



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Registration Division (7505T)
 1200 Pennsylvania Ave., N.W.
 Washington, D.C. 20460

EPA Reg. Number:

71085-95

Date of Issuance:

2/18/26

NOTICE OF PESTICIDE:

Registration
 Reregistration
 (under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

CH-PRO-5823

Name and Address of Registrant (include ZIP Code):

CHEMAGCO LLC
 15401 WESTON PARKWAY, SUITE 170
 CARY, NC 27513

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

Continues page 2

Signature of Approving Official:

Manjula Unnikrishnan
 Product Manager 21 Fungicide
 Branch Registration Division (7505T)
 Office of Pesticide Programs

Date:

2/18/26

2. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 71085-OL."
 - Add an appropriate EPA Establishment Number and Net Contents information.
3. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The record for this product currently contains the following CSF(s):

- Basic CSF dated 08/01/2025

If you have any questions, please contact Senedu Alemu at alemu.senedu@epa.gov.

Enclosure

ACCEPTED
 02/18/2026
 Under the Federal Insecticide, Fungicide
 and Rodenticide Act as amended, for the
 pesticide registered under
 EPA Reg. No. 71085-95

TRIFLUMIZOLE	GROUP	3	FUNGICIDE
--------------	-------	----------	-----------

CH-PRO-5823

Agricultural Fungicide

COMPOSITION

Active Ingredient:
 Triflumizole
 1-[1-[[4-chloro-2-(trifluoromethyl)phenyl]imino]-2- propoxyethyl]-1H-imidazole....42.14%
 Other Ingredients:57.86%
 Total: 100.00%
 Contains 4 lb Triflumizole per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by the poison control center or doctor. • DO NOT give anything by mouth to an unconscious person.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. Contact the Rocky Mountain Poison and Drug Safety at 1-866-673-6671 for emergency medical treatment information.</p>	

See inside booklet for additional First Aid, Precautionary Statements, and Directions for Use

CHEMAGCO LLC
 15401 Weston Parkway, Suite 170
 Cary, NC 27513

EPA Reg. No. 71085-XX
 EPA Est. No.

Net Contents: _____

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or the toilet. Remove contaminated clothing and wash clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemically-resistant to this product are listed below.

Applicators and Other Handlers Must Wear:

- A long-sleeved shirt & long pants;
- chemical-resistant gloves made of any waterproof material;
- shoes plus socks.
- For applicators using open cab air blast equipment in pome fruit, and cherries, a chemical resistant hat is also required.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

When handlers use closed systems or enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling the product. Wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when cleaning equipment or disposing of equipment washwater or rinsate. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire label directions before using. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. **DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. REI is 24 hours (1 day) for girdling of table and raisin grapes.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water, is:

- coveralls;
- chemical-resistant gloves made of any waterproof material; and
- shoes plus socks.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry, secure location.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. **DO NOT** reuse or refill this container.

[For containers equal to or less than 5 gallons in size:]

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

If pressure rinsing: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about

40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Then offer container for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Recycling: Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer or contact the Ag Container Recycling Council (ACRC) at 1-877-952-2272 (toll free) or www.acrecycle.org.

PRODUCT INFORMATION

CH-PRO-5823 is a broad-spectrum foliar fungicide for control of certain diseases of listed fruits and vegetables. CH-PRO-5823 belongs to the demethylation inhibitor (DMI) group of fungicides classified as Group 3 and acts as a protectant and eradicant by preventing disease symptoms before and after infection has occurred. Its anti-sporulant activity reduces spores after lesions become visible.

Read the entire label and directions before using CH-PRO-5823.

RESISTANCE MANAGEMENT: Populations of fungal pathogens are known to develop resistance to fungicides with the same mode of action when used repeatedly. Thus, Macdermid Agricultural Solutions encourages responsible resistance management to ensure an effective long-term control of diseases. Resistance management strategies include alternating and/or tank mixing with other labeled

fungicides having different modes of action or limiting the total number of applications per year.

APPLICATION INSTRUCTIONS

Apply CH-PRO-5823 in sufficient water to ensure complete and thorough coverage of foliage, flowers and fruit.

Application rates are best determined by the user based on disease pressure, weather conditions, varietal susceptibility to specific diseases, and planned timing of applications versus the start of infection periods. In general, use rates in the higher listed rate range for heavy disease pressure, more susceptible varieties, conditions that favor disease development and conditions in which thorough coverage is more difficult.

Consult your State Agricultural Experimental Station or Extension Service Specialist for guidance on selection of specific dosage rates and timing of application appropriate to your specific crop and area conditions. State regulations may contain additional restrictions or requirements.

DO NOT use less than the specified application rate. CH-PRO-5823 can be tank mixed with other fungicides. CH-PRO-5823 can be applied with dilute or concentrate ground equipment. Aerial applications for tree fruits can be made in accordance with use directions noted for each crop. Equipment should be frequently checked for calibration.

PRODUCT RESTRICTIONS

DO NOT allow livestock to graze on treated orchard ground cover. Aerial application of CH-PRO-5823 is allowed unless prohibited under a specific crop use direction. **DO NOT** apply this product through any type of irrigation system.

RESISTANCE MANAGEMENT RESTRICTIONS:

DO NOT make more than two sequential applications of CH-PRO-5823 before alternating to a registered fungicide with a different mode of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

ROTATIONAL CROPS RESTRICTIONS: DO NOT plant leafy or fruiting vegetables within 30 days after application. **DO NOT** plant root and tuber vegetables within 60 days after application. **DO NOT** plant cotton, small cereal grains and all other crops not registered on the label for CH-PRO-5823, within one year after application.

POME FRUIT (CROP GROUP 11-10)

CROP	DISEASE	RATE FL OZ PER ACRE (LB AI/A)	APPLICATION INSTRUCTIONS
Apples Azarole Crabapple Loquat Mayhaw Medlar Pear Pear, Asian Quince Quince, Chinese Quince, Japanese Tejocote (including all cultivars, varieties and or hybrids of these)	Powdery Mildew <i>(Podosphaera leucotricha)</i>	8 to 16 (0.25 to 0.5 lb ai)	Initiate spraying at first sign of mildew, usually at the 1/2" green tip stage.
	Scab <i>(Venturia inaequalis)</i> Cedar Apple Rust <i>(Gymnosporangium juniperi-virginianae)</i>	8 to 16 (0.25 to 0.5 lb ai)	Standard Spray Schedule: Initiate spraying at 1/2" green tip stage continue through second cover or until primary scab sporulation is complete. Eradicant Spray Schedule (Post Infection Schedule): Initiate applications within 96 hours of the start of an infection period. Under low to moderate disease pressure conditions, rates in the lower range can be applied within 48 hours after the start of an infection period. Use rates in the higher range when disease pressure is high or if the time interval after the start of infection is greater than 48 hours. For improved scab control, apply CH-PRO-5823 in combination with a protectant fungicide. After first cover, CH-PRO-5823 can be used in a standard cover spray program in tank mixes with appropriate protectant fungicides.

RESTRICTIONS:

DO NOT apply more than a total of 4 applications of CH-PRO-5823 at reduced rates per crop per year.
DO NOT apply more than a total of 64 fl oz (2 lb ai) of CH-PRO-5823 per crop per year.
DO NOT apply less than 14 days apart. Wait at least 21 days between the second and third applications.
DO NOT apply within 14 days of harvest.
DO NOT allow livestock to graze on treated orchard ground cover.
Single maximum application rate: 16 fl oz (0.5 lb ai)
Minimum RTI: DO NOT apply less than 14 days apart. Wait at least 21 days between the second and third application.
 When making aerial applications, apply 12 - 16 fl oz (0.375 - 0.5 lb ai) CH-PRO-5823 in a minimum of 10 gallons of water per acre.
 Not registered for use by AIR in California.
 Ground applications – apply in a minimum of 100 gallons per acre.

BRASSICA LEAFY GREENS (CROP SUBGROUP 5B)

CROP	DISEASE	RATE FL OZ PER ACRE (LB AI/A)	APPLICATION INSTRUCTIONS
Broccoli raab Chinese cabbage (bok choy) Collards Kale Mizuna Mustard spinach Rape greens Turnip greens (Including all cultivars and/or hybrids of these)	Powdery Mildew <i>(Erysiphe polygoni)</i> Black Spot <i>(Alternaria spp.)</i>	6 to 8 (0.19 to 0.25 lb ai)	Begin application at the first sign of disease. A single repeat application can be made 14 days or more after the first application.

RESTRICTIONS:

DO NOT apply more than a total of 2 applications of PROCURE 480 SC per crop per year.

DO NOT apply more than a total of 16 fl oz (0.5 lb ai/A) of PROCURE 480 SC per crop per year.

DO NOT apply within 1 day of harvest.

Use a minimum of 10 gallons per acre spray volume for ground applications.

Use a minimum of 2 gallons per acre spray volume for aerial applications.

Single maximum application rate: 8 fl oz (0.25 lb ai)

Minimum RTI: 14 days.

CHERRIES (SWEET & TART)

CROP	DISEASE	RATE FL OZ PER ACRE (LB AI/A)	APPLICATION INSTRUCTIONS
Cherries (Sweet & Tart)	Powdery Mildew <i>(Podosphaera clandestina)</i>	8 to 16 (0.25 to 0.50 lb ai)	Initiate treatments at early popcorn and continue until terminal growth ceases, timed to maintain coverage on susceptible tissue throughout the possible infection period.
	Blossom Blight & Fruit Rot <i>(Monilinia spp.)</i>	10 to 16 (0.31 to 0.50 lb ai)	Blossom Blight Control: Apply between early popcorn and petal fall. Complete fungicide coverage must be maintained throughout the bloom period. Fruit Rot Control: Depending on disease pressure, applications can also be directed to the fruit during the cover spray period.
	Leaf Spot <i>(Blumeriella jaapii)</i>	10 to 16 (0.31 to 0.50 lb ai)	Applications made to control powdery mildew will suppress the development of leaf spot. Tank mix with other products registered for leaf spot control for control.

RESTRICTIONS:

DO NOT apply more than a total of 4 applications of CH-PRO-5823 at reduced rate per crop per year.

DO NOT apply more than a total of 56 fl oz (1.75 lb ai) of CH-PRO-5823 per crop per year.

DO NOT apply less than 14 days apart. Wait at least 21 days between the second and third application.

DO NOT apply within 1 day of harvest.

DO NOT allow livestock to graze on treated orchard ground cover.

Single maximum application rate: 16 fl oz (0.5 lb ai)

When making aerial applications, apply 12 - 16 fl oz (**0.5 lb ai**) CH-PRO-5823 in a minimum of 10 gallons of water per acre.

Ground applications – apply in a minimum of 100 gallons per acre.

CUCURBIT VEGETABLES (CROP SUBGROUP 9)

CROP	DISEASE	RATE FL OZ PER ACRE (LB AI/A)	APPLICATION INSTRUCTIONS
Cucumbers Gherkins Gourds Muskmelons (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon) Pumpkins Squash (summer and winter) Watermelons	Powdery Mildew <i>(Erysiphe cichoracearum, Podosphaera xanthii)</i>	4 to 8 (0.125 to 0.25 lb ai)	Begin applications at the start of vining or at first indication of disease. Repeat applications can be made at not less than 7-day intervals.

RESTRICTIONS:

If applying to a transplanted crop, **DO NOT** apply more than a total of 3 applications of CH-PRO-5823 per crop per year.
 If applying to a crop grown from seed, **DO NOT** apply more than a total of 4 applications of CH-PRO-5823 per crop per year at the 4 – 6 oz (0.125-0.19) per acre rate or 3 applications at the 8 oz (0.25 lbs ai) per acre rate.
DO NOT apply more than a total of 24 fl oz (0.75 lb ai) of CH-PRO-5823 per crop per year.
 Applications can be made up to and including the day of harvest.
 Single maximum application rate: 8 fl oz (0.25 lbs ai)
 Yearly max use rate: 24 fl oz (0.75 lb ai)
 Minimum RTI: 7 day

GRAPES, SMALL FRUIT VINE CLIMBING, EXCEPT FUZZY KIWIFRUIT (CROP SUBGROUP 13-07F)

CROP	DISEASE	RATE FL OZ PER ACRE (LB AI/A)	APPLICATION INSTRUCTIONS
Amur River Grape Grape Gooseberry Maypop Schisandra berry (including all cultivars, varieties, and/or hybrids of these)	Powdery Mildew <i>(Uncinula necator)</i>	4 to 8 (0.125 to 0.25 lb ai)	Start protectant applications before bloom and continue on a 14 to 21-day interval. When disease pressure is high or for the most susceptible varieties, use 6 to 8 fl oz (0.19-0.25 lb ai) /acre and/or 14-day spray intervals. When disease pressure is low or for least susceptible varieties, use 4 to 6 fl oz (0.125-0.19 lb ai)/acre and/or 21-day spray intervals. Select spray volume, nozzle adjustments and ground speed to ensure complete coverage of fruit clusters and foliage. Use a minimum of 50 gallons of water per acre for early season applications. Sulfur at specified use rates can be applied before, with, and/or between CH-PRO-5823 applications. When applied on a powdery mildew spray schedule CH-PRO-5823 will enhance the activity of registered fungicides used for the control of Botrytis
			bunch rot. Applications must be made on a 14-day spray schedule for suppression of Botrytis bunch rot.

RESTRICTIONS:

DO NOT apply more than a total of 4 applications of CH-PRO-5823 per crop per year.
DO NOT apply more than a total of 32 fl oz (1.0 lb ai) of CH-PRO-5823 per crop per year.
DO NOT apply within 7 days of harvest.
 The Restricted Entry Interval (REI) for girdling table and raisin grapes is 24 hours (1-day). Maintain 12 hours for wine grapes.
DO NOT apply by air.
 Single maximum application rate: 8 fl oz (0.25 lbs ai)
 Yearly maximum use rate: 32 fl oz (1.0 lb ai)
 Minimum RTI: 14 days

HAZELNUTS (FILBERTS)

CROP	DISEASE	RATE FL OZ PER ACRE (LB AI/A)	APPLICATION INSTRUCTIONS
Hazelnuts (Filberts)	Eastern Filbert Blight <i>(Anisogramma anomala)</i>	4 to 6 (0.125 to 0.19 lb ai)	Apply at budbreak (when the first green tissue is visible) as a preventative spray. One repeat application can be made no less than 10 days later if weather conditions are conducive to disease development, and to protect the expanding shoot tip during bud break and rapid shoot elongation. CH-PRO-5823 may be applied as a tank mix or alternated with non-DMI fungicides as a resistance management strategy. Tank mixing the lowest labeled rate of a spray surfactant with CH-PRO-5823 may improve spray coverage and penetration.

RESTRICTIONS:

DO NOT apply more than a total of 2 applications of CH-PRO-5823 per crop per year.
DO NOT apply more than a total of 12 fl oz (0.375 lb ai) of CH-PRO-5823 per crop per year.
DO NOT apply within 18 days of harvest.
DO NOT allow livestock to graze on treated orchard ground cover.
 Ground applications – apply in a minimum of 100 gallons per acre.
 Single maximum application rate: 6 fl oz (0.19 lbs ai)
 Minimum RTI: 10 days
 For aerial applications, use only the 6 fl oz/A (0.19 lbs ai) rate - apply in a minimum of 15 gallons per acre.

HEAD AND STEM BRASSICA (CROP SUBGROUP 5A)

CROP	DISEASE	RATE FL OZ PER ACRE (LB AI/A)	APPLICATION INSTRUCTIONS
Head & Stem Brassicas: Broccoli Broccoli, Chinese (gai lon) Brussels sprouts Cabbage Cabbage, Chinese (napa) Cauliflower Cavalo broccolo Kohlrabi Mustard, Chinese (gai choy)	Powdery Mildew <i>(Erysiphe polygoni)</i> Black Spot / Alternaria leaf spot <i>(Alternaria spp.)</i>	6 to 8 (0.19 to 0.25 lb ai)	Begin applications at the first indication of disease. A single repeat application can be made no less than 14 days after the first application. Use the higher specified rate when disease pressure is high.

RESTRICTIONS:

DO NOT apply more than a total of 2 applications of CH-PRO-5823 per crop per year.
DO NOT apply more than a total of 16 fl oz (0.5 lb ai) of CH-PRO-5823 per crop per year.
DO NOT apply within 1 day of harvest.
 Single maximum application rate: 8 fl oz (0.25 lbs ai)
 Minimum RTI: 14 days
 Use a minimum of 10 gallons per acre spray volume for ground applications.
 Use a minimum of 2 gallons per acre spray volume for aerial applications.

HOPS

CROP	DISEASE	RATE FL OZ PER ACRE (LB AI/A)	APPLICATION INSTRUCTIONS
Hops	Powdery Mildew <i>(Podosphaera macularis)</i>	12 (0.375 lb ai)	Begin applications prior to disease development or at the first indication of the disease. One repeat application can be made not less than 14 days after the first application.

RESTRICTIONS:

DO NOT apply more than a total of 2 applications of CH-PRO-5823 per crop per year.
DO NOT apply more than a total of 24 fl oz (0.75 lb ai) of CH-PRO-5823 per crop per year.
DO NOT apply within 7 days of harvest.
 Single maximum application rate: 12 fl oz (0.375 lbs ai)
 Minimum RTI: 14 days
 Use a minimum of 50 gallons per acre spray volume for ground applications.
 Use a minimum of 10 gallons per acre spray volume for aerial applications.
DO NOT graze livestock in treated areas, and **DO NOT** harvest crops grown in treated areas for silage or for hay.

LEAFY GREENS EXCEPT SPINACH (CROP SUBGROUP 4A)

CROP	DISEASE	RATE FL OZ PER ACRE (LB AI/A)	APPLICATION INSTRUCTIONS
Leafy Greens (except spinach) Amaranth Arugula Chervil Chrysanthemum, edible-leaved & garland Cilantro Corn salad Cress, garden & upland Dandelion Dock Endive Lettuce (head and leaf,) Cos (Romaine) Butterhead varieties (both field and greenhouse) Parsley Purslane, garden and winter Radicchio Swiss chard (including all cultivars and/or hybrids of these)	Powdery Mildew <i>(Erysiphe polygoni)</i> Black Spot / Alternaria leaf spot (<i>Alternaria</i> <i>spp.</i>)	6 to 8 (0.19 to 0.25 lb ai)	Begin applications prior to disease development or at the first indication of the disease. A single repeat application can be made no less than 14 days after the first application.

RESTRICTIONS:

DO NOT apply more than a total of 2 applications of CH-PRO-5823 per crop per year.
DO NOT apply more than a total of 16 fl oz (0.5 lb ai) of CH-PRO-5823 per crop per year.
 Applications can be made up to and including day of harvest.
 Single maximum application rate: 8 fl oz (0.25 lbs ai)
 Minimum RTI: 14 days
 Use a minimum of 10 gallons per acre spray volume for ground applications.
 Use a minimum of 2 gallons per acre spray volume for aerial applications.

PAPAYA

CROP	DISEASE	RATE FL OZ PER ACRE (LB AI/A)	APPLICATION INSTRUCTIONS
Canistel Mango Papaya Sapodilla Sapote, black Sapote, mammey Star apple	Black Spot <i>(Asperisporium caricae)</i> Powdery Mildew <i>(Oidium caricae)</i>	8 (0.25 lb ai)	Apply prior to disease development or at the first indication of the disease. An approved surfactant may be used to enhance coverage to the foliage.

RESTRICTIONS:

DO NOT apply more than a total of 1 application of CH-PRO-5823 per crop per year.

DO NOT apply more than a total of 8 fl oz (0.25 lb ai) of CH-PRO-5823 per crop per year.

Single maximum application rate: 8 fl oz (0.25 lbs ai)

Minimum RTI: Not Applicable

Application can be made up to and including day of harvest.

Use a minimum of 75 gallons per acre spray volume for ground applications.

Use a minimum of 2 gallons per acre spray volume for aerial applications.

PINEAPPLES

CROP	DISEASE	RATE FL OZ PER ACRE (LB AI/A)	APPLICATION INSTRUCTIONS
Pineapples (pre-plant seed piece) dip treatment	Butt Rot <i>(Chalara paradoxa)</i>	4 to 8 (0.125 to 0.25 lb ai)	Apply only as a pre-plant seed piece dip treatment. Freshly removed seed piece material from the parent plant must be immersed immediately for thorough wetting and coverage. Store treated plant material with butt ends exposed to the sun prior to planting for disease control.

RESTRICTIONS:

DO NOT use treated green forage or green fodder for animal feed.

Single maximum application rate: 8 fl oz (0.25 lbs ai)

LOW-GROWING BERRIES EXCEPT CRANBERRIES (CROP SUBGROUP 13-07G)

CROP	DISEASE	RATE FL OZ PER ACRE (LB AI/A)	APPLICATION INSTRUCTIONS
Bearberry Bilberry Blueberry, lowbush Cloudberry Lingonberry Muntries Partridgeberry Strawberry (including all cultivars, varieties and/or hybrids of these)	Powdery Mildew <i>(Sphaerotheca macularis f. sp. fragariae)</i>	4 to 8 (0.125 to 0.25 lb ai)	Begin applications at first indication of disease. Repeat applications can be made at no less than 14 day intervals while conditions favor disease development. Protection of leaves before bloom will reduce in-season disease pressure and result in greater control of flower and fruit infections. Apply with dilute or concentrate ground equipment or by air. Use higher specified application rates and shorter spray intervals on more susceptible varieties and/or under higher disease pressure.

RESTRICTIONS:

DO NOT apply more than a total of 4 applications of CH-PRO-5823 per crop per year.

DO NOT apply more than a total of 32 fl oz (1.0 lb ai) of CH-PRO-5823 per crop per year.

DO NOT apply within 1 day of harvest.

Single maximum application rate: 8 fl oz (0.25 lbs ai)

Minimum RTI: 14 days

Use a minimum of 75 gallons per acre spray volume for ground applications.

Use a minimum of 2 gallons per acre spray volume for aerial applications.

**IMPORTANT INFORMATION
READ BEFORE USING PRODUCT**

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product reflect the opinion of experts based on field use and tests, and must be followed carefully. It is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of CHEMAGCO LLC or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of CHEMAGCO LLC and Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold CHEMAGCO LLC and Seller harmless for any claims relating to such factors.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, CHEMAGCO LLC AND SELLER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ON THIS LABEL.

To the extent consistent with applicable law, CHEMAGCO LLC or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product and **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF CHEMAGCO LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF CHEMAGCO LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

CHEMAGCO LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of CHEMAGCO LLC.