

71096-4

2/14/2001

1/7

(Front Panel)

# SLUG-FEST

## ALL WEATHER FORMULA

*Lasts longer under wet conditions*

**Active Ingredient:**

Metaldehyde (2,4,6,8-tetramethyl-1,3,5,7-tetraoxycyclo-octane) .....	25%
Other Ingredients: .....	75%
Total .....	100%

*Contains 2.1 pounds Metaldehyde per gallon.*

KEEP OUT OF REACH OF CHILDREN

WARNING AVISO

Si Usted no entiende la etiqueta, busque a alguien para que se la explique a Usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)


See side panels for additional precautionary statements.

All applicable directions, restrictions and precautions on the label are to be followed. See Directions for Use in Booklet.

**SHAKE WELL****NET CONTENTS**

EPA Reg. No. 71096-4  
EPA Est. No. 52251-OR-005

**OR-CAL, Inc.**  
29454 Meadowview Road  
Junction City, OR 97448  
(541)689-4413



## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**WARNING.** May be fatal if swallowed. Avoid breathing spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after use. Keep out of reach of children or domestic animals.

**IMPORTANT:** *KEEP CHILDREN AND PETS OUT OF TREATED AREA UNTIL SPRAY HAS DRIED. THIS PESTICIDE MAY BE FATAL TO DOGS OR OTHER PETS IF ANIMAL DRINKS. DO NOT ALLOW PUDDLES TO FORM ON GROUND WHEN MIXING OR CLEANING APPLICATION EQUIPMENT.*

#### FIRST AID

**If Swallowed:** Call a physician or Poison Control Center. Give 3-4 glasses of water but DO NOT induce vomiting. If vomiting does occur, give fluids again.

**If In Eyes:** Wash eyes immediately with fresh water for at least 15 minutes. See a doctor if irritation persists.

**If On Skin:** Wash skin with plenty of soap and water.

**If Inhaled:** Remove person from exposure area.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

*Applicators and handlers must wear:* Coveralls over long-sleeved shirt and long pants, chemical-resistant or waterproof gloves, chemical resistant footwear plus socks, protective eyewear, chemical resistant apron when cleaning equipment, mixing, or loading. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### USER SAFETY RECOMMENDATIONS

#### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water or areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water.

### STORAGE AND DISPOSAL

**Prohibitions:** Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

**Storage:** Keep pesticides in original container in a locked place out of reach of children. Protect from freezing.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Do not reuse container. Dispose of container in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### AGRICULTURAL USE

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**REENTRY:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

**EARLY ENTRY PPE:** PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: Coveralls over long-sleeved shirt and long pants, chemical resistant or waterproof gloves, chemical resistant footwear plus socks, and protective eyewear.

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of the product that are NOT within the scope of the Worker Protection Standard, for agricultural pesticides 40 CFR Part 170. The WPS applies only when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

**USE PRECAUTIONS**

Read entire label before using. Apply directly to the soil area beneath plants as listed on this label. Do not allow spray to come in contact with parts of plant that may be used for food or feed purposes. The grower is responsible for residues on his crops, as well as for damages caused by drift from his property to that of others. Do not tank mix this product with other chemicals. Do not allow drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops.

**MIXING:** Shake SLUG-FEST ALL WEATHER FORMULA well before mixing. Thoroughly mix spray solution before application. Use entire mixed solution at one time. Do not store mixed solution.

**LAWNS AND ORNAMENTALS:** Such as Geraniums, Salvia, Primroses, Marigolds, Petunias, Pansies, Dahlias, Begonias, Coleus, Ajuga and Dichondra.

**HOSE-ON TYPE APPLICATOR:** Thoroughly water areas to be treated. Use one ounce of SLUG-FEST ALL WEATHER FORMULA for each two gallons of solution delivered by your applicator to cover approximately 200 sq. ft.

**SPRINKLING CAN:** Thoroughly water areas to be treated. Use one ounce of SLUG-FEST ALL WEATHER FORMULA to each two gallons of water and sprinkle over areas to be treated.

**WHERE SNAILS ARE A PROBLEM:** We suggest double rate of application on Ornamentals only.

**BEST RESULTS:** Apply SLUG-FEST ALL WEATHER FORMULA in the cool of the evening.

**TERRESTRIAL FOOD CROPS:** Alfalfa, Almond, Anise, Artichoke, Asparagus, Barley, Beans, Beets, Blackeyed Peas, Broccoli, Brussels Sprouts, Cabbage, Cantaloupes, Caraway, Carrots, Catnip, Cauliflower, Celery, Chicory, Chive, Clover, Collards, Coriander, Corn, Cowpeas, Cucumbers, Currant, Dill, Eggplants, Endive, Garlic, Ginseng, Horseradish, Kale, Kohlrabi, Leeks, Lentils, Lettuce, Melons, Muskmelons, Mustard, Mustard Greens, Oats, Okra, Onions, Oregano, Parsley, Parsnips, Peas, Peppermint, Peppers, Pimentos, Potatoes, Pumpkins, Radishes, Rhubarb, Rutabagas,

Page 5 – (SLUG-FEST ALL WEATHER FORMULA)  
(Side Panel or Booklet)

Sage, Salsify, Savory, Shallots, Spearmint, Spinach, Squash, Sugar Beets, Sweet Potatoes, Swiss Chard, Tomatoes, Turnips, Watermelon, Wheat, Seed Crops such as alfalfa, beans, clover, corn (sweet), grasses, peas, potato.

**SLUGS AND SNAILS:** Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage OR 5 to 10 oz. in ten gallons of water per treated 1,000 sq. ft. OR 1 to 2 oz. in two gallons of water per 200 sq. ft. Where Snails are a problem we suggest the higher rate of application.

Thoroughly water areas to be treated before application. Apply only to soil surface around plants and in rows between plants. Use equipment that applies directly to the soil surface. Do not apply in such a manner as to cause drift of product onto plants. Do not apply to edible parts of plants. For best results, apply in the evening. Repeat as needed for control.

**TREE CROPS:** Apples, Apricots, Avocados, Bananas, Cherries, Citrus fruits, Grapefruits, Lemons, Limes, Nectarines, Oranges, Peaches, Pears, Pecans, Plums, Stone fruits, Walnuts.

**SLUGS AND SNAILS:** Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage OR 5 to 10 oz. in ten gallons of water per treated 1,000 sq. ft. OR 1 to 2 oz. in two gallons of water per treated 200 sq. ft. Where snails are a problem we suggest the higher rate of application.

Make application when foliage and fruit are completely dry but soil is wet following an irrigation. Apply under trees, around irrigation pipes, farm buildings and any other place where slugs and snails may congregate. Use equipment that applies directly to soil surface. Do not apply in such a manner as to cause drift of product onto trees. Do not apply directly to trees or fruit. For best results, apply in the evening. Repeat as needed for control.

**BERRY CROPS:** Blackberries, Blueberries, Boysenberries, Caneberries, Dewberries, Gooseberries, Loganberries, Raspberries, Strawberries.

**SLUGS AND SNAILS:** Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage OR 5 to 10 oz. in ten gallons of water per treated 1,000 sq. ft. OR 1 to 2 oz. in two gallons of water per treated 200 sq. ft. Where snails are a problem we suggest the higher rate of application.

Thoroughly water areas to be treated before application. Apply only to soil surface around plants and in rows between plants. Use equipment that applies directly to soil surface. Do not apply in such a manner as to cause drift of product onto plants. Do not

Page 6 -- (SLUG-FEST ALL WEATHER FORMULA)  
(Side Panel or Booklet)

apply to edible parts of plants. For best results, apply in the evening. Repeat as needed for control.

**TERRESTRIAL NONFOOD CROPS**

**SEED CROPS:** Beets, Cabbage, Carrots, Collards, Kale, Kohlrabi, Lettuce, Mustard, Radish, Spinach, Swiss Chard, Tobacco, Turnips.

*SLUGS AND SNAILS:* Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage.

Thoroughly water areas to be treated before application. Apply only to soil surface around plants and in rows between plants. Use equipment that applies directly to soil surface. Do not apply in such a manner as to cause drift of product onto plants. For best results, apply in the evening. Repeat as needed for control.

**GREENHOUSE FOOD CROP:** Asparagus, Beans, Beets, Cabbage, Carrots, Celery, Corn, Cucumbers, Eggplant, Endive, Lettuce, Melons, Onions, Peas, Peppers, Potatoes, Radish, Spinach, Squash, Tomatoes, Turnips.

*SLUGS AND SNAILS:* Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage OR 5 to 10 oz. in ten gallons water per treated 1,000 sq. ft. OR 1 to 2 oz. in two gallons of water per treated 200 sq. ft. Where snails are a problem we suggest the higher rate of application.

Water or irrigate areas to be treated before application. Apply only to soil surface around plants, in rows between plants and under benches in greenhouses and around plants; any place where slugs and snails may congregate. Use equipment that applies directly to soil surface. Do not apply in such a manner as to cause drift of product onto plants. Do not apply to plants or edible parts of plants. For best results, apply in the evening. Repeat as needed for control.

**GREENHOUSE NONFOOD CROPS:** Orchids, Ornamental Plants

*SLUGS AND SNAILS:* Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage OR 5 to 10 oz. in ten gallons of water per treated 1,000 sq. ft. OR 1 to 2 oz. in two gallons of water per treated 200 sq. ft. Where snails are a problem we suggest the higher rate of application.

Water or irrigate areas to be treated before application. Apply only to soil surface around plants, in rows between plants and under benches in greenhouse and around plants; any place where snails and slugs might congregate. Do not apply directly to plants unless prior experience has shown the plant to be tolerant to this product. Most plants are tolerant, however, some slight tip burning has been noticed on extremely tender orchids.

Page 7 – (SLUG-FEST ALL WEATHER FORMULA)  
(Side Panel or Booklet)

**DOMESTIC AND NONDOMESTIC OUTDOOR:** Domestic Dwellings, Farm Buildings, Lath Houses, Ornamental Plants

*SLUGS AND SNAILS:* Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage OR 5 to 10 oz. in ten gallons of water per treated 1,000 sq. ft. OR 1 to 2 oz. in 2 gallons of water per treated 200 sq. ft. Where snails are a problem we suggest the higher rate of application.

Thoroughly water areas to be treated before application. Use equipment that applies directly to the soil surface. Apply evenly over infested areas and along flower beds adjacent to border or ground cover plants. Also apply to ground at base of fences and walls in shady, protected locations and any other place where slugs and snails may congregate.

Do not apply in such a manner as to cause drift of product onto edible plants or plant parts. Do not apply directly to plants unless prior experience has shown the plant to be tolerant, however, some slight tip burning has been noticed on extremely tender orchids. For best results, apply in the evening. Repeat as needed for control.

**LAWNS (INCLUDING DICHONDRA), ORNAMENTAL GRASSES, TURF, NEWLY SEEDED GRASS GROWN FOR SEED** (prior to canopy closure).

*SLUGS AND SNAILS:* Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage OR 5 to 10 oz. in ten gallons of water per treated 1,000 sq. ft. OR 1 to 2 oz. in two gallons of water per treated 200 sq. ft. Where snails are a problem we suggest the higher rate of application.

Uniformly broadcast directly on lawn or turf. Withhold heavy watering for at least 2 days after treatment. Keep children and pets off the treated area until spray has dried.