

71096-4

12/13/2007

1/14

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460



OFFICE OF
PREVENTION, PESTICIDES,
AND TOXIC SUBSTANCES

December 13, 2007

Irene Boone
OR-CAL, Inc.
29454 Meadowview Road
Junction City, OR 97448

Dear Ms. Boone:

Subject: Submission of amended label to add non-crop use sites on master label
Slug-Fest All Weather Formula
EPA Registration No. 71096-4
Your Submission Dated May 21, 2007

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, is acceptable with the following comments:

- On page 8, add the following, bolded statement to the NON CROP AREAS section:

NON CROP AREAS:

This product may be used only on the following use sites in response to State and/or Federal invasive mollusk eradication operations. Broadcast applications of this product may be made to the following use sites for State and/or Federal invasive mollusk eradication operations:

Fallow land; barrier strips; uncultivated non-agricultural areas; recreational areas, etc...


This product may be applied for State and/or Federal invasive mollusk eradication operations at an individual application rate of no more than 2 lb. of active ingredient per acre (0.95 gallons of Slug-Fest product per acre), and at no more than six applications per year.

- On page 11, remove the NON CROP AREAS section from the Home Garden portion of the label.

2/14

A stamped copy of the label is enclosed for your records. Submit two copies of your final printed labeling before you release the product for shipment. If you have any questions, please Julie Chao at (703) 308-8735 or chao.julie@epa.gov.

Regards,



Venus Eagle
Insecticide-Rodenticide Branch
Registration Division (7505P)

Enclosure

(Front Panel – AG Label)

SLUG-FEST

ALL WEATHER FORMULA

Lasts longer under wet conditions

Active Ingredient:

Metaldehyde (2,4,6,8-tetramethyl-1,3,5,7-tetraoxocyclo-octane) 25%

Other Ingredients: 75%

Total 100%

Contains 2.1 pounds Metaldehyde per gallon.

KEEP OUT OF REACH OF CHILDREN
WARNING AVISO

Si Usted no entiende la etiqueta, busque a alguien para que se la explique a Usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See side panels for additional precautionary statements.

All applicable directions, restrictions and precautions on the label are to be followed. See Directions for Use in Booklet.

SHAKE WELL

NET CONTENTS

EPA Reg. No. 71096-4
EPA Est. No. 52251-OR-005

OR-CAL, Inc.
29454 Meadowview Road
Junction City, OR 97448
(541)689-4413

ACCEPTED
with COMMENTS
In EPA Letter Dated:
DEC 13 2007
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
71096-4

4/14

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Harmful if swallowed. Avoid breathing spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after use. Keep out of reach of children or domestic animals.

IMPORTANT: *KEEP CHILDREN AND PETS OUT OF TREATED AREA UNTIL SPRAY HAS DRIED. THIS PESTICIDE MAY BE FATAL TO DOGS OR OTHER PETS IF ANIMAL DRINKS. DO NOT ALLOW PUDDLES TO FORM ON GROUND WHEN MIXING OR CLEANING APPLICATION EQUIPMENT.*

FIRST AID

If Swallowed-Call a poison control center or doctor immediately for treatment advice.

- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

If in eyes -Hold eye open and rinse slowly and gently with water for 15-20 minutes.
-Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

If inhaled -Move person to fresh air.
-If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
-Call a poison control center or doctor for further treatment advice.

If on skin or clothing -Take off contaminated clothing.
-Rinse skin immediately with plenty of water for 15-20 minutes.
-Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Telecommunications Network at 1-800-858-7378 for emergency treatment.

Note to Physician

Probable mucosal damage may contraindicate gastric lavage.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and handlers must wear: Coveralls over long-sleeved shirt and long pants, chemical-resistant gloves made of a waterproof material, chemical resistant footwear plus socks, protective eyewear, chemical resistant apron when cleaning equipment, mixing, or loading. Follow manufacturer’s instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water or areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water.

STORAGE AND DISPOSAL

Prohibitions: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

Storage: Keep pesticides in original container in a locked place out of reach of children. Protect from freezing.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Do not reuse container. Dispose of container in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

6/14

AGRICULTURAL USE

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

REENTRY: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

EARLY ENTRY PPE: PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: Coveralls over long-sleeved shirt and long pants, chemical resistant gloves made of a waterproof material, chemical resistant footwear plus socks and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of the product that are NOT within the scope of the Worker Protection Standard, for agricultural pesticides 40 CFR Part 170. The WPS applies only when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

USE PRECAUTIONS

Read entire label before using. Apply directly to the soil area beneath plants as listed on this label. Do not allow spray to come in contact with parts of plant that may be used for food or feed purposes. **(AGRICULTURAL LABEL ONLY)** The grower is responsible for residues on his crops, as well as for damages caused by drift from his property to that of others. Do not tank mix this product with other chemicals. Do not allow drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops.

MIXING: Shake SLUG-FEST ALL WEATHER FORMULA well before mixing. Thoroughly mix spray solution before application. Use entire mixed solution at one time. Do not store mixed solution.

LAWNS AND ORNAMENTALS: Such as Geraniums, Salvia, Primroses, Marigolds, Petunias, Pansies, Dahlias, Begonias, Coleus, Ajuga and Dichondra.

HOSE-ON TYPE APPLICATOR: Thoroughly water areas to be treated. Use one ounce of SLUG-FEST ALL WEATHER FORMULA for each two gallons of solution delivered by your applicator to cover approximately 200 sq. ft.

SPRINKLING CAN: Thoroughly water areas to be treated. Use one ounce of SLUG-FEST ALL WEATHER FORMULA to each two gallons of water and sprinkle over areas to be treated.

WHERE SNAILS ARE A PROBLEM: We suggest double rate of application on Ornamentals only.

BEST RESULTS: Apply SLUG-FEST ALL WEATHER FORMULA in the cool of the evening.

TERRESTRIAL FOOD CROPS: Alfalfa, Almond, Anise, Artichoke, Asparagus, Barley, Beans, Beets, Blackeyed Peas, Broccoli, Brussels Sprouts, Cabbage, Cantaloupes, Caraway, Carrots, Catnip, Cauliflower, Celery, Chicory, Chive, Clover, Collards, Coriander, Corn, Cowpeas, Cucumbers, Currant, Dill, Eggplants, Endive, Garlic, Ginseng, Horseradish, Kale, Kohlrabi, Leeks, Lentils, Lettuce, Melons, Muskmelons, Mustard, Mustard Greens, Oats, Okra, Onions, Oregano, Parsley, Parsnips, Peas, Peppermint, Peppers, Pimentos, Potatoes, Pumpkins, Radishes, Rhubarb, Rutabagas, Sage, Salsify, Savory, Shallots, Spearmint, Spinach, Squash, Sugar Beets, Sweet Potatoes, Swiss Chard, Tomatoes, Turnips, Watermelon, Wheat, Seed Crops such as alfalfa, beans, clover, corn (sweet), grasses, peas, potato.

SLUGS AND SNAILS: Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage OR 5 to 10 oz. in ten gallons of water per treated 1,000 sq. ft. OR 1 to 2 oz. in two gallons of water per 200 sq. ft. Where Snails are a problem we suggest the higher rate of application.

Thoroughly water areas to be treated before application. Apply only to soil surface around plants and in rows between plants. Use equipment that applies directly to the soil surface. Do not apply in such a manner as to cause drift of product onto plants. Do not apply to edible parts of plants. For best results, apply in the evening. Repeat as needed for control.

TREE CROPS: Apples, Apricots, Avocados, Bananas, Cherries, Citrus fruits, Grapefruits, Lemons, Limes, Nectarines, Oranges, Peaches, Pears, Pecans, Plums, Stone fruits, Walnuts.

SLUGS AND SNAILS: Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage OR 5 to 10 oz. in ten gallons of water per treated 1,000 sq. ft. OR 1 to 2 oz. in two gallons of water per treated 200 sq. ft. Where snails are a problem we suggest the higher rate of application.

Make application when foliage and fruit are completely dry but soil is wet following an irrigation. Apply under trees, around irrigation pipes, farm buildings and any other place where slugs and snails may congregate. Use equipment that applies directly to soil surface. Do not apply in such a manner as to cause drift of product onto trees. Do not apply directly to trees or fruit. For best results, apply in the evening. Repeat as needed for control.

BERRY CROPS: Blackberries, Blueberries, Boysenberries, Caneberries, Dewberries, Gooseberries, Loganberries, Raspberries, Strawberries.

SLUGS AND SNAILS: Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage OR 5 to 10 oz. in ten gallons of water per treated 1,000 sq. ft. OR 1 to 2 oz. in two gallons of water per treated 200 sq. ft. Where snails are a problem we suggest the higher rate of application.

Thoroughly water areas to be treated before application. Apply only to soil surface around plants and in rows between plants. Use equipment that applies directly to soil surface. Do not apply in such a manner as to cause drift of product onto plants. Do not apply to edible parts of plants. For best results, apply in the evening. Repeat as needed for control.

TERRESTRIAL NONFOOD CROPS

SEED CROPS: Beets, Cabbage, Carrots, Collards, Kale, Kohlrabi, Lettuce, Mustard, Radish, Spinach, Swiss Chard, Tobacco, Turnips.

SLUGS AND SNAILS: Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage OR 5 to 10 oz. in ten gallons of water per treated 1,000 sq. ft. OR 1 to 2 oz. in two gallons of water per treated 200 sq. ft. Where snails are a problem we suggest the higher rate of application.

Thoroughly water areas to be treated before application. Apply only to soil surface around plants and in rows between plants. Use equipment that applies directly to soil surface. Do not apply in such a manner as to cause drift of product onto plants. For best results, apply in the evening. Repeat as needed for control.

9/14

Page 7-(SLUG-FEST ALL WEATHER FORMULA-AG Label)

GREENHOUSE FOOD CROP: Asparagus, Beans, Beets, Cabbage, Carrots, Celery, Corn, Cucumbers, Eggplant, Endive, Lettuce, Melons, Onions, Peas, Peppers, Potatoes, Radish, Spinach, Squash, Tomatoes, Turnips.

SLUGS AND SNAILS: Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage OR 5 to 10 oz. in ten gallons water per treated 1,000 sq. ft. OR 1 to 2 oz. in two gallons of water per treated 200 sq. ft. Where snails are a problem we suggest the higher rate of application.

Water or irrigate areas to be treated before application. Apply only to soil surface around plants, in rows between plants and under benches in greenhouses and around plants; any place where slugs and snails may congregate. Use equipment that applies directly to soil surface. Do not apply in such a manner as to cause drift of product onto plants. Do not apply to plants or edible parts of plants. For best results, apply in the evening. Repeat as needed for control.

GREENHOUSE NONFOOD CROPS: Orchids, Ornamental Plants

SLUGS AND SNAILS: Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage OR 5 to 10 oz. in ten gallons of water per treated 1,000 sq. ft. OR 1 to 2 oz. in two gallons of water per treated 200 sq. ft. Where snails are a problem we suggest the higher rate of application.

Water or irrigate areas to be treated before application. Apply only to soil surface around plants, in rows between plants and under benches in greenhouse and around plants; any place where snails and slugs might congregate. Do not apply directly to plants unless prior experience has shown the plant to be tolerant to this product. Most plants are tolerant, however, some slight tip burning has been noticed on extremely tender orchids.

DOMESTIC AND NONDOMESTIC OUTDOOR: Domestic Dwellings, Farm Buildings, Lath Houses, Ornamental Plants

SLUGS AND SNAILS: Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage OR 5 to 10 oz. in ten gallons of water per treated 1,000 sq. ft. OR 1 to 2 oz. in 2 gallons of water per treated 200 sq. ft. Where snails are a problem we suggest the higher rate of application.

Thoroughly water areas to be treated before application. Use equipment that applies directly to the soil surface. Apply evenly over infested areas and along flower beds adjacent to border or ground cover plants. Also apply to ground at base of fences and walls in shady, protected locations and any other place where slugs and snails may congregate.

10/14

Page 8- Slug Fest All Weather Formula AG label

Do not apply in such a manner as to cause drift of product onto edible plants or plant parts. Do not apply directly to plants unless prior experience has shown the plant to be tolerant, however, some slight tip burning has been noticed on extremely tender orchids. For best results, apply in the evening. Repeat as needed for control.

NON CROP AREAS:

Slugs and Snails: Fallow land; barrier strips; uncultivated non-agricultural areas; recreational areas; non food and non feed brush, weed, or dense vegetative areas; railroad, pipeline, highway, power and telephone rights-of-way and road sides; guardrails and fences; lumberyards; storage areas, industrial plant sites, parking areas; parks, golf courses, other public areas; airport and similar industrial non-crop areas: Apply 2 to 4 quarts in sufficient water per acre. Repeat treatment as needed. Use the higher rate for snail control.

LAWNS (INCLUDING DICHONDRA), ORNAMENTAL GRASSES, TURF, NEWLY SEEDED GRASS GROWN FOR SEED (prior to canopy closure).

SLUGS AND SNAILS: Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage OR 5 to 10 oz. in ten gallons of water per treated 1,000 sq. ft. OR 1 to 2 oz. in two gallons of water per treated 200 sq. ft. Where snails are a problem we suggest the higher rate of application.

Uniformly broadcast directly on lawn or turf. Withhold heavy watering for at least 2 days after treatment. Keep children and pets off the treated area until spray has dried.

Non-AG (Home Garden) label

SLUG-FEST

ALL WEATHER FORMULA

Lasts longer under wet conditions

Active Ingredient:

Metaldehyde (2,4,6,8-tetramethyl-1,3,5,7-tetraoxycyclo-octane)	25%
Other Ingredients:	75%
Total	100%

Contains 2.1 pounds Metaldehyde per gallon.

11/14

KEEP OUT OF REACH OF CHILDREN
WARNING AVISO

Si Usted no entiende la etiqueta, busque a alguien para que se la explique a Usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See side panels for additional precautionary statements.

All applicable directions, restrictions and precautions on the label are to be followed. See Directions for Use in Booklet.

SHAKE WELL

NET CONTENTS

EPA Reg. No. 71096-4
EPA Est. No. 52251-OR-005

OR-CAL, Inc.
29454 Meadowview Road
Junction City, OR 97448
(541)689-4413

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Harmful if swallowed. Avoid breathing spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after use. Keep out of reach of children or domestic animals. Wear chemical resistant gloves made of any waterproof material when handling and applying product.

IMPORTANT: *KEEP CHILDREN AND PETS OUT OF TREATED AREA UNTIL SPRAY HAS DRIED. THIS PESTICIDE MAY BE FATAL TO DOGS OR OTHER PETS IF ANIMAL DRINKS. DO NOT ALLOW PUDDLES TO FORM ON GROUND WHEN MIXING OR CLEANING APPLICATION EQUIPMENT.*

12/14

FIRST AID

If Swallowed-Call a poison control center or doctor immediately for treatment advice.
-Have person sip a glass of water if able to swallow.
-Do not induce vomiting unless told to do so by the poison control center or doctor.
-Do not give anything by mouth to an unconscious person.

If in eyes -Hold eye open and rinse slowly and gently with water for 15-20 minutes.
-Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

If inhaled -Move person to fresh air.
-If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
-Call a poison control center or doctor for further treatment advice.

If on skin or clothing -Take off contaminated clothing.
-Rinse skin immediately with plenty of water for 15-20 minutes.
-Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Telecommunications Network at 1-800-858-7378 for emergency treatment.

Note to Physician

Probable mucosal damage may contraindicate gastric lavage.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water or areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water.

STORAGE AND DISPOSAL
Prohibitions: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.
Storage: Keep pesticides in original container in a locked place out of reach of children. Protect from freezing.
Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash.
Container Disposal: Do not reuse container. Rinse thoroughly before discarding in trash.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes. For use on plants intended for anesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or golf courses or lawns and grounds.

USE PRECAUTIONS

Read entire label before using. Apply directly to the soil area beneath plants as listed on this label. Do not allow spray to come in contact with parts of plant that may be used for food or feed.

MIXING: Shake **SLUG-FEST ALL WEATHER FORMULA** well before mixing. Thoroughly mix spray solution before application. Use entire mixed solution at one time. Do not store mixed solution.

LAWNS AND ORNAMENTALS: Such as Geraniums, Salvia, Primroses, Marigolds, Petunias, Pansies, Dahlias, Begonias, Coleus, Ajuga and Dichondra.

NON CROP AREAS: Fallow land; barrier strips; uncultivated non-agricultural areas; recreational areas; non food and non feed brush, weed, or dense vegetative areas; fences; storage areas.

HOSE-ON TYPE APPLICATOR: Thoroughly water areas to be treated. Use one ounce of **SLUG-FEST ALL WEATHER FORMULA** for each two gallons of solution delivered by your applicator to cover approximately 200 sq. ft.

14/14

Page 12
NON-AG (Home Garden) label - continued

SPRINKLING CAN: Thoroughly water areas to be treated. Use one ounce of SLUG-FEST ALL WEATHER FORMULA to each two gallons of water and sprinkle over areas to be treated.

WHERE SNAILS ARE A PROBLEM: We suggest double rate of application on Ornamentals only.

BEST RESULTS: Apply SLUG-FEST ALL WEATHER FORMULA in the cool of the evening.