



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Biopesticides and Pollution Prevention Division (7511P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

71096-24

Date of Issuance:

4/8/2022

NOTICE OF PESTICIDE:

☒ Registration
☐ Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Slugger 360

Name and Address of Registrant (include ZIP Code):

OR-CAL, Inc.
29454 Meadowview Road
Junction City, OR 97448

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Biopesticides and Pollution Prevention Division prior to use of the label in commerce. In any correspondence on this product, always refer to the above EPA Registration Number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA or the Act).

Registration is in no way to be construed as an endorsement or recommendation of this product by the U.S. Environmental Protection Agency (EPA). In order to protect health and the environment, the Administrator, on his or her motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under the Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration or registration review of your product when the EPA requires all registrants of similar products to submit such data.
2. Submit storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) data as these data requirements are not satisfied. A one-year study is required to satisfy these data requirements. You have 18 months from the date of this registration to provide these data to the EPA.

Signature of Approving Official:

Gina Burnett, Senior Regulatory Advisor
Biochemical Pesticides Branch
Biopesticides and Pollution Prevention Division (7511M)
Office of Pesticide Programs

Date:

4/8/2022

3. Make the following labeling change before you release this product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 71096-24."
4. Submit one (1) copy of the final printed labeling for the record before you release this product for shipment.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains statements or claims substantially differing from statements or claims made in connection with obtaining a FIFRA section 3 registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6. A stamped copy of the labeling is enclosed for your records. Please also note that the record for this product currently contains the following acceptable Confidential Statement of Formula (CSF):

- Basic CSF dated 10/07/2021

If you have any questions, please contact Alex Horansky via email at Horansky.alex@epa.gov.

Sincerely,




Gina Burnett, Senior Regulatory Advisor
Biochemical Pesticides Branch
Biopesticides and Pollution
Prevention Division (7511M)
Office of Pesticide Programs

Enclosure

MASTER LABEL**SLUGGER 360**

Slug & Snail Bait



Active Ingredient:

Iron phosphate 1.0%

Other Ingredients 99.0%

Total 100.0%

SUBLABEL A Home and Garden Use**SUBLABEL B Commercial Agriculture Use**

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side/back/booklet for additional precautionary statements

Home & Garden Use: NET WEIGHT: 1 oz., 13 oz., 26 oz, 0.5 lb, 1 lb, 1.25 lbs, 1.5 lbs. 2 lbs, 2.5 lbs., 3 lbs., 3.2 lbs., 3.5 lbs., 4.25 lbs., 5 lbs., 5.5 lbs, 6 lbs., 7 lbs., 8.5 lbs., 10 lbs., 20 lbs., 25 lbs. and (equivalent metric weights)

Commercial Agriculture Use: NET WEIGHT: 10 lbs, 20 lbs., 25 lbs., 40 lbs., 50 lbs., 55 lbs., 100 lbs., 2000 lbs.

OR-CAL, Inc.

29454 Meadowview Road

Junction City, OR 97448

541-689-4413

EPA Reg. No 71096-

EPA Est. 66876-OR-1

SUBLABEL A Home and Garden Use**SLUGGER 360**

Slug & Snail Bait

Active Ingredient:

Iron phosphate 1.0%

Other Ingredients 99.0%

Total 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side/back/booklet for additional precautionary statements

Home & Garden Use: NET WEIGHT: 1 oz., 13 oz., 26 oz, 0.5 lb, 1 lb, 1.25 lbs, 1.5 lbs. 2 lbs, 2.5 lbs., 3 lbs., 3.2 lbs., 3.5 lbs., 4.25 lbs., 5 lbs., 5.5 lbs, 6 lbs., 7 lbs., 8.5 lbs., 10 lbs., 20 lbs., 25 lbs. and (equivalent metric weights)

OR-CAL, Inc.

29454 Meadowview Road

Junction City, OR 97448

541-689-4413

EPA Reg. No 71096-

EPA Est. 66876-OR-1

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wear protective eyewear and clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

FIRST AID	
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For general information on this product, contact the National Pesticide Information Center (NPIC) at 1-800-858-7378, Monday through Friday, 8 AM to 12 PM PST, or at http://npic.orst.edu</p>	

ENVIRONMENTAL HAZARDS

To protect the environment do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway sidewalk or street back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

HOW TO APPLY: Scatter the slug and snail bait granules on the soil around or near the plants to be protected. For broadcast application use standard broadcast spreaders. For row application use standard granular spreaders. Apply bait evenly at the rate of 0.5-1 lb per 1000 square feet or for smaller areas 0.15 oz or 1 level teaspoon per square yard and reapply as the bait is consumed or at least every two weeks. Do not place in piles. If the ground is dry, wet it before applying bait. The product works best when soil is moist but with little or no standing water. Reapply as the bait is consumed

or at least every two weeks. Apply more heavily if the infestation is severe if the area is heavily watered or after long periods of heavy rain. See specific directions for different plant types and for inside greenhouses.

WHEN TO APPLY: Apply the bait during the evening, as slugs and snails travel and feed mostly by night, or early morning.

WHERE TO APPLY: Treat all likely areas of infestation, especially around the perimeter of garden plots, because these pests travel into plant areas from daytime refuges. They favor damp places around vegetable plants such as beans, tomatoes, lettuce, cabbage, celery, and squash. Other favorite areas are flower gardens, rockeries, hedges, dichondra, lawns, citrus groves, ivy patches, and other ground cover where they obtain shelter by day.

Vegetables

Use the bait to protect any vegetables from slug and snail damage including (but not limited to) **artichokes, asparagus, beans, root and tuber vegetables** (beets, carrots, ginger, parsnip, potato, radish, rutabaga, sweet potato, sugar beet, taro, turnip, yam), **bulb vegetables** (dry bulb onion, garlic, great-headed elephant garlic, green onion, leek, onion, shallot, Welsh onion), **leafy vegetables and leaves of root and tuber and legume vegetables** (arugula, beet, blackeyed pea, celery, lettuce, pea, rhubarb, spinach, Swiss chard, turnip greens, watercress), **cole crops and brassica leafy vegetables** (broccoli, broccoli raab, Brussels sprouts, cabbage, canola, cauliflower, cavalo, Chinese broccoli, Chinese cabbage (bok choy), Chinese cabbage (napa), Chinese mustard cabbage (gai choy), collards, kale, kohlrabi), **legumes** (beans and peas, soybean), **cucurbit vegetables** (cantaloupe, cucumber, edible gourd, honeydew, melon, muskmelon, pumpkin, summer squash, watermelon, winter squash), **fruiting vegetables** (eggplant, ground cherry, okra, pepino, pepper, tomatillo, tomato, **corn** (field and sweet)).

Scatter the bait around the perimeter of the vegetable plot at the rate of 0.5-1 lb per 1000 square feet or for smaller areas 0.15 oz or 1 level teaspoon per square yard to provide a protective barrier for slugs and snails entering the garden plot. If slugs or snails are inside the rows then scatter the bait on the soil around the base of the plants and between the rows.

Herbs

Use the bait to protect herbs from slugs and snails, such as angelica, balm, basil, borage, burnet, chamomile, catnip, chervil, chive, clary, costmary, cilantro, curry, dillweed, horehound, hyssop, lavender, lemongrass, lovage, marigold, nasturtium, parsley, pennyroyal, rosemary, rue, sage, savory, sweet bay, tansy, tarragon, thyme, wintergreen, woodruff and wormwood. Scatter the bait around the perimeter of the herb plot at the rate of 0.5-1 lb per 1000 square feet or for smaller areas 0.15 oz or 1 level teaspoon per square yard. If slugs or snails are inside the rows, then scatter the bait on the soil around the base of the plants and between the rows.

Fruits Including Citrus, and Nuts

Use the bait to protect fruits from slugs and snails including (but not limited to) **banana, plantain, fig, grape, tropical tree fruits** (avocado, guava, mango, papaya, passion fruit), **citrus fruits** (grapefruit, lemon, lime, orange, tangerine), **pome fruits** (apple, crabapple, loquat, mayhaw, pear, quince), **stone fruits** (apricot, cherry, nectarine, peach, plum, prune), **melons, tree nuts** (almond, cashew, chestnut, filbert/hazelnut, macadamia nut, pecan, walnut), **peanuts** (not for use in California), and **pistachios**. For seedlings spread the bait around the base of the stem. Apply at 0.15 oz or 1 level teaspoon per square yard in a 6-inch circular band around the base of the plants to be protected. For older trees spread the bait around the base of the tree to intercept slugs and snails traveling to the trunk. Apply the bait at the rate of 0.5-1 lb per 1000 square feet for orchards using standard fertilizer granular spreaders.

Berries

Use the bait to protect berries from slugs and snails including (but not limited to) **bushberries** (blueberry, currant, elderberry, gooseberry, huckleberry, juneberry, ligonberry, salal), **caneberries** (blackberry, boysenberry, loganberry, red and black raspberry), **cranberry, strawberry**. Spread the bait around the perimeter of the plot to intercept slugs and snails migrating toward the berries. Use a rate of 0.5-1 lb per 1000 square feet or for smaller areas 0.15 oz or 1 level teaspoon per square yard and scatter by hand or with granular spreaders. If slugs and snails are already in the plots then carefully spread bait between the furrows near the base of the plants. For small plots treat around the base of the plants to be protected. Do not spread over the entire area but apply selectively.

Outdoor Ornamentals

Scatter bait in a 6-inch circular band around the base of the plants to be protected at 0.15 oz or 1 level teaspoon per square yard. If plants are next to a grassy area spread the bait between the ornamentals and the grass. Slugs and snails traveling to the plants will encounter the bait before reaching the plant. In these situations scatter the bait around the perimeter of the plot at the rate of 1 lb per 1000 square feet to intercept snails and slugs traveling to the plot.

Greenhouses

Where slugs or snails are a problem in greenhouses scatter the bait in the plant pots of plants being damaged or around pots on greenhouse benches. Apply 1/2 teaspoon per 9-inch pot.

Lawns

Use the bait on lawns as a buffer zone around gardens. When slugs or snails are detected scatter the bait at a rate of 0.15 oz or 1 level teaspoon per square yard. Scatter by hand or with a granular spreader where the slugs or snails are observed.

STORAGE AND DISPOSAL

Do not contaminate water food or feed by storage or disposal.

PESTICIDE STORAGE: Store this product in its original container and keep in a secure storage area out of reach of children and domestic animals.

CONTAINER DISPOSAL: Nonrefillable container.

If empty: Do not reuse or refill this container. Offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

BATCH CODE

[The registrant may use any or none of these optional statements]

WARRANTY

To the extent consistent with applicable law the seller warrants that this product conforms to the chemical description on this label and is reasonably fit for purposes stated on this label only when used in accordance with directions for use. This warranty does not extend to use of this product contrary to label directions or under abnormal use conditions or under conditions not reasonably foreseeable to seller. To the extent consistent with applicable law seller makes no other warranties either expressed or implied.

OR

OUR COMMITMENT TO YOUR SATISFACTION

If you are not happy with the way this product works in your garden please let us know. We will be happy to replace the product or refund your money. Simply send your name address and proof of purchase to [insert name and ' address of supplemental registrant].

OR

The (insert company name) Guarantee

If for any reason you are not satisfied with this product mail us proof of purchase to obtain a full refund of your purchase price.

OR

The [brand name] promise: Satisfaction guaranteed or your money back. If you apply this product as directed and are not completely satisfied send proof of purchase from this package and the original cash register receipt to the address at right (below). We will refund the purchase price you paid.

[The following claims and product information may or may not be presented on the Product's label and labeling]

Quick Facts

Product Facts

What it does

Kills snails and slugs

For use around vegetables, fruit trees, citrus, berries, ornamentals, shrubs, flowers, trees, lawns, gardens, and in greenhouses

Read the Label First

Sold by weight not by volume

Contents may have settled during shipment

This container is made from XX% recycled materials

Resealable waterproof package

The highly compressed granules (pellets) are easy to use, clean to handle ,and economical

Easy to use (ready to use) (RTU) granular (pellet) formulation

Easy to apply granules

Kills snails & (and) slugs

Treats (will treat) x,xxx sq ft

Remains effective after rain or sprinkling

Proven snail & (and) slug killer (kill control)

Convenient

Easy to use

Requires no mixing spraying or special applicators

SATISFACTION GUARANTEED

Money Back Guarantee. If you are not satisfied with this product we will gladly refund your original purchase price.

Can be used in vegetable gardens

For use around vegetables, fruit trees, citrus, berries, ornamentals, shrubs, flowers, trees, lawns, gardens, and in greenhouses

Use around (flowers) (gardens) (vegetables) and (fruit trees)

Can be used around domestic animals (pets) and wildlife

Read Entire Container Label Before Using This Product

The active ingredient in this product is exempt from the requirement for a tolerance when used as a molluscicide in or on all food commodities

For broadcast application standard broadcast applicators may be used such as (but not limited to) Cyclone

For row application standard granular spreaders may be used such as (but not limited to) Gandy and Clampco

Unconditionally guaranteed by [insert name of supplemental registrant]

Baits and Kills

Baits and Kills Slugs and Snails

For household home garden use

The slugs and snails controlled by this product include (but are not limited to)

Deroceras reticulatum (Field slug), *Deroceras laeve* (Smooth slug), *Anon subfuscus* (Dusky slug), *Anon circumscriptus* (Gray garden slug), *Anon hortensis* (Black field slug), *Anon rufus* (Large red slug), *Anon ater* (Large black slug), *Limax flavus* (Spotted garden slug), *Limax tenellus* (Slender slug), *Anohmax columbianus* (Banana slug), *Helix* spp, *Hehcella* spp, and *Cepaea* spp

Kills a wide variety of slugs and snails

Effective against a wide variety of slugs and snails
 No slime trails
 No messy cleanup
 Dead snails no slime trails
 Rain Proof Formula
 Guaranteed (Results)
 Protecting your garden and landscapes from snail and slug damage has never been easier
 Easy (to use) Shaker Applicator
 Easy (to use) (Pour) Spout
 May be used up to and including day of harvest
 Does not contain metaldehyde
 Iron phosphate is a component of soil
 Iron phosphate occurs in soil
 Iron phosphate is a plant nutrient
 Iron phosphate is a plant nutrient that occurs in soils
 Iron phosphate is a plant nutrient that occurs in soil and is used as an ingredient in fertilizers
 Contains iron phosphate a plant nutrient
 Contains iron phosphate a fertilizer
 Contains iron phosphate which is found in soil and which is a plant nutrient
 Contains iron phosphate which is a fertilizer
 Contains iron phosphate which is found in soil
 Contains iron phosphate which is a plant nutrient
 Unconsumed bait will degrade and become part of soil
 Exclusive smaller particle size makes it easier for pests to find
 Questions comments (or medical information) X XXX XXX XXXX
 Kills/controls Giant African Land Snail

GENERAL INFORMATION (WHY SLUG AND SNAIL BAIT IS EFFECTIVE)

This product contains a unique combination of iron phosphate which is a plant nutrient that occurs in soils, with slug and snail bait additives. The bait, which is not ingested by snails and slugs, will degrade and become a part of the soil.

The bait is ingested by slugs and snails when they travel from their hiding places to Plants. Ingestion even in small amounts will cause them to cease feeding. This physiological effect of the bait gives immediate protection to the plants even though the slugs and snails may remain in the area. After eating the bait the slugs and snails cease feeding, become less mobile, and begin to die within three to six days. Dead slugs and snails may not be visible as they often crawl away to secluded places to die. Plant protection will be observed in the decrease in plant damage. This product is effective against a wide variety of slugs and snails and will give protection to home lawns, gardens, greenhouses, outdoor ornamentals, vegetable gardens, fruits, berries, citrus, and crop plants. The bait can be scattered on the lawn or on the soil around any vegetable plants, flowers, or fruit trees or bushes to be protected.

SUBLABEL B Commercial Agriculture Use**SLUGGER 360**

Slug & Snail Bait

Active Ingredient:

Iron phosphate	1.0%
Other Ingredients	<u>99.0%</u>
Total	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side/back/booklet for additional precautionary statements

Commercial Agriculture Use: NET WEIGHT: 10 lbs, 20 lbs., 25 lbs., 40 lbs., 50 lbs., 55 lbs., 100 lbs., 2000 lbs.

OR-CAL, Inc.
29454 Meadowview Road
Junction City, OR 97448
541-689-4413
EPA Reg. No 71096-
EPA Est. 66876-OR-1

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wear protective eyewear and clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

FIRST AID	
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For general information on this product, contact the National Pesticide Information Center (NPIC) at 1-800-858-7378, Monday through Friday, 8 AM to 12 PM PST, or at http://npic.orst.edu	

Personal Protective Equipment (PPE) Requirements

Applicators and other handlers must wear long sleeved shirts and long pants and shoes plus socks. Follow manufacturer's instructions for maintaining PPE. If no such instructions for washables use detergent and hot water. Keep and wash PPE separately from other laundry'

User Safety Recommendations

- Users should wash hands before eating drinking chewing gum using tobacco or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible wash thoroughly and change into clean Clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its

labeling.

Do not apply this product in a manner that will contact workers or other persons either directly or through drift. Only protected workers may be in the area during application. For any requirements specific to your State or Tribe consult the State or Tribal agency representative responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product in accordance with its labeling and with the Worker Protection Standard (40 CFR part 170). This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE), the restricted entry interval, and notification to workers. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS). For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- coveralls,
- chemical resistant gloves made of any water proof material,
- and shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Do not enter or allow others to enter the treated area until applied material is no longer visible.

COMMERCIAL AGRICULTURE

HOW TO APPLY: Scatter the slug and snail bait granules on the soil around or near the plants to be protected.

For broadcast application use standard broadcast spreaders.

For row application use standard granular spreaders.

For aerial application, use aerial equipment calibrated to give correct dosage within the application rates on this label; do not apply when weather conditions favor drift from treated areas. Do not allow dust to drift from the application site and contact people, structures people occupy at any time and the associated property, parks and recreation areas, nontarget crops, aquatic and wetland areas, woodlands, pastures, rangelands, or animals. For aerial applications, use upwind swath displacement and apply only when wind speed is 3 -- 10 mph as measured by an anemometer. If application includes a no-spray zone, do not release product at a height greater than 10 feet above the ground or the crop canopy. The applicator also must use all other measures necessary to control drift.

Apply bait evenly at the rate of 20-44 lbs per acre (0.5-1 lb per 1000 square feet or for smaller areas 0.15 oz or 1 level teaspoon per square yard) and reapply as the bait is consumed or at least every two weeks. Do not place in piles. If the ground is dry, wet it before applying bait. The product works best when soil is moist but with little or no standing water. Reapply as the bait is consumed or at least every two weeks. Apply more heavily if the infestation is severe if the area is heavily watered or after long periods of heavy rain. See specific directions for different plant types and for inside greenhouses.

WHEN TO APPLY: Apply the bait during the evening, as slugs and snails travel and feed mostly by night, or early morning.

WHERE TO APPLY: Treat all likely areas of infestation, especially around the perimeter of garden plots, because these pests travel into plant areas from daytime refuges. They favor damp places around vegetable plants such as beans, tomatoes, lettuce, cabbage, celery, and squash. Other favorite areas are flower gardens, rockeries, hedges, dichondra, lawns, citrus groves, ivy patches, and other ground cover where they obtain shelter by day.

Vegetables

Use the bait to protect any vegetables from slug and snail damage including (but not limited to) **artichokes, asparagus, beans, root and tuber vegetables** (beets, carrots, ginger, parsnip, potato, radish, rutabaga, sweet potato, sugar beet, taro, turnip, yam), **bulb vegetables** (dry bulb onion, garlic, great-headed elephant garlic, green onion, leek, onion, shallot, Welch onion), **leafy vegetables and leaves of root and tuber and legume vegetables** (arugula, beet, blackeyed pea, celery, lettuce, pea, rhubarb, spinach, Swiss chard, turnip greens, watercress), **cole crops and brassica leafy vegetables** (broccoli, broccoli raab, Brussels sprouts, cabbage, canola, cauliflower, cavalo, Chinese broccoli, Chinese cabbage (bok choy), Chinese cabbage (napa), Chinese mustard cabbage (gai choy), collards, kale, kohlrabi), **legumes** (beans and peas, soybean), **cucurbit vegetables** (cantaloupe, cucumber, edible gourd, honeydew, melon, muskmelon, pumpkin, summer squash, watermelon, winter squash), **fruiting vegetables** (eggplant, ground cherry, okra, pepino, pepper, tomatillo, tomato, **corn**

(field and sweet). Scatter the bait around the perimeter of the vegetable plot at the rate of 20-44 lbs per acre (0.5-1 lb per 1000 square feet or for smaller areas 0.15 oz or 1 level teaspoon per square yard) to provide a protective barrier for slugs and snails entering the garden plot. If slugs or snails are inside the rows then scatter the bait on the soil around the base of the plants and between the rows.

Herbs

Use the bait to protect herbs from slugs and snails, such as angelica, balm, basil, borage, burnet, chamomile, catnip, chervil, chive, clary, costmary, cilantro, curry, dillweed, horehound, hyssop, lavender, lemongrass, lovage, marigold, nasturtium, parsley, pennyroyal, rosemary, rue, sage, savory, sweet bay, tansy, tarragon, thyme, wintergreen, woodruff and wormwood. Scatter the bait around the perimeter of the herb plot at the rate of 20-44 lbs per acre (0.5-1 lb per 1000 square feet or for smaller areas 0.15 oz or 1 level teaspoon per square yard). If slugs or snails are inside the rows, then scatter the bait on the soil around the base of the plants and between the rows.

Fruits Including Citrus, and Nuts

Use the bait to protect fruits from slugs and snails including (but not limited to) **banana, plantain, fig, grape** (currant, table and wine), **tropical tree fruits** (avocado, guava, mango, papaya, passion fruit), **citrus fruits** (grapefruit, lemon, lime, orange, tangerine), **pome fruits** (apple, crabapple, loquat, mayhaw, pear, quince), **stone fruits** (apricot, cherry, nectarine, peach, plum, prune), **melons, tree nuts** (almond, cashew, chestnut, filbert/hazelnut, macadamia nut, pecan, walnut), **peanuts** (not for use in California), and **pistachios**. For seedlings spread the bait around the base of the stem. Apply at 0.15 oz or 1 level teaspoon per square yard in a 6-inch circular band around the base of the plants to be protected. For older trees spread the bait around the base of the tree to intercept slugs and snails traveling to the trunk. Apply the bait at the rate of 20-44 lbs per acre (0.5-1 lb per 1000 square feet) for orchards using standard fertilizer granular spreaders.

Berries

Use the bait to protect berries from slugs and snails including (but not limited to) **bushberries** (blueberry, currant, elderberry, gooseberry, huckleberry, junberry, ligonberry, salal), **caneberries** (blackberry, boysenberry, loganberry, red and black raspberry), **cranberry, strawberry**. Spread the bait around the perimeter of the plot to intercept slugs and snails migrating toward the berries. Use a rate of 20-44 lbs per acre (0.5-1 lb per 1000 square feet or for smaller areas 0.15 oz or 1 level teaspoon per square yard) and scatter by hand or with granular spreaders. If slugs and snails are already in the plots then carefully spread bait between the furrows near the base of the plants. For small plots treat around the base of the plants to be protected. Do not spread over the entire area but apply selectively.

Outdoor Ornamentals

Scatter bait in a 6-inch circular band around the base of the plants to be protected at 0.15 oz or 1 level teaspoon per square yard. If plants are next to a grassy area spread the bait between the ornamentals and the grass. Slugs and snails traveling to the plants

will encounter the bait before reaching the plant. In these situations scatter the bait around the perimeter of the plot at the rate of 20-44 lbs per acre (1 lb per 1000 square feet) to intercept snails and slugs traveling to the plot.

Greenhouses

Where slugs or snails are a problem in greenhouses scatter the bait in the plant pots of plants being damaged or around pots on greenhouse benches. Apply 1/2 teaspoon per 9-inch pot.

Outdoor Container-Grown Nursery Plants

Where slugs or snails are a problem in outdoor nurseries, scatter the bait in the plant containers at the rate of 1 tablespoon per container of plants being damaged, or scatter on the soil near the containers at the rate of 1 teaspoon per square yard (24-44 lbs. per acre).

Lawns

Use the bait on lawns as a buffer zone around gardens. When slugs or snails are detected scatter the bait at a rate of 22-44 lbs per acre (0.54 lb per 1000 sq ft or 0.15 oz or 1 level teaspoon per square yard). Scatter by hand or with a granular or standard broadcast spreader where the slugs or snails are observed.

Field Crops

Use the bait to protect field and seed crops from slugs and snails including (but not limited to) beans, field corn, sweet corn, soybeans, sugarbeets, sugarcane, asparagus, beets, broccoli, Brussels sprouts, cabbage, carrots, cauliflower, cucumbers, lettuce, onions, peas, peppers, potatoes, radishes, strawberries, tomatoes, turnips, alfalfa, cotton, flax, hops, peanuts, small grains (barley, oats, rye, wheat), sorghum. At the seedling and later stages apply the bait between the rows and around the perimeter of the field. Scatter pellets at a rate of 20-44 lbs per acre (0.5-1 lb per 1000 square feet). Use the higher dosage rate for heavy infestations.

Artichokes

Use the bait to protect artichokes from slugs and snails. At the seedling and later stages apply the bait within the rows and around the perimeter of the field. Scatter pellets at a rate of 10-44 lbs per acre (0.23-1 lb per 1000 square feet). Use the higher dosage rate for heavier infestations.

Grass and Legumes Grown for seed production, Forage, Sugarcane, and Cereal Crops (such as: wheat, barley, oats and rye): Use the bait on grass grown for seed production, grass crops, sugarcane, legume forage grown for seed production (such as clover and alfalfa), and small cereal grains (such as barley buckwheat, corn, grain amaranth, milo, oats, pearl millet, proso millet, rice, rye, sorghum, triticale, wheat). When broadcast or spread the bait at a rate of approximately 10-44 lbs per acre (0.23-1 lb per 1000 square feet). Use the higher rate for heavy infestations. Scatter by hand or with a granular or standard broadcast spreader where the slugs or snails are observed.

Non-Crop Areas

The bait can be used to protect the following non-crop areas: fallow land; barrier strips; non-food or non-feed brush; weed or dense vegetative areas; railroad, pipeline, highway, power and telephone rights of way and road sides; guardrails and fences; lumberyards; storage areas; industrial facility sites, including yards and walkways around industrial building; parking areas, parks; airports. Apply at a rate of approximately 20-44 lbs. per acre (0.5-1 lb. per 1,000 square feet). Use the higher rate for heavy infestations.

Commercial Turf, Sod, Golf Courses and Grass Buffers around Gardens, Crop Areas and Ornamentals

Use the bait on turf and sod for resale, golf courses and on lawns as a buffer zone around gardens and agricultural crop areas. When slugs or snails are detected scatter the bait at a rate of 22-44 lbs per acre (0.54-1 lb per 1000 square feet). Use the higher rate for heavy infestations. Scatter by hand or with a granular spreader where the slugs or snails are observed.

STORAGE AND DISPOSAL

Do not contaminate water food or feed by storage or disposal.

PESTICIDE STORAGE: Store this product in its original container and keep in a secure storage area out of reach of children and domestic animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Completely empty metal drum, plastic bag, box, or plastic tote into application equipment. Then dispose of empty metal drum, plastic bag, box or plastic tote in a sanitary landfill or by incineration or if allowed by State and local authorities by burning. If burned stay out of smoke.

BATCH CODE

[The registrant may use any or none of these optional statements]

WARRANTY

To the extent consistent with applicable law the seller warrants that this product conforms to the chemical description on this label and is reasonably fit for purposes stated on this label only when used in accordance with directions for use. This warranty does not extend to use of this product contrary to label directions or under abnormal use conditions or under conditions not reasonably foreseeable to seller. To the extent consistent with applicable law seller makes no other warranties either expressed or implied.

OR

OUR COMMITMENT TO YOUR SATISFACTION

If you are not happy with the way this product works in your garden please let us know. We will be happy to replace the product or refund your money. Simply send your name address and proof of purchase to [insert name and ' address of supplemental registrant].

OR

The (insert company name) Guarantee

If for any reason you are not satisfied with this product mail us proof of purchase to obtain a full refund of your purchase price.

OR

The [brand name] promise: Satisfaction guaranteed or your money back. If you apply this product as directed and are not completely satisfied send proof of purchase from this package and the original cash register receipt to the address at right (below). We will refund the purchase price you paid.

[The following claims and product information may or may not be presented on the Product's label and labeling]

Quick Facts

Product Facts

What it does

Kills snails and slugs

For use around vegetables, fruit trees, citrus, berries, ornamentals, shrubs, flowers, trees, lawns, gardens, and in greenhouses

Read the Label First

Sold by weight not by volume

Contents may have settled during shipment

This container is made from XX% recycled materials

Resealable waterproof package

The highly compressed granules (pellets) are easy to use, clean to handle ,and economical

Easy to use (ready to use) (RTU) granular (pellet) formulation

Easy to apply granules

Kills snails & (and) slugs

Treats (will treat) x,xxx sq ft

Remains effective after rain or sprinkling

Proven snail & (and) slug killer (kill control)

Convenient

Easy to use

Requires no mixing spraying or special applicators

SATISFACTION GUARANTEED

Money Back Guarantee. If you are not satisfied with this product we will gladly refund your original purchase price.

Can be used in vegetable gardens

For use around vegetables, fruit trees, citrus, berries, ornamentals, shrubs, flowers, trees, lawns, gardens, and in greenhouses

Use around (flowers) (gardens) (vegetables) and (fruit trees)

Can be used around domestic animals (pets) and wildlife

Read Entire Container Label Before Using This Product

The active ingredient in this product is exempt from the requirement for a tolerance when used as a molluscicide in or on all food commodities

For broadcast application standard broadcast applicators may be used such as (but not limited to) Cyclone

For row application standard granular spreaders may be used such as (but not limited to) Gandy and Clampco

Unconditionally guaranteed by [insert name of supplemental registrant]

Baits and Kills

Baits and Kills Slugs and Snails

For household home garden use

The slugs and snails controlled by this product include (but are not limited to)

Deroceras reticulatum (Field slug), *Deroceras laeve* (Smooth slug), *Anon subfuscus* (Dusky slug), *Anon circumscriptus* (Gray garden slug), *Anon hortensis* (Black field slug), *Anon rufus* (Large red slug), *Anon ater* (Large black slug), *Limax flavus* (Spotted garden slug), *Limax tenellus* (Slender slug), *Anohmax columbianus* (Banana slug), *Helix* spp, *Hehcella* spp, and *Cepaea* spp

Kills a wide variety of slugs and snails

Effective against a wide variety of slugs and snails

No slime trails

No messy cleanup

Dead snails no slime trails

Rain Proof Formula

Guaranteed (Results)

Protecting your garden and landscapes from snail and slug damage has never been easier

Easy (to use) Shaker Applicator

Easy (to use) (Pour) Spout

May be used up to and including day of harvest

Does not contain metaldehyde

Iron phosphate is a component of soil

Iron phosphate occurs in soil

Iron phosphate is a plant nutrient

Iron phosphate is a plant nutrient that occurs in soils

Iron phosphate is a plant nutrient that occurs in soil and is used as an ingredient in fertilizers

Contains iron phosphate a plant nutrient

Contains iron phosphate a fertilizer

Contains iron phosphate which is found in soil and which is a plant nutrient

Contains iron phosphate which is a fertilizer

Contains iron phosphate which is found in soil

Contains iron phosphate which is a plant nutrient
 Unconsumed bait will degrade and become part of soil
 Exclusive smaller particle size makes it easier for pests to find
 Questions comments (or medical information) X XXX XXX XXXX
 Kills/controls Giant African Land Snail

GENERAL INFORMATION (WHY SLUG AND SNAIL BAIT IS EFFECTIVE)

This product contains a unique combination of iron phosphate which is a plant nutrient that occurs in soils, with slug and snail bait additives. The bait, which is not ingested by snails and slugs, will degrade and become a part of the soil.

The bait is ingested by slugs and snails when they travel from their hiding places to Plants. Ingestion even in small amounts will cause them to cease feeding. This physiological effect of the bait gives immediate protection to the plants even though the slugs and snails may remain in the area. After eating the bait the slugs and snails cease feeding, become less mobile, and begin to die within three to six days. Dead slugs and snails may not be visible as they often crawl away to secluded places to die. Plant protection will be observed in the decrease in plant damage. This product is effective against a wide variety of slugs and snails and will give protection to home lawns, gardens, greenhouses, outdoor ornamentals, vegetable gardens, fruits, berries, citrus, and crop plants. The bait can be scattered on the lawn or on the soil around any vegetable plants, flowers, or fruit trees or bushes to be protected.