

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

March 20, 2024

Maria F. Abildgaard Regulatory Affairs Manager AgroFresh Inc 510-530 Walnut Steet, Suite 1350 Philadelphia, PA 19106

Subject: Non-PRIA (Pesticide Registration Improvement Act) Labeling Amendment -

Amendment to Tank Mix Directions to Include Adjuvant; Amendment to Simplify

Ground and Aerial Application

Product Name: AF-701

EPA Registration Number: 71297-17

EPA Receipt Date: 6/1/2023 Action Case Number: 00452629

Dear Ms. Abildgaard:

The amended labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable.

This approval does not affect any terms or conditions that were previously imposed on this registration. You continue to be subject to existing terms or conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release this product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR § 152.3.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the U.S. Environmental Protection Agency (EPA). If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or

Page 2 of 2 EPA Reg. No. 71297-17 Action Case No. 00452629

misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains statements or claims substantially differing from statements or claims made in connection with obtaining a FIFRA section 3 registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Chris Pfeifer of my team by phone at (202) 566-1599 or via email at <u>pfeifer.chris@epa.gov</u>.

Sincerely,

James Parker, Team Leader Biochemical Pesticides Branch Biopesticides and Pollution Prevention Division (7511M) Office of Pesticide Programs

Enclosure

[Text in brackets is optional]

AF-701

Protects fruit and vegetable crops from the various effects of ethylene. [Alternate Brand Name: HarvistaTM 1.3 SC]

ACTIVE INGREDIENT:	% By WT.
1-Methylcyclopropene	1.3%
OTHER INGREDIENTS	98.7%
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION/ PRECAUCION

Contains 0.143 lbs 1-MCP/gallon

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

[See[front][back][side][inside] panel for First Aid Instructions and [Leaflet][Booklet] for complete Precautionary Statements and Directions for Use.]

FIRST AID			
If in eyes	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. 		
If on skin or clothing	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. 		
If inhaled	 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. 		
If swallowed	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. 		

POISON CONTROL CENTER 1-800-222-1222

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For general information on product use, etc., call the National Pesticides Information Center at 1-800-858-7378. For emergencies, call the poison control center 1-800-222-1222.

Lot No. on container

EPA Reg. No. 71297-17

EPA Est. No.

Net Contents (on container):

[1 gallon] [4 gallons] [55 gallons] [2 gallons] [5 gallons] [275 Gallons]

[2.5 gallons] [15 gallons] [330 Gallons] [3 gallons] [500 Gallons]

Produced for: AgroFresh Inc., 510-530 Walnut Street, Suite 1350

Philadelphia PA 19106, USA, Tel: 1-866-206-1001

NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions and Storage and Disposal Instructions. If the Conditions of Sale and Warranty are not acceptable, return the product unused within thirty days of purchase to the place of purchase.

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to Supplemental Labeling entitled AGRICULTURAL USE REQUIREMENTS in the DIRECTIONS FOR USE section of the labeling for information about this standard.

GENERAL INFORMATION

Harvista is a specialized formulation of 1-MCP (Methylcyclopropene) used to counteract the undesirable effects of ethylene on a wide range of crops. Benefits may include:

- Delayed ripening and associated benefits
- Management of harvest timing
- Increased retention of fruit firmness
- Increased yields and fruit size
- Reduced pre-harvest fruit drop
- Reduced ethylene production
- Improved color development
- Maintenance of fruit firmness and acid levels
- Increase in soluble solids
- Delayed starch hydrolysis
- Delayed onset and incidence of watercore
- Enhanced storage potential
- Enhanced fruit set

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS - CAUTION Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with eyes, skin or clothing. Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking chewing gum using tobacco or using the toilet. Remove contaminated clothing and wash before reuse. Avoid breathing vapor.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves (made of any waterproof material)
- Protective eyewear (such as goggles, safety glasses, or face shield)

Follow manufacturer's instructions for cleaning / maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing or other absorbent materials that have been fogged or heavily contaminated with the concentrated product. Do not reuse them.

USER SAFETY RECOMMENDATIONS: Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing / PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing

ENVIRONMENTAL HAZARDS

For terrestrial use: Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate surface water when disposing of equipment wash water or rinsate. This pesticide is moderately toxic to fish and aquatic invertebrates.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Workers Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the **restricted-entry interval (REI) of 4 hours** unless wearing the appropriate PPE.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Long -sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves (made of any waterproof material)
- Protective eyewear (such as goggles, safety glasses, or face shield)

Read the entire label before using this product.

APPLICATION INSTRUCTIONS

AF-701 is packaged in plastic containers, such as pails and totes. While the package is opened, stand upwind from the container. DO NOT ADD ANY WATER TO THE CONTAINER.

Rate, time and mode of application: Refer to instructions in crop tables.

Plant Safety: This product has been tested on numerous plant varieties with no phytotoxic effects. However, since this product has not been tested on all plant varieties or in combination with all available adjuvants or tank mixes, the manufacturer recommends testing on a small number of plants to check for adverse plant effects before applying on a larger number of plants.

Incompatibility: Do not mix or allow contact with products containing copper.

For treatment of FRUITS AND VEGETABLES.

CROP	RATE fl oz/acre	Restrictions
Apple	48 to 256 fl oz	Do not apply more than 256 fl oz/acre per crop.
Pear		

Blueberry	48 to 256 fl oz	Do not apply more than 256 fl oz/acre per crop.
Kiwifruit	12 to 48 fl oz	Do not apply more than 48 fl oz /acre per crop.
BRASSICA LEAFY VEGETABLES (Crop Group 5):[*] Broccoli[*]; Broccoli, Chinese[*]; Broccoli raab[*]; Brussels sprouts[*]; Cabbage[*]; Cabbage, Chinese (bok choy) [*]; Cabbage, Chinese (napa) [*]; Cabbage, Chinese mustard[*]; Cauliflower[*]; Cavalo broccoli[*]; Collards[*]; Kale[*]; Kohlrabi[*]; Mustard Greens[*]; Mustard Spinach[*]; Rape Greens[*].	12 to 256 fl oz	Do not apply more than 256 fl oz /acre per crop.
CUCURBIT VEGETABLES (Crop Group 9) [*]: Melon[*]: Cantaloupe[*], Citron melon[*], Honeydew[*], Muskmelon[*] and Watermelon[*]; Cucumber[*]; Gherkin[*]; Summer Squash[*]; cultivars, varieties, and/or hybrids of these[*].	121 to 256 fl oz	Do not apply more than 256 fl oz / acre per crop.
Persimmon[*].		Do not apply more than 256 fl oz / acre per crop.
CROP	RATE fl oz/acre	Restrictions

STONE FRUIT (Crop Group 12-12A)[*]: Capulin[*]; Cherry, black[*]; Cherry, Nanking[*]; Cherry, sweet[*]; Cherry, tart[*]; cultivars, varieties, and/or hybrids of these.[*]	121 to 256 fl oz	Do not apply more than 256 fl oz / acre per crop.
STONE FRUIT (Crop Group 12-12B)[*]: Peach[*]; Nectarine[*]	121 to 256 fl oz	Do not apply more than 256 fl oz / acre per crop.
STONE FRUIT (Crop Group 12-12C) [*]: Apricot[*]; Apricot, Japanese[*]; Jujube, Chinese[*]; Plum[*]; Plum, American[*]; Plum, beach[*]; Plum, Canada[*]; Plum, cherry[*]; Plum, Chickasaw[*]; Plum, Damson[*]; Plum, Japanese[*]; Plum, Klamath[*]; Plumcot[*]; Plum, prune[*]; Sloe[*]; cultivars, varieties, and/or hybrids of these.[*]		
Table grapes [*]	12 to 256 fl oz	Do not apply more than 256 fl oz /acre per crop.

[*Not for Use in California]

GROUND APPLICATION

AF-701 must be applied using either an inline co-injection system or as a tank mix with an approved Cycloamylose adjuvant such as Havista Mix AF14050.

In-line co-injection application

AF-701 can be applied from a separate nurse tank through an in-line injection into the water spray line of a conventional spray equipment. Applicators should follow the use rate recommendations in the table given above for a specific crop. The co-injected water spray volume should be at least 2 gal/ac and as high as needed for desired application performance. Tank mix adjuvants such as surfactants and spray oils may be added into the water tank for better wetting, spreading, or other desirable results. Other agricultural products may be added into the water tank as co-application partners for convenience and time saving, but an AgroFresh authorized professional must be consulted to ensure there are no compatibility issues.

Tank mix spray application

Alternatively, AF-701 can be applied as an aqueous tank mix product by a conventional spray equipment using an approved Cycloamylose adjuvant or equivalent such as Havista Mix AF14050. AF14050 must be used at a minimum of 5% by weight of the spray water to ensure the spray stability and performance. An example of spray solution preparation procedure may include: 1) add water into the spray tank in the amount of ~80% of the final spray volume; 2) add AF14050 into the tank at a rate of 41.7 lbs per 100 gal water in the final spray volume, and mix till no solid visible; 3) add AF-701 at the use rate recommended in the table above; 4) top off with addition water needed to make the final spray volume. The spray volume should be at least 5 gal/ac and is recommended to be as low as practical for better efficiency and productivity. Tank mix adjuvants such as surfactants and spray oils may be added into the tank prior to the addition of AF-701 for better wetting, spreading, or other desirable results. Other agricultural products may also be added into the tank prior to the addition of AF-701 as co-application partners for convenience and time saving, but an AgroFresh authorized professional must be consulted to ensure there are no compatibility issues.

AERIAL AND DRONE APPLICATION

AF-701 must be applied using either an inline co-injection system or as a tank mix with an approved Cycloamylose adjuvant such as Havista Mix AF14050.

Spray at a height of fifteen (15) feet or less above the tree canopy unless a greater height is required for safety.

Nozzles must always point down, perpendicular to the ground.

In-line co-injection application

AF-701 can be applied from a separate nurse tank through an in-line injection into the water spray line of a typical aerial or drone spray equipment. Applicators should follow the use rate recommendations in the table given above for a specific crop. The co-injected water spray volume should be at least 2 gal/ac and up to 20 gal/ac. Tank mix adjuvants such as surfactants and spray oils may be added into the water tank for better wetting, spreading, or other desirable results. Other agricultural products may be added into the water tank as co-application partners for convenience and time saving, but an AgroFresh authorized professional must be consulted to ensure there are no compatibility issues.

Tank mix spray application

Alternatively, AF-701 can be applied as an aqueous tank mix product by a typical aerial or drone spray equipment using adjuvant, AF14050. AF14050 must be used at a minimum of 5% by weight of the spray water to ensure the

spray stability and performance. An example of spray solution preparation procedure may include: 1) add water into the spray tank in the amount of ~80% of the final spray volume; 2) add AF14050 into the tank at a rate of 41.7 lbs per 100 gal water in the final spray volume, and mix till no solid visible; 3) add AF-701 at the use rate recommended in the table above; 4) top off with addition water needed to make the final spray volume. The spray volume should be at least 5 gal/ac and is recommended to be as low as practical for better efficiency and productivity. Tank mix adjuvants such as surfactants and spray oils may be added into the tank prior to the addition of AF-701 for better wetting, spreading, or other desirable results. Other agricultural products may also be added into the tank prior to the addition of AF-701 as co-application partners for convenience and time saving, but an AgroFresh authorized professional must be consulted to ensure there are no compatibility issues.

IMPORTANT CONSIDERATIONS FOR USE OF AF-701

- Do not apply when temperatures exceed 95 degrees.
- Do not use overhead mist cooling or irrigation equipment for at least 1 hour following application.
- Do not apply if rain is expected within 1 hour of application.
- Combination effects and potential interactions between AF-701 and products containining ethephon have not been thoroughly evaluated. Use of these products on crops previously treated with AF-701 may reduce the benefits provided by AF-701 at harvest or after storage.
- Do not apply in winds exceeding 10mph
- Do not allow product to contact any copper or mix with products containing copper.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original packaging in a cool, dry place.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or

at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill container smaller than 2 (two)

gallons. Offer for recycling, if available.

For non refillable container, triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with concentrated salt water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure a second time with salt water and a third time with water.

For containers greater than 50 lbs: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with concentrated salt water. Replace and tighten closures. Tip container on its side and roll it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure a second time with salt water and a third time with water.

Then offer for recycling if available, or puncture and dispose of in a sanitary landfill or by incineration. Do not burn unless allowed by the State and local ordinances.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of AGROFRESH INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of AGROFRESH INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, AGROFRESH INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL, THIS

PRODUCT IS SOLD "AS IS" TO THE EXTENT CONSISTENT WITH APPLICABLE LAW. AGROFRESH INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT AGROFRESH INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL, BUYER OR USER MUST SEND WRITTEN NOTICE OF SUCH CLAIM TO THE FOLLOWING ADDRESS, TO THE EXTENT REQUIRED BY APPLICABLE LAW:

AGROFRESH INC.

ATTENTION: LAW DEPARTMENT

510-530 WALNUT STREET - SUITE 1350

PHILADELPHIA. PA 19106

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, OR STRICT LIABILITY.

Harvista is a registered trademark of AgroFresh Inc. [Patent No. XXXXX]