

71368-11

3-30-2004

1/13

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 11-30-93

(A) EPA	United States Environmental Protection Agency Office of Pesticide Programs (H7505C) Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number _____
	Application for Pesticide:		Please Attach An Original Form

Section I

1. Company/Product Number 71368-11	2. EPA Product Manager J. Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Nufarm Weedone LV 6 Broadleaf Herbicide	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Nufarm Inc. 1333 Burr Ridge Parkway, Suite 125A Burr Ridge, IL 60521 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section II

<input type="checkbox"/> Amendment - Explain Below	<input type="checkbox"/> Final printed labels in response to Agency letter dated	NOTIFICATION MAR 30 2004
<input type="checkbox"/> Resubmission in response to Agency letter dated	<input type="checkbox"/> "Me Too" Application	
<input checked="" type="checkbox"/> Notification - Explain below	<input type="checkbox"/> Other - explain below	


Explanation: Use additional page(s) if necessary. (For Section I and Section II).
Notification of addition of California restrictions.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Sections 12 and 14 of FIFRA.

Section III

1. Material This Product Will Be Packaged In:			
Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No <small>*Certification must be submitted.</small>	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No If "Yes," Unit Package Wt. No. Per Container	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No If "Yes," Package Wt. No. Per Container	2. Type of Container <input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label or <input checked="" type="checkbox"/> Container		4. Size(s) of Retail Container 1 qt.; 1/2, 1, 2 1/2, 5, 30, 55, 110, 220 gal.; bulk	5. Location of Label Directions <input checked="" type="checkbox"/> On Label or <input checked="" type="checkbox"/> On Labeling accompanying product
6. Manner In Which Label Is Affixed To Product <input checked="" type="checkbox"/> Lithograph <input checked="" type="checkbox"/> Other (<u>Self-Adhesive Integrated Label/Booklet</u>) <input checked="" type="checkbox"/> Paper Glued <input type="checkbox"/> Stenciled			

Section IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Theodore D. Head	Title Product Registration Manager / Agriculture	Telephone No. (Include Area Code) (708) 754-3330
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Product Registration Manager / Agriculture	
4. Typed Name Theodore D. Head	5. Date 3-16-04	

2/13

NOTIFICATION
MAR 30 2004



Weedone[®] LV6 EC

Broadleaf Herbicide

Weed And Brush Control In Small Grains, Field Corn, Rangeland, Pastures,
Roadsides And Fencerows

ACTIVE INGREDIENT:	
2,4-Dichlorophenoxyacetic acid, isooctyl ester	86.6%*
INERT INGREDIENTS:	13.4%
	100.0%

*2,4-Dichlorophenoxyacetic acid equivalent 57.5% by weight or 5.4 lbs./gal. Isomer specific by AOAC method No. 978.05

EPA REG. NO. 71368-11

EPA EST. NO. 228-IL-1

KEEP OUT OF REACH OF CHILDREN
CAUTION

See Inside for Additional Precautionary Statements

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300.
For Medical Emergencies Only, Call 877-325-1840.

NET CONTENTS 2.5 GALLONS (9.46L)

Manufactured by:
Nufarm, Inc.
Burr Ridge, IL

071368-00011. 20040303.wlv6ec

3/13

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Get medical attention, preferably an ophthalmologist.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.</p>	
NOTE TO PHYSICIAN	
<p>No specific antidote is available. All treatments should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.</p> <p>This product contains petroleum distillates. If large amounts, greater than 1 mg/kg body weight have been ingested, the stomach should be evacuated by gastric intubation with the aid of a cuffed endotracheal tube to prevent aspiration of petroleum distillates. After removal of stomach contents, wash stomach by instilling 30 to 50 grams of activated charcoal in 3 to 4 ounces of water through the stomach tube and again remove stomach contents. Avoid oily laxatives.</p>	

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid breathing vapors or spray mist. Do not get in eyes, on skin or on clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other handlers must wear: long-sleeved shirt and long pants, waterproof gloves, shoes plus socks and protective eye wear.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE (personal protective equipment) may be reduced or modified as specified in the WPS.

For containers over 1 gallon but less than 5 gallons, mixers and loaders who do not use a mechanical system (probe and pump) to transfer the contents of this container must wear coveralls or chemical-resistant apron in addition to the other required PPE.

For containers of 5 gallons or more, a mechanical system (probe and pump) must be used for transferring the contents of the container. If the contents of a non-refillable pesticide containers are emptied, the probe must be rinsed before removal. If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4)] the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

4/13

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Do not apply when weather conditions favor drift from treated areas. Do not use the same spray equipment for other purposes unless thoroughly cleaned.

Do not contaminate water used for irrigation or domestic purposes.

Do not apply this product directly to, or permit spray mist to drift onto cotton, okra, grapes, tomatoes, fruit trees, vegetables, flowers or other desirable crop or ornamental plants which are susceptible to 2,4-D herbicide. Do not apply near susceptible plants since very small quantities of the 2,4-D will cause severe injury during the growing or dormant periods. Crops contacted by sprays or spray drift may be killed or suffer significant stand loss with extensive quality and yield reduction.

Do not apply when a temperature air inversion exists. Such a condition is characterized by little or no air movement and an increase in air temperature with an increase in height. In humid regions a fog or mist may form. An inversion may be detected by producing a smoke column and checking for a layering effect. If questions exist pertaining to the existence of an inversion, consult with local weather services before making an application.

Use coarse sprays to minimize drift. Do not apply with hollow cone type insecticide or other nozzles that produce fine spray droplets.

Drift from aerial or ground application may be reduced by: (1) applying as near to the target as possible in order to obtain coverage; (2) by increasing the volume of spray mix per acre; (3) by decreasing the pounds of pressure at the nozzle tips; (4) by using nozzles which produce a coarse spray pattern; (5) by not applying when wind is blowing toward susceptible valuable plants.

At high air or ground surface temperatures, vapors from this product may injure susceptible plants.

MIXING AND LOADING: Most cases of ground water contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of ground water supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent ground water contamination.

DIRECTIONS FOR USE

It is a Violation of Federal Law To Use This Product In A Manner Inconsistent With Its Labeling. Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil or water is: coveralls, waterproof gloves, shoes plus socks and protective eye wear.

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food or feed by storage or disposal. Store in original container in a dry, secured storage area. Keep container tightly closed when not in use.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate ground water. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse or equivalent. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

5/13

GENERAL CAUTIONS AND RESTRICTIONS

Do not apply this product through any type of irrigation system.
Do not use in or near a greenhouse.

MIXING INSTRUCTIONS

Add 1/2 the required amount of water to the spray tank, then add the Nufarm WEEDONE® LV6 EC Broadleaf Herbicide with agitation, and finally, the balance of the water with continued agitation. This material forms an emulsion in water, not a solution. This tends to separate on standing. Provide agitation to prevent such separation and insure uniform spray mixtures.

COMPATIBILITY

If this herbicide is to be tank mixed with fertilizers or with other pesticides, compatibility should be tested prior to mixing. To test for compatibility, use a small container and mix a small amount (0.5 to 1 quart) of spray, combining all ingredients in the same ratio as the anticipated use. If any indications of physical incompatibility develop, do not use this mixture for spraying. Indications of incompatibility usually will appear within 5 to 15 minutes after mixing.

Read and follow the label of each tank mix product used for precautionary statements, directions for use, geographic and other restrictions.

COMBINATION WITH LIQUID NITROGEN FERTILIZER

Use 1/3 to 1/2 pint of Nufarm WEEDONE® LV6 EC Broadleaf Herbicide per acre for weeding and feeding corn, small grains or grass pastures as directed on this label.

Use fertilizer at rates recommended by supplier or Extension Service Specialist.

Fill the spray tank about half full with the liquid fertilizer, then add the Nufarm WEEDONE® LV6 EC Broadleaf Herbicide with vigorous agitation, and complete filling the tank with fertilizer. Apply immediately and continue agitation in the spray tank during application. Application during very cold weather (near freezing) is not advisable. Do not allow mixture to stand overnight. Incompatibility may be encountered with some fertilizer brands or under some environmental conditions. If in doubt, test a small sample in the dilution ratio planned for application.

NOTE: Fertilizers can increase foliage contact burn of herbicides. Reducing the fertilizer rate and concentration will reduce the hazard of leaf burn.

GENERAL INFORMATION

Nufarm WEEDONE® LV6 EC Broadleaf Herbicide contains the iso-octyl ester of 2,4-D, the original and one of the best low volatile esters. In cropland, this herbicide is more effective than amines for controlling hard-to-kill weeds such as bindweed, thistle, smartweeds, wild garlic, curled dock, tansy ragwort and wild onions.

INJURY TO CROPS FROM THIS HERBICIDE MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY, DO NOT USE THIS PRODUCT.

Crop varieties vary in response to 2,4-D and some are easily injured. Apply this herbicide to varieties known to be tolerant to 2,4-D. If you are uncertain concerning tolerant varieties or local use situations that may affect crop tolerance to 2,4-D, consult your seed company, state Agricultural Extension Service or qualified crop consultant for advice.

GENERAL WEED LIST

ANNUAL AND BIENNIAL WEEDS

annual yellow sweetclover
*beggarticks
bull thistle
coffeeweed
*common broomweed
common burdock
common cocklebur
common evening primrose
common lambsquarters
croton (Texas or woolly)
hairy galinsoga
jimsonweed

*knotweed
*mallow (venice, dwarf, little)
marshelder
morning glory (common, ivy, woolly)
mustards (except blue mustard)
pepperweeds (except perennial)
**pigweeds (Amaranthus spp.)
*prickly lettuce
ragweed (common, giant)
rough fleabane
*Russian thistle

salsify (western or common)
*smartweeds (annual species)
sowthistle (annual or spiny)
sunflower
velvetleaf
*vervains
vetches
*wild carrot
wild lettuce
wild parsnips
wild radish

PERENNIAL WEEDS

*alfaifa	*dogbanes	*nettles (including stinging)
*bindweed (hedge, field & European)	*goldenrod	plantains
blue lettuce	*ground ivy	sowthistle (perennial)
*Canada thistle	*hawkweed (orange)	*tansy ragwort
catnip	healall	*vervains
chicory	*hoary cress	*western ironweed
dandelion	Jerusalem - artichoke	*wild garlic
*docks	*many-flowered aster	*wild onion

*These species may require repeat applications and/or use of the higher rate recommended on this product label even under ideal conditions for applications.

**Control of pigweeds in the High Plains area of Texas and Oklahoma may not be satisfactory with this product.

SPECIFIC USE DIRECTIONS

CEREAL GRAINS: WHEAT, BARLEY AND RYE (not underseeded with legumes)

WEEDS	AMOUNT OF WEEDONE® LV6 EC PER ACRE	DIRECTIONS
Postemergence Annual and biennial weeds Perennial broadleaf weeds	1/3 to 1-1/3 pints*	Apply after grain is well tillered (usually about 4 to 8 inches high) but before boot. Do not spray grain in boot to dough stage.
Wild onion or garlic	1 to 1-1/3 pints*	Apply 1 pint when grain is well tillered and wild garlic or onion plants are small. Apply 1-1/3 pints after harvest in the crop stubble. For control of new fall growth of wild onion or garlic, refer to FALLOWLAND use directions.
Emergency Weed Control in Wheat Perennial broadleaf weeds	2 pints*	Apply when weeds are approaching bud stage, but do not spray grain during the boot to dough stage. The 2 pints per acre application can produce injury to wheat. Balance the severity of your weed problem against the possibility of crop damage. Where perennial weeds are scattered, spot treatment is suggested to minimize the extent of crop injury.
Preharvest	2/3 to 1-1/3 pints*	Apply when grains are in the hard dough stage to suppress large weeds that may interfere with harvest. Best results will be obtained when soil moisture is sufficient to induce succulent weed growth.

*Use the lower rate if small annual and biennial weeds are the major problem. Use the higher rate if perennial weeds or annual and biennial weeds are present which are in the hard-to-kill categories as determined by local experience. The higher rates increase the risk of grain injury and should be used only where the weed control problem justifies the risk of grain damage. Do not apply to grain in the seedling stage.

Note: Maximum rate 1 pint/acre from jointing to before boot stage. Treatment at this growth stage (jointing to before boot) should be used where such increased risk of injury to crop is acceptable. Higher rates listed on this label are more likely to cause crop injury.

RESTRICTIONS AND LIMITATIONS FOR USE ON CEREAL GRAINS

- For aerial application on grain: Apply this product in 2 or more gallons of water per acre.
- For ground application: A minimum of 10 to 15 gallons of water per acre is recommended for proper spray coverage.
- Do not mix with oil for crop uses.
- Do not permit dairy animals or meat animals being finished for slaughter to forage treated grain fields within 2 weeks after treatment.
- Do not feed treated straw to livestock if a preharvest treatment or emergency treatment as described above is applied.

7/13

FIELD CORN

WEEDS	AMOUNT OF WEEDONE® LV6 EC PER ACRE	DIRECTIONS
Preplant	2/3 to 1-1/3 pints	To control emerged broadleaf weed seedlings or existing cover crops prior to planting corn, apply 7 to 14 days before planting. Do not use on light, sandy soil, or where soil moisture is inadequate for normal weed growth. Use high rate for control of less susceptible weeds or cover crops such as alfalfa.
Preemergence	1-1/3 to 2 pints	To control broadleaf weeds and suppress annual grasses, apply 3 to 5 days after planting, but before corn emerges. Use high rate on soil high in organic matter. Do not use on light, sandy soil, or where soil moisture is inadequate for normal weed growth.
Postemergence Annual broadleaf weeds Perennial broadleaf weeds	1/3 pint 1/2 pint	Avoid spraying just after corn leaves unfold as injury may occur. Apply when weeds are small and corn is less than 8 inches tall. If corn more than 8 inches tall must be sprayed, use drop pipes. Spray when weeds are in the bud to bloom stage. If corn is more than 8 inches tall, use drop pipes to keep spray off corn leaves. Do not spray corn from tassel to dough stage. 2,4-D may make corn brittle. Winds or cultivation may cause stalk breakage while brittle.
Preharvest	2/3 to 1-1/3 pints	After the hard dough (denting) stage, to suppress weeds that interfere with harvest such as bindweed, cocklebur, dogbane, jimsonweed, ragweed, sunflower and velvetleaf, and to decrease production of weed seeds, spray with air or ground equipment. The high rate will be needed for tough weeds under stress.

RESTRICTIONS AND LIMITATIONS FOR USE ON FIELD CORN

- Do not forage or feed corn fodder for 7 days following application.

GRAIN SORGHUM (milo)

WEEDS	AMOUNT OF WEEDONE® LV6 EC PER ACRE	DIRECTIONS
Annual broadleaf weeds Perennial broadleaf weeds	1/2 pint 3/4 pint	Treat when sorghum is 6 to 15 inches tall (to top of canopy). If sorghum is taller than 8 inches to top of canopy, use drop nozzles to keep spray off crop foliage. Do not treat during boot, flowering, or early dough stages.

RESTRICTED AND LIMITATIONS FOR USE ON SORGHUM

- Do not forage or feed fodder for 7 days following application.
- A minimum application volume of 5 gallons per acre by air or 10 gallons per acre by ground is recommended.

SOYBEANS (Preplant Only)*

WEEDS	AMOUNT OF WEEDONE® LV6 EC PER ACRE	DIRECTIONS
Postemergence	1/2 to 2/3 pint	Apply not less than 7 days prior to planting soybeans, when weeds are small and actively growing. Use the higher rate on larger weeds and when perennials are present. Some weeds may require repeat treatment for adequate control (see GENERAL WEED LIST and below).
	> 2/3 to 1-1/3 pints	Apply not less than 30 days prior to planting soybeans, when weeds are actively growing. Some weeds may require repeat treatment for adequate control (see GENERAL WEED LIST and below).

(continued)

8/13

SOYBEANS (Preplant Only) (continued)

WEEDS	AMOUNT OF WEEDONE® LV6 EC PER ACRE	DIRECTIONS
Postemergence		<p>In addition to those weeds found on the GENERAL WEED LIST, this product will suppress or control the following broadleaf weeds frequently encountered in reduced tillage soybean production systems: bullnettle, smallflowered bittercress, Carolina geranium, smallflowered buttercup, common and rough cinquefoil, red clover* horseweed or maretail, mousetail, wild mustard, field pennycress, cutleaf evening primrose, common purslane, speedwell, and Virginia copperleaf.</p> <p>* These weeds are only partially controlled.</p> <p>Apply no more than 1-1/3 pints of this herbicide in one season prior to planting soybeans. After applying, plant soybean seed as deep as practical or at least 1-1/2 to 2 inches deep. Adjust the planter press wheel, if necessary, to ensure that planted seed is completely covered.</p> <p>If desired, this product may be applied pre-plant to soybeans in tank mixtures with other herbicides such as Poast®, Poast Plus®, Roundup®, Roundup D-Pak®, Honcho®, Gramoxone® Extra, Prowl®, Pursuit Plus®, Scepter®, Scepter 70 DG®, Squadron® and others that are registered for pre-plant soybean use.</p> <p>NOTE: Unacceptable injury to soybeans planted in fields previously treated with this herbicide may occur. Whether or not soybean injury occurs and the extent of the injury will depend on weather and agronomic factors such as the amount of weed vegetation and previous crop residue present that may be in effect between the time of herbicide application and the emergence of the soybean plant.</p>

RESTRICTIONS AND LIMITATIONS FOR USE IN SOYBEANS (PRE-PLANT)

- Do not apply this product prior to planting soybeans if you are not prepared to accept the results of soybean injury including possible loss of stand and yield.
- Do not apply this product when weather conditions such as temperature, air inversions, or wind favor drift from treated areas to susceptible plants.
- Apply no more than 1-1/3 pints of this herbicide per acre in one season prior to planting soybeans.
- Do not mow or cultivate weeds prior to treating with this herbicide as poor control may result.
- Do not feed treated hay, forage, or fodder or graze treated soybeans to livestock.
- Only one application of this product may be made prior to planting soybeans per growing season.
- Do not feed or graze treated cover crops to livestock.
- * Not currently registered in California.

GRASSES GROWN FOR SEED PRODUCTION

WEEDS	AMOUNT OF WEEDONE® LV6 EC PER ACRE	DIRECTIONS
Annual broadleaf weeds	2/3 to 1 pint	Apply to established stands in the spring before the seed head comes into the boot stage. Do not spray in boot stage of growth. In seedling grasses apply in the spring after the grass has tillered or has at least 5 leaves but before the boot stage. Use only the low rate on seedling grasses.
Biennial and Perennial broadleaf weeds	1-1/3 to 2 pints	DO NOT USE IN CALIFORNIA

RESTRICTIONS AND LIMITATIONS FOR USE ON GRASSES FOR SEED PRODUCTION

- Do not graze dairy animals or cut forage for hay within 7 days of application.

9/13

**WEEDS AND BRUSH IN PASTURES, RANGELAND AND FALLOWLAND,
ESTABLISHED GRASS PASTURES AND RANGELAND**

WEEDS IN CROP	AMOUNT OF WEEDONE® LV6 EC PER ACRE	DIRECTIONS
Established Grasses and Rangeland Annual broadleaf weeds Perennial and biennial broadleaf weeds	1-1/3 pints 2 to 2-2/3 pints	Apply when weeds are small and actively growing and prior to bud stage. Do not apply to newly seeded areas until grass is well established. Do not apply when grass is in the boot to milk stage if grass seed production is desired. Bentgrass and legumes may be injured by this treatment.
Sand sage, sagebrush, shinnery oak and other 2,4-D susceptible woody species	2/3 to 2-3/4 pints	Apply aerially to brush in 2 or more gallons of water per acre. One gallon of fuel oil may be included in the mixture. Consult state or local brush control specialists for most effective rate, volume and timing of spray applications.

FALLOW LAND AND CROP STUBBLE

WEEDS	AMOUNT OF WEEDONE® LV6 EC PER ACRE	DIRECTIONS
Annual broadleaf weeds	2/3 to 1-1/3 pints	Use the lower rate when weeds are small (2 to 3 inches tall) and growing actively. Use the higher rate on older and drought-stressed plants.
Biennial broadleaf weeds	2/3 to 1-2/3 pints	Spray while musk thistles or other biennial species are in the seedling to rosette stage, and before flower stalks are initiated. The lower rate can be used in spring during rosette stage. In fall or after flower stalks have developed, use the highest rate.
Perennial broadleaf weeds	1-1/3 to 4 pints	Spray weeds in bud to bloom stage, or in good vegetative growth. Do not disturb treated area for at least 2 weeks after treatment, or until weed tops are dead.
Wild garlic or onions	2-2/3 to 4 pints	Apply to new regrowth of wild onion or garlic which occurs in the fall following harvest of small grains, soybeans, corn or grain sorghum.

RESTRICTIONS AND LIMITATIONS FOR USE ON PASTURES, RANGELAND, AND FALLOW LAND

- For ground application, a minimum of 10 to 15 gallons of water per acre is recommended for proper spray coverage.
- For aerial application, use a minimum of 2 gallons of water per acre.
- Do not graze animals on treated areas within 7 days after treatment.
- Do not cut grass for hay within 30 days after application.
- Do not permit dairy animals or meat animals being finished for slaughter to forage treated fields within 3 days of slaughter.
- Do not plant treated fallow land until 3 months after treatment or until chemical has disappeared from soil.

FENCEROWS, ROADSIDES, VACANT LOTS, AIRFIELDS, RAILROAD, HIGHWAY AND UTILITY RIGHTS-OF-WAY, AND OTHER NON-CROP AREAS.

WEEDS	AMOUNT OF WEEDONE® LV6 EC PER ACRE	DIRECTIONS
Annual broadleaf weeds	1-1/3 to 2-2/3 pints	Apply when most weeds are still young and growing vigorously.
Perennial and biennial broadleaf weeds	2 to 4 pints	Apply when weeds are actively growing and near the bud stage, but before flowering. For best results on tansy ragwort and musk thistle, treat in rosette stage, before bolting. A second application is usually needed for best results on thistle, nettle and bindweed. Treat wild onion or garlic in early spring and in fall when they are young and growing actively.

(continued)

10/13

FENCEROWS, ROADSIDES, VACANT LOTS, AIRFIELDS, RAILROAD, HIGHWAY AND UTILITY RIGHTS-OF-WAY, AND OTHER NON-CROP AREAS. (continued)

WEEDS	AMOUNT OF WEEDONE® LV6 EC PER ACRE	DIRECTIONS
Cattails and tules	1 to 1-2/3 gallons	Mix 1-1/3 quarts of this product in 2 quarts kerosene or diesel oil then add this mixture to 100 gallons of water. Apply 300 to 500 gallons of spray per acre, depending on the stand. The addition of a wetting agent such as SURFEL® spray adjuvant is suggested.
Woody plants	2/3 to 1-1/3 gallons	Apply in 30 to 100 gallons of spray per acre, preferably through low volume equipment such as DIRECTA-SPRA®, WOBBLER®, MINI-WOBBLER® or SPIROMETER®. Treat when 2,4-D susceptible species are in full leaf and growing actively. Repeated applications, oil or wetting agent and/or high volume spraying (300 to 500 gallons of spray per acre) may be needed for acceptable control of some species or if brush is dense.

RESTRICTIONS AND LIMITATIONS FOR USE IN NON-CROP AREAS

- For aerial application to solid stands of susceptible brush, use 1-1/3 to 2-2/3 quarts in 3 to 12 gallons volume per acre. Two to 4 quarts of fuel oil may be included in this mixture.
- Applications will be less effective when conditions such as deficient soil moisture reduce brush growth.

**TANK MIXTURES FOR NON-CROP AREAS
WEEDONE® LV6 EC and GARLON® 4 or GARLON® 3A Tank Mixtures for Non-Crop Areas**

WEEDS	AMOUNT OF WEEDONE® LV6 EC PER ACRE	DIRECTIONS
Broadleaf weed control	1-1/3 to 2-2/3 pints WEEDONE® LV6 EC plus 2 to 6 pints GARLON-4 or 3 to 8 pints GARLON-3A	For wider spectrum control of broadleaf weeds and woody plants: Apply as a broadcast spray in enough water to deliver 20 to 100 gallons total spray per acre. Apply when broadleaf weeds are actively growing.
Woody plant control Broadcast foliar spray	2/3 to 1-1/3 gallons WEEDONE® LV6 EC plus 1-1/2 to 3 quarts GARLON-4 or 2 to 4 quarts GARLON-3A	Apply as a broadcast spray in enough water to deliver 20 to 100 gallons total spray per acre. Apply when woody plants are actively growing.
Woody plant control High volume leaf-stem treatment with ground equipment	2/3 to 5-1/3 quarts WEEDONE® LV6 EC plus 1-1/2 to 12 pints GARLON-4 or 2 to 16 pints GARLON-3A	Mix 2/3 to 1-1/3 quarts of Nufarm WEEDONE® LV6 EC Broadleaf Herbicide plus 1-1/2 to 3 pints GARLON-4 or 2 to 4 pints GARLON-3A in enough water to make 100 gallons of spray. Apply at a volume of 100 to 400 gallons of total spray per acre depending on size and density of woody plants. Thoroughly wet all leaves, stems and root collars of plants to be controlled.
Woody plant control Aerial application (helicopter only)	2/3 to 1-1/3 gallons WEEDONE® LV6 EC plus 3 to 4 quarts GARLON-4 or 4 to 6 quarts GARLON-3A	Apply in a total spray volume of 10 to 30 gallons per acre using drift control equipment such as the MICRO-FOIL® boom or an effective drift control agent such as LO-DRIFT® spray additive. Use the higher rates and volumes when plants are dense or under drought conditions.

11/13

WEEDONE® LV6 EC and BANVEL® Herbicide Tank Mixtures For Non-Crop Areas

WEEDS	AMOUNT OF WEEDONE® LV6 EC PER ACRE	DIRECTIONS
Annual broadleaf weeds	1-1/3 to 2-2/3 pints WEEDONE® LV6 EC plus 1/2 to 1-1/2 pints BANVEL	For wider spectrum control of broadleaf weeds and woody plants: Apply as a broadcast spray in enough water to deliver 20 to 100 gallons total spray per acre. Apply when broadleaf weeds are actively growing. Use the higher rates when treating dense or tall vegetative growth.
Perennial and biennial broadleaf weeds	2 to 4 pints WEEDONE® LV6 EC plus 1/2 to 6 pints BANVEL	Apply as a broadcast spray in enough water to deliver 20 to 100 gallons total spray per acre. Apply when broadleaf weeds are actively growing but prior to flowering. Use the lower rates for biennials less than 3 inches rosette diameter. Use the higher rates for perennial weeds or for biennial weeds past the 3-inch rosette stage.
Woody plant control Broadcast, high volume, stem foliage or aerial application	2/3 to 1-1/3 gallons WEEDONE® LV6 EC plus 2 to 8 quarts BANVEL	Apply as broadcast spray in enough water to deliver 20 to 100 gallons total spray per acre or apply as a high volume stem foliage spray in enough volume to thoroughly wet leaves, stems and root collars (100 to 400 gallons per acre) or apply aerially in enough water to deliver total spray volume of 10 to 30 gallons per acre using drift control equipment such as the MICRO-FOIL boom or an effective drift control agent such as LO-DRIFT spray additive. Use the higher rates and volumes when plants are dense or under drought conditions.

SMALL AREA APPLICATIONS FOR NON-CROP USE

For control of broadleaf weeds in small non-crop areas with hand held or back-pack sprayers mix 2-2/3 fluid ounces of this herbicide per gallon of water. Thoroughly wet all weed foliage. Maintain agitation of mixture to prevent separation.

FOREST MANAGEMENT*

CONIFER RELEASE

To control alder, apply 2/3 to 2 quarts Nufarm WEEDONE® LV6 EC Broadleaf Herbicide in 9-1/2 to 15 gallons water per acre as a foliage spray. Usually 1-1/3 quarts of this herbicide per acre is sufficient to provide good alder control with minimum conifer injury. Treat when 3/4 of the brush foliage has attained full size leaves and before new conifer growth reaches 2" in length. This is usually between early May and mid-June. Adjust treatment date depending on stage of growth of conifers and brush species. This may cause leader deformation on exposed firs, but they should overcome this during the second year after spraying.

To control tanoak, madrone, ceanothus, canyon live oak, and manzanita, and to release Douglas fir, hemlock, Sitka spruce or grand fir, apply 2 quarts Nufarm WEEDONE® LV6 EC Broadleaf Herbicide plus 3 quarts fuel oil in 8-1/2 to 15 gallons water per acre before new growth on Douglas fir is 2" long. To control manzanita and ceanothus in ponderosa pine, apply 3 quarts Nufarm WEEDONE® LV6 EC Broadleaf Herbicide before pine growth begins in spring.

To control hazel brush in the Lake states, apply 1-2/5 quarts, Nufarm WEEDONE® LV6 EC Broadleaf Herbicide in 6 to 25 gallons of water per acre when new shoot growth of hazel is complete (usually mid-July).

In northern areas, if possible conifer injury can be tolerated, 1 to 2 quarts of Nufarm WEEDONE® LV6 EC Broadleaf Herbicide applied aerially in 8 to 25 gallons of water per acre after conifers such as jack pine, red pine, black spruce, and white spruce have hardened off (usually mid-July) will provide control of competing hardwoods including alder, aspen, birch, hazel and willow.

SITE PREPARATION

Budbreak Spray: To control alder and other susceptible species before planting forest seedlings, apply 1-2/5 to 2-3/5 quarts Nufarm WEEDONE® LV6 EC Broadleaf Herbicide in 9 to 15 gallons of water per acre plus 2 quarts of fuel oil per acre after alder buds break, but before foliage is 1/4 full size.

Foliage Spray: To control alder before planting forest tree seedlings, apply 1-2/5 quarts Nufarm WEEDONE® LV 6 EC Broadleaf Herbicide plus 2 quarts fuel oil in 9 to 15 gallons water per acre after most alder leaves are full size.

Consult your regional or extension forester or state herbicide specialist for recommendations to fit local conditions.

12/13

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants that this product conforms to the chemical description on the label; that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESSED OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS, OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.


NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

Manufactured by:
Nufarm, Inc.
Burr Ridge, IL

BANVEL® , POAST® and POAST PLUS® are Registered Trademarks of BASF Corp.
GARLON® is a trademark of Dow AgroSciences
GRAMOXONE EXTRA® is a Registered Trademark of a Syngenta Crop Protection.
MICRO-FOIL® , DIRECTA-SPRAY® , WOBBLER® , MINI WOBBLER® , SPIROMETER® , SURFEL® and LO-DRIFT® are Registered Trademarks of Aventis.
PROWL® , PURSUIT PLUS® , SCEPTER® and SQUADRON® are Registered Trademarks of BASF.
ROUNDUP® , ROUNDUP D-PAK® and HONCHO® are Registered Trademarks of Monsanto Co.
WEEDONE® is a Registered Trademark of Nufarm, Inc.

071368-00011. 200300421.wlv6ec





13/13

Nufarm Americas Inc.
1333 Burr Ridge Parkway, Suite 125A
Burr Ridge, IL 60527-0866
Phone: 708.754.3330 Fax: 708.754.0314
www.ag.us.nufarm.com

March 16, 2004

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504C)
U. S. Environmental Protection Agency
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202-4501

Attention: Joanne Miller, PM 23
Herbicide Branch

Subject: Label Notification Addition of Similar Application Methods
EPA Reg. No. 71368-11 Weedone LV6 EC.

Dear Joanne:

Enclosed please find EPA Form 8570-1 and one copy of our label, which includes changes recently requested by California. The California changes are highlighted on pages 5,6 and 7. Please note that no rates or application sites have changed.

If you should have any questions, please contact me at the address and telephone number above or by e-mail to <ted.head@us.nufarm.com>. Thank you for your cooperation.

Sincerely,

A handwritten signature in black ink, appearing to read 'Ted Head', written over the word 'Sincerely,'.

Theodore D. Head
Product Registration Manager

