

# SUPPLEMENTAL LABELING

READ AND FOLLOW THE ENTIRE LABEL BOOKLET FOR NUFARM Weedone LV6 EC BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for this product and this supplement.



## Weedone LV 6 EC Broadleaf Herbicide

EPA Reg. No. 71368-11

### Maximum Use Rates For Forestry and Non-Crop Sites

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

All applicable directions and precautions in the label booklet for this product must be followed.

Also refer to additional "GENERAL INFORMATION" and "WEEDS CONTROLLED" in the label booklet. INJURY TO CROPS FROM THIS HERBICIDE MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY, DO NOT USE THIS PRODUCT. Product should not be used in or near greenhouses.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR, Part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

#### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. For Turf use, do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

#### NON-CROPLAND

DITCHES, ROADSIDES, VACANT LOTS, AIRFIELDS, RAILROAD, HIGHWAY AND UTILITY RIGHTS-OF-WAY, AND OTHER NON-CROP AREAS.

WEEDS	AMOUNT OF WEEDAR 64 PER ACRE	DIRECTIONS
Annual broadleaf weeds	1-1/3 to 2-2/3 pints	Apply when most weeds are still young and growing vigorously.
Perennial and biennial broadleaf weeds	2 to 3 pints	Apply when weeds are actively growing and near the bud stage, but before flowering. For best results on tansy ragwort and musk thistle, treat in rosette stage, before bolting. A second application is usually needed for best results on thistle, nettle and bindweed. Treat wild onion or garlic in early spring and in fall when they are young and growing actively.

ACCEPTED

OCT 20 2005

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

71368-11

**NON-CROPLAND - Continued**  
**FENCEROWS, ROADSIDES, VACANT LOTS, AIRFIELDS,**  
**RAILROAD, HIGHWAY AND UTILITY RIGHTS-OF-WAY, AND**  
**OTHER NON-CROP AREAS.**

**Tank Mixtures for Non-Crop Areas**

**FOR NON-CROP AREAS WEEDONE® LV6 EC AND**  
**GARLON® 4 OR GARLON® 3A**

WEEDS	AMOUNT OF WEEDAR 64 PER ACRE	DIRECTIONS
Cattails and tules	1-1/3 quarts	Mix 1-1/3 quarts of this product per acre in 2 quarts kerosene or diesel oil then add this mixture to 100 gallons of water. This may require 300 to 500 gallons of water per acre, depending on the stand. The addition of a wetting agent such as SURFEL® spray adjuvant is suggested.
Woody plants	2/3 to 3/4 gallons	Apply preferably through low volume equipment such as DIRECTA-SPRA®, WOBBLER®, MINI-WOBBLER® or SPIROMETER®. Treat when 2,4-D susceptible species are in full leaf and growing actively. Repeated applications, oil or wetting agent and/or high volume spraying (300 to 500 gallons of water per acre) may be needed for acceptable control of some species or if brush is dense.

**Woody plant control**

Broadcast foliar spray ¾ gallons per acre plus 1-1/2 to 3 quarts GARLON-4 Or 2 to 4 quarts GARLON-3A. Apply as a broadcast spray in enough water to wet all parts of the brush foliage, stem and bark. This may require 20 to 100 gallons of water per acre. Apply when woody plants are actively growing.

**Woody plant control** High volume leaf-stem treatment with plus ground equipment 3 quarts WEEDONE® LV6 EC per acre plus 1-1/2 to 12 pints GARLON-4 or 2 to 16 pints GARLON-3A

Mix 2/3 to 1-1/3 quarts of Nufarm WEEDONE® LV6 EC Broadleaf Herbicide per acre plus 1-1/2 to 3 pints GARLON-4 or 2 to 4 pints GARLON-3A in enough water to wet all parts of the brush foliage, stem and bark. This may require 100 – 400 gallons of water per acre depending on size and density of woody plants. Thoroughly wet all leaves, stems and root collars of plants to be controlled.

**Woody plant control** Aerial application (helicopter only) ¾ gallons WEEDONE® LV6 EC per acre plus 3 to 4 quarts GARLON-4 or 4 to 6 quarts GARLON-3A. Apply in enough water to wet all parts of the brush foliage, stem and bark. This may require 10 to 30 gallons of water per acre using drift control equipment such as the MICRO-FOIL® boom or an effective drift control agent such as LO-DRIFT® spray additive. Use the higher rates and volumes when plants are dense or under drought conditions.

**WEEDONE® LV6 EC AND BANVEL® Herbicide Tank**  
**Mixtures For Non-Crop Areas**

Perennial and biennial broadleaf weeds 2 to 3 pints WEEDONE® LV6 EC per acre Plus 1/2 to 6 pints BANVEL. Apply as a broadcast spray in enough water to wet all parts of the brush foliage, stem and bark. This may require 20 to 100 gallons total spray per acre. Apply when broadleaf weeds are actively growing but prior to flowering. Use the lower rates for biennials less than 3 inches rosette diameter. Use the higher rates for perennial weeds or for biennial weeds past the 3-inch rosette stage.

**WEEDONE® LV6 EC AND BANVEL® Herbicide Tank  
Mixtures For Non-Crop Areas**

Woody plant control Broadcast, high volume, stem foliage or aerial application  $\frac{3}{4}$  gallons WEEDONE® LV6 EC per acre plus 2 to 8 quarts BANVEL. Apply as broadcast spray in enough water to thoroughly wet leaves, stems and root collars, or apply as a high volume stem foliage spray in enough volume to thoroughly wet leaves, stems and root collars (100 to 400 gallons of water per acre) or apply aerially in enough water (10 to 30 gallons per acre) using drift control equipment such as the MICRO-FOIL boom or an effective drift control agent such as LO-DRIFT spray additive. Use the higher rates and volumes when plants are dense or under drought conditions.

This product will either kill, control or suppress the weeds listed in the label booklet for this product. Some of these species may require repeat spot applications even under ideal conditions.

\* Maximum Seasonal Application Rate Rate to non-crop and forestry sites is 4 pounds 2,4-D Acid equivalent per acre per application site.

**NOTICE**

Before buying or using this product, read "WARRANTY LIMITATIONS AND DISCLAIMER" in the label booklet. If terms are not acceptable, return unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by the Buyer or any other User constitutes acceptance of the terms under the "WARRANTY LIMITATIONS AND DISCLAIMER".

**Manufactured By  
Nufarm Inc.  
Burr Ridge, IL 60521**

09/05