


PLEASE NOTE

**This image contains more than one label
approved for this product on this date.**

SUPPLEMENTAL LABELING

READ AND FOLLOW THE ENTIRE LABEL BOOKLET FOR NUFARM CREDIT HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Nufarm Credit Herbicide and this supplement.



Credit®

Herbicide

EPA Reg. No. 71368-20

PREHARVEST AND SPOT-TREATMENTS OF WEEDS IN DRY BEANS

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

All applicable directions and precautions in the Nufarm Credit Herbicide label booklet must be followed.

See the "GENERAL INFORMATION" and "MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS" sections of the Nufarm Credit Herbicide label booklet for essential product performance information.

AVOID CONTACT WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

BROADCAST SPRAY: This product may be applied as an over the top broadcast spray to control labeled weeds prior to the harvest of dry beans. Apply up to 32 fluid ounces in 3 to 20 gallons of water per acre at the hard dough stage of the legume seed (30 percent grain moisture or less). Either ground broadcast or aerial applications may be made.

PRECAUTIONS RESTRICTIONS:

- Apply at least 7 days before harvest.
- Only one application per year may be made; do not combine a preharvest spray with a spot treatment on the same crop area.
- Employ at least a 30 day plant-back interval between treatment and replanting for any crop not listed in the Credit Systemic herbicide label.
- Preharvest application is not recommended for dry beans grown for seed, as a reduction in germination or vigor may occur.
- Do not feed treated vines and hay from these crops to livestock. Do not apply this product through any type of irrigation system.
- Do not treat field cowpeas, since these are considered to be grown as livestock feed.

SPOT TREATMENTS

This product may be applied as spot treatment to control troublesome weeds such as Canada thistle, quackgrass, mayweed (dog fennel), and milkweed in dry beans. Apply up to 26 fluid ounces in 10 to 20 gallons of water through ground spray equipment or use a 2 percent solution in a handheld sprayer. For best results, applications should be made at or beyond the bud stage of

growth. The crop receiving spray in treated areas will be killed.

PRECATIONS, RESTRICTIONS:

- Apply at least 14 days before harvest.
- Only one application per year may be made; do not combine a preharvest spray with a spot treatment on the same crop area.
- Employ at least a 30 day plant-back interval between treatment and replanting for any crop not listed in the Credit Systemic herbicide label.
- Do not feed treated vines and hay from these crops to livestock. Do not apply this product through any type of irrigation system.
- Do not treat field (feed) peas, since these are considered to be grown as livestock feed.

NOTICE

Read the "LIMIT of WARRANTY AND LIABILITY" in the label booklet for Nufarm Credit Herbicide before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Manufactured By

Nufarm, Inc.

Burr ridge, IL 60527

Credit® is a registered trademark of Nufarm, Inc.


NEXT

LABEL

SUPPLEMENTAL LABELING

READ AND FOLLOW THE ENTIRE LABEL BOOKLET FOR NUFARM CREDIT HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Nufarm Credit Herbicide and this supplement.



Credit®

Herbicide

EPA Reg. No. 71368-20

PREHARVEST AND SPOT-TREATMENTS OF WEEDS IN DRY PEAS, LENTILS, AND CHICKPEAS

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

All applicable directions and precautions in the Nufarm Credit Herbicide label booklet must be followed.

See the "GENERAL INFORMATION" and "MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS" sections of the Nufarm Credit Herbicide label booklet for essential product performance information.

AVOID CONTACT WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

BROADCAST SPRAY: This product may be applied as an over the top broadcast spray to control labeled weeds prior to the harvest of dry peas, lentils, and chickpeas. Apply up to 24 fluid ounces in 3 to 20 gallons of water per acre at the hard dough stage of the legume seed (30 percent grain moisture or less). Either ground broadcast or aerial applications may be made.

PRECAUTIONS RESTRICTIONS:

- Apply at least 14 days before harvest.
- Only one application per year may be made; do not combine a preharvest spray with a spot treatment on the same crop area.
- Employ at least a 30 day plant-back interval between treatment and replanting for any crop not listed in the Credit Systemic herbicide label.
- Preharvest application is not recommended for peas, lentils, or chickpeas grown for seed, as a reduction in germination or vigor may occur.
- Do not feed treated vines and hay from these crops to livestock. Do not apply this product through any type of irrigation system.
- Do not treat field (feed) peas, since these are considered to be grown as livestock feed.

SPOT TREATMENTS

This product may be applied as spot treatment to control troublesome weeds such as Canada thistle, quackgrass, mayweed (dog fennel), and milkweed in dry peas, lentils, and chickpeas. Apply up to 20 fluid ounces in 10 to 20 gallons of water through ground spray equipment or use a 2 percent solution in a handheld sprayer. For best results,

applications should be made at or beyond the bud stage of growth. The crop receiving spray in treated areas will be killed.

PRECAUTIONS, RESTRICTIONS:

- Apply at least 14 days before harvest.
- Only one application per year may be made; do not combine a preharvest spray with a spot treatment on the same crop area.
- Employ at least a 30 day plant-back interval between treatment and replanting for any crop not listed in the Credit Systemic herbicide label.
- Do not feed treated vines and hay from these crops to livestock. Do not apply this product through any type of irrigation system.
- Do not treat field (feed) peas, since these are considered to be grown as livestock feed.

NOTICE

Read the "LIMIT of WARRANTY AND LIABILITY" in the label booklet for Nufarm Credit Herbicide before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

**Manufactured By
Nufarm, Inc.**

Burr ridge, IL 60527

Credit® is a registered trademark of Nufarm, Inc.