



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Mr. George Meindl
Regulatory Affairs Manager
Nufarm America, Inc.
150 Harvester Drive suite 200
Burr Ridge, IL 60525

AUG 14 2008

RE: Notification of Alternate Brand Name: "Weedmaster® Herbicide"

EPA Registration Number: 71368-34
Date of Submission: July 15, 2008

Dear Mr. Meindl:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated July 15, 2008, for the product Nufarm Kambamaster® Herbicide. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Joyce Edwards of my staff at 703-308-5479.

Sincerely,

A handwritten signature in black ink, appearing to read "Linda Arrington".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs



United States
Environmental Protection Agency
 Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 71368-34	2. EPA Product Manager Joanne Miller	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Weedmaster Herbicide	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Nufarm Inc. 150 Harvester Drive Suite 200 Burr Ridge, IL 60527 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. NOTIFICATION Product Name AUG 14 2008	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Label notification (adding an alternate brandname) consistent with PRN98-10, see cover letter for detailed explanation. This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make false statements to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
				<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
* Certification must be submitted	If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	Other (Specify) _____	
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name George Meindl george.meindl@us.nufarm.com	Title Regulatory Affairs Manager	Telephone No. (Include Area Code) 630.455.2017
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		8. Date Application Received (Stamped)
2. Signature 	3. Title Regulatory Affairs Manager	
4. Typed Name George Meindl	5. Date 7/15/2008	

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER - PELIGRO**

Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear goggles. Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with skin. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other handlers must wear a long-sleeved shirt and long pants, shoes plus socks and protective eyewear.

Mixers and loaders who do not use a mechanical system (probe and pump) must wear: Coveralls, chemical-resistant apron. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not re-use them. Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

ENGINEERING CONTROLS STATEMENT:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agriculture pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

For containers of 5 gallons or more, do not open pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

For containers greater than 1 gallon but less than 5 gallons, when handlers use a mechanical system (probe and pump), enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS	
Users Should:	
<ul style="list-style-type: none"> • Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. • Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water. • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. 	

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.</p>

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

ENDANGERED SPECIES CONCERNS

The use of any pesticide in a manner that may kill or otherwise harm an endangered species or adversely modify their habitat is a violation of federal law.

DIRECTIONS FOR USE

It Is A Violation Of Federal Law To Use This Product In A Manner Inconsistent With Its Labeling.

Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Unless otherwise directed in supplemental labeling, all applicable directions, restrictions, and precautions are to be followed. Labeling must be in the user's possession during application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is: coveralls over short-sleeved shirt and short pants, waterproof gloves, chemical-resistant footwear plus socks, chemical-resistant headgear for overhead exposure and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

For ornamental turf uses (golf courses, cemeteries, parks and other turf grass areas), do not enter treatment areas until sprays have dried. Do not allow people (other than applicator) or pets on treatment area during application.

Steps to be taken in case material is released or spilled:

Dike and contain the spill with inert material (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal. Remove contaminated clothing, and wash affected skin areas with soap and water. Wash clothing before re-use. Keep the spill out of all sewers and open bodies of water.

I. GENERAL INFORMATION

This product is a selective postemergence herbicide for controlling a wide spectrum of annual, biennial, and perennial broadleaf weeds and brush in grass forages and selected row crops.

Mode of Action

This product contains two active ingredients: dicamba and 2,4-D. This herbicide is readily absorbed by plants through shoot and root uptake, translocates throughout the plant's system, and accumulates in areas of active growth. This product interferes with the plant's growth hormones (auxins) resulting in death of many broadleaf weeds.

Cleaning Spray Equipment

Clean application equipment thoroughly by using a strong detergent or commercial sprayer cleaner according to the manufacturer's directions and then triple rinsing the equipment before and after applying this product.

II. APPLICATION PROCEDURES

Apply this product at the rates and growth stages listed in Tables 1 and 2 as follows unless instructed differently by Section VI or VII. (Food/Feed Crop Specific Information or Non-Food/Feed Use Specific Information). Applications can be made to actively growing weeds as aerial, broadcast, band, or spot spray applications. This product may be applied using water or sprayable fluid fertilizer as a carrier. Sprayable fluid fertilizer may be used as the carrier in preplant or pre-emergence uses for all crops listed on this label. Postemergence uses with sprayable fluid fertilizer may be made on pasture, hayland, or wheat crops only.

The most effective application rate and timing varies based on the target weed species (refer to Table 1). In mixed populations of weeds the correct rate is determined by the weed species requiring the highest rate. Delaying application permits weeds to exceed the maximum size stated and will prevent adequate control.

IRRIGATION

In irrigated areas, it may be necessary to irrigate before treatment to ensure active weed growth.

SPRAY COVERAGE

Weeds must be thoroughly covered with spray. Dense leaf canopies shelter smaller weeds and can prevent adequate spray coverage.

SENSITIVE CROP PRECAUTIONS

This product may cause injury to desirable trees and plants, particularly beans, cotton, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers, tobacco, tomatoes and other broadleaf plants when contacting their roots, stems or foliage. These plants are most sensitive to this product during their development or growing stage.

Do not treat areas where either possible downward movement into the soil or surface washing may cause contact of this product with the roots of desirable plants such as trees and shrubs.

- Avoid making applications when spray particles may be carried by air currents to areas where sensitive crops and plants are growing. Do not spray near sensitive plants if wind is gusty or in excess of 5 mph and moving in the direction of nearby sensitive crops or if a temperature inversion exists. However, always make applications when there is some air movement to determine the direction and distance of possible spray drift. Leave an adequate buffer zone between area to be treated and sensitive plants. Coarse sprays are less likely to drift out of the target area than fine sprays. Agriculturally-approved drift-reducing additives may be used.
- Do not use aerial equipment or apply this product when sensitive crops and plants are growing in the vicinity of area to be treated.

SPRAY DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information.

Importance of Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversion section of this label).

Controlling Droplet Size

Volume: Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure: Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of nozzles: Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation: Orienting nozzles so that the spray is released backwards, parallel to the airstream will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.

Nozzle Type: Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.

Boom Length: For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

Application: Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment

When applications are made with a cross-wind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

Wind

Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion, because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a connected cloud

(under low wind conditions) indicates an inversion, while smoke that moves upwards and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

AERIAL APPLICATION METHODS AND EQUIPMENT

Water Volume: Use 3 to 10 gallons of water per acre. Use the higher spray volume when treating dense or tall vegetation.

Application Equipment: Select nozzles designed to produce minimal amounts of fine spray particles. Make applications at the lowest safe height to reduce the exposure of spray droplets to evaporation and wind. The applicator must follow the most restrictive use cautions to avoid drift hazards, including those found in this labeling as well as applicable state and local regulations and ordinances.

Do not use aerial equipment if spray particles can be carried by the wind into areas where sensitive crops or plants are growing or when temperature inversions exist.

TABLE 1. APPLICATION RATE AND TIMING - ANNUAL WEEDS

Weeds Controlled (including ALS - and triazine-resistant)	Weedmaster Rate Per Acre (according to weed growth stage)					
	0.5 pint	1 pint	1.5 pints	2 pints	3 pints	4 pints
Beebalm, Spotted	-	-	-	pre-bloom	postbloom	-
Broomweed	1 to 3"	3" branching	-	branching	-	after branching
Buckwheat, Wild	-	1 to 6"	-	-	-	-
Buffalobur	-	-	-	1 to 6"	-	flowering
Burdock	-	pre-flower	-	-	-	-
Buttercup	-	pre-flower	-	early bloom	late bloom	-
Chickweed, Common	-	seedling	1 to 3"	-	-	-
Cockle, Cow	-	< 3"	-	-	-	-
Cocklebur, Common	-	1 to 6"	6 to 12"	12 to 18"	-	-
Coreopsis, Plains	-	1 to 6"	-	-	-	-
Croton, Woolly	1 to 4"	4 to 12"	12 to 30"	-	-	-
Devils-claw	-	-	-	< 8"	-	-
Dogfennel	-	-	-	10 to 15"	-	-
Evening Primrose	-	< 2"	-	2 to 6"	-	-
Flax	-	< 2"	-	-	-	-
Fleabane, Annual	-	1 to 4"	4 to 8"	8"	-	-
Flixweed	-	< 3"	-	-	-	-
Henbit	-	-	preflower	-	flower	-
Knotweed Spp.	-	< 3" runners	-	> 3" runners	-	actively growing
Kochia	-	1 to 6"	6 to 10"	10 to 20"	-	actively growing
Lambsquarters, Common	-	1 to 6"	6 to 10"	10 to 20"	-	actively growing
Mallow, Common	-	< 3"	-	-	-	-
Morningglory, Ivyleaf , Tail	-	pre-flower pre-flower	-	post-flower early bolt	-	-
Mustards, Annual , Tansy	-	rosette < 3"	-	-	-	-
Pennycress, Field	-	-	-	rosette	-	-
Pepperweed, Virginia	-	-	1 to 3"	3 to 6"	after branching	-
Pigweed, Prostrate , Redroot , Smooth , Tumble	-	< 3" < 3" < 3" < 3"	3 to 10" - - -	- - - mature	- - - -	- - - -
Poojoe	-	prior to flower	-	-	-	actively growing
Purslane, Common	-	< 3"	3 to 8"	-	-	-
Ragweed, Common Western, Lanceleaf	1 to 3"	3 to 6"	6 to 10"	> 10" actively growing	-	-
Sedge ¹	-	-	-	-	-	-
Shepherdspurse	-	rosette	-	-	-	-
Smartweed, Pennsylvania	-	< 4"	-	-	4 to 12"	-
Sneezeweed, Bitter	-	1 to 4"	prior to flower	flower	-	-
Sowthistle	-	rosette	-	bolting	-	-
Sunflower	-	1 to 3"	3 to 6"	6 to 24"	-	-
Thistle, Russian	-	-	-	rosette	-	-
Velvetleaf	-	< 6"	6 to 20"	> 20"	-	-

¹ For use in non-food/feed crop only. Adding crop oil concentrate has shown to improve performance on actively-growing annual sedge.

TABLE 2. APPLICATION RATE AND TIMING - BIENNIAL AND PERENNIAL WEEDS

Weeds Controlled	Weedmaster Rate Per Acre (according to weed growth stage)					
	0.5 pint	1 pint	1.5 pints	2 pints	3 pints	4-6 pints
Bindweed, Field	-	-	-	-	-	actively growing
Bittercress	-	2 to 3"	-	-	-	-
Buckeye species ¹	-	-	-	-	full leaf	-
Bullnettle ²	-	-	-	flower	-	-
Chicory	-	-	-	-	early bolting	-
Clover, Bur	-	-	pre-flower	-	-	-
Dandelion, Common	-	rosette	-	bolting	-	-
Dewberry, Southern ¹	-	-	-	-	-	spring or fall
Dock, Curly	-	-	prior to bolting	-	after bolting	-
Elderberry ²	-	-	-	-	-	actively growing
Goldenrod, Missouri	-	-	-	3 to 15"	flower	-
Goldenweed, Common	-	-	-	-	-	actively growing
Groundsel, Texas	-	rosette	post bolting	-	-	-
Honeysuckle, Hairy	-	-	-	-	spring or fall	-
Horsenettle, Carolina ¹	-	-	-	-	-	flower or berry
Ivy, Poison	-	-	-	after bloom	-	-
Knapweed, Black ²	-	-	-	-	-	actively growing
Russian ²	-	-	-	-	-	actively growing
Spotted	-	-	-	-	-	actively growing
Marsheld	-	-	-	< 12"	12"/prebloom	-
Mesquite	-	-	-	-	-	45 to 90 days after bud-break
Milkweed	-	-	-	-	-	-
Antelopehorn ¹	-	-	-	pre-flower	-	flower
Nightshade, Silverleaf ¹	-	-	-	full flower	-	-
Black ¹	-	-	-	full flower	-	actively growing
Persimmon, Eastern ²	-	-	-	-	-	actively growing
Prickly Lettuce	-	-	-	rosette	-	actively growing
Rabbitbrush ²	-	-	-	-	-	-
Ragwort, Tansy	-	-	-	rosette	-	actively growing
Redvine ²	-	-	-	-	-	actively growing
Sagebrush, Fringed ²	-	-	-	-	-	actively growing
Smartweed	-	-	-	-	-	-
Sorrel, Red	-	-	rosette	bolting	flower	actively growing
Sowthistle ²	-	-	-	-	-	actively growing
Spurge, Leafy ²	-	-	-	-	-	full leaf
Tallow Tree, Chinese ²	-	-	-	-	-	-
Thistle, Bull	-	-	rosette	bolting	-	actively growing
Canada ²	-	-	-	-	-	-
Musk	-	-	-	rosette/bolting	-	-
Plumeless	-	-	rosette	bolting	-	-
Vetch, Hairy	-	1 to 4"	4 to 8"	8" full flower	-	-
Yankeeeweed	-	-	-	10 to 18"	-	rosette
Yellow Starthistle ¹	-	-	-	-	-	-

¹May require repeat applications.

²Recommended rate will provide top growth suppression only.

³For improved root kill or weedy species such as mesquite and eastern persimmon, spray 4 pints of this product per acre each year for 3 consecutive years. For increased control of weeds such as blackberry and dewberry, this product may be tank mixed with Ally® herbicide (0.1 to 0.2 ounces per acre), if labeled for the use site.

⁴Under dense populations, a second application may be needed the following growing season.

TABLE 4. ADDITIVE RATE PER ACRE

Additive	Rate Per Acre
Nonionic Surfactant	2 to 4 pints per 100 gallons
Sprayable liquid fertilizers (28-0-0, 32-0-0) Crop Oil Concentrate	2 to 4 quarts 1 quart*

*see manufacturer's label for specific rate recommendations.

IV. GENERAL TANK MIXING INFORMATION

Tank Mix Partners/Components

The following products may be tank mixed with this product according to the specific tank mixing instructions in this label and respective product labels.

- Aim™ (carfentrazone-ethyl)
- Ally® (metsulfuron-methyl)
- Amber® (triasulfuron)
- Asulox® (asulam)
- Atrazine
- Banvel® (dicamba)
- Basagran® (bentazon)
- Bronate® (bromoxynil + MCPA)
- Buctril® (bromoxynil)
- Canvas® (thifensulfuron + tribenuron + metsulfuron)
- Clarity® (dicamba)
- Curtail® (clorpyralid + 2,4-D)
- Cyclone® (paraquat)
- Dakota® (fenoxaprop-p-ethyl + MCPA)
- Distinct® (diflufenzopyr)
- Evik® (ametryn)
- Express® (thifensulfuron + tribenuron-methyl)
- Fallowmaster® (glyphosate + dicamba)
- Finesse® (chlorsulfuron + metsulfuron-methyl)
- Glean® (chlorsulfuron)
- Gramoxone® Extra (paraquat)
- Harmony® Extra (thifensulfuron + tribenuron-methyl)
- Karmex® (diuron)
- Kerb® (pronamide)
- Laddok® S-12 (bentazon + atrazine)
- Landmaster® (glyphosate + 2,4-D)
- Lexone® (metribuzin)
- MCPA
- Paramount® (quinclorac)
- Peak® (prosulfuron)
- Permit® (halosulfuron-methyl)
- Rave™ (dicamba + triasulfuron)
- Roundup Ultra® (glyphosate)
- Sencor® (metribuzin)
- Sinbar® (terbacil)
- Stinger® (clopyralid)
- Tiller® (fenoxaprop-p-ethyl + 2,4-D + MCPA)
- Tordon® (picloram)
- Touchdown® (sulfosate)
- 2,4-D

See section VI. Crop-Specific Information for more details. Read and follow the applicable Restrictions and Limitations and Directions For Use on all product involved in tank mixing. The most restrictive labeling applies to tank mixes. Physical incompatibility, reduced weed control, or crop injury may result from mixing this product with other pesticides (fungicides, herbicides, insecticides, or miticides), additives, or fertilizers. NUFARM does not recommend using tank mixes other than those listed on NUFARM labeling. Local agricultural authorities may be a source of information when using other than NUFARM recommended tank mixes.

COMPATIBILITY TEST FOR MIX COMPONENTS

Before mixing components, always perform a compatibility jar test.

For 20 gallons per acre spray volume, use 3.3 cups (800 ml) of water. For other spray volumes, adjust accordingly. Only use water from the intended source at the source temperature.

Add components in the sequence indicated in the Mixing Order using 2 teaspoons for each pound or 1 teaspoon for each pint of recommended label rate per acre.

Always cap the jar and invert 10 cycles between component additions.

When the components have all been added to the jar, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent. If the solution is then compatible, use the compatibility agent as directed on its label. If the solution is still incompatible, do not mix the ingredients in the same tank.

MIXING ORDER

If an inductor is used, rinse it thoroughly after each component has been added. Maintain constant agitation during application.

- 1) **Water***. Begin by agitating a thoroughly clean sprayer tank half full of clean water.
- 2) **Agitation**. Maintain constant agitation throughout mixing and application.
- 3) **Products in PVA bags**. Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.

- 4) **Water-dispersible products** (such as dry flowables, wettable powders, suspension concentrates, or suspo-emulsions).
- 5) **Water-soluble products.** (such as this product)
- 6) **Emulsifiable concentrates** (such as oil concentrate when applicable).
- 7) **Water-soluble additives** (such as liquid fertilizers (28-0-0, 32-0-0) when applicable).
- 8) **Remaining quantity of water.**

* If sprayable fluid fertilizer is used as the carrier, This product must be diluted with a minimum of 5 parts water to 1 part this product. Then add 0.25 to .05% volume/volume of a nonionic surfactant to the dilution before adding it to the sprayable fluid fertilizer to reduce the concern for compatibility problems with this mix. Always perform the Compatibility Test before mixing into the spray tank. Also, when using a sprayable fluid fertilizer as the carrier, any product contained in PVA bags must first be completely dissolved in water before the contents can be added to the fertilizer mix.

V. RESTRICTIONS AND LIMITATIONS

- Maximum seasonal use rate: Refer to Crop-Specific Restrictions and Limitations table below.
- Preharvest Interval (PHI): Refer to section VI. Food/Feed Crop-Specific Information
- Restricted Entry Interval (REI): 48 hours
- Crop rotational Restrictions:

The interval between application and planting rotational crop is given below. Always exclude counting days when the ground is frozen. Planting at intervals less than specified below may result in crop injury. Moisture is essential for the degradation of this herbicide in soil. If dry weather prevails, use cultivation to allow herbicide contact with moist soil.

- **Planting/replanting restrictions for Weedmaster herbicide applications of 6 pints per acre or less:** No rotational cropping restrictions apply at 120 days or more following application. Additionally, for annual crop uses in this label including sorghum, follow the preplant use directions in section VI. Food/Feed Crop-Specific Information. For barley, oat, wheat, and other grass seedings, the interval between application and planting is 10 days per pint per acre.


- **Planting/replanting restrictions for applications of more than 6 pints and up to 8 pints of this product per acre:** Corn, sorghum, cotton (east of the Rocky Mountains) and all other crops grown in areas with 30" or more of annual rainfall may be planted 120 days or more after application. Barley, oat, wheat, and other grass seedings, may be planted if the interval from application to planting is 10 days per pint per acre east of the Mississippi River and 15 days per pint per acre west of the Mississippi River. For all other crops in areas with less than 30" of annual rainfall, the interval between application and planting is 180 days or more.

- **Rainfast period:** Rainfall or irrigation occurring within 4 hours after postemergence applications may reduce the effectiveness of this product
- **Stress:** Do not apply to crops under stress such as stress due to lack of moisture, hail damage, flooding, herbicide injury, mechanical injury, or widely fluctuating temperatures, as unsatisfactory control may result.
- Do not apply to crops that show injury (leaf phytotoxicity or plant stunting) produced by any other prior herbicide applications, because this injury may be enhanced or prolonged.
- Do not apply through any type of irrigation equipment. Do not contaminate irrigation ditches or water used for domestic purposes.
- This product cannot be used to formulate or reformulate any other pesticide product.

Table 5. CROP-SPECIFIC RESTRICTIONS AND LIMITATIONS

Crop	Maximum Rate Per Acre Per Application	Maximum Rate Per Acre Per Season	Livestock Grazing or Feeding ¹	Aircraft Application
Between Crop Applications	6 pints	8 pints	Yes	Yes
Pasture, Hay, Silage	4 pints	8 pints	Yes	Yes
Sugarcane	6 pints	16 pints	Yes	Yes
Sorghum	1 pint	1 pint	Yes	Yes
Wheat	2 pints	3.33 pints	Yes	Yes

¹ refer to Section VI. Food/Feed Crop-Specific Information for grazing and feeding restrictions.



SUGARCANE

Applications of this product can be made any time after the weeds have emerged and are actively growing but prior to the close-in stage of sugarcane. When possible, direct the spray beneath the sugarcane canopy in order to minimize the likelihood of crop injury. The use of directed sprays will also aid in maximizing spray coverage of weed foliage. Application rates and timing are given below. Use the higher level of listed rate ranges when treating dense vegetative growth.

Rate:

- For control of listed annual broadleaf weeds, apply 2 pints of this product per treated acre.
- For suppression of listed perennial weeds, apply 1 to 6 pints of this product per treated acre. Retreatments may be made as needed, however, do not exceed 16 pints of this product per treated acre during a growing season.

SUGARCANE TANK MIXES

This product may be applied in tank mixes with one or more of the following herbicides:

- Asulox®
- Atrazine
- Evik®
- Lexone®
- Sencor®
- Sinbar®

WHEAT

(Fall and Spring - Seeded)

If small grains are grown for pasture or hay only, refer to Pastures, Rangeland and Grass (Hay, Silage). Do not graze or harvest for livestock feed prior to crop maturity. Do not use this product in wheat underseeded with legumes.

EARLY SEASON APPLICATIONS: Apply 0.5 to 1 pint of this product per acre to wheat unless using one of the wheat specific programs below. Early season applications to spring-seeded wheat must be made after tillering and before wheat reaches the 6-leaf stage.

Early season applications to fall-seeded wheat must be made after tillering and prior to the jointing stage. Care should be taken in staging early developing wheat varieties such as TAM 107, Madison, or Wakefield to be certain that the application occurs prior to the jointing stage.

SPECIFIC USE PROGRAMS FOR FALL-SEEDED WHEAT ONLY: Up to 1.33 pints of this product per acre may be applied on fall-seeded wheat after the wheat begins to tiller for suppression of perennial weeds, such as field bindweed. Applications may be made in the fall following a frost but before a killing freeze. Periods of extended stress such as cold and wet weather may enhance the possibility of crop injury. For fall applications only, do not use if the potential for crop injury is not acceptable.

PREHARVEST APPLICATIONS: This product can be used to control weeds that may interfere with harvest of wheat. Apply up to 2 pints of this product per acre as a broadcast or spot treatment to annual broadleaf weeds when wheat is in the hard dough stage and the green color is gone from the nodes (joints) of the stem. Best results will be obtained if application can be made when weeds are actively growing but before weeds canopy. A waiting interval of 7 days is required before harvest. Do not use preharvest-treated wheat for seed unless a germination test is performed on the seed with an acceptable result of 95% germination or better. For control of additional broadleaf weeds or grasses, This product may be tank mixed with other herbicides such as Ally or Roundup® Ultra that are registered for preharvest use in wheat.

Preharvest use of this product is not registered for use in California.

Table 7. WHEAT TANK MIXES

Tank Mix Partner	Rate Per Acre
Aim™	0.3 ounce
Ally®	0.05 to 0.1 ounce ¹
Amber®	0.14 to 0.28 ounce ¹
Bronate®	0.75 to 1.5 pints
Buctril®	1 to 1.5 pints
Canvas®	0.2 to 0.4 ounce ¹
Curtail®	2 to 2.67 pints
Dakota® ²	16 fluid ounces
Express®	0.083 to 0.167 ounce ¹
Finesse®	0.167 to 0.33 ounce ¹
Glean®	0.167 ounce ¹
Harmony® Extra	0.167 to 0.33 ounce ¹
Karmex® ³	0.5 to 1.5 pounds
2,4-D amine	4 to 20 fluid ounces ¹

(continued)

Table 7. WHEAT TANK MIXES (continued)

Tank Mix Partner	Rate Per Acre
Metribuzin [®] (Sencor [®] , Lexone [®])	0.25 to 0.375 pound a.i.
Peak [®]	0.25 to 0.38 ounce
Stinger [®]	4 to 5.33 fluid ounces
Tiller [®]	1 to 1.7 pints

- 1 Do not use low rates of sulfonylurea herbicides, such as Ally, Amber, Canvas, Express, Finesse, Glean, Harmony Extra, and Peak on more mature weeds or on dense vegetative growth.
- 2 Do not use this product as a tank mix treatment with Dakota or Tiller on Durum wheat. Do not tank mix with Tiller if wild oat is the target weed.
- 3 Tank mixes with Karmex and metribuzin are for use in fall-seeded wheat only.
- 4 This product contains 0.36 pounds a.e. of 2,4-D per pint. When tank mixing with 2,4-D, do not exceed a combined total of 1.0 pound a.e. per acre of 2,4-D and do not exceed 0.5 pound a.e. of 2,4-D unless injury to wheat is acceptable.

**BETWEEN CROP APPLICATIONS, CONSERVATION RESERVE PROGRAMS,
GENERAL FARMSTEAD AND FALLOW SYSTEMS**

These uses are considered Food/Feed Crops when harvested, grazed or foraged. Consult Section III. for adjuvant restrictions and Section VII. on Non-Food/Feed Use for specific use directions.

Section VII. NON-FOOD/FEED USE (Land not Harvested, Grazed or Foraged) - Specific Information

BETWEEN CROP APPLICATIONS

PREPLANT DIRECTIONS (POSTHARVEST, FALLOW, CROP STUBBLE, SET-ASIDE) FOR BROADLEAF WEED CONTROL: This product can be applied either postharvest in the fall, spring, or summer during the fallow period or to crop stubble/set-aside acres. Apply this product as a broadcast or spot treatment to emerged and actively growing weeds after crop harvest (postharvest) and before a killing frost or in the fallow cropland or crop stubble the following spring or summer.

See Crop Rotational Restrictions in Section V. General Restrictions and Limitations for the recommended interval between application and planting to prevent crop injury.


Rates and Timings: Apply 0.5 to 6 pints of this product per acre. Refer to Table 1 to determine use rates for specific targeted weed species. Retreatments may be made as needed; however, do not exceed a total of 8 pints of this product per treated acre during a growing season. For best performance, apply this product when annual weeds are less than 6" tall, when biennial weeds are in the rosette stage and to perennial weed regrowth in late summer or fall following a mowing or tillage treatment. The most effective control of upright perennial broadleaf weeds such as Canada thistle and Jerusalem artichoke occurs if this product is applied when the majority of weeds have at least 4 to 6" of regrowth or for weeds such as field bindweed and hedge bindweed that are in or beyond the full bloom stage.

Avoid disturbing treated areas following application. Treatments may not kill weeds that develop from seed or underground plant parts such as rhizomes or bulblets, after the effective period for this product. For seedling control, a follow-up program or other cultural practices could be instituted.

BETWEEN CROP TANK MIXES

In tank mixes with one or more of the following herbicides, apply 0.5 to 2 pints of this product per acre for control of annual weeds, or 2 to 8 pints of this product per acre for control of biennial and perennial weeds:

- Aim™
- Ally[®]
- Amber[®]
- Atrazine
- Bladex[®]
- Curtail[®]
- Cyclone[®]
- Distinct[®]
- Fallowmaster[®]
- Finesse[®]
- Glyphosate
- Gramoxone[®] Extra
- Kerb[®]
- Landmaster[®] BW
- Paramount[®]
- Sencor[®]
- Tordon[®] 22K
- Touchdown[®]
- 2,4-D



CONSERVATION RESERVE PROGRAMS AND GENERAL FARMSTEAD

This product herbicide is recommended for use for Conservation Reserve Programs, general farmstead (non-cropland only), weed and brush control, or use in State Recognized Noxious Weed areas (noncropland areas).

Refer to Tables 1 and 2 for rate selection based on targeted weed or brush species. Some weed species will require tank mixes for adequate control.

Rates above 4 pints of this product per acre are for spot treatments only. Retreatments may be made as needed; however, do not exceed a total of 8 pints of this product per treated acre during a growing season.

FARMSTEAD AND FENCEROW TREATMENT APPLICATION INSTRUCTIONS

This product may be applied using water or oil and water emulsions in spot application to control undesirable vegetation using handgun or similar types of application equipment. In addition to weed species listed in Tables 1 and 2, these treatments may be used to control or suppress woody plant species listed in Table 6.

To prepare oil and water emulsions, mix in the order and proportions indicated below.

The solution should remain milky colored without an oily layer on top when under agitation. If an oily layer forms, increase the amount of emulsifier or change to a more effective emulsifier.

Do not exceed 40 gallons of spray solution per treated acre per application. Forty gallons of spray solution contains 1.0 pound acid equivalent of dicamba and 2.87 pounds acid equivalent of 2,4-D. Spray plants to wet. Do not allow this spray mix to contact desirable vegetation.

To control brush, briars, and weeds along fencerows surrounding pasture and ranch lands, and fallow fields, use a tank mix of 2.5% of this product, 87.5% water, 10% diesel oil, and sufficient emulsifier (to mix the diesel and emulsifier). The diesel oil in this tank mix will damage or kill desirable grasses and should not be used in pastures or where damage to desirable species cannot be tolerated.

- 1) **Water:** Begin by agitating a thoroughly clean sprayer tank with the desired quantity of clean water. Maintain constant agitation during complete mixing procedure.
- 2) **Emulsifier:** Add 0.5% volume to volume.
- 3) **This product:** Add 2.5 gallons per 100 gallons of total intended solution.
- 4) **Diesel Oil:** Add 10 gallons per 100 gallons of total intended solution.

Maintain constant agitation during application. Under good agitation, the spray solution should be milky white with no oil layer on top. If an oil layer forms, increase the amount of emulsifier or change to a more effective emulsifier.

FOR SPRAYING FOLIAR APPLICATIONS:

1. Spray when leaves have reached full size but have not hardened due to drought or maturity.
2. Spray individual plants to wet with handgun.
3. For larger stems (up to 3" in diameter) and hard to control species, direct spray stream to base of stems to wet the stem at soil surface in addition to wetting the foliage.
4. Do not apply under drip line of desirable trees or adjacent to desirable vegetation.

FOR DORMANT BASAL APPLICATIONS:

1. Increase diesel oil content to 15% or 15 gallons of diesel oil per 100 gallons of total solution.
2. Spray in late winter and early spring before plants break dormancy.
3. Spray the bottom 24" of the target stem to wet on all sides.
4. For larger stems (up to 3" in diameter) and hard to kill species direct the spray solution to the base of target stems to wet the soil at the stem/soil junction in addition to wetting the stem.
5. Do not apply under drip line of desirable trees or adjacent to desirable vegetation.

FOR CUT SURFACE TREATMENTS:

Apply this product in an undiluted state as a cut surface treatment to control unwanted trees and prevent sprouts of cut trees.

- **Frill or Girdle Treatments:** Make a continuous cut or a series of overlapping cuts using an axe to girdle tree trunk. Spray or paint the cut surface with this product.
- **Stump Treatments:** Spray or paint freshly cut surface with this product. The cambium layer (the area adjacent to the bark) should be thoroughly wet. Treat stumps within 6 hours after cutting.




Table 6. The following list of trees and vines can be controlled on farmsteads and fencerows as foliar, basal, or cut surface treatments:

Alder	Dogwood	Kudzu	Rose, McCartney
Ash	Elm	Locust, Black	Rose, Multiflora
Aspen	Grape	Maple	Sagebrush, Fringe
Basswood	Greenbriar	Mesquite	Sassafras
Beech	Hawthorn (Thornapple)	Oak	Spruce
Blackberry	Hemlock	Oak, Poison	Sumac
Blackgum	Hickory	Olive, Russian	Sweetgum
Cedar	Honeylocust	Persimmon, Eastern	Sycamore
Cherry	Honeysuckle	Pine	Tarbrush
Chinquapin	Hornbeam	Plum, Sand (Wild Plum)	Willow
Cottonwood	Huckleberry	Poplar	Witchhazel
Creosotebush	Huisache	Rabbitbrush	Yaupon
Dewberry	Ivy, Poison	Redcedar, Eastern	Yucca

WEEDS LISTED IN THIS LABEL:

ANNUALS		BIENNIALS AND PERENNIALS	
Common Name	Scientific Name	Common Name	Scientific Name
Beebalm, Spotted	<i>Monarda punctata</i>	Bindweed, Field	<i>Convolvulus arvensis</i>
Broomweed, Common	<i>Gutierrezia dracunculoides</i>	Bittercross	<i>Cardamine spp.</i>
Buckwheat, Wild	<i>Polygonum convulvulus</i>	Buckeye	<i>Aesculus spp.</i>
Buffalobur	<i>Solanum rostratum</i>	Bullnettle	<i>Cnidoscopus stimulosus</i>
Burdock	<i>Arctium spp.</i>	Chicory	<i>Cichorium intybus</i>
Buttercup, Corn	<i>Ranunculus arvensis</i>	Clover, Hop	<i>Trifolium aureum</i>
Chickweed, Common	<i>Stellaria media</i>	Dandelion	<i>Taraxacum officinale</i>
Cockle, Corn	<i>Agrostemma githago</i>	Dock, Curly	<i>Rumex crispus</i>
Cocklebur, Common	<i>Xanthium strumarium</i>	Elderberry	<i>Sambucus canadensis</i>
Coreopsis, Plains	<i>Coreopsis tinctoria</i>	Goldenrod, Missouri	<i>Solidago missouriensis</i>
Croton, Woolly	<i>Croton capitatus</i>	Goldenweed, Common	<i>Isocoma coronopifolia</i>
Devilsclaw	<i>proboscidea louisianica</i>	Groundsel	<i>Senecio vulgaris</i>
Dogfennel(Cypressweed)	<i>Eupatorium capillifolium</i>	Honeysuckle, Hairy	<i>Lonicera</i>
Eveningprimrose, Cutleaf	<i>Oenothera lacinata</i>	Horsenettle	<i>Solanum carolinense</i>
Flax	<i>Linum catharticum</i>	Ivy, Poison	<i>Rhus radicans</i>
Fleabane, Annual	<i>Erigeron annuus</i>	Knapweed, Black	<i>Centaurea nigra</i>
Flixweed	<i>Descurainia sophia</i>	, Russian	<i>Centaurea repens</i>
Henbit	<i>Lamium amplexicaule</i>	, Spotted	<i>Centaurea maculosus</i>
Knotweed, Prostrate	<i>Polygonum aviculare</i>	Marshelder	<i>Ina annua</i>
Kochia	<i>Kochia scoparia</i>	Mesquite	<i>Prosopis juliflora</i>
Lambsquarters, Common	<i>Chenopodium album</i>	Milkweed, Antelopehorn	<i>Asclepius</i>
Lettuce, Prickly	<i>Lactuca scariola</i>	Nightshade, Silverleaf	<i>Solanum elaeagnifolium</i>
Mallow, Common	<i>Malva neglecta</i>	, Black	<i>Solanum nigrum</i>
Morningglory, Ivyleaf	<i>Ipomea hederacea</i>	Persimmon, Eastern	<i>Diospyros virginiana</i>
, Tall	<i>Ipomea purpurea</i>	Rabbitbrush	<i>Chrysanthemum pulchellus</i>
Mustard, Annual	<i>Brassica spp.</i>	Ragwort, Tansy	<i>Senecio jacobia</i>
, Tansy	<i>Descurainia pinnata</i>	Redvine	<i>Brunnichia ovata</i>
Pennycress, Field	<i>Thlaspi arvense</i>	Sagebrush, Fringed	<i>Artemisia frigida</i>
Pepperweed, Virginia	<i>Lepidium virginicum</i>	Smartweed, Swamp	<i>Polygonum coccineum</i>
Pigweed, Prostrate	<i>Amaranthus blitoides</i>	Sorrel, Red (Sheep Sorrel)	<i>Rumex acetosella</i>
, Redroot	<i>Amaranthus retroflexus</i>	Sowthistle, Perennial	<i>Sonchus arvensis</i>
, Smooth	<i>Amaranthus hybridus</i>	Spurge, Leafy	<i>Euphorbia esula</i>
, Tumble	<i>Amaranthus albus</i>	Starthistle, Yellow	<i>Centaurea solstitialis</i>
Poorjoe	<i>Diodia teres</i>	Tallow Tree, Chinese	<i>Sapium sebiferum</i>
Purslane, Common	<i>Portulaca oleracea</i>	Thistle, Bull	<i>Cirsium vulgare</i>
Ragweed, Common	<i>Ambrosia artemisiifolia</i>	, Canada	<i>Cirsium arvense</i>
, Lance-Leaf	<i>Ambrosia bidentata</i>	, Musk	<i>Carduus nutans</i>
, Western	<i>Ambrosia psilostachya</i>	, Plumeless	<i>Carduus acanthoides</i>
Sedge	<i>Cyperus compressus</i>	Vetch	<i>Vicia spp.</i>
Shepherdspurse	<i>Capsella bursa-pastoris</i>	Yankee weed	<i>Eupatorium compositifolium</i>
Smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>		
Sneezeweed, Bitter	<i>Helianthus amurum</i>		
Sunflower, Common (Wild)	<i>Helianthus annuus</i>		
Thistle, Russian	<i>Salsola iberica</i>		
Velvetleaf	<i>Abutilon theophrasti</i>		

FOOD/FEED CROP USES

This product can be used on the following:

- * Conservation Reserve Program Land
- * Fallow Systems (Between Crop Applications)
- * General Farmstead
 - Grain Sorghum
 - Grass (Hay or Silage)
 - Pastures
 - Rangeland
 - Sugarcane
 - Wheat

* These crops are considered Food/Feed crops only when harvested, grazed or foraged. Otherwise, they are considered as non-Food/Feed uses.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Do not store below 32°F or above 100°F. Store in original container in a well-ventilated area separately from fertilizer, feed, and foodstuffs. Avoid cross-contamination with other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility. Improper disposal of excess pesticide, spray mix, or rinsate is a violation of federal law. If these wastes cannot be disposed of according to label instructions, contact the state agency responsible for pesticide regulation or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL - NON-RETURNABLE PLASTIC: Triple rinse (or equivalent) and add rinsate to spray tank. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

RETURNABLE - REFILLABLE CONTAINERS: After use, return the container to the point of purchase or designated locations. This container must only be refilled with this product. **DO NOT REUSE THE CONTAINER FOR ANY OTHER PURPOSE.** Prior to refilling, inspect thoroughly for damage such as cracks, punctures, abrasions and damaged or worn out threads on closure devices. Do not refill or transport damaged or leaking containers. Check for leaks after refilling and before transportation. If the container is not being refilled, return it to the point of purchase.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(NEW)

