



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

17 APR 2009

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Nufarm Americas, Inc
150 Harvester Drive Suite 200
Burr Ridge, IL 60527

Subject: Nufarm Kambamaster Herbicide
Supplemental label for preplant applications to cropland rotated to wheat
EPA Reg. No. 71368-34
Application dated January 27, 2009

Dear Mr. Meindl:

The supplemental labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable, provided you make the following changes before you release the product for shipment.

1. Replace the text "recommended" with "listed" in the sentence "KAMBAMASTER, when applied at the **recommended** rates, will control ANNUAL and BIENNIAL weeds and suppress the PERENNIAL weeds listed below."
2. Add the following text after the 'Rates and Timing' section in order to be substantially similar with EPA Reg. No. 1381-202 and in compliance with the 2,4-D RED.

"Cropland Rotated to Wheat (Post-Harvest/ Fallow/ Stubble/ Set-Aside)

Restrictions:

Plant only labeled crops within 29 days following application.

Limited to 2 applications per year.

Maximum of 2.0 lbs. 2,4-D ae/acre per application.

Minimum of 30 days between applications."


3. In the sentence above the 'Tank Mix Treatments' section, revise "Retreatments may be made as needed; however, do not exceed a total of 8 pints of KAMBAMASTER per treated acre" to read:
"Retreatment may be made 30 days after the initial treatment; however, do not exceed a total of 8 pints of KAMBAMASTER per treated acre per year."

Page 2
EPA Reg. No. 71368-34

At your next label printing, or within eighteen months of the date of this letter, whichever comes first, you must incorporate this supplemental labeling (with revisions) into the main product labeling.

A stamped copy of the label is enclosed for your records. Please submit one copy of your final printed label before you release the product for shipment.

Sincerely,



Joanne Miller
Product Manager 23
Herbicide Branch
Registration Division (7505P)

SUPPLEMENTAL LABELING

Nufarm Kambamaster Herbicide EPA Reg. No. 71368-34

READ AND FOLLOW THE ENTIRE LABEL BOOKLET FOR WEEDMASTER HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING

"Label" as used in this supplemental labeling for Nufarm Kambamaster Herbicide and this supplement.

PREPLANT APPLICATION DIRECTIONS FOR BROADLEAF CONTROL IN CROPLAND ROTATED TO WHEAT (POST-HARVEST / FALLOW / STUBBLE / SET-ASIDE)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application. All application directions and precautions in the Nufarm Kambamaster Herbicide label booklet must be followed.

Before using this product, read and carefully observe all application directions, restrictions, and precautionary statements in this booklet and on the container label.

WEEDS CONTROLLED

KAMBAMASTER®, when applied at the recommended rates, will control the ANNUAL and BIENNIAL weeds and suppress the PERENNIAL weeds listed below.

ANNUALS		
Buckwheat, Wild	Mustards	Salsify, Western
Cockle, Cow	Nightshade, Black	Smartweed, Pennsylvania
Cocklebur, Common	Pigweed, Redroot (Carelessweed)	Sowthistle, Annual
Knotweed	Pigweed, Rough	Sunflower
Kochia	Purslane, Common	Tansymustard
Lambsquarters, Common	Ragweed, Common	Thistle, Russian
Mallow, Common	Sage, Lanceleaf	Velvetleaf
BIENNIALS		
Carrot, Wild	Starthistle, Yellow	Thistle, Musk
Ragwort, Tansy	Thistle Bull	Thistle, Plumeless
PERENNIALS		
Bindweed, Field	Dock, Curly	Thistle, Canada

RATES AND TIMING

ACCEPTED
with COMMENTS
in EPA Letter Dated

17 APR 2009

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

71368-34

Application may be made to fallow land, wheat stubble or land to be rotated to wheat. Application should be made to emerged and actively growing weeds. Use higher rate when treating dense vegetative growth. Avoid disturbing treated areas for seven days following application.

Wheat injury may occur if the interval between application and planting is less than 10 days for each pint per acre of KAMBAMASTER® used. Exclude days when ground is frozen.

Weed Type & Stage	Broadcast Rate per Treated Acre Amount
Annual	
Small, actively growing (less than 4 inches)	1.0 – 1.5 pints
Established weed growth (greater than 4 inches)	1.5 – 3.0 pints
Biennial	
Rosette diameter (3 inches or less)	1.5 – 2.0 pints
(3 inches or more)	2.0 – 4.0 pints
Greater than 4 inches, tillering	4.0 pints
Bolted or flowering	
Perennial	
Suppression or top growth control	2.0 – 4.0 pints
Seasonal Control	4.0 – 8.0 pints

Add 0.5% v/v of an agriculturally approved surfactant to KAMBAMASTER® when used alone or in a tank mix. The addition of a surfactant will enhance spray coverage and the herbicide's penetration of weed foliage.

Retreatments may be made as needed; however, do not exceed a total of 8 pints of KAMBAMASTER® per treated acre.

TANK MIX TREATMENTS

KAMBAMASTER® may be tank mixed with one or more of the following herbicides for control of grasses or additional broadleaf weeds. Read and follow the label of each tank mix product used for precautionary statements, directions for use, rates and timings, weeds controlled, geographic or other restrictions. Add 0.5% v/v of an agriculturally approved surfactant to all tank mixes.

Herbicide	Rate per Treated Acre (lbs a.i.)
Atrazine	½ to 3.0
Chlorsulfuron	0.016 to 0.024

Glyphosate	¼ to 2.0
Metribuzin	1/3 to ¾
Paraquat	½ to 1.0

NOTICE

Read the WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY in the label booklet for Nufarm Kambamaster Herbicide before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

**Manufactured By
Nufarm Inc.
Burr Ridge, IL 60527**