

71368-38

1/15/2009

10f26



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Ms. Lizbeth Rea
Registration Specialist
Nufarm Americas Inc.
150 Harvester Drive Suite 200
Burr Ridge, ILL 60527

JAN 15 2009

RE: Notification change in Primary Brand Name: "Purestand Selective Herbicide"
EPA Registration Number: 71368-38
Date of Submission: November 14, 2008

Dear Ms. Rea:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated November 14, 2008, for the product Purestand Selective Herbicide. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Joyce Edwards of my staff at 703-308-5479.

Sincerely,

A handwritten signature in black ink, appearing to read "Linda Arrington".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

2026



United States
Environmental Protection Agency
 Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 71368-38	2. EPA Product Manager Jim Tompkins	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Purestand Dry Flowable Herbicide	PM# 25	
5. Name and Address of Applicant (Include ZIP Code) Nufarm Americas, Inc. 150 Harvester Drive Suite 200 Burr Ridge, IL 60527 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is NOTIFICATION composition and labeling to: EPA Reg. No. <u>JAN 15 2009</u> Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)
 Primary brand name was changed from Purestand Dry Flowable Herbicide to Purestand Selective Herbicide.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Metal Plastic Glass Paper Other (Specify) _____		
* Certification must be submitted		If "Yes" Unit Packaging wgt.			
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Lizbeth Rea	Title Registration Manager	Telephone No. (Include Area Code) 919/655-0701
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Registration Manager	
4. Typed Name Lizbeth Rea	5. Date 11/14/2008	



30236

Nufarm Americas, Inc.
Lizbeth Rea
Registration Manager
150 Harvester Drive, Suite 200
Burr Ridge, IL 60527
Phone: 919.655.0701 Fax: 919.342.5176
liz.rea@us.nufarm.com

November 14, 2008

Via Overnight Courier

Jim Tompkins (PM-25)
Document Processing Desk (AMEND)
Office of Pesticide Programs (7504P)
U. S. Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 S. Crystal Drive
Arlington, VA 22202-4501

Subject Purestand Dry Flowable Herbicide
: EPA Reg. No. 71368-38
Change in Primary Brand Name

Dear Mr. Tompkins:

The following documents are attached in support of our request to change the primary brand name of Purestand Dry Flowable Herbicide to Purestand Selective Herbicide.


- Application for a Product Registration (EPA Form 8570-1)
- One (1) copy of the Purestand Dry Flowable Herbicide label with changes tracked.
- Five (5) copies of the Purestand Dry Flowable Herbicide label with changes incorporated.

The primary brand name is the only change made to the enclosed label. No other changes to the label were made.

Please call me at 919/655-0701 if you have any questions and/or comments regarding this submission.

Sincerely,

Lizbeth Rea
Registration Manager
Nufarm Americas Inc.



4 of 36

this product ~~Dry Flowable~~ Purestand Selective Herbicide

For Use on Wheat, Barley, Fallow, Pastures and Rangeland

ACTIVE INGREDIENT:

Metsulfuron Methyl Methyl 2-[[[(4-methoxy-6-methyl -1,3,5-triazin-2y0amino) Carbonyl] amino]sutfonyl]benzoate..... 60.0%

INERT INGREDIENTS..... 40.0%

Total..... 100.0%

E.P.A. Reg. No. 71368-38

E.P.A. Est. No. 22844

NOTIFICATION

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

JAN 15 2009

Si usted no endende la etiqueta, busque a algulen para que se la explique a usted en details.
(If you do not understand the label, find someone to explain it to you in detail.)

See Inside for Additional Precautionary Statements.

**For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300.
For Medical Emergencies Only, Call 877-325-1840.**

Manufactured By:
Nufarm, Inc.
Burr Ridge, IL

NET CONTENTS:

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none"> • Call a doctor or poison control center immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Takeoff contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment for advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after first 5 minutes, rinsing eye. • Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
<p align="center">HOT LINE NUMBER</p> <p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.</p>	
<p align="center">NOTE TO PHYSICIAN</p> <p>This product may pose an aspiration pneumonia hazard. Contains petroleum distillates:</p>	

50724

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Causes eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust or spray mist.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard [(40 CFR Part 170)] must wear:

- Long-sleeved shirt and long pants.
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE, if no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, and chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

IMPORTANT INFORMATION

PESTICIDE HANDLING

- Calibrate sprayers only with clean water away from the well site.
- Make scheduled checks of spray equipment.
- Assure accurate measurement of pesticides by all operation employees.
- Mix only enough product for the job at hand.
- Avoid overfilling of spray tank.
- Do not discharge excess material on the soil at a single spot in the field or mixing/loading station.
- Dilute and agitate excess solution and apply at labeled rates/uses.
- Avoid storage of pesticides near well sites.
- When triple rinsing the pesticide container, be sure to add the rinsate to the spray mix.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls.
- Shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Weed control in pastures and rangeland is not within the scope of the Worker Protection Standard.

Use ~~Purestand~~this product only in accordance with recommendations on this label or in separate published Nufarm recommendations.

Nufarm will not be responsible for losses or damages resulting from the use of this product in any manner not specifically recommended by Nufarm.

Do not apply this product through any type of irrigation system.

GENERAL INFORMATION

Nufarm ~~Purestand~~this product herbicide is recommended for use on land primarily dedicated to the production of wheat, barley, fallow, pasture, and rangeland.

~~Purestand~~This product is recommended for use on wheat, barley, fallow, pasture and rangeland in most states, check with your state extension or Dept. of Agriculture before use, to be certain ~~Purestand~~this product is registered in your state.

~~Purestand~~This product is not registered for use in Alamosa, Conejos, Costilla, Rio Grande, and Saquache counties of Colorado.

~~Purestand~~This product is a dry-flowable granule that controls weeds in wheat (including durum), barley, pasture, rangeland grasses, and fallow. ~~Purestand~~This product is mixed in water or can be pre-slurried in water and added to liquid nitrogen carrier solutions and applied as a uniform broadcast spray. A surfactant should be used in the spray mix unless otherwise specified on this label. ~~Purestand~~this product is noncorrosive, nonflammable, nonvolatile, and does not freeze.

~~Purestand~~This product controls weeds by postemergence activity. For best results, apply ~~Purestand~~this product to young, actively growing weeds. The use rate depends upon the weed spectrum and size of weeds at application. The degree and duration of control may depend on the following factors:

- weed spectrum and infestation intensity
- weed size at application
- environmental condition at and following treatment

Environmental Conditions and Biological Activity

~~Purestand~~This product is absorbed through the foliage of broadleaf weeds, rapidly inhibiting their growth. Leaves of susceptible plants appear chlorotic from 1 to 3 weeks after application and the growing point subsequently dies.

Application of ~~Purestand~~this product provides the best control in vigorously growing crops that shade competitive weeds. Weed control in areas of thin crop stand or seeding skips may not be as satisfactory. However, a crop canopy that is too dense at application can intercept spray and reduce

weed control.

Nufarm Purestand This product may injure crops that are stressed from adverse environmental conditions (such as extreme temperatures or moisture), abnormal soil conditions, or cultural practices. In addition, different varieties of the crop may be sensitive to treatment with Purestand this product under otherwise normal conditions. Treatment of such varieties may injure crops.

In warm, moist conditions, the expression of herbicide symptoms is accelerated in weeds; in cold, dry conditions, expression of herbicide symptoms is delayed. In addition, weeds hardened-off by drought stress are less susceptible to Purestand this product.

Weed control may be reduced if rainfall or snowfall occurs soon after application.

APPLICATION INFORMATION

Use Rates

Wheat (including durum) and Barley

Apply 1/10 oz Purestand this product per acre to wheat or barley.

Pasture and Rangeland

Apply 1/10 to 4/10 oz Purestand this product per acre as a broadcast treatment to pasture and rangeland. For spot applications, use 1 oz per 100 gal of water. Do not exceed 3/4 oz Purestand this product per acre.

Harvest Aid

Apply 1/10 oz Purestand this product per acre in combination with 2,4-D or Roundup to aid in dry down of many broadleaved weeds, thereby aiding grain harvest.

Fallow

Apply Purestand this product at 1/10 oz per acre.

Application Timing—Wheat and Barley

Dryland Wheat and Barley

(Except Durum or Wampum Variety)

Make applications after the crop is in the 2-leaf stage but before boot.

Durum and Wampum Variety Spring Wheat

Make applications after the crop is tillering but before boot. Applications to durum and wampum varieties should be made in combination with 2,4-D.

Irrigated Wheat and Barley

Make applications after the crop begins tillering but before boot. First post-treatment irrigation should be delayed for at least 3 days after treatment and should not exceed 1 in. of water

Wheat and Barley--Harvest Aid

Make applications after the crop has reached the hard dough stage, but no later than 10 days before harvest.

See section on Harvest Aid tank mixtures.

Fallow

Purestand This product may be used as a fallow treatment, in the spring or fall when the majority of weeds have emerged and are actively growing.

Do not apply during boot or early heading, as crop injury may result.

Application Timing—Pasture Grasses

Purestand This product may be used on some native grasses such as bluestems and grama, and on other pasture grasses such as bermudagrass, bluegrass, orchardgrass, bromegrass, fescue and timothy. Specific application information on several of these pasture grasses follows:

Pasture Grass	Minimum time from grass establishment to Purestand application
Bermudagrass	2 months
Bluegrass, brome grass, and orchardgrass	6 months
Timothy	12 months
Fescue	24 months

Fescue Precautions:

Note that Purestandthis product may temporarily stunt fescue, cause it to turn yellow, or cause seed head suppression. To minimize these symptoms, take the following precautions:

- tank mix Purestandthis product with 2,4-D
- use the lowest recommended rate for target weeds
- use surfactant at 1/2 to 1 pt per 100 gal of spray solution (1/16 to 1/8% v/v)
- make application later in the spring after the new growth is 5 to 6 inches tall, or in the fall.
- Do not use surfactant when liquid nitrogen is used as a carrier.

The first cutting yields may be reduced due to seed head suppression resulting from treatment with Purestandthis product.

Timothy Precautions:

Timothy should be at least 6" tall at application and be actively growing. Applications of Purestandthis product to timothy under any other conditions may cause crop yellowing and/or stunting. To minimize these symptoms, take the following precautions:

- tank mix Purestandthis product with 2,4-D
- use the lowest recommended rate for target weeds
- use surfactant at 1/2 pt per 100 gal (1/16% v/v)
- make applications in the late summer or fall
- Do not use surfactant when liquid nitrogen is used as a carrier.

Ryegrass Pastures (Italian or perennial): Do not apply Purestandthis product as injury to or loss of the pasture may result.

Other Pastures: Varieties and species of pasture grasses differ in their tolerance to herbicides. When using Purestandthis product on a particular grass for the first time, limit use to one container. If no injury occurs throughout the season, larger acreage may be treated the following season.

Broadleaf pasture species, such as alfalfa and clover are highly sensitive to Purestandthis product and will be severely stunted or injured by Purestandthis product.

WEEDS CONTROLLED

Unless otherwise directed, treat when weeds are less than 4" tall or in diameter and are actively growing.

Effectiveness may be reduced if rainfall occurs within 4 hrs after application.

Cereals, Pasture, Rangeland, and Fallow
1/10 oz per acre

Blue/purple mustard*	Filaree	Pigweed (redroot, smooth, tumble)	Snow speedwell
Bur buttercup (testiculate)	Flixweed*	Plains coreopsis	Tansymustard*
Coast fiddleneck (tarweed)	Groundsel	Prickly lettuce*	Treacle mustard (Bushy Wallflower)
Common chickweed	Henbit	Russian thistle*	Tumble/Jim Hill
Common purslane	Kochia*	Shepherd's purse	Volunteer sunflower
Conical catchfly	Lambsquarters (common, slimleaf)	Smallseed falseflax	Waterpod
Cowcockle	Mayweed chamomile	Smartweed (green, ladysthumb, pale)	Wild Mustard
False chamomile	Miners lettuce		
Field pennycress (fanweed)			

Additional Weeds in Pasture/Rangeland Only

1/10 to 2/10 oz per acre

Bitter sneezeweed	Common broomweed	Dandelion	Wild garlic*
Buttercup	Common mullein	Marestail	Woolly croton*
Carolina geranium	Curly dock	Plantain	

2/10 to 3/10 oz per acre

Annual marshelder	Burclover	Horsemint (beebalm)	Purple scabious
Blackeyed-Susan	Common yarrow	Musk thistle*	Western snowberry‡
Buckbrush†	Dogfennel	Pensacola bahiagrass*	Wild carrot

4/10 oz per acre

Serecia lespedeza*

Weeds Suppressed‡*

Cereals, Pastures, Rangeland, and Fallow

1/10 oz per acre

Canada thistle*	Corn groomwell*	Sowthistle (annual)*
Common sunflower*	Knotweed (prostrate)*	Wild buckwheat*

Brush Suppressed‡

3/10 oz per acre

Blackberry	Dewberry	Multiflora rose*
------------	----------	------------------

Weeds/Brush Suppressed with Spot Application

(Pasture/Rangeland only)

1 oz per 100 gal of water

Blackberry*	Canada thistle*	Dewberry*	Multiflora rose*
-------------	-----------------	-----------	------------------

* See the **Specific Weed Problems** section.

‡ Weed suppression is a reduction in weed competition (reduced population and/or vigor) as visually compared to an untreated area. The degree of suppression varies with the rate used, the size of the weeds, and the environmental conditions following treatment.

Specific Weed Problems

Note: Thorough spray coverage of all weed species listed below is very important.

Blue Mustard, Flixweed, and Tansymustard: For best results, apply ~~Nufarm Purestand~~ this product tank mixtures with 2,4-D or MCPA postemergence to mustards, but before bloom.

Canada Thistle and Sowthistle: Apply either ~~Purestand~~ this product plus surfactant or ~~Purestand~~ this product plus 2,4-D or MCPA in the spring after the majority of thistles have emerged and are small (rosette stage to 6" elongating stems) and actively growing. The application will inhibit the ability of emerged thistles to compete with the crop.

For Spot applications to Canada thistle in pasture and rangeland, apply as a foliar spray once plant is fully leafed. Apply to runoff and include a surfactant in the spray mix at 1 to 2 qt per 100 gal of spray solution. Complete coverage of all foliage and stems is required for control. On tall, dense stands, it is often necessary to spray from both sides to obtain adequate coverage.

Corn Cromwell and Prostrate Knotweed: Apply ~~Purestand~~ this product plus surfactant when weeds are actively growing are no larger than 2" tall, and when crop canopy will allow thorough coverage. Tank mixing 2,4-D or MCPA with ~~Purestand~~ this product can improve results.

Kochia, Russian thistle, and Prickly lettuce: Naturally occurring resistant biotypes of these weeds are known to occur. For best results, use ~~Purestand~~ this product in a tank mix with "Banvel"/"Banvel" SGF and 2,4-D, or bromoxynil and 2,4-D (such as 3/4 - 1 pt "Buctril" + 1/4-3/8 lb active 2,4-D ester). ~~Purestand~~ This product should be applied in the spring when kochia, Russian thistle, and prickly lettuce are less than 2" tall or 2" across and are actively growing (refer to the Tank Mixtures section of this label for additional details).

Sunflower (common/volunteer): Apply either Nufarm Purestandthis product plus surfactant or Purestandthis product plus 2,4-D or MCPA after the majority of sunflowers have emerged, are 2" to 4" tall and are actively growing. Use spray volumes of at least 3 gal by air or 5 gal by ground (10 gal by ground in pastures).

Wild Buckwheat: For best results, apply Purestandthis product plus 2,4-D or MCPA when plants have no more than 3 true leaves (not counting the cotyledons). If plants are not actively growing, delay treatment until environmental conditions favor active weed growth.

Musk Thistle: Apply Purestandthis product at 2/10 to 3/10 oz per acre in the spring or early summer prior to flowering or in the fall after newly emerged plants have reached the rosette stage of growth. Fall applications should be made before the soil freezes.

Multiflora Rose: For best control, apply Purestandthis product as a broadcast application when multiflora rose is less than 3' tall. Application should be made in the spring, soon after multiflora rose is fully leafed.

For Spot applications in pasture and rangeland, apply as a foliar spray once plant is fully leafed. Apply to runoff and include a surfactant in the spray mix at 1 to 2 qt per 100 gal of spray solution. Complete coverage of all foliage and stems is required for control. On tall, dense stands, it is often necessary to spray from both sides to obtain adequate coverage.

Blackberry and Dewberry: For Spot applications in pasture and rangeland, apply as a foliar spray once plant is fully leafed. Apply to runoff and include a surfactant in the spray mix at 1 to 2 qt per 100 gal of spray solution. Complete coverage of all foliage and stems is required for control. On tall, dense stands, it is often necessary to spray from both sides to obtain adequate coverage.

Pensacola bahiagrass control in established Bermudagrass pasture: Apply Purestandthis product at 3/10 oz per acre plus surfactant Apply after green-up in the spring but before bahiagrass seedhead formation.

Application should be made when moisture is sufficient to enhance grass growth. PurestandThis product is very effective for removal of bahiagrass from bermudagrass pastures. In highly infested pastures, the use of Purestandthis product can clear the areas of useful forage until the bermudagrass has time to cover the area.

Therefore, Purestandthis product treatments should be spread out over a period of years. Do not apply to an entire farm or ranch in one year. Fertilization (particularly with nitrogen and potassium) and/or replanting may accelerate the process of reestablishment of bermudagrass.

Under heavy bahiagrass pressure, grazing pressure, or adverse weather conditions (heat and drought), bahiagrass regrowth may occur.

Note: PurestandThis product should not be used for the control of common or Argentine bahiagrass. Also, Purestandthis product should not be applied in liquid fertilizer solutions for Pensacola bahiagrass control, as poor control and/or regrowth may occur.

Serecia lespedeza: Apply Purestandthis product at 4/10 oz per acre plus a surfactant at 1 to 2 qt per 100 gal of total spray solution. For best results, make applications to serecia lespedeza beginning at flower bud initiation through the full bloom stage of growth.

Note: Do not make applications if drought conditions exist at intended time of application.

Wild Garlic: Apply 1/10 to 2/10 oz per acre of Purestandthis product in the early spring when wild garlic is less than 12" tall with 2" to 4" of new growth.

Woolly Croton: Apply 1/10 to 2/10 oz per acre of Purestandthis product in the late spring or early summer at preemergence through 2 true leaf stage.

Surfactants

Unless otherwise specified, add a Nufarm recommended nonionic surfactant having at least 80% active ingredient at 1 to 2 qt per 100 gal of spray solution (0.25 to 0.5% v/v).

Exceptions: (1) On all spring wheat and spring or winter barley use 1/2 to 1 qt per 100 gals; (2) on Fescue pastures use 1/4 to 1/2 qt per 100 gals; (3) on Timothy pastures use 1/4 qt per 100 gals.

Consult your agricultural dealer, applicator, or Nufarm representative for a listing of recommended surfactants.

Antifoaming agents may be used if needed.

Do not use low rates of liquid fertilizer as a substitute for surfactant.

Ground Application

To obtain optimum spray distribution and thorough coverage, use flat-fan or low-volume flood nozzles.

For flood nozzles on 30" spacings, use at least 10 gallons per acre (GPA), flood nozzles no larger than 10 (or equivalent), and a pressure of at least 30 pounds per square inch (psi). For 40" nozzle spacings, use at least 13 GPA; for 60" spacings, use at least 20 GPA. It is essential to overlap the nozzles 100% for all spacings.

With "Raindrop RA" nozzles, use at least 30 GPA and ensure that nozzle spray patterns overlap 100%.

For flat-fan nozzles, use at least 3 CPA for applications to wheat or barley. Use at least 10 GPA for applications to pasture or rangeland.

Use 50-mesh screens or larger.

Aerial Application

Use nozzle types and arrangements that provide optimum spray distribution and maximum coverage.

Wheat, Barley and Fallow--use 1 to 5 GPA. Use at least 3 GPA in Idaho, Oregon, or Utah.

Pasture and Rangeland--use 2 to 5 GPA.

When applying Purestand this product by air in areas adjacent to sensitive crops, use solid stream nozzles oriented straight back. Adjust the swath to avoid spray drift damage to sensitive crops downwind and/or use ground equipment to treat the border edge of fields. See the **Spray Drift Management** section of this label.

Product Measurement

Purestand This product is measured using the Purestand volumetric measuring cylinder. The degree of accuracy of this cylinder varies by +/- 7.5%. For more precise measurement, use scales calibrated in ounces.

TANK MIXTURES

Nufarm Purestand This product may be tank mixed with other suitable registered herbicides to control weeds listed under **Weeds Suppressed**, weeds resistant to Purestand this product, or weeds not listed under **Weeds Controlled**. Read and follow all manufacturer's label recommendations for the companion herbicide. If those recommendations conflict with this label, do not tank mix the herbicide with Purestand this product.

Tank Mixtures in Cereals (Wheat and Barley)

With 2,4-D (amine or ester) or MCPA (amine or ester)

Purestand This product can be used as a tank-mix treatment with 2,4-D or MCPA (ester formulations provide best results) herbicides after weeds have emerged. For best results, use 1/10 oz of Purestand this product per acre; add 2,4-D or MCPA herbicides to the tank at 1/4 to 1/2 lb active ingredient. Surfactant may be added to the mixture at 1/2 to 1 qt per 100 gal of spray solution; however, adding surfactant may increase the potential for crop injury.

Apply Purestand this product plus MCPA after the 3 to 5-leaf stage but before boot (with Durum and Wampum varieties do not apply before tillering). Apply Purestand this product plus 2,4-D after tillering (refer to appropriate 2,4-D manufacturer's label), but before boot.

With "Banvel"/"Banvel" SGF

For best results, apply Purestand this product at 1/10 oz per acre; add 1/16 to 1/8 lb active ingredient "Banvel"/"Banvel" SGF. Surfactant may be added to the mixture at 1/2 to 1 qt per 100 gal of spray solution; however, adding surfactant may increase the potential for crop injury. Also refer to "Banvel"/"Banvel" SGF labels for application timing and restrictions

With 2,4-D (amine or ester) and "Banvel"

Purestand This product may be applied in a 3-way tank mix with formulations of "Banvel" and 2,4-D. Observe all applicable directions, restrictions and precautions on labels of all products used.

Make applications at 1/10 oz of Purestand this product + 2 - 3 oz "Banvel" (4 - 6 oz "Banvel" SGF) + 4 - 6 oz active 2,4-D Ester or Amine per acre. Use higher rates when weed infestation is heavy. Add 1-2 pt of surfactant to the 3 way mixture, where necessary, as deemed by local recommendations. Use of additional surfactant may not be needed with the higher phenoxy rates and ester phenoxy formulations. Consult the specific 2,4-D or "Banvel" label, or local recommendations for more information.

Apply this 3-way combination to winter wheat after the crop is tillering and prior to jointing (first node). In Spring Wheat (including Durum wheat) apply after the crop is tillering and before it exceeds the 5-leaf stage. Do not apply this 3-way mixture at high rates more than once a year or more than twice per year at the low rates.

With bromoxynil (such as "Buctril", "Bronate")

~~Purestand~~This product may be tank mixed with bromoxynil containing herbicides registered for use on wheat, barley, or fallow. For best results, add bromoxynil containing herbicides to the tank at 3 to 6 oz active ingredient per acre (such as "Bronate" or "Buctril" at 3/4 - 1 1/2 pt per acre).
Read and follow all label instructions on timing, precautions, and warnings for these herbicides before using these tank mixtures. Follow the most restrictive labeling.

With grass control products

Tank mixtures of ~~Purestand~~this product and grass control products may result in poor grass control. Nufarm recommends that you first consult your state experiment station, university, or extension agent, Agricultural dealer, or Nufarm representative as to the potential for antagonism before using the mixture. If no information is available, limit the initial use of ~~Purestand~~this product and the grass product to a small area.
To control wild oat, tank mix ~~Purestand~~this product with "Avenge" or "Assert".
When tank mixing ~~Purestand~~this product with "Assert", always include 2,4-D ester, MCPA ester, or Bromoxynil containing products (such as "Buctril", or "Bronate"). Tank-mixed applications of ~~Purestand~~this product plus "Assert" may cause temporary crop discoloration, stunting, or injury when heavy rainfall occurs shortly after application.
Do not tank mix ~~Purestand~~this product with "Hoelon" 3EC, as grass control may be reduced.

EXPRESS®

~~Purestand~~This product may be tank mixed with Express based on local recommendations.
Read and follow all label instructions on timing, precautions, and warnings for these herbicides before using this tank mixture.

HARMONY® EXTRA

~~Purestand~~this product may be tank mixed with Harmony Extra based on local recommendations.
Read and follow all label instructions on timing, precautions, and warnings for these herbicides before using this tank mixture.

With Insecticides and Fungicides

~~Purestand~~This product may be tank-mixed or used sequentially with insecticides and fungicides registered for use on cereal grains.
However, under certain conditions (drought stress, cold weather, or if the crop is in the 2-4 leaf stage), tank mixes or sequential applications of ~~Purestand~~this product with organophosphate insecticides (such as parathion, "Di-Syston") may produce temporary crop yellowing or, in severe cases, crop injury.
The potential for crop injury is greatest when wide fluctuations in day/night temperatures occur just prior to or soon after application.
Test these mixtures in a small area before treating large areas.
Do not apply ~~Purestand~~this product within 60 days of crop emergence where an organophosphate insecticide (such as "Di-Syston") has been applied as an in-furrow treatment, as crop injury may result.
Do not use ~~Purestand~~this product plus Malathion, as crop injury will result.

With Liquid Nitrogen Solution Fertilizer

Liquid nitrogen fertilizer solutions may be used as a carrier in place of water. Run a tank mix compatibility test before mixing ~~Purestand~~this product in fertilizer solution..
~~Purestand~~This product must first be slurried with water and then added to liquid nitrogen solutions (e.g., 28-0-0, 32-0-0). Ensure that the agitator is running while the ~~Purestand~~this product is added. Use of this mixture may result in temporary crop yellowing and stunting.
If using low rates of liquid nitrogen fertilizer in the spray solution (less than 50% of the spray solution volume), the addition of surfactant is necessary. Add surfactant at 1/2 pt-1 qt per 100 gal of spray solution (0.06 - 0.25% v/v) based on local recommendations.

When using high rates of liquid nitrogen fertilizer in the spray solution, adding surfactant increases the risk of crop injury. Consult your agricultural dealer, consultant, fieldman, or Nufarm representative for a specific recommendation before adding an adjuvant to these tank mixtures.

If 2,4-D or MCPA is included with Nufarm Purestandthis product and fertilizer mixture, ester formulations tend to be more compatible (See manufacturer's label). Do not add surfactant when using Purestandthis product in tank mix with 2,4-D ester or MCPA ester and liquid nitrogen fertilizer solutions.

Do not use low rates of liquid fertilizer as a substitute for a surfactant.

Do not use with liquid fertilizer solutions with a pH less than 3.0.

Tank Mixtures in Harvest Aid

A tank mix of Purestandthis product plus 2,4-D and surfactant, or "Roundup", will typically aid in dry down of many broadleaved weeds, thereby aiding grain harvest. Postemergence application should be made to actively growing weeds after the crop is in the hard dough stage. If weeds are not dry within 10 days after application, delay harvest until weeds are dry.

See weeds listed in Weeds Controlled chart of this label.

With 2,4-D.

Use 1/10 oz Purestandthis product plus 1/4 to 1/2 lb active ingredient 2,4-D per acre on moderate weed infestations; higher rates of 2,4D may be used on large weeds if permitted by the 2,4-D brand labeling. Include 1 to 2 qt surfactant per 100 gal spray solution.

In addition to the weeds listed in Weeds Controlled chart of this label, the 2,4-D combination will also dry down common cocklebur, marestail, puncturevine and common and wild sunflower. In areas where 2,4-D use is restricted, apply Purestandthis product with surfactant only; however, this treatment may be less effective.

With "Roundup"

Use 1/10 oz Purestandthis product plus the locally recommended rate of "Glyphosate" (see "Glyphosate" label for maximum seasonal rate). Purestandthis product requires the use of an adjuvant for optimum activity - Consult the "Glyphosate" label or local recommendations for the amount of adjuvant to include.

Tank Mixtures in Fallow

PurestandThis product may be used as a fallow treatment, and may be tank mixed with other herbicides that are registered for use in fallow.

Read and follow all manufacturer's label recommendations for the companion herbicide. If those recommendations conflict with this label, do not tank mix the herbicide with Purestandthis product.

Tank Mixtures in Pastures or Rangeland

PurestandThis product can be applied in a tank-mix combination with "Grazon P+D", "Tordon" 22K, 2,4-D, "Banvel", or "Weedmaster" in states where these products are labeled for postemergence control of the following weeds:

Annual marshelder	Common cocklebur	Giant ragweed	Western ragweed
Burclover	Common milkweed	Prickly lettuce	
Carolina horsenettle	Common ragweed	Sunflower	

For best results, apply Purestandthis product at 1/10 to 2/10 oz per acre with one of the following products:

Product	Rate (oz /A)
"Grazon P+D"	8 to 32
"Tordon" 22K	4 to 16
2,4-D	16 to 32
"Banvel"	4 to 32
"Weedmaster"	8 to 32
"Remedy"	8
"Amber"	0.35 *

* For suppression of Ragweed In Phenoxy Restricted and Herbicide Regulated Counties

With Liquid Nitrogen Solution Fertilizer

Liquid nitrogen fertilizer solutions may be used as a carrier in place of water. Run a tank mix compatibility test before mixing Purestandthis product in fertilizer solution.

PurestandThis product must first be slurried with water and then added to liquid nitrogen solutions (e.g., 28-0-0, 32-0-0). Ensure that the agitator is running while the Purestandthis product is added. Use of this mixture may result in temporary crop yellowing and stunting.

If using low rates of liquid nitrogen fertilizer in the spray solution (less than 50% of the spray solution volume), the addition of surfactant is necessary. Add surfactant at 1/4 pt per 100 gal of spray solution (0.03% v/v).

When using high rates of liquid nitrogen fertilizer in the spray solution, adding surfactant increases the risk of crop injury. Consult your agricultural dealer, consultant, fieldman, or Nufarm representative for a specific recommendation before adding an adjuvant to these tank mixtures.

If 2,4-D or MCPA is included with Purestandthis product and fertilizer mixture, ester formulations tend to be more compatible (See manufacturer's label). Do not add surfactant when using Purestandthis product in tank mix with 2,4-D ester and liquid nitrogen fertilizer solutions.

Do not use low rates of liquid fertilizer as a substitute for a surfactant.

Do not use with liquid fertilizer solutions with a pH less than 3.0.

CROP ROTATION

Before using Nufarm Purestandthis product, carefully consider your crop rotation plans and options. For rotational flexibility, do not treat all of your wheat, barley, fallow, pasture, or rangeland acres at the same time.

Minimum Rotational intervals

Minimum rotation intervals* are determined by the rate of breakdown of Purestandthis product applied. PurestandThis product breakdown in the soil is affected by soil pH, presence of soil microorganisms, soil temperature, and soil moisture. Low soil pH, high soil temperature, and high soil moisture increase Purestandthis product breakdown in soil, while high soil pH, low soil temperature, and low soil moisture slow Purestandthis product breakdown.

Of these 3 factors, only soil pH remains relatively constant. Soil temperature, and to a greater extent, soil moisture, can vary significantly from year to year and from area to area. For this reason, soil temperatures and soil moisture should be monitored regularly when considering crop rotations.

* The minimum rotation interval represents the period of time from the last application to the anticipated date of the next planting.

Soil pH Limitations

PurestandThis product should not be used on soils having a pH above 7.9, as extended soil residual activity could extend crop rotation intervals beyond normal. Under certain conditions, Purestandthis product could remain in the soil for 34 months or more, injuring wheat and barley. In addition, other crops planted in high-pH soils can be extremely sensitive to low concentrations of Purestandthis product

Checking Soil pH

Before using Purestandthis product, determine the soil pH of the areas of intended use. To obtain a representative pH value for the test area, take several 0" to 4" samples from different areas of the field and analyze them

separately. Consult local extension publications for additional information on recommended soil sampling procedures.

Rotational Intervals for Cereals

All Areas - Following Use of Purestand at 1/10 oz per Acre

Crop	Soil pH	Minimum Cumulative Precipitation (inches)	Minimum Rotation Interval
Winter and spring wheat	7.9 or lower	No restrictions	1
Durum wheat, barley, spring/winter oat	7.9 or lower	No restrictions	10

Rotation Intervals for Crops in Non-Irrigated Land

Following Use of Purestand at 1/10 oz per Acre on Wheat, Barley, Fallow or Pasture

Location		Crop	Soil pH	Minimum Cumulative Precipitation (inches)	Minimum Rotation Interval (months)
State	County or Area				
Colorado	Statewide	Grain sorghum, Proso millet	7.9 or lower	No restrictions	10
	Generally N. of 1-70	Flax, Safflower, Sunflower	7.9 or lower	No restrictions	22
		Field corn	7.9 or lower	15	12
Idaho	Southern Idaho	Flax, Safflower, Sunflower	7.9 or lower	No restrictions	22
	Statewide	Peas Lentils Canola	6.8 or lower	18	10
		Peas	6.9 to 7.9	18	15
		Lentils	6.9 to 7.9	18	34
		Canola	6.9 to 7.9	18	22

Continued on next page

Rotation Intervals For Crops in Non-Irrigated Land (continued)
Following Use of Nufarm Purestand at 1/10 oz per Acre on Wheat, Barley, Fallow or Pasture

Location		Crop	Soil pH	Minimum Cumulative Precipitation (inches)	Minimum Rotation Interval (months)
State	County or Area				
Kansas	Statewide	Grain sorghum, Proso millet	7.9 or lower	No restrictions	10
		Flax, Safflower, Sunflower	7.9 or lower	No restrictions	22
	Central and Western Kansas (West of the Flint Hills)	Field corn	7.9 or lower	15	12
	Western Kansas W. of Hwy. 183	Soybeans	7.5 or lower 7.6-7.9	22 33	22 34
	Central Kansas; generally E. of Hwy. 183 and W. of the Flinthills	Soybeans	7.9 or lower	15	12
Montana	Statewide	Grain sorghum, Proso millet, Field corn	7.9 or lower	22	22
		Alfalfa (hay only)	7.6-7.9	No restrictions	34
			7.5 or lower	No restrictions	22
		Flax, Safflower, Sunflower	7.9 or lower	No restrictions	22
Nebraska	Statewide	Grain sorghum, Proso millet	7.9 or lower	No restrictions	10
		Flax, Safflower, Sunflower	7.9 or lower	No restrictions	22
	Generally W. of Hwy. 77 and E. of the Panhandle	Field corn	7.9 or lower	15	12
		Soybeans	7.5 or lower 7.6-7.9	22 33	22 34
New Mexico	Statewide	Grain sorghum, Proso millet	7.9 or lower	No restrictions	10
		Flax, Safflower, Sunflower	7.9 or lower	No restrictions	22
	Eastern New Mexico	Cotton (dryland only)	7.9 or lower	30	22
North Dakota	W. of Hwy. 1	Grain sorghum, Proso millet, Field corn, Dry beans, Flax, Safflower, Sunflower	7.9 or lower	22	22
	E. of Hwy. 1	Grain sorghum, Proso millet, Field corn, Dry beans, Flax, Safflower, Sunflower	7.9 or lower	34	34

Continued on next page

Rotation Intervals For Crops in Non-Irrigated Land (continued)
Following Use of Nufarm Purestand at 1/10 oz per Acre on Wheat, Barley, Fallow or Pastures

Location		Crop	Soil pH	Minimum Cumulative Precipitation (inches)	Minimum Rotation Interval (months)
State	County or Area				
Oklahoma	Statewide	Grain sorghum, Proso millet	7.9 or lower	No restrictions	10
		Flax, Safflower, Sunflower	7.9 or lower	No restrictions	22
		Field corn	7.9 or lower	15	12
	Panhandle	Cotton (dryland only)	7.9 or lower	30	22
	E. of the Panhandle	Cotton (dryland only)	7.9 or lower	25	14
Oregon	Statewide	Peas Lentils Canola	6.8 or lower	18	10
		Peas	6.9 to 7.9	18	15
		Lentils	6.9 to 7.9	18	34
		Canola	6.9 to 7.9	18	22
South Dakota	Statewide	Flax, Safflower, Sunflower	7.9 or lower	No restrictions	22
	S. of Hwy. 212 & E. of the Missouri River, & S. of Hwy. 34 & W. of Missouri River	Grain sorghum, Proso millet	7.9 or lower	13	12
	Generally E. of Missouri River & S. of Hwy. 14, & W. of Missouri River	Field corn	7.9 or lower	15	12
Texas	Statewide	Grain sorghum, Proso millet	7.9 or lower	No restrictions	10
		Flax, Safflower, Sunflower	7.9 or lower	No restrictions	22
	Panhandle	Field corn	7.9 or lower	15	12
		Cotton (dryland only)	7.9 or lower	30	22
	N. Central Texas*	Field corn	7.9 or lower	15	12
		Cotton (dryland only)	7.9 or lower	25	14
	* The counties of N. Central Texas are: Archer, Baylor, Bell, Bosque, Bowie, Callahan, Camp, Cass, Clay, Collin, Cooke, Coryell, Dallas, Delta, Denton, Eastland, Ellis, Falls, Fannin, Foard, Franklin, Grayson, Hardeman, Haskell, Hill, Hood, Hopkins, Hunt, Jack, Johnson, Kaufman, Knox, Lamar, Limestone, McLennan, Milam, Montague, Morris, Nafarro, Palo Pinto, Parker, Rains, Red River, Robertson, Rockwall, Shackelford, Somervell, Stephens, Tarrant, Throckmorton, Titus, Upshur, Van Zandt, Wilbarger, Wichita, Williamson, Wise.				
Washington	Statewide	Peas Lentils Canola	6.8 or lower	18	10
		Peas	6.9 to 7.9	18	15
		Lentils	6.9 to 7.9	18	34
		Canola	6.9 to 7.9	18	22

Continued on next page

Rotation Intervals For Crops in Non-Irrigated Land (continued)

Following Use of Nufarm Purestand at 1110 oz per Acre on Wheat, Barley, Fallow or Pasture

Location		Crop	Soil pH	Minimum Cumulative Precipitation (inches)	Minimum Rotation Interval (months)
State	County or Area				
Utah	Statewide	Flax, Safflower, Sunflower	7.9 or lower	No restrictions	22
Wyoming	Statewide	Flax, Safflower, Sunflower	7.9 or lower	No restrictions	22
	Southern Wyoming	Grain sorghum,	7.9 or lower	No restrictions	10
	Southern Wyoming (Goshen, Laramie, and Platte counties only)	Field corn	7.9 or lower	15	12
	Northern Wyoming	Grain sorghum, Proso millet, Field corn	7.9 or lower	22	22

Rotation Intervals not covered above - The minimum rotation interval is 34 months with at least 28" of cumulative precipitation during the period:

- to any major field crop not listed (See the Rotation Intervals table)
- if the soil pH is not in the specified range
- if the use rate applied is not specified in the table
- or if the minimum cumulative precipitation has not occurred since application.

To rotate to a major field crop at an interval shorter than recommended, a field bioassay must be successfully completed to that crop. A field bioassay must be successfully completed before rotation to any minor crops (as determined by the USDA criteria). See section on Field Bioassay for further information.

Rotation Intervals in Pasture or Rangeland for Overseeding and Renovation

Location	Crop	Maximum Purestand Rate on Pasture (oz)	Minimum Rotation Interval (months)
AL, AR, FL, GA, KY, LA, MS, NC, OK, SC, TN, TX, VA, WV	Alfalfa, red clover, white clover, sweet clover, bermudagrass, bluegrass, orchardgrass, bromegrass, ryegrass, fescue, timothy	1/10 to 3/10	4
	Wheat (except durum)	1/10 to 3/10	1
	Durum, barley, oat	1/10 to 3/10	10
ALL AREAS NOT INCLUDED ABOVE*	Red clover, white clover, and sweet clover	1/10 to 2/10	12
	Bermudagrass, bluegrass, orchardgrass, bromegrass, ryegrass, timothy	1/10 to 2/10	6
	Fescue	WO to 2/10	18
	Wheat (except durum)	1/10 to 2/10	1
	Durum, barley, oat	1/10 to 2/10	10

Rotation Intervals not covered above - The minimum rotation interval is 34 months with a least 28" of cumulative precipitation during the period :

- to any major field crop or pasture crop not listed (See the Rotation Intervals table)
- if the use rate applied is not specified in the table

To rotate to a major field crop at an interval shorter than recommended, a field bioassay must be successfully completed to that crop. A field bioassay must be successfully completed before rotation to any minor crops (as determined by the USDA criteria). See section on Field Bioassay for further information.

BIOASSAY

A field bioassay must be completed before rotating to any crop not listed (See the Rotation Intervals table), or if the soil pH is not in the specified range, or if the use rate applied is not specified in the table, or if the minimum cumulative precipitation has not occurred since application.

Field Bioassay

To conduct a field bioassay, grow test strips of the crop or crops you plan to grow the following year in fields previously treated with Purestandthis product. Crop response to the bioassay will indicate whether or not to rotate to the crop(s) grown in the test strips.

If a field bioassay is planned, check with your local Agricultural dealer or Nufarm representative for information detailing the field bioassay procedure.

GRAZING

There are no grazing restrictions on Purestandthis product.

IMPORTANT PRECAUTIONS

Treated vegetation may be cut for forage or hay. Coveralls, shoes plus socks must be worn if cutting within 4 hours of treatment.

MIXING INSTRUCTIONS

1. Fill the tank 1/4 to 1/3 full of water (If using liquid nitrogen fertilizer solution in place of water, see Tank Mixtures sections for additional details).
2. While agitating, add the required amount of Purestandthis product.
3. Continue agitation until the Purestandthis product is fully dispersed, at least 5 minutes.
4. Once the Purestandthis product is fully dispersed, maintain agitation and continue filling tank with water. PurestandThis product should be thoroughly mixed with water before adding any other material.
5. As the tank is filling, add tank mix partners (if desired) then add the necessary volume of nonionic surfactant. Always add surfactant last.
6. If the mixture is not continuously agitated, settling will occur. If settling occurs, thoroughly re-agitate before using.
7. Apply Purestandthis product spray mixture within 24 hours of mixing to avoid product degradation.
8. If Purestandthis product and a tank mix partner are to be applied in multiple loads, pre-slurry the Purestandthis product in clean water prior to adding to the tank. This will prevent the tank mix partner from interfering with the dissolution of the Purestandthis product.

Do not use Purestandthis product with spray additives that reduce the pH of the spray solution to below 3.0.

SPRAY EQUIPMENT

For specific application equipment, refer to the manufacturer's recommendations for additional information on GPA, pressure, speed, nozzle types and arrangements, nozzle heights above the target canopy, etc.

Be sure to calibrate air or ground equipment properly before application. Select a spray volume and delivery system that will ensure thorough coverage and a uniform spray pattern with minimum drift. Use higher spray volumes to obtain better coverage when the crop canopy is dense. Avoid

swath overlapping, and shut off spray booms while starting, turning, slowing, or stopping to avoid crop injury.

Do not make applications using equipment and/or spray volumes or under weather conditions that might cause spray to drift onto non-target sites. For additional information on spray drift, refer to the **Spray Drift Management** section of the label.

Continuous agitation is required to keep Purestandthis product in suspension.

SPRAYER CLEANUP

Spray equipment must be cleaned before Purestandthis product is sprayed. Follow the cleanup procedures specified on the labels of previously applied products. If no directions are provided, follow the six steps outlined in **After Spraying Purestand** section of this label.

At the End of the Day

When multiple loads of Purestandthis product herbicide are applied, it is recommended that at the end of each day of spraying, the interior of the tank be rinsed with fresh water and then partially filled, and the boom and hoses flushed. This will prevent the buildup of dried pesticide deposits that can accumulate in the application equipment.

After Spraying Purestandthis product and Before Spraying Crops Other Than Wheat, Barley, Fallow, Pasture, or Rangeland

To avoid subsequent injury to desirable crops, thoroughly clean all mixing and spray equipment immediately following applications of Purestandthis product as follows:

1. Drain tank; thoroughly rinse spray tanks, boom, and hoses with clean water. Loosen and physically remove any visible deposits.
2. Fill the tank with clean water and 1 gal of household ammonia* (contains 3% active) for every 100 gal of water. Flush the hoses, boom, and nozzles with the cleaning solution. Then add more water to completely fill the tank. Circulate the cleaning solution through the tank and hoses for at least 15 min. Flush the hoses, boom, and nozzles again with the cleaning solution, and then drain the tank.
3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
4. Repeat step 2.
5. Rinse the tank, boom, and hoses with clean water.
6. If only Ammonia is used as a cleaner, the rinsate solution may be applied back to the crop(s) recommended on this label. Do not exceed the maximum labeled use rate. If other cleaners are used, consult the cleaner label for rinsate disposal instructions. If no instructions are given dispose of the rinsate on site or at an approved waste disposal facility.

* Equivalent amounts of an alternate-strength ammonia solution or a Nufarm-approved cleaner can be used in the cleanout procedure. Carefully read and follow the individual cleaner instructions. Consult your agricultural dealer, applicator, or Nufarm representative for a listing of approved cleaners.

Notes:

1. **Attention:** Do not use chlorine bleach with ammonia, as dangerous gases will form. Do not clean equipment in an enclosed area.
2. Steam-cleaning aerial spray tanks is recommended prior to performing the above cleanout procedure to facilitate the removal of any caked deposits.
3. When ~~Nufarm~~ Purestandthis product is tank mixed with other pesticides, all required cleanout procedures should be examined and the most rigorous procedure should be followed.
4. In addition to this cleanout procedure, all precleanout guidelines on subsequently applied products

220f26

should be followed as per the individual labels.

- 5. Where routine spraying practices include shared equipment frequently being switched between applications of Purestandthis product and applications of other pesticides to Purestandthis product - sensitive crops during the same spray season, it is recommended that a sprayer be dedicated to Purestandthis product to further reduce the chance of crop injury.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

IMPOR TANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets (>150 - 200 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!** See **Wind, Temperature and Humidity, and Temperature Inversions** sections of this label.

Controlling Droplet Size – General Techniques

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.**
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

Controlling Droplet Size - Aircraft

- **Number of Nozzles** - Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations.
- **Nozzle Type** - Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.
- **Boom Length** - The boom length should not exceed 3/4 of the wing or rotor length - longer booms increase drift potential.
- **Application Height** - Application more than 10 ft above the canopy increases the potential for spray drift.

BOOM HEIGHT

Setting the boom at the lowest labeled height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.

WIND

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. **AVOID GUSTY OR WINDLESS CONDITIONS.**

Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can redact the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

AIR ASSISTED (AIR BLAST) FIELD CROP SPRAYERS

Air assisted field crop sprayers carry droplets to the target via a downward directed air stream. Some may reduce the potential for drift, but if a sprayer is unsuitable for the application and/or set up improperly, high drift potential can result. It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application, is configured properly, and that drift is not occurring.

Note: Air assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration. Consult the application equipment section of this label to determine if use of an air assisted sprayer is recommended.

WEED RESISTANCE

Biotypes of certain weeds listed on this label are resistant to ~~Nufarm Purestand~~this product and other herbicides with the same mode of action*, even at exaggerated application rates. Biotypes are naturally occurring individuals of a species that are identical in appearance but have slightly different genetic compositions; the mode of action of an herbicide is the chemical interaction that interrupts a biological process necessary for plant growth and development.

If weed control is unsatisfactory, it may be necessary to retreat problem areas using a product with a different mode of action, such as postemergence broadleaf and/or grass herbicides.

If resistant weed biotypes such as kochia, prickly lettuce, and Russian thistle are suspected or known to be present use a tank-mix partner with ~~Purestand~~this product to help control these biotypes, or use a planned herbicide rotation program where other residual broadleaf herbicides having different modes of action are used.

* Naturally occurring weed biotypes that are resistant to ALS inhibitor herbicides (such as "Amber" herbicide, "Pursuit" herbicide, FINESSE® herbicide, or HARMONY® EXTRA herbicide) may also be resistant to ~~Purestand~~this product.

INTEGRATED PEST MANAGEMENT

To better manage weed resistance when using ~~Purestand~~this product, use a combination of tillage, and tank-mix partners or sequential herbicide applications that have a different mode of action than ~~Purestand~~this product, to control escaped weeds. Do not let weed escapes go to seed.

Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative herbicide recommendations available in your area.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes.

PRECAUTIONS

- Injury to or loss of desirable trees or vegetation may result from failure to observe the following:

-Do not apply, drain, or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.

-Do not use on lawns, walks, driveways, tennis courts, golf courses, athletic fields, commercial sod operations, or other high-maintenance, fine turfgrass areas, or similar areas.

-Do not use on grasses grown for seed.

- Do not apply to irrigated land where tailwater will be used to irrigate crops other than wheat and barley.
- Do not apply to frozen ground as surface runoff may occur.
- Do not apply to snow-covered ground.
- Wheat and barley varieties may differ in their response to various herbicides. Nufarm recommends that you first consult your state experiment station, university, or extension agent as to sensitivity to any herbicide. If no information is available, limit the initial use of Purestandthis product to a small area.
- Under certain conditions such as heavy rainfall, prolonged cold weather, or wide fluctuations in day/night temperatures prior to or soon after Purestandthis product application, temporary discoloration and/or crop injury may occur. Purestandthis product should not be applied to wheat or barley that is stressed by severe weather conditions, drought, low fertility, water-saturated soil, disease, or insect damage, as crop injury may result. Risk of injury is greatest when crop is in the 2 to 5- leaf stage. Severe winter stress, drought, disease, or insect damage following application also may result in crop injury.
- The combined treatment effects of Purestandthis product postemergence preceded by preemergence wild oat herbicides may cause crop injury to spring wheat when crop stress (soil crusting, planting too deep, prolonged cold weather, or drought) causes poor seedling vigor.
- In the Pacific Northwest, to prevent cold weather-related crop injury, avoid making applications during winter months when weather conditions are unpredictable and can be severe.
- Do not apply to wheat, barley or pastures undersown with legumes, as injury to the forage may result.
- To reduce the potential for movement of treated soil due to wind erosion, do not apply to powdery dry or light sandy soils until they have been stabilized by rainfall, trashy mulch, reduced tillage, or other cultural practices. Injury to immediately adjacent crops may occur when treated soil is blown onto land used to produce crops other than cereal grains or pasture/rangeland.
- For ground applications applied to weeds when dry, dusty field conditions exist, control of weeds in wheel track areas may be reduced. The addition of 2,4-D or MCPA should improve weed control under these conditions.
- Preplant or preemergence applications of 2,4-D or herbicides containing 2,4-D made within 2 weeks of planting spring cereals may cause crop injury when used in conjunction with early postemergence applications of Nufarm Purestandthis product. For increased crop safety, delay Purestandthis product treatment until crop tillering has begun.

STORAGE AND DISPOSAL

Storage: Store product in original container only. Do not contaminate water, other pesticides, fertilizer, food or feed in storage. Store in a cool, dry place.

Product Disposal: Do not contaminate water, food, or feed by disposal. Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: For Plastic Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. **For Fiber Sacks:** Completely empty fiber sack by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into manufacturing or application equipment. Then dispose of sack in a sanitary landfill or by incineration if allowed by State and local authorities. **For Fiber Drums With Liners:** Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by State and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner. **For Bags Containing Water Soluble Packets:** Do not reuse the outer box or the resealable plastic bag. When all water-soluble packets are used, the outer packaging should be clean and may be disposed of in a sanitary landfill or by incineration, or if allowed by State and local authorities, by open burning. If burned, stay out of smoke. If the resealable plastic bag contacts the formulated product in any way, the bag must be triple-rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer wrap as described above. **For Metal Containers (non aerosol):** Triple rinse (or equivalent) the container. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. **For Paper and Plastic Bags:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

EXPRESS®, HARMONY® and FINESSE® are trademarks or registered trademarks of E. I. duPont de Nemours and Company

"Raindrop RA" is a trademark of Dclavan Corp. "Amber" is a trademark of Novartis Crop Protection Inc.

"Assert", "Avenge", "Banvel", "Pursuit", "Weedmaster" are trademarks of BASF Corp.

"Grazon P+D", "Remedy", "Tordon" are trademarks of Dow AgroSciences

"Buctril", "Bronate", "Hoelon" are trademarks of Aventis CropScience "Di-Syston" is a trademark of Bayer Crop Protection

LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read This Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of Nufarm. These risks can cause: ineffectiveness of the product; crop injury, or; injury to non-target crops or plants.

Nufarm does not agree to be an insurer of these risks. **WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.**

Nufarm warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

NUFARM MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO EVENT SHALL NUFARM OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF NUFARM OR SELLER, FOR ANY

AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF NUFARM OR SELLER, THE REPLACEMENT OF THE PRODUCT.

Nufarm or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify Nufarm or a Nufarm Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

NOTIFICATION

JAN 15 2009