



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (7505C)  
401 "M" St., S.W.  
Washington, D.C. 20460

EPA Reg.  
Number:

71368-50

Date of Issuance: JUN 30 2004

Term of Issuance:

Conditional

Name of Pesticide Product:

Weedhawk Dry Flowable  
Herbicide

NOTICE OF PESTICIDE:

Registration  
 Reregistration  
(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Nufarm, Inc.  
1333 Burr Ridge Parkway, Suite 125A  
Burr Ridge, IL 60521-0866

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit data.
2. Make the following label changes listed below before you release the product for shipment:
  - a. Add the phrase, "EPA Reg. No.71368-50".
  - b. Please note that 40 CFR 156.10(i)(1)(ii) states that only the Directions For Use may appear on printed or graphic matter which accompanies the pesticide. The referral statement "See Inside For Additional Precautionary Statements" implies that the precautionary statements are inside a booklet. If all of the Precautionary Statements and First Aid Statements cannot appear on the outside container label due to size constraints, you should formally submit a request for a size exemption from the regulations.

Signature of Approving Official:

/s/

Date:

JUN 30 2004

page 2

EPA Reg. No. 71368-50

c. In the Recommended Tank Mixes Table specify the Weedhawk application rate for each crop in the tank mix. Specify a Buctril application rate or delete this column. Assure that the crop and application rate is identical to a currently registered product.

3. Submit one (1) copy of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

4. Submit the results of storage stability (GRN 830.6317) and corrosion characteristics (GRN 830.6320) studies to EPA upon completion.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Joanne I. Miller  
Product Manager (23)  
Herbicide Branch  
Registration Division (7505C)

3/13

# WEEDHAWK

## DRY FLOWABLE HERBICIDE

### ACTIVE INGREDIENT:

\*Dimethylamine salt of 2,4-Dichlorophenoxyacetic Acid ..... 7.50%

INERT INGREDIENTS ..... 92.50%

TOTAL ..... 100.00%

\*2,4-Dichlorophenoxyacetic acid equivalent 6.2% by weight.

EPA Reg. No.

EPA Est. No.

### CAUTION

### KEEP OUT OF REACH OF CHILDREN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

#### FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

NOTICE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact: 1-877-325-1840/1840 for emergency medical treatment information.

SEE INSIDE FOR ADDITIONAL PRECAUTIONARY STATEMENTS

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if absorbed through skin. Causes moderate eye damage. Avoid contact with skin, eyes or clothing.

### PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- \* Coveralls over long-sleeved shirt and long pants.
- \* Chemical-resistant (such as nitrile or butyl) gloves.
- \* Chemical-resistant footwear plus socks.
- \* Protective eyewear.
- \* Chemical-resistant apron when mixing or loading.
- \* For exposures outdoors, a dust/mist filtering respirator (MSHA/NIOH approval number prefix TC-21C).

Net Contents: 8 X 9 oz. packets.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated  
JUN 30 2004

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

71368-50

4/13

- \* During aerial application, human flag-waver must be in enclosed cabs.
- \* Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacture's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and Wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

#### ENVIRONMENT HAZARD STATEMENTS

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water marks. Do not contaminate water when disposing of equipment wash waters. Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides as such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

#### PHYSICAL OR CHEMICAL HAZARDS

Combustible. Do not use or store near heat or open flame.

5/13

**ENGINEERING CONTROL STATEMENTS**

When handlers use enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR part 170.240 (d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

**USER SAFETY RECOMMENDATIONS**

**USER SHOULD:** Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.  
Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handler may be in the area during application.  
For requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170.

This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouse, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by Worker Protection Standard.

Do not enter or allow workers entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- \* Coveralls over long-sleeved shirt and long pants.
- \* Chemical-resistant (such as nitrile or butyl) gloves.
- \* Chemical-resistant footwear plus socks.
- \* Protective eyewear.

**STORAGE AND DISPOSAL**

**STORAGE:** Do not store near heat or open flame. Keep upright in a cool place in original container.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposal of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose if in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**IN CASE OF SPILLS:** Recover all material practical for use. Soak up balance with sand, dirt or absorbent, and haul to approved land fill, check local regulations.

6/13

**Active Ingredient 2,4-D acid per volume of Weedhawk:**

Weedhawk	Amount of 2,4-dichlorophenoxyacetic acid
9 oz (dry)	0.035 lb.
18 oz (dry)	0.07 lb.
36 oz (dry)	0.14 lb.

This product is recommended for use in most geographic locations of the United States, regardless of humidity or lack of rain fall and at elevations. It is recommended for use on grapes, pome fruits (apples and pears), nut-orchard floors, small grains (barley, oats, rye and wheat), stone fruits, sugarcane, and certain non-crop uses (golf courses, parks, turfgrass, railroads, and drainage ditches). Use appropriate spray nozzles to obtain the best coverage with minimum distance from target weeds. Apply to weed growth so that coverage is uniform and complete, but not to the point of run-off. Do not apply during rain. Apply when conditions allow spray to dry on foliage.

**PRECAUTIONS FOR VULNERABLE CROPS**

This product should not be used on vulnerable crops such as cotton and pomegranates. Please be reminded that this product contains 2,4-D and should be used only as recommended. It may prove harmful when misused. Avoid spraying near crops, plants, or trees if wind moves at or above 5 mph, or in their direction. Avoid herbicide spraying if crops and planting are under stress conditions. **Weedhawk tank mix** is for use in State where approved for spraying. Consult local and state authorities concerning restrictions.

**GENERAL RESTRICTIONS AND LIMITATIONS**

- Do not apply more than 36 (dry) ounces per season.
- Do not apply to crops listed on this label that have been subjected to stress conditions such as drought, flooding, extreme diseases, nematodes, insects, where crops injury could result. Likewise with non-actively thriving weeds, control may be reduced.
- Do not apply if rain is threatening. Rainfall soon after application can reduce effectiveness.
- Do not apply through any type of irrigation system.
- Do not use treated plants for livestock feed.
- Do not use tank mixes other than listed on a label or technical bulletins. After mixing, spray within 8 hours. Do not cultivate or disturb sprayed ground for at least 7 days after spray application. Avoid applying where spray can reach young tender growth.

**MIXING**

Fill mixing tank 1/2 full with clean water. Drop entire water soluble packet into tank and begin agitation. Allow packets to dissolve. If tank mixing add partner herbicide while continuing agitation Allow to thoroughly mix. Add remaining water and continue aggitation. Add anti-foaming agent if desired. One to two 9 oz packets of Weedhawk per acre is reccomended. (see application rates and timing.)

7/13

**Notice:** To avoid subsequent injury to other crops, clean the sprayer before and after application.  
**Cleaning the Sprayer:** Fill with clean water. Add commercial spray tank cleaner, as recommended on its label. Circulate and clean all parts, drain the tank and rinse total system as many times as needed to be thoroughly clean. Do not contaminate water when disposing of equipment wash waters.

**GENERAL WEEDS LIST:**

**Weedhawk** herbicide combined with other herbicides gives control or growth suppression on following:

**Broadleaf Weeds:**

African Daisy	Hairy fleabane	Mustard	Puncturevine
Carrot, wild	Lambquarters	Nettles	Spurge
Common groundsel	Lettuce, wild	Nightshade, hairy	Sowthistle
Fiddleneck	Malva	Nutsedge	Thistle, bull
Filaree	Marestail	Pigweed	Thistle, russian
Kochia	Morning-glory, annual	Prickly. lettuce	Tumbleweed

**APPLICATIONS**

**Weedhawk** herbicide should be applied at a rate of 9 - 18 (dry) oz. per acre when weeds are small (e.g., 2- leaf stage) and growing actively before the bud stage. Do not exceed 36 (dry) oz. / acre / season. Apply at the rate of 5 to 100 gallons of water per acre application and a minimum of 20 to 30 psi pressures for adequate coverage. Spray when weeds are actively growing.  
 • Always read and follow all label directions for tank mixes. Refer to individual use sections for additional Precautions, Restrictions, Application rates and timings.

**GRAPES:**

**Weedhawk** is effective at 9-18 dry.oz. of product per acre per application. (Equivalent 0.035-0.070 pounds of acid/acre). Make one directed application. Grapes are highly sensitive to 2,4-D. Do not allow spray to contact fruit, foliage, exposed roots, stems or lower limbs of vines. Do not spray bare ground. Use

8/13

**POME FRUITS (APPLE AND PEAR):**

Weedhawk is effective at 9-18 dry oz. of product per acre per application. (Equivalent 0.035-0.070 pounds of acid /acre) as a broadcast treatment to the orchard floor. Do not allow spray to contact fruit, foliage, exposed roots, stems or lower limbs of trees. Do not spray bare ground. Use flat, fan-type nozzles and low pressures. Do not apply during periods of extremely high temperatures. Apply after irrigation and allow the maximum time before the next irrigation (at least three days). Do not apply to newly established or young orchards. Trees should be at least one (1) year or older. Avoid applying where spray can reach bloom. A maximum of two (2) broadcast application may be made per crop. Allow at least 75 days between applications. Do not harvest within forty (40) days of application. Do not graze or feed cover crops from treated orchards to livestock.

**NUT ORCHARD FLOORS:**

Weedhawk is effective at 9 -18 dry oz. of product per acre per application. (Equivalent 0.035-0.070 pounds of acid /acre) as a broadcast treatment to the orchard floor. Do not allow spray to contact fruit, foliage, exposed roots, stems or lower limbs of trees. Do not spray bare ground. Use flat, fan-type nozzles and low pressures. Do not apply during periods of extremely high temperatures. Apply after irrigation and allow the maximum time before the next irrigation (at least three days). Do not apply to newly established or young orchards. Trees should be at least one (1) year or older. Avoid applying where spray can reach bloom. A maximum of two (2) broadcast application may be made per crop. Do not harvest these tree nuts within sixty (60) days of applications. Do not graze or feed cover crops from treated orchards to livestock.

**SMALL GRAINS (Barley, Oats, Rye and Wheat):**

Weedhawk is effective at 9-18 dry oz. of product per acre per application. (Equivalent 0.035-0.070 pounds of acid /acre) as a broadcast treatment to the grains. Apply after crop begins tillering and before the boot stage. Do not apply to grains underseeded with legumes. Do not apply before tiller stage or from early boot to milk stage. Do not spray grain in the boot to dough stage. Do not graze or feed forage from treated fields within fourteen (14) days after treatment. A maximum of two (2) application may be made per crop. Do not harvest these grain within seven (7) days of application.

**STONE FRUITS:**

Weedhawk is effective at 9-18 dry oz. of product per acre per application. (Equivalent 0.035-0.070 pounds of acid /acre) as a broadcast treatment to the orchard floor. Do not allow spray to contact fruit, foliage, exposed roots, stems or lower limbs of trees. Do not spray bare ground. Use flat, fan-type nozzles and low pressures. Do not apply during periods of extremely high temperatures. Apply after irrigation and allow the maximum time before the next irrigation (at least three days). Do not apply to newly established or young orchards. Trees should be at least one (1) year or older. Avoid applying where spray can reach bloom. A maximum of two (2) broadcast application may be made per crop. Do not harvest these tree nuts within forty (40) days of applications. Do not graze or feed cover crops from treated orchards to livestock.

**SUGARCANE:**

Weedhawk is effective at 9-18 dry oz. of product per acre per application. (Equivalent 0.035-0.070 pounds of acid /acre) as a broadcast treatment to the sugarcane. Apply through layby. A maximum of two (2) application may be made per crop. Do not harvest these sugarcane within seven (7) days of application.



9/13

**Corn: (Field and Sweet)**

Weedhawk is effective at 9-18 dry ounces of product per acre per application. (Equivalent 0.035-0.070 pounds of acid / acre) as a broadcast treatment to corn. Apply after crop begins tillering and before the boot stage. Do not apply to corn underseeded with legumes. Corn is susceptible to injury at time of emergence and shortly after unfolding of leaves. Do not spray during this period nor after first tassels appear. Do not spray corn in tassel. 2,4-D may cause brittleness to corn. Certain single cross corn hybrids may be more susceptible to 2,4-D injury than others. Do not graze or feed forage from treated fields within 14 days after treatment. A maximum of 2 applications may be made per crop. Do not harvest corn within 7 days following application.

**Postemergence: (Annual broadleaf weeds and Perennial broadleaf weeds.)**

Weedhawk is effective at 9-18 dry ounces / acre per application. (Equivalent 0.035-0.070 pounds of acid / acre) as a broadcast treatment to corn. Apply when most weeds have germinated. Corn is susceptible to injury at time of emergence and shortly after unfolding of leaves. Do not spray during this period nor after first tassels appear. Use drop nozzles when corn is 10 inches tall to place spray below its tops. Do not cultivate soon after spraying as corn may be brittle. Do not forage or feed corn fodder for seven days following application.

**Grass Seed Crops**

Weedhawk is effective at 9-18 dry ounces per acre per application. (Equivalent 0.035-0.070 pounds of acid / acre) as a broadcast treatment to the grasses. Apply after crop begins tillering and before the boot stage. Do not apply to grasses undreseeded with legumes. Do not apply before tiller stage or from early boot to milk stage. Do not spray grain in the boot to dough stage. Do not graze or feed forage from treated fields within 14 days after treatment. A maximum of 2 application may be made per crop. Do not harvest these grasses within 7 days of application. Do not graze dairy cattle for at least 7 days after application. Do not allow meat animals to enter treated fields for at least 3 days prior to slaughter. Do not cut grass for hay for at least 30 days after application.

**Sorghum**

Weedhawk is effective at 9-18 dry ounces per acre per application. (Equivalent 0.035-0.070 pounds of acid / acre) as a broadcast treatment to sorghum. Apply after crop begins tillering and before the boot stage. Do not apply to sorghum underseeded with legumes. Do not apply before tiller stage or from early boot to milk stage. Do not graze or feed forage from treated fields within 14 days after treatment. A maximum of 2 applications may be made per crop. Do not harvest sorghum within 7 days of application.

**Pistachio Orchards**

Weedhawk is effective at 9-18 dry ounces per acre per application. (Equivalent 0.035-0.070 pounds of acid / acre) as a broadcast treatment to the orchard floor. Do not allow spray to contact fruit, foliage, exposed roots, stems or lower limbs. of trees. Do not spray bare ground. Use flat, fan type nozzles and low pressure. Do not apply during periods of extremely high temperature. Apply after irrigation and allow the maximum time before the next irrigation (at least 3 days). Do not apply to newly established or young orchards. Trees should be at least 1 year or older. Avoid applying where spray can reach bloom. A maximum of 2 applications may be made per crop. Do not harvest these tree nuts within 60 days of application. Do not graze or feed cover crops from treated orchards to livestock.

10/13

**RECOMMENDED TANK MIXES**

- Always read and follow all label directions for tank mixes in combination with other herbicides. Refer to individual herbicide use sections for additional Precautions, Restrictions, and Application rates.
- **Weedhawk for weed control in herbicide combination with the following crops:**

Crops	Weedhawk dry.oz/acre	Credit Systemic lbs per acre	Buctril
<b>Pome fruits:</b> Apple, and Pear	9-18	1.0-3.0 lbs	
<b>Nut orchard floors</b>		1.0-3.0 lbs	
<b>Rye</b>		1.0-3.0 lbs	
<b>Stone fruits</b>	9-18	1.0-3.0 lbs	
<b>Pistachi Orchards</b>		1.0-3.0 lbs	

**NOTE**  
This product can be mixed with Nufarm Credit Herbicide and Buctril for use on the crops listed in this table, in accordance with the most restrictive label limitations and precautions. No label dosage rate should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

**FOR FALLOW LAND AND STUBBLE**

Apply Weedhawk to annual weeds at the rate of 9-18 dry ounces of product per acre per application. Apply Weedhawk to perennial weeds at the rate of 18- 36 dry ounces of product per acre per application. Apply Weedhawk when weeds are actively growing. Do not apply more than 36 dry ounces per acre per season. Weedhawk can be applied as a tank mix with Glyphosate and Banvel to broaden the spectrum of weed control. In order to assure maximum safety and weed control, follow all precautions and limitations on the Weedhawk label and on the labels of products used in tank mixtures with Weedhawk. Do not make applications within 90 days of planting, or until chemical has disappeared from soil.

Weedhawk 9-18 oz/acre	Glyphosate lbs/ acre 1/4 to 1 pound	Banvel Pints/Acre
9-18 oz/acre		1-2 pints

11/13

### RECOMMENDED TANK MIXTURE FOR PASTURE AND RANGELAND

Weedhawk can be applied as a tank mix with Banvel and Tordon to broaden the spectrum of weed control. In order to assure maximum safety and weed control, follow all precautions and limitations on the Weedhawk label and the labels of products used in tank mixtures with Weedhawk. Apply the amounts listed below in 5-100 gallons of water per acre as a broadcast treatment.

Weedhawk dry ounces per acre	Banvel pint/acre	Tordon pint/acre
9-18	1-2 pints	
9-18		1/2 to 2 pints

#### Pasture and Rangeland Restrictions and Limitations.

Do not allow dairy animals to graze on treated areas for at least 7 days after treatment.

Do not cut forage or graze treated fields within 30 days after treatment.

Do not allow livestock to enter treated fields within 3 days of slaughter.

12/13

**FORESTRY USE (FOREST SITE PREPARATION)**

Use 9.18 dry ounces of product in 5 to 100 gallons of water per acre. Dosage rates per acre depend on the density of brush and/or weeds. For small broadleaf weeds, use the lower rate. Heavy dense stands of brush require the higher rate with higher water volume. To effectively control brush, all leaves, stems, and suckers should be thoroughly wetted to the ground. Apply when plants come into full leaf (spring) to the time plants begin to go dormant. Best results are obtained when brush and broadleaf weeds are young and actively growing. Do not cut brush until the herbicide has translocated throughout the plant causing root death. Do not apply as a stand release or cover spray to established conifers as injury may result.

**RECOMMENDED TANK MIXES FOR FOREST SITE PREPARATION**

- Always read and follow all label directions for tank mixes in combination with other herbicides. Refer to individual herbicide use sections for additional Precautions, Restrictions, and Application rates.
- **Weedhawk for weed control in herbicide combination with the following:**

Application	Weedhawk dry.oz/acre	Glyphosate lbs/acre	Water
Aerial	dry.oz/acre	1 to 3	5 to 25 gal/acre
Ground	dry oz/acre	1 to 3	10 to 100 gal/acre

**AERIAL SPRAY (FOR FORESTRY USE ONLY)**

Use 5 to 25 gallons of spray solution per acre as a broad cast spray. WHILE SPRAYING, A BUFFER ZONE MUST BE MAINTAINED TO PROTECT UNINTENDED VEGETATION: VARIOUS CONDITIONS, INCLUDING WINDY CONDITIONS AND AIR INVERSIONS, MAY CAUSE DRIFT AND WILL CAUSE DAMAGE TO ANY CONTACTED VEGETATION. USE STRAIGHT STREAM NOZZLES DIRECTED STRAIGHT BACK AND USE A SPRAY BOOM NO LONGER THAN ¾ THE WINGSPAN OF THE AIRCRAFT.

**SPOT TREATMENT SPRAY DIRECTIONS**

- Weedhawk may be used as a spot spray application to control weeds in golf courses, parks, turfgrass, railroads, and drainage ditches. Spray to wet target weeds thoroughly.
- Control spray drift using nozzles that provide the largest size droplet consistent with adequate and uniform coverage. When applying as a spot spray use sufficient material to obtain thorough coverage of weeds.
- Always read and follow all label directions for tank mixes in combination with other herbicides. Refer to individual herbicide use sections for additional Precautions, Restrictions and Application rates.

**FOR AERIAL APPLICATION**

For applications to Small Grains, Pasture, Rangeland and Sugarcane. Make sure the aerial equipment for application provides a maximum spray distribution and maximum coverage. Recommend a minimum spray volume of 5 GPA and a maximum pressure of 40 PSI per acre. To avoid off target movement, increase spray volume per acre, do not apply in winds above 10 m.p.h.

13/13

**SPECIFICATIONS FOR WARRANTY AND SALE**

There is no warranty expressed or implied, concerning the use of this product other than indicated on the label. The buyer assumes all risk of liability for any and all claims, losses, damages, or injuries resulting from the use, handling and storage that are contrary to label instructions. The Directions for Use of this product are believed to be reliable and should be followed carefully. All such risks will be assumed by Buyer and User, and Buyer and User agree to hold Seller harmless for any claims relating to such factors. Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label, subject to the inherent risks referred to above, when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller and Buyer and Users assume the risk of any such use.