

(A) <b>EPA</b>	United States Environmental Protection Agency Office of Pesticide Programs (H7505C) Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number  Please Attach An Original Form
----------------	--	--	--

**Application for Pesticide:**

Section I

1. Company/Product Number 71368-50	2. EPA Product Manager J. Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Nufarm Weedhawk Dry Flowable Herbicide	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Nufarm Inc 1333 Burr Ridge Parkway, Suite 125A Burr ridge, IL 60527 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to:  EPA Reg. No. _____ Product Name _____	

Section II

Amendment - Explain Below  Final printed labels in response to Agency letter dated NOTIFICATION  
 Resubmission in response to Agency letter dated \_\_\_\_\_  "Me Too" Application NOV 17 2004  
 Notification - Explain below  Other - explain below

Explanation: Use additional page(s) if necessary. (For Section I and Section II).  
Updated label. Language left off final printed label.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Sections 12 and 14 of FIFRA.

Section III

1. Material This Product Will Be Packaged In:

Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No If "Yes," Unit Package Wt. No. Per Container	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No If "Yes," Package Wt. No. Per Container	2. Type of Container <input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
--	---	---	---

\*Certification must be submitted.

3. Location of Net Contents Information <input checked="" type="checkbox"/> Label or <input checked="" type="checkbox"/> Container	4. Size(s) of Retail Container 1 qt.; 1/2, 1, 2 1/2, 5, 30, 55, 110, 220 gal.; bulk	5. Location of Label Directions <input checked="" type="checkbox"/> On Label or <input checked="" type="checkbox"/> On Labeling accompanying product
---	--	--

6. Manner in Which Label Is Affixed To Product  Lithograph  Other ( Self-Adhesive Integrated Label/Booklet )  
 Paper Glued  
 Stenciled

Section IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Theodore D. Head	Title Product Registration Manager / Agriculture	Telephone No. (Include Area Code) (708) 754-3330
--------------------------	---	---

**Certification**

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

Signature: *Theodore D. Head*

3. Title Product Registration Manager / Agriculture

4. Typed Name Theodore D. Head

5. Date 10-28-04

Date Application Received (Stamp)

2/9

NOTIFICATION

NOV 17 2004

# Nufarm

# WEEDHAWK™

## DRY FLOWABLE HERBICIDE

**ACTIVE INGREDIENT:**

*Dimethylamine salt of 2,4-Dichlorophenoxyacetic Acid .....	7.50%
---	-------

**OTHER INGREDIENTS:**

92.50%

<b>TOTAL:</b> .....	100.00%
---------------------	---------

\*2,4-Dichlorophenoxyacetic acid equivalent 6.2% by weight.

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

PRECAUCION AL USUARIO: Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
 (If you do not understand the label, find someone to explain it to you in detail.)

**SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS**

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300.  
 For Medical Emergencies Only, Call (877) 325-1840.

EPA REG. NO. 71368-50  
 EPA EST. NO. 228-JL-1

MANUFACTURED BY  
 NUFARM INC.  
 BURR RIDGE, IL 60527-0866



NET CONTENTS 8 x 9 OUNCE PACKETS

071368-00050.092804.WDFH

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION - CAUCION**

Harmful if absorbed through skin. Causes moderate eye damage. Avoid contact with skin, eyes or clothing.

**PERSONAL PROTECTIVE EQUIPMENT (PPE):**

**Applicators and other handlers must wear:** Coverall over long-sleeved shirt and long pants, chemical-resistant (such as nitrile or butyl) gloves, chemical-resistant footwear plus socks, protective eyewear, chemical-resistant apron when mixing or loading, and for exposures outdoors, a dust/mist filtering respirator (MSHA/NIOH approval number prefix TC-21C). During aerial application, human flag-waver must be in enclosed cabs.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacture's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and Wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

**ENGINEERING CONTROL STATEMENTS**

When handlers use enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR part 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

<b>USER SAFETY RECOMMENDATIONS</b>	
<b>Users Should:</b>	
<ul style="list-style-type: none"> <li>• Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.</li> <li>• Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.</li> <li>• Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.</li> </ul>	

<b>FIRST AID</b>	
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Call poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF INHALED</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>HOT LINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	
<b>NOTE TO PHYSICIAN</b>	
Probable mucosal damage may contraindicate the use of gastric lavage.	

**ENVIRONMENTAL HAZARDS**

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water, to areas where surface waters are present or to intertidal areas below the mean high water marks. Do not contaminate water when disposing of equipment wash waters. Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides as such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

**PHYSICAL OR CHEMICAL HAZARDS**

Combustible. Do not use or store near heat or open flame.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls over long-sleeved shirt and long pants, chemical-resistant (such as nitrile or butyl) gloves, chemical-resistant footwear plus socks and protective eyewear.

**GENERAL RESTRICTIONS AND LIMITATIONS**

- Do not apply more than 36 (dry) ounces per season.
- Do not apply to crops listed on this label that have been subjected to stress conditions such as drought, flooding, extreme diseases, nematodes, insects, where crops injury could result. Likewise with non-actively thriving weeds, control may be reduced.
- Do not apply if rain is threatening. Rainfall soon after application can reduce effectiveness.
- Do not apply through any type of irrigation system.
- Do not use treated plants for livestock feed.
- Do not use tank mixes other than listed on a label or technical bulletins. After mixing, spray within 8 hours. Do not cultivate or disturb sprayed ground for at least 7 days after spray application. Avoid applying where spray can reach young tender growth.

**MIXING**

Fill mixing tank 1/2 full with clean water. Drop entire water soluble packet into tank and begin agitation. Allow packets to dissolve. If tank mixing add partner herbicide while continuing agitation. Allow to thoroughly mix. Add remaining water and continue agitation. Add anti-foaming agent if desired. One to two 9 ounce packets of this product per acre is recommended. (See application rates and timing.)

**Notice:** To avoid subsequent injury to other crops, clean the sprayer before and after application.

**Cleaning the Sprayer:** Fill with clean water. Add commercial spray tank cleaner, as recommended on its label. Circulate and clean all parts, drain the tank and rinse total system as many times as needed to be thoroughly clean. Do not contaminate water when disposing of equipment wash waters.

**ACTIVE INGREDIENT 2,4-D ACID PER VOLUME OF THIS PRODUCT**

THIS PRODUCT	AMOUNT OF 2,4-DICHLOROPHENOXYACETIC ACID
9 ounces (dry)	0.035 pound
18 ounces (dry)	0.07 pound
36 ounces (dry)	0.14 pound

This product is recommended for use in most geographic locations of the United States, regardless of humidity or lack of rain fall and at elevations. It is recommended for use on grapes, pome fruits (apples and pears), nut-orchard floors, small grains (barley, oats, rye and wheat), stone fruits, sugarcane, and certain non-crop uses (golf courses, parks, turfgrass, railroads, and drainage ditches). Use appropriate spray nozzles to obtain the best coverage with minimum distance from target weeds. Apply to weed growth so that coverage is uniform and complete, but not to the point of run-off. Do not apply during rain. Apply when conditions allow spray to dry on foliage.

**PRECAUTIONS FOR VULNERABLE CROPS**

This product should not be used on vulnerable crops such as cotton and pomegranates. Please be reminded that this product contains 2,4-D and should be used only as recommended. It may prove harmful when misused. Avoid spraying near crops, plants, or trees if wind moves at or above 5 mph, or in their direction. Avoid herbicide spraying if crops and planting are under stress conditions. This product tank mix is for use in States where approved for spraying. Consult local and State authorities concerning restrictions.

**GENERAL WEED LIST**

This product combined with other herbicides gives control or growth suppression on the following:

**Broadleaf Weeds:**

African Daisy	Kochia	Marestail	Nutsedge	Sowthistle
Carrot, wild	Hairy fleabane	Morning-glory, annual	Pigweed	Thistle, bull
Common groundsel	Lambquarters	Mustard	Prickly, lettuce	Thistle, Russian
Fiddleneck	Lettuce, wild	Nettles	Puncturevine	Turnleweed
Filaree	Malva	Nightshade, hairy	Spurge	

**APPLICATIONS**

This product should be applied at a rate of 9 - 18 (dry) oz. per acre when weeds are small (e.g., 2- leaf stage) and growing actively before the bud stage. Do not exceed 36 (dry) ounces / acre / season. Apply at the rate of 5 to 100 gallons of water per acre application and a minimum of 20 to 30 psi pressures for adequate coverage. Spray when weeds are actively growing.

- Always read and follow all label directions for tank mixes. Refer to individual use sections for additional Precautions, Restrictions, Application rates and timings.

**CORN (Field and Sweet)**

This product is effective at 9-18 dry ounces of product per acre per application. (Equivalent 0.035-0.070 pounds of acid/acre) as a broadcast treatment to corn. Apply after crop begins tillering and before the boot stage. Do not apply to corn underseeded with legumes. Corn is susceptible to injury at time of emergence and shortly after unfolding of leaves. Do not spray during this period nor after first tassels appear. Do not spray corn in tassel. 2,4-D may cause brittleness to corn. Certain single cross corn hybrids may be more susceptible to 2,4-D injury than others. Do not graze or feed forage from treated fields within 14 days after treatment. A maximum of 2 applications may be made per crop. Do not harvest corn within 7 days following application.

**Postemergence: (Annual broadleaf weeds and Perennial broadleaf weeds.)**

This product is effective at 9-18 dry ounces/acre per application. (Equivalent 0.035-0.070 pounds of acid/acre) as a broadcast treatment to corn. Apply when most weeds have germinated. Corn is susceptible to injury at time of emergence and shortly after unfolding of leaves. Do not spray during this period nor after first tassels appear. Use drop nozzles when corn is 10 inches tall to place spray below its tops. Do not cultivate soon after spraying as corn may be brittle. Do not forage or feed corn fodder for seven days following application.

**GRAPES**

This product is effective at 9-18 dry ounces of product per acre per application. (Equivalent 0.035-0.070 pounds of acid/acre). Make one directed application. Grapes are highly sensitive to 2,4-D. Do not allow spray to contact fruit, foliage, exposed roots, stems or lower limbs of vines. Do not spray bare ground. Use flat, fan-type nozzles and low pressures. Do not apply during periods of extremely high temperatures. Apply after irrigation and allow the maximum time before the next irrigation (at least three days). Grape vineyards should be at least three (3) years old or older and well established. Do not apply to newly established or young vineyards. Avoid applying where spray can reach bloom. Do not harvest these grapes within forty (40) days of application.

**GRASS SEED CROPS**

This product is effective at 9-18 dry ounces of product per acre per application. (Equivalent 0.035-0.070 pounds of acid/acre) as a broadcast treatment to the grasses. Apply after crop begins tillering and before the boot stage. Do not apply to grasses underseeded with legumes. Do not apply before tiller stage or from early boot to milk stage. Do not spray grain in the boot to dough stage. Do not graze or feed forage from treated fields within 14 days after treatment. A maximum of 2 applications may be made per crop. Do not harvest these grasses within 7 days of application. Do not graze dairy cattle for at least 7 days after application. Do not allow meat animals to enter treated fields for at least 3 days prior to slaughter. Do not cut grass for hay for at least 30 days after application.

**NUT-ORCHARD FLOORS**

This product is effective at 9-18 dry ounces of product per acre per application. (Equivalent 0.035-0.070 pounds of acid/acre) as a broadcast treatment to the orchard floor. Do not allow spray to contact fruit, foliage, exposed roots, stems or lower limbs of trees. Do not spray bare ground. Use flat, fan-type nozzles and low pressures. Do not apply during periods of extremely high temperatures. Apply after irrigation and allow the maximum time before the next irrigation (at least three days). Do not apply to newly established or young orchards. Trees should be at least one (1) year or older. Avoid applying where spray can reach bloom. A maximum of two (2) broadcast applications may be made per crop. Do not harvest these tree nuts within sixty (60) days of applications. Do not graze or feed cover crops from treated orchards to livestock.

**PISTACHIO ORCHARDS**

This product is effective at 9-18 dry ounces of product per acre per application. (Equivalent 0.035-0.070 pounds of acid/acre) as a broadcast treatment to the orchard floor. Do not allow spray to contact fruit, foliage, exposed roots, stems or lower limbs of trees. Do not spray bare ground. Use flat, fan-type nozzles and low pressure. Do not apply during periods of extremely high temperatures. Apply after irrigation and allow the maximum time before the next irrigation (at least three days). Do not apply to newly established or young orchards. Trees should be at least one (1) year or older. Avoid applying where spray can reach bloom. A maximum of two (2) broadcast applications may be made per crop. Do not harvest these tree nuts within sixty (60) days of applications. Do not graze or feed cover crops from treated orchards to livestock.

**POME FRUITS (Apple and Pear)**

This product is effective at 9-18 dry ounces of product per acre per application. (Equivalent 0.035-0.070 pounds of acid/acre) as a broadcast treatment to the orchard floor. Do not allow spray to contact fruit, foliage, exposed roots, stems or lower limbs of trees. Do

not spray bare ground. Use flat, fan-type nozzles and low pressures. Do not apply during periods of extremely high temperatures. Apply after irrigation and allow the maximum time before the next irrigation (at least three days). Do not apply to newly established or young orchards. Trees should be at least one (1) year or older. Avoid applying where spray can reach bloom. A maximum of two (2) broadcast applications may be made per crop. Allow at least seventy-five (75) days between applications. Do not harvest within forty (40) days of application. Do not graze or feed cover crops from treated orchards to livestock.

**SMALL GRAINS (Barley, Oats, Rye and Wheat)**

This product is effective at 9-18 dry ounces of product per acre per application. (Equivalent 0.035-0.070 pounds of acid/acre) as a broadcast treatment to the grains. Apply after crop begins tillering and before the boot stage. Do not apply to grains underseeded with legumes. Do not apply before tiller stage or from early boot to milk stage. Do not spray grain in the boot to dough stage. Do not graze or feed forage from treated fields within fourteen (14) days after treatment. A maximum of two (2) applications may be made per crop. Do not harvest these grains within seven (7) days of application.

**SORGHUM**

This product is effective at 9-18 dry ounces of product per acre per application. (Equivalent 0.035-0.070 pounds of acid/acre) as a broadcast treatment to sorghum. Apply after crop begins tillering and before the boot stage. Do not apply to sorghum underseeded with legumes. Do not apply before tiller stage or from early boot to milk stage. Do not graze or feed forage from treated fields within fourteen (14) days after treatment. A maximum of two (2) applications may be made per crop. Do not harvest sorghum within seven (7) days of application.

**STONE FRUITS**

This product is effective at 9-18 dry ounces of product per acre per application. (Equivalent 0.035-0.070 pounds of acid/acre) as a broadcast treatment to the orchard floor. Do not allow spray to contact fruit, foliage, exposed roots, stems or lower limbs of trees. Do not spray bare ground. Use flat, fan-type nozzles and low pressures. Do not apply during periods of extremely high temperatures. Apply after irrigation and allow the maximum time before the next irrigation (at least three days). Do not apply to newly established or young orchards. Trees should be at least one (1) year or older. Avoid applying where spray can reach bloom. A maximum of two (2) broadcast applications may be made per crop. Do not harvest these tree nuts within forty (40) days of application. Do not graze or feed cover crops from treated orchards to livestock.

**SUGARCANE**

This product is effective at 9-18 dry ounces of product per acre per application. (Equivalent 0.035-0.070 pounds of acid/acre) as a broadcast treatment to the sugarcane. Apply through layby. A maximum of two (2) applications may be made per crop. Do not harvest these sugarcane within seven (7) days of application.

**RECOMMENDED TANK MIXES**

- Always read and follow all label directions for tank mixes in combination with other herbicides. Refer to individual herbicide use sections for additional Precautions, Restrictions, and Application rates.
- This product for weed control in combination with the following crops:

CROPS	THIS PRODUCT DRY OUNCES/ACRE	CREDIT SYSTEMIC POUNDS/ACRE	BUCTRIL PINTS/ACRE
Pome fruits: Apple, and Pear	9-18	1.0-3.0 lbs	
Nut-orchard floors	9-18	1.0-3.0 lbs	
Rye	9-18	1.0-3.0 lbs	1/2 to 1-1/2 pts.
Stone fruits	9-18	1.0-3.0 lbs	
Pistachio Orchards	9-18	1.0-3.0 lbs	

**NOTE:** This product can be mixed with Nufarm Credit Herbicide and Buctril for use on the crops listed in this table, in accordance with the most restrictive label limitations and precautions. No label dosage rate should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

**FALLOW LAND AND STUBBLE**

Apply this product to annual weeds at the rate of 9-18 dry ounces of product per acre per application. Apply this product to perennial weeds at the rate of 18-36 dry ounces of product per acre per application. Apply this product when weeds are actively growing. Do not apply more than 36 dry ounces per acre per season. This product can be applied as a tank mix with Glyphosate and Banvel to broaden the spectrum of weed control. In order to assure maximum safety and weed control, follow all precautions and limitations on this product label and on the labels of products used in tank mixtures with this product. Do not make applications within 90 days of planting, or until chemical has disappeared from soil.

THIS PRODUCT	GLYPHOSATE POUNDS/ACRE	BANVEL PINTS/ACRE
9-18 ounces/acre	1/4 to 1 pound	
9-18 ounces/acre		1-2 pints

**RECOMMENDED TANK MIXTURE FOR PASTURE AND RANGELAND**

This product can be applied as a tank mix with Banvel and Tordon to broaden the spectrum of weed control. In order to assure maximum safety and weed control, follow all precautions and limitations on this label and the labels of products used in tank mixtures with this product. Apply the amounts listed below in 5-100 gallons of water per acre as a broadcast treatment.

THIS PRODUCT OUNCES/ACRE	BANVEL PINTS/ACRE	TORDON PINTS/ACRE
9-18 ounces/acre	1-2 pints	
9-18 ounces/acre		1/2 to 2 pints

**Pasture and Rangeland Restrictions and Limitations.**

- Do not allow dairy animals to graze on treated areas for at least 7 days after treatment.
- Do not cut forage or graze treated fields within 30 days after treatment.
- Do not allow livestock to enter treated fields within 3 days of slaughter.

**FORESTRY USE (FOREST SITE PREPARATION)**

Use 9-18 dry ounces of product in 5 to 100 gallons of water per acre. Dosage rates per acre depend on the density of brush and/or weeds. For small broadleaf weeds, use the lower rate. Heavy dense stands of brush require the higher rate with higher water volume. To effectively control brush, all leaves, stems, and suckers should be thoroughly wetted to the ground. Apply when plants come into full leaf (spring) to the time plants begin to go dormant. Best results are obtained when brush and broadleaf weeds are young and actively growing. Do not cut brush until the herbicide has translocated throughout the plant causing root death. Do not apply as a stand release or cover spray to established conifers as injury may result.

**RECOMMENDED TANK MIXES FOR FOREST SITE PREPARATION**

- Always read and follow all label directions for tank mixes in combination with other herbicides. Refer to individual herbicide use sections for additional Precautions, Restrictions, and Application rates.
- This product for weed control in combination with the following:

APPLICATION	THIS PRODUCT DRY OUNCES/ACRE	GLYPHOSATE POUNDS/ACRE	WATER
Aerial	dry ounce/acre	1-3 pounds	5 to 25 gallons/acre
Ground	dry ounce/acre	1-3 pounds	10 to 100 gallons/acre

**AERIAL SPRAY (FOR FORESTRY USE ONLY)**

Use 5 to 25 gallons of spray solution per acre as a broad cast spray. WHILE SPRAYING, A BUFFER ZONE MUST BE MAINTAINED TO PROTECT UNINTENDED VEGETATION: VARIOUS CONDITIONS, INCLUDING WINDY CONDITIONS AND AIR INVERSIONS, MAY CAUSE DRIFT AND WILL CAUSE DAMAGE TO ANY CONTACTED VEGETATION. USE STRAIGHT STREAM NOZZLES DIRECTED STRAIGHT BACK AND USE A SPRAY BOOM NO LONGER THAN 3/4 THE WINGSPAN OF THE AIRCRAFT.

**SPOT TREATMENT SPRAY DIRECTIONS**

- This product may be used as a spot spray application to control weeds in golf courses, parks, turfgrass, railroads, and drainage ditches. Spray to wet target weeds thoroughly.
- Control spray drift using nozzles that provide the largest size droplet consistent with adequate and uniform coverage. When applying as a spot spray use sufficient material to obtain thorough coverage of weeds.
- Always read and follow all label directions for tank mixes in combination with other herbicides. Refer to individual herbicide use sections for additional Precautions, Restrictions and Application rates.

**FOR AERIAL APPLICATION**

For applications to Small Grains, Pasture, Rangeland and Sugarcane. Make sure the aerial equipment for application provides a maximum spray distribution and maximum coverage. Recommend a minimum spray volume of 5 GPA and a maximum pressure of 40 PSI per acre. To avoid off target movement, increase spray volume per acre, do not apply in winds above 10 mph.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Do not store near heat or open flame. Keep upright in a cool place in original container.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposal of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent) adding rinsate to spray tank. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### SPECIFICATIONS FOR WARRANTY AND SALE

There is no warranty expressed or implied, concerning the use of this product other than indicated on the label. The buyer assumes all risk of liability for any and all claims, losses, damages, or injuries resulting from the use, handling and storage that are contrary to label instructions. The Directions for Use of this product are to be reliable and should be followed carefully. All such risks will be assumed by Buyer and User, and Buyer and User agree to hold Seller harmless for any claims relating to such factors. Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label, subject to the inherent risks referred to above, when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller and Buyer and Users assume the risk of any such use.

(RV "Date of stamped or notification label")

- Barvel® is a Registered Trademark of BASF Corp.
- Buctril® is a Registered Trademark of Bayer CropScience
- Credit® Systemic is a registered Trademark of Nufarm, Inc.
- Tordon® is a Registered Trademark of Dow AgroSciences
- Weedhawk™ is a Trademark of Nufarm, Inc.





9/9  
Nufarm Americas Inc.  
1333 Burr Ridge Parkway, Suite 125A  
Burr Ridge, IL 60527-0866  
Phone: 708.754.3330 Fax: 708.754.0314  
www.us.nufarm.com

October 28, 2004

Document Processing Desk (NOTIF)  
Office of Pesticide Programs (7504C)  
U. S. Environmental Protection Agency  
Room 266A, Crystal Mall 2  
1801S Bell Street  
Arlington, VA 22202-4501

Attention: Ms. Joanne Miller, PM 23  
Herbicide Branch

Subject: Label Correction

Dear Joanne:

In reviewing this label I noticed that there was verbiage missing from the grapes section of our final printed label. Accordingly, I am submitting a revised label with the language that was inadvertently left out.

If you should have any questions please do not hesitate to contact me at the letterhead address and telephone number.

Sincerely,

  
Theodore D. Head  
Product Registration Manager  
North America

