

71368-50

3-18-2008

1/11



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
 Registration Division (7505P)
 1200 Pennsylvania Ave., N.W.
 Washington, D.C. 20460

NOTICE OF PESTICIDE:

Registration
 Reregistration

(under FIFRA, as amended)

EPA Reg. Number:

71368-50

Date of Issuance:

18 MAR 2008

Term of Issuance:

Name of Pesticide Product:

Weedhawk Dry Flowable
Herbicide

Name and Address of Registrant (include ZIP Code):

Nufarm Inc.
 150 Harvester Drive
 Burr Ridge, IL 60527

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is reregistered in accordance with FIFRA section 4(g)(2)(C) provided you agree in writing to:

1) Per the acute toxicity review and the 2,4-D RED label table, the PPE section must be revised to read:

“Some materials that are chemical-resistant to this product are made of any waterproof material. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

All mixers, loaders, applicators, flaggers, and other handlers must wear:
 Long-sleeved shirt and long pants,

Shoes plus socks,
Chemical-resistant gloves (except for pilots), and
Chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

See engineering controls for additional requirements.”

2) Per the RED, the following text must be added to the engineering control section of the label:

“Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40CFR 170.240 (d)(6)].”

3) Delete the text “except as noted on appropriate labels” from the text “Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels” appearing in the Environmental Hazards section.

4) In the spray drift section, reference to chemigation must be deleted from the label because the product prohibits application through any type of irrigation system.

Also, the text “Avoid spraying near crops, plants, or trees if wind moves at or above 5 mph, or in their directions” [Page 4] and “To avoid off target movement, increase spray volume per acre, do not apply in winds above 10 mph” [Page 7] conflicts with required spray drift text and must be deleted from the label.

5) The following revisions are needed to the Directions for Use:

The text “Do not apply more than 36 (dry) ounces per season” appearing in the “General Restrictions and Limitations” section is incomplete because the area to be treated and crops are not specified. This text should be deleted from the label.

Field Corn:

If this product is intended for preplant or preemergence use, the following text must be added to the label:

“Preplant or preemergence-

Limited to one preplant or preemergence application per crop cycle.”

Sweet Corn:

If this product is also intended for preplant or preemergence use, the following text must be added to the label:

“The minimum retreatment interval is 21 days.

Limited to one postemergence application per crop cycle.”

Grass seed crop:

3/11

"The minimum retreatment interval is 21 days."

Nut-Orchard Floors and Pistachios:


"The minimum retreatment interval is 30 days."

Forestry Use:

No application rates were provided in the forestry rate table on Page 7 of the label. Rates must be listed in the table and the rates must comply with those listed on Page 139 of the RED.

6) On page 5, change "should be applied" to "is to be applied".

Signature of Approving Official:



Joanne Miller, Product Manager (23)
Herbicide Branch, Registration Division (7505P)

Date:

18 MAR 2006

EPA Form 8570-6

You will submit one copy of your final printed labeling before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). If you have any questions please call Erik Kraft at 703-308-9358.

4/11

WEEDHAWK™ DRY FLOWABLE HERBICIDE

ACTIVE INGREDIENT:

*Dimethylamine salt of 2,4-Dichlorophenoxyacetic Acid 7.50%

OTHER INGREDIENTS: 92.50%

TOTAL: 100.00%

*2,4-Dichlorophenoxyacetic acid equivalent 6.2% by weight.

**KEEP OUT OF REACH OF CHILDREN
CAUTION - PRECAUCION**

PRECAUCION AL USUARIO: Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300.
For Medical Emergencies Only, Call (877) 325-1840.

ACCEPTED
with COMMENTS
in EPA Letter Dated

18 MAR 2008

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

71368-50

EPA REG. NO. 71368-50
EPA EST. NO. 228-IL-1

MANUFACTURED BY
NUFARM INC.
150 HARVESTER DRIVE
BURR RIDGE, IL 60527



NET CONTENTS 8 x 9 OUNCE PACKETS

071368-00050.20071030.PENDING RED

5/11

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION - PRECAUCION**

Harmful if absorbed through skin. Causes moderate eye damage. Avoid contact with skin, eyes or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

All mixers, loaders, applicators, flaggers, and other handlers must wear:

- long-sleeved shirt and long pants,
- shoes and socks, plus
- chemical-resistant gloves, when applying postharvest dips or sprays to citrus, applying with any handheld nozzle or equipment, mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.
- chemical-resistant apron when applying postharvest dips or sprays to citrus, mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.
- protective eyewear

See engineering controls for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements:

When handlers use enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR part 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS. Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(6)].

Water-soluble packets when used correctly qualify as a closed loading system under the WPS. Mixers and loaders using water-soluble packets (1) must wear the PPE specified above for mixers and loaders and (2) must be provided, have immediately available for use in an emergency, such as a broken package, spill, or equipment breakdown a NIOSH-approved dust mist filtering respirator with MSHA/NIOSH approval number prefix TC-21C or a NIOSH-approved respirator with any N², R, P, or HE filter.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage.

6/11

ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

PHYSICAL OR CHEMICAL HAZARDS

Combustible. Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, chemical-resistant gloves made of any waterproof material, shoes plus socks and protective eyewear.

GENERAL RESTRICTIONS AND LIMITATIONS

- Do not apply more than 36 (dry) ounces per season.
- Do not apply to crops listed on this label that have been subjected to stress conditions such as drought, flooding, extreme diseases, nematodes, insects, where crops injury could result. Likewise with non-actively thriving weeds, control may be reduced.
- Do not apply if rain is threatening. Rainfall soon after application can reduce effectiveness.
- Do not apply through any type of irrigation system.
- Do not use treated plants for livestock feed.
- Do not use tank mixes other than listed on a label or technical bulletins. After mixing, spray within 8 hours. Do not cultivate or disturb sprayed ground for at least 7 days after spray application. Avoid applying where spray can reach young tender growth.

MIXING

Fill mixing tank 1/2 full with clean water. Drop entire water soluble packet into tank and begin agitation. Allow packets to dissolve. If tank mixing add partner herbicide while continuing agitation. Allow to thoroughly mix. Add remaining water and continue agitation. Add anti-foaming agent if desired. One to two 9 ounce packets of this product per acre is recommended. (See application rates and timing.)

Notice: To avoid subsequent injury to other crops, clean the sprayer before and after application.

Cleaning the Sprayer: Fill with clean water. Add commercial spray tank cleaner, as recommended on its label. Circulate and clean all parts, drain the tank and rinse total system as many times as needed to be thoroughly clean. Do not contaminate water when disposing of equipment wash waters.

ACTIVE INGREDIENT 2,4-D ACID PER VOLUME OF THIS PRODUCT

THIS PRODUCT	AMOUNT OF 2,4-DICHLOROPHENOXYACETIC ACID
9 ounces (dry)	0.035 pound
18 ounces (dry)	0.07 pound
36 ounces (dry)	0.14 pound

This product is recommended for use in most geographic locations of the United States, regardless of humidity or lack of rain fall and at elevations. It is recommended for use on grapes, pome fruits (apples and pears), nut-orchard floors, small grains (barley, oats, rye

7/11

and wheat), stone fruits, sugarcane, and certain non-crop uses (golf courses, parks, turfgrass, railroads, and drainage ditches). Use appropriate spray nozzles to obtain the best coverage with minimum distance from target weeds. Apply to weed growth so that coverage is uniform and complete, but not to the point of run-off. Do not apply during rain. Apply when conditions allow spray to dry on foliage.

PRECAUTIONS FOR VULNERABLE CROPS

This product should not be used on vulnerable crops such as cotton and pomegranates. Please be reminded that this product contains 2,4-D and should be used only as recommended. It may prove harmful when misused. Avoid spraying near crops, plants, or trees if wind moves at or above 5 mph, or in their direction. Avoid herbicide spraying if crops and planting are under stress conditions. This product tank mix is for use in States where approved for spraying. Consult local and State authorities concerning restrictions.

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a Coarse or coarser spray, apply only as a Coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a Medium or more fine spray, apply only as a Medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

Additional requirements for aerial applications:

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. This requirement does not apply to forestry or rights-of-way applications.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this by adjusting the path of the aircraft upwind.

Additional requirements for ground boom application:

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

GENERAL WEED LIST

This product combined with other herbicides gives control or growth suppression on the following:

Broadleaf Weeds:

- | | | | | |
|------------------|----------------|-----------------------|------------------|------------------|
| African Daisy | Kochia | Marestail | Nutsedge | Sowthistle |
| Carrot, wild | Hairy fleabane | Morning-glory, annual | Pigweed | Thistle, bull |
| Common groundsel | Lambquarters | Mustard | Prickly, lettuce | Thistle, Russian |
| Fiddleneck | Lettuce, wild | Nettles | Puncturevine | Tumbleweed |
| Filaree | Malva | Nightshade, hairy | Spurge | |

8/11

APPLICATIONS

This product should be applied at a rate of 9 - 18 (dry) oz. per acre when weeds are small (e.g., 2- leaf stage) and growing actively before the bud stage. Do not exceed 36 (dry) ounces / acre / season. Apply at the rate of 5 to 100 gallons of water per acre application and a minimum of 20 to 30 psi pressures for adequate coverage. Spray when weeds are actively growing.

- Always read and follow all label directions for tank mixes. Refer to individual use sections for additional Precautions, Restrictions, Application rates and timing.

CORN (Field and Sweet)

This product is effective at 9-18 dry ounces of product per acre per application. (Equivalent 0.035-0.070 pounds of acid/acre) as a broadcast treatment to corn. Apply after crop begins tillering and before the boot stage. Do not apply to corn underseeded with legumes. Corn is susceptible to injury at time of emergence and shortly after unfolding of leaves. Do not spray during this period nor after first tassels appear. Do not spray corn in tassel. 2,4-D may cause brittleness to corn. Certain single cross corn hybrids may be more susceptible to 2,4-D injury than others.

Postemergence: (Annual broadleaf weeds and Perennial broadleaf weeds.)

This product is effective at 9-18 dry ounces/acre per application. (Equivalent 0.035-0.070 pounds of acid/acre) as a broadcast treatment to corn. Apply when most weeds have germinated. Corn is susceptible to injury at time of emergence and shortly after unfolding of leaves. Do not spray during this period nor after first tassels appear. Use drop nozzles when corn is 10 inches tall to place spray below its tops. Do not cultivate soon after spraying as corn may be brittle.

Limitations for Use on Field Corn:

Do not use treated crops as fodder for 7 days following application. The preharvest interval (PHI) is 7 days. Limited to one postemergence and one preharvest application per crop cycle.

Limitations for Use on Sweet Corn:

Do not use treated crop as fodder for 7 days following application. The preharvest interval (PHI) is 45 days. Limited to one postemergence application per crop cycle.

GRAPES (California Only)

This product is effective at 9-18 dry ounces of product per acre per application. (Equivalent 0.035-0.070 pounds of acid/acre). Make one directed application. Grapes are highly sensitive to 2,4-D. Do not allow spray to contact fruit, foliage, exposed roots, stems or lower limbs of vines. Do not spray bare ground. Use flat, fan-type nozzles and low pressures. Do not apply during periods of extremely high temperatures. Apply after irrigation and allow the maximum time before the next irrigation (at least three days). Grape vineyards should be at least three (3) years old or older and well established. Do not apply to newly established or young vineyards. Do not harvest these grapes within 100 days of application. Limited to one application per crop cycle.

GRASS SEED CROPS

This product is effective at 9-18 dry ounces of product per acre per application. (Equivalent 0.035-0.070 pounds of acid/acre) as a broadcast treatment to the grasses. Apply after crop begins tillering and before the boot stage. Do not apply to grasses underseeded with legumes. Do not apply before tiller stage or from early boot to milk stage. Do not spray grain in the boot to dough stage. A maximum of 2 applications may be made per year.

NUT-ORCHARD FLOORS

This product is effective at 9-18 dry ounces of product per acre per application. (Equivalent 0.035-0.070 pounds of acid/acre) as a broadcast treatment to the orchard floor. Do not allow spray to contact fruit, foliage, exposed roots, stems or lower limbs of trees. Do not spray bare ground. Use flat, fan-type nozzles and low pressures. Do not apply during periods of extremely high temperatures. Apply after irrigation and allow the maximum time before the next irrigation (at least three days). Do not apply to newly established or young orchards. Trees should be at least one (1) year or older. Avoid applying where spray can reach bloom. A maximum of two (2) broadcast applications may be made per year. Do not harvest these tree nuts within sixty (60) days of applications. Do not cut orchard floor forage for hay within 7 days of application.

PISTACHIO ORCHARDS

This product is effective at 9-18 dry ounces of product per acre per application. (Equivalent 0.035-0.070 pounds of acid/acre) as a broadcast treatment to the orchard floor. Do not allow spray to contact fruit, foliage, exposed roots, stems or lower limbs of trees. Do not spray bare ground. Use flat, fan-type nozzles and low pressure. Do not apply during periods of extremely high temperatures. Apply after irrigation and allow the maximum time before the next irrigation (at least three days). Do not apply to newly established or young orchards. Trees should be at least one (1) year or older. Avoid applying where spray can reach bloom. A maximum of two (2) broadcast applications may be made per year. Do not harvest these tree nuts within sixty (60) days of applications. Do not cut orchard floor forage for hay within 7 days of application.

POME FRUITS (Apple and Pear)

This product is effective at 9-18 dry ounces of product per acre per application. (Equivalent 0.035-0.070 pounds of acid/acre) as a broadcast treatment to the orchard floor. Do not allow spray to contact fruit, foliage, exposed roots, stems or lower limbs of trees. Do not spray bare ground. Use flat, fan-type nozzles and low pressures. Do not apply during periods of extremely high temperatures. Apply after irrigation and allow the maximum time before the next irrigation (at least three days). Do not apply to newly established or young orchards. Trees should be at least one (1) year or older. Avoid applying where spray can reach bloom. A maximum of two (2) broadcast applications may be made per crop cycle. Allow at least seventy-five (75) days between applications. Do not harvest within 14 days of application. Do not cut orchard floor forage for hay within 7 days of application.

9/11

SMALL GRAINS (Barley, Oats, Rye and Wheat)

This product is effective at 9-18 dry ounces of product per acre per application. (Equivalent 0.035-0.070 pounds of acid/acre) as a broadcast treatment to the grains. Apply after crop begins tillering and before the boot stage. Do not apply to grains underseeded with legumes. Do not apply before tiller stage or from early boot to milk stage. Do not spray grain in the boot to dough stage. Do not graze or feed forage from treated fields within fourteen (14) days after treatment. Limited to one preharvest and one postemergence application per crop cycle. Do not harvest these grains within 14 days of application.

SORGHUM

This product is effective at 9-18 dry ounces of product per acre per application. (Equivalent 0.035-0.070 pounds of acid/acre) as a broadcast treatment to sorghum. Apply after crop begins tillering and before the boot stage. Do not apply to sorghum underseeded with legumes. Do not apply before tiller stage or from early boot to milk stage. Do not permit meat or dairy animals to consume treated crop as fodder or forage for 30 days following application. Limited to one postemergence application per crop cycle. Do not harvest sorghum within 30 days of application.

STONE FRUITS

This product is effective at 9-18 dry ounces of product per acre per application. (Equivalent 0.035-0.070 pounds of acid/acre) as a broadcast treatment to the orchard floor. Do not allow spray to contact fruit, foliage, exposed roots, stems or lower limbs of trees. Do not spray bare ground. Use flat, fan-type nozzles and low pressures. Do not apply during periods of extremely high temperatures. Apply after irrigation and allow the maximum time before the next irrigation (at least three days). Do not apply to newly established or young orchards. Trees should be at least one (1) year or older. Avoid applying where spray can reach bloom. A maximum of 2 broadcast applications may be made per crop cycle. Allow a minimum of 75 days between applications. Do not harvest these tree nuts within 40 days of application. Do not cut orchard floor forage for hay within 7 days of application.

SUGARCANE

This product is effective at 9-18 dry ounces of product per acre per application. (Equivalent 0.035-0.070 pounds of acid/acre) as a broadcast treatment to the sugarcane. Apply through layby. Limited to one preemergence and one postemergence application per crop cycle. Do not harvest cane prior to crop maturity.

RECOMMENDED TANK MIXES

- Always read and follow all label directions for tank mixes in combination with other herbicides. Refer to individual herbicide use sections for additional Precautions, Restrictions, and Application rates.
- **This product for weed control in combination with the following crops:**

CROPS	THIS PRODUCT DRY OUNCES/ACRE	GLYPHOSATE POUNDS/ACRE	MAESTRO 2EC PINTS/ACRE
Pome fruits: Apple, and Pear	9-18	1.0-3.0 lbs	
Nut-orchard floors	9-18	1.0-3.0 lbs	
Rye	9-18	1.0-3.0 lbs	1/2 to 1-1/2 pts.
Stone fruits	9-18	1.0-3.0 lbs	
Pistachio Orchards	9-18	1.0-3.0 lbs	

NOTE: This product can be mixed with Glyphosate and Maestro 2EC for use on the crops listed in this table, in accordance with the most restrictive label limitations and precautions. No label dosage rate should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

FALLOW LAND AND STUBBLE

Apply this product to annual weeds at the rate of 9-18 dry ounces of product per acre per application. Apply this product to perennial weeds at the rate of 18-36 dry ounces of product per acre per application. Apply this product when weeds are actively growing. Do not apply more than 36 dry ounces per acre per season. This product can be applied as a tank mix with Glyphosate and Diablo to broaden the spectrum of weed control. In order to assure maximum safety and weed control, follow all precautions and limitations on this product label and on the labels of products used in tank mixtures with this product. Plant only labeled crops within 29 days following application. A maximum of 2 applications may be made per year. Allow a minimum of 30 days between applications.

THIS PRODUCT	GLYPHOSATE POUNDS/ACRE	DIABLO PINTS/ACRE
9-18 ounces/acre	1/4 to 1 pound	
9-18 ounces/acre		1-2 pints

10/11

RECOMMENDED TANK MIXTURE FOR PASTURE AND RANGELAND

This product can be applied as a tank mix with Diablo and Tordon to broaden the spectrum of weed control. In order to assure maximum safety and weed control, follow all precautions and limitations on this label and the labels of products used in tank mixtures with this product. Apply the amounts listed below in 5-100 gallons of water per acre as a broadcast treatment.

THIS PRODUCT OUNCES/ACRE	DIABLO PINTS/ACRE	TORDON PINTS/ACRE
9-18 ounces/acre	1-2 pints	
9-18 ounces/acre		1/2 to 2 pints

Pasture and Rangeland Restrictions and Limitations.

Do not cut forage for hay within 7 days of application.

A maximum of 2 applications may be made per year.

Allow a minimum of 30 days between applications.

If grass is to be cut for hay, Agricultural Use Requirements for the Worker Protection Standard are applicable.

FORESTRY USE (FOREST SITE PREPARATION)

Use 9-18 dry ounces of product in 5 to 100 gallons of water per acre. Dosage rates per acre depend on the density of brush and/or weeds. For small broadleaf weeds, use the lower rate. Heavy dense stands of brush require the higher rate with higher water volume. To effectively control brush, all leaves, stems, and suckers should be thoroughly wetted to the ground. Apply when plants come into full leaf (spring) to the time plants begin to go dormant. Best results are obtained when brush and broadleaf weeds are young and actively growing. Do not cut brush until the herbicide has translocated throughout the plant causing root death. Do not apply as a stand release or cover spray to established conifers as injury may result. A maximum of one application may be made per year.

RECOMMENDED TANK MIXES FOR FOREST SITE PREPARATION

- Always read and follow all label directions for tank mixes in combination with other herbicides. Refer to individual herbicide use sections for additional Precautions, Restrictions, and Application rates.
- **This product for weed control in combination with the following:**

APPLICATION	THIS PRODUCT DRY OUNCES/ACRE	GLYPHOSATE POUNDS/ACRE	WATER
Aerial	dry ounce/acre	1-3 pounds	5 to 25 gallons/acre
Ground	dry ounce/acre	1-3 pounds	10 to 100 gallons/acre

AERIAL SPRAY (FOR FORESTRY USE ONLY)

Use 5 to 25 gallons of spray solution per acre as a broadcast spray. WHILE SPRAYING, A BUFFER ZONE MUST BE MAINTAINED TO PROTECT UNINTENDED VEGETATION: VARIOUS CONDITIONS, INCLUDING WINDY CONDITIONS AND AIR INVERSIONS, MAY CAUSE DRIFT AND WILL CAUSE DAMAGE TO ANY CONTACTED VEGETATION. USE STRAIGHT STREAM NOZZLES DIRECTED STRAIGHT BACK AND USE A SPRAY BOOM NO LONGER THAN 3/4 THE WINGSPAN OF THE AIRCRAFT.

SPOT TREATMENT SPRAY DIRECTIONS

- This product may be used as a spot spray application to control weeds in golf courses, parks, turfgrass, railroads, and drainage ditches. Spray to wet target weeds thoroughly.
- Control spray drift using nozzles that provide the largest size droplet consistent with adequate and uniform coverage. When applying as a spot spray use sufficient material to obtain thorough coverage of weeds.
- Always read and follow all label directions for tank mixes in combination with other herbicides. Refer to individual herbicide use sections for additional Precautions, Restrictions and Application rates.

FOR AERIAL APPLICATION

For applications to Small Grains, Pasture, Rangeland and Sugarcane. Make sure the aerial equipment for application provides a maximum spray distribution and maximum coverage. Recommend a minimum spray volume of 5 GPA and a maximum pressure of 40 PSI per acre. To avoid off target movement, increase spray volume per acre, do not apply in winds above 10 mph.

11/11

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Do not store near heat or open flame. Keep upright in a cool place in original container.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent) adding rinsate to spray tank. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV103007A)

Diablo and Maestro are Registered Trademarks and Weedhawk is a Trademark of Nufarm Americas Inc.

Tordon is a Registered Trademark of Dow AgroSciences