



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

Michael A. Peplowski  
ISK Biosciences Corporation  
7470 Auburn Road, Suite A  
Concord, Ohio 44077

JAN 31 2011

Subject: Technical Fluazinam Fungicide; 71512-8  
Decision 412724 (new use on apple)  
Decision 411457 (new use on carrot)

Omega 500F; 71512-1  
Decision 411455 - new use on carrot  
Decision 412726 - new use on apple

Decision 411458 - Petition 9E7570 - carrot  
Decision 412727 - Petition 9F7571 - apple

Dear Mr. Peplowski:

The labels referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, to add apple and carrot, are acceptable provided the following conditions are satisfied:

Submit to the Agency the following conditional data or items by the stated due date:

1. As per the prior EPA acceptance letters dated 3/27/08 and 5/28/10 for these products, submit the conditional data listed in those letters (eg. immunotoxicity study by December 31, 2011, etc.). Also with the immunotoxicity or next conditional data to be submitted below, provide a status summary of what was required in prior EPA letters vs what was submitted with copies of the EPA letters.
2. By 1/31/2012, submit multiresidue method testing data for the regulated animal metabolites, AMPA and DAPA
3. By 1/31/2012 a study that analyzes samples from the lettuce field trial for residues of AMGT. The residue data must be supported by data depicting the frozen


storage stability of AMGT in the lettuce samples. This requirement was in the EPA letter dated 5/28/2010 but no due date was listed.

**EFED BASED REQUIRED CONDITIONAL DATA (see EFED review dated 1/28/2011 for details). Submit all by 1/31/2012**

4. 835.6100, 835.6200 Environmental Chemistry Methods and ILVs for soil, water, and sediment.
5. 850.2100 - Avian acute oral toxicity using a passerine sp.
6. 850.1400 - Fish Early Life Stage for Estuarine/Marine Fish
7. 850.1350 - Aquatic invertebrate Life-Cycle Marine Study
8. 850.4150 - Tier I terrestrial plant growth- Veg. Vigor
9. 850.4400 - Tier I Aquatic Plant Growth- duckweed
10. 850.5400 - Tier I Aquatic Plant Growth - green algae, freshwater and marine diatom
11. 850.1790 - Whole Sediment Chronic Toxicity Invert.
12. LABEL CHANGES- Within 45 days submit two copies of the final printed labeling for each product and for the carrot restrictions on Omega 500F change '6 pints' to '4 pints' to match the HED review basis of 2.08 lb ai./season, and delete the duplicate lettuce section on page 9.

If you have any questions about this letter, please contact me at (703)308-9443 or kish.tony@epa.gov.

Sincerely yours,



Tony Kish  
Product Manager (22)  
Fungicide Branch  
Registration Division (7504P)

Attachments: Label stamped "ACCEPTED WITH COMMENTS"

OMEGA<sup>®</sup> 500F

## AGRICULTURAL FUNGICIDE

ACTIVE INGREDIENT: Fluazinam\* ..... 40.0%  
 OTHER INGREDIENTS: ..... 60.0%  
 Total ..... 100.0%

\*3-chloro-*N*-[3-chloro-2,6-dinitro-4-(trifluoromethyl)phenyl]-5-trifluoromethyl-2-pyridinamine (CA)

Contains 4.17 pounds fluazinam per gallon (500 grams per liter)

**KEEP OUT OF REACH OF CHILDREN**

## WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See side panel for additional precautionary statements.

Read entire label carefully and use only as directed

ISK Biosciences Corporation  
 7470 Auburn Road, Suite A  
 Concord, Ohio 44077

EPA Reg. No. 71512-1

Formulated in USA

Net Contents: 2.5 Gallons

ACCEPTED  
 with COMMENTS  
 In Final Form 2011  
 JAN 31 2011  
 Under the Federal Insecticide,  
 Fungicide, and Rodenticide Act  
 as amended for the pesticide  
 registered under EPA Reg. No.

71512-1

30613

## FIRST AID

<b>If on skin</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>

## NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

## HOT LINE NUMBER

For 24-Hour Medical Emergency Assistance  
 Call 1-888-484-7546.

For Chemical Emergency, Spill, Leak, Fire or Accident, Call  
 CHEMTREC 1-800-424-9300.

---

## PRECAUTIONARY STATEMENTS

---

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING

Causes skin irritation. Harmful if absorbed through skin. Causes moderate eye irritation. Harmful if inhaled or swallowed. Do not get on skin or on clothing. Avoid contact with eyes. Avoid breathing spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before use. Do not take internally.

#### Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical-resistance category selection chart.

Applicators, flaggers, and other handlers must wear coveralls worn over long-sleeved shirt, long pants, socks and chemical resistant footwear, chemical resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride, and protective eyewear. When mixing and loading, or when cleaning equipment, also wear a chemical resistant apron. Mixer/loaders must wear a dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C), or a NIOSH approved respirator with an N, R, P, or HE filter.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. Do not allow contact of contaminated clothing with unprotected skin.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**IMPORTANT:** When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment break-down. Do not allow contact between contaminated sprayer parts and unprotected skin. Ensure sprayer is washed down daily.

#### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove contaminated clothing and wash clothing before reuse.

---

### ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash waters or rinsate. Do not apply when weather conditions favor drift from treated areas. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas.

---

### STORAGE AND DISPOSAL

---

DO NOT contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

**PESTICIDE STORAGE:** Store in original container, in a secured, dry place separate from food and feed.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, pesticide spray or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Nonrefillable container. DO NOT reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the

40413

flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

## **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours. Refer to use directions for each crop to see additional REI restrictions for high exposure activities (i.e., hand harvesting/pruning/pinching/training) greater than 48 hours.

PPE required for early entry to the treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls worn over long-sleeved shirt and long pants, socks and chemical-resistant footwear, chemical resistant gloves made of any waterproof material, and protective eyewear.

Omega 500F may cause allergic skin reactions in a small number of sensitive individuals. To prevent the potential for an allergic reaction:

when entering treated crops, wear protective clothing (coveralls, socks and shoes) to avoid contact of unprotected skin with foliage; wash all protective clothing (coveralls) regularly, preferable daily; remove PPE immediately after leaving treated area, wash thoroughly, as soon as possible, and change into clean clothing; keep and wash PPE separately from other laundry; when entering treated crops, avoid contact of unprotected skin with treated foliage. People who have been sensitized to Omega 500F should not use or have further contact with the product.

**FAILURE TO FOLLOW THE USE DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONTROL.**

Omega 500F may be applied with all types of spray equipment normally used for ground applications. Aerial application or application through sprinkler irrigation systems is not allowed unless specific directions are given for a crop. See the crop table, and application and calibration instructions below.

Do not cultivate within 25 feet of permanent water bodies (lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, and estuaries) so as to allow growth of a vegetative filter strip.

Do not apply OMEGA 500F within 25 feet of permanent water bodies (lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, and estuaries). Do not apply OMEGA 500F by aerial equipment within 150 feet of marine/estuarine areas.

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making decisions. Where states have more stringent regulations, they must be observed.

## **MIXING AND SPRAYING**

OMEGA 500F can be used effectively in dilute or concentrate sprays. Thorough, uniform coverage is essential for disease control.

Apply OMEGA 500F in sufficient water to obtain adequate coverage of the foliage. Gallonage to be used will vary with crop and amount of plant growth. Spray volume will usually range from 20 to 100 gallons per acre

50413

for dilute sprays, and 5 to 10 gallons per acre for concentrate ground and aerial sprays. For aerial applications, apply OMEGA 500F in a minimum of 5 gallons of water per acre.

Dosage rates on this label indicate pints of OMEGA 500F per acre, unless otherwise stated. Under conditions that favor disease development, the high rate specified and the shortest application interval should be used.

NOTE: Slowly invert container several times to assure uniform mixture.

The required amount of OMEGA 500F should be added slowly into the spray tank during filling. With concentrate sprays, premix the required amount of OMEGA 500F in a clean container and add to the spray tank as it is being filled. Keep agitator running when filling spray tank and during spray operations.

DO NOT allow spray mixture to stand overnight or for prolonged periods. Prepare only the amount of spray required for immediate use. Spraying equipment should be thoroughly cleaned immediately after the application.

#### **TANK MIX COMPATIBILITY**

OMEGA 500F is physically compatible (no nozzle or screen blockage) with many products recommended for control of diseases and insects on vegetable crops. Read and follow all manufacturers' label recommendations for the tank mix companion product. It is the applicator's responsibility to ensure that the companion product is EPA approved for use on the intended crop. OMEGA 500F is generally compatible with other insecticides, fungicides, fertilizers and micronutrient products provided sufficient free water is available for dispersion of all the tank mix products. However, the physical compatibility of OMEGA 500F with tank mix partners should be evaluated before use. A jar test should be conducted with intended tank-mix pesticides prior to preparation of large volumes. Use the following procedure: 1) Pour the recommended proportions of the products into a suitable container of water, 2) Mix thoroughly and 3) Allow to stand 5 minutes. If the combination remains mixed or can be re-mixed readily, it is considered physically compatible. Any physical incompatibility in the jar test indicates that OMEGA 500F should not be used in the tank-mix.

#### **ROTATIONAL CROP (PLANTBACK) RESTRICTIONS**

Areas treated with Omega 500F may be replanted with crops on this label immediately after the last treatment. All other crops can be planted 30 days after the last application.

#### **FIELD AND ROW CROPS:**

Apply OMEGA 500F in sufficient water to obtain adequate coverage of foliage. Gallonage to be used will vary with crop and amount of plant growth. Spray volume usually will range from 20 to 60 gallons per acre (200 to 600 liters per hectare) for dilute sprays and 5 to 10 gallons per acre (50 to 100 liters per hectare) for concentrate ground sprays. Application through sprinkler irrigation systems is not allowed unless specific directions are given for a crop. See application and calibration instruction below.

#### **INTEGRATED PEST MANAGEMENT**

OMEGA 500F is an excellent disease control agent when used according to label directions for control of a broad spectrum of plant diseases. OMEGA 500F is recommended for use as part of an Integrated Pest Management (IPM) program, which may include the use of disease resistant crop varieties, cultural practices, biological control agents, pest scouting and disease forecasting systems aimed at preventing economic pest damage. Practices known to reduce disease development should be followed. Consult your state cooperative extension service or local agricultural authorities for additional IPM strategies established in your area. OMEGA 500F may be used in State Agricultural Extension advisory (disease forecasting) programs that recommend application timing based on environmental factors which favor disease development.

#### **RESISTANCE MANAGEMENT**

Some plant pathogens are known to develop resistance to products used repeatedly for disease control. OMEGA 500F is effective for strategic use in programs that attempt to minimize disease resistance to fungicides. OMEGA 500F has a multi-site mode of action that disrupts the energy production in the fungus. It is listed in FRAC code 29, as an uncoupler of oxidative phosphorylation. Some other fungicides, which are at risk from disease resistance, exhibit a single-site mode of fungicidal action. OMEGA 500F, with its multi-site mode of action, may be used to delay or prevent the development of resistance to single-site fungicides. Consult with your Federal or State Cooperative Extension Service representatives for guidance on the proper use of OMEGA 500F in programs that seek to minimize the occurrence of disease resistance to other fungicides. No known resistance has developed to OMEGA 500F and thus it is an excellent partner for those products that specify the use of a protectant or other fungicide that has a different mode of action.

60413

## **APPLICATION AND CALIBRATION TECHNIQUES FOR SPRINKLER IRRIGATION**

Apply this product only through center pivot, motorized lateral move, traveling gun, solid set or portable (wheel move, side roll, end tow, or hand move) irrigation system(s). DO NOT apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

DO NOT apply OMEGA 500F through irrigation systems connected to a public water system. "Public water system" means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days per year.

Controls for both irrigation water and pesticide injection systems must be functionally interlocked, so as to automatically terminate pesticide injection when the irrigation water pump motor stops. A person knowledgeable of the irrigation system and responsible for its operation shall be present so as to discontinue pesticide injection and make necessary adjustments, should the need arise.

The irrigation water pipeline must be fitted with a functional, automatic, quick-closing check valve to prevent the flow of treated irrigation water back toward the water source. The pipeline must also be fitted with a vacuum relief valve and low-pressure drain, located between the irrigation water pump and the check valve, to prevent back-siphoning of treated irrigation water into the water source.

**Always inject OMEGA 500F into irrigation water after it discharges from the irrigation pump and after it passes through the check valve. Never inject pesticides into the intake line on the suction side of the pump.**

Pesticide injection equipment must be fitted with a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump. Interlock this valve to the power system, so as to prevent fluid from being withdrawn from the chemical supply tank when the irrigation system is either automatically or manually turned off.

The pesticide injection pipeline must contain a functional, automatic,

quick-closing check valve to prevent the flow of fluid back toward the injection pump. The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur. DO NOT apply when wind speed favors drift beyond the area intended for treatment.

OMEGA 500F may be used through two basic types of sprinkler irrigation systems as outlined in Sections A and B below. Determine which type of system is in place, then refer to the appropriate directions provided for each type.

### **A. Center Pivot, Motorized Lateral Move and Traveling Gun Irrigation Equipment**

For injection of pesticides, these continuously moving systems must use a positive displacement injection pump of either diaphragm or piston type, constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock and capable of injection at pressures approximately 2-3 times those encountered within the irrigation water line. Venturi applicator units cannot be used on these systems.

Thoroughly mix recommended amount of this product for acreage to be covered into the same amount of water used during calibration and inject into system continuously for one revolution or run. Mixture in the chemical supply tank must be continuously agitated during the injection run. Shut off injection equipment after one revolution or run, but continue to operate irrigation system until this product has been cleared from the last sprinkler head.

### **B. Solid Set and Portable (Wheel Move, Side Roll, End Tow, or Hand Move) Irrigation Equipment**

With stationary systems, an effectively designed in-line venturi applicator unit is preferred which is constructed of materials that are compatible with pesticides; however, a positive-displacement pump can also be used.

Determine acreage covered by sprinkler. Fill tank of injection equipment with water and adjust flow to use contents over a 30 to 45 minute period. Mix desired amount of OMEGA 500F for acreage to be covered with water so that the total mixture of this product plus water in the injection tank is equal to the quantity of water used during calibration. Agitation is

70413

recommended. OMEGA 500F can be injected at the beginning or end of the irrigation cycle or as a separate application. Stop injection equipment

after treatment is completed and continue to operate irrigation system until this product has been cleared from last sprinkler head.

### DIRECTIONS FOR USE

Crop	Diseases	Rate Per Acre	Instructions
Apples	<b>Diseases Controlled</b> Apple Scab ( <i>Venturia inaequalis</i> )	10 to 13.8 fl oz	Omega 500F should be applied as a broadcast spray on a preventative basis. For scab control begin applications at green tip or when conditions are favorable for primary scab development. Repeat applications at 7- to 10-day intervals. The high rate and shortest intervals should be used for more susceptible varieties and heavy disease pressure.
	Flyspeck ( <i>Zygophiala jamaicensis</i> ) Sooty blotch ( <i>disease complex</i> )	10 to 13.8 fl oz	For control of flyspeck and sooty blotch begin applications before disease occurs and continue on a 7- to 10-day schedule. Use the higher rate and shorter interval when disease pressure is high.
	Bitter rot ( <i>Colletotrichum cingulata</i> ) Black rot ( <i>Botryosphaeria obtusa</i> ) Brooks spot ( <i>Mycosphaerella pomi</i> ) Cedar apple rust ( <i>Gymnosporangium juniperi-virginianae</i> )	13.8 fl oz	For control of bitter rot, black rot, Brooks spot, cedar apple rust, two-spotted spider mite and European red mite begin applications before disease occurs or mites are present, continue on a 7- to 10-day schedule and shorten application intervals when disease pressure or mite infestations are high. When Omega 500F is used as a cover spray, initiate the applications at petal fall and continue applications on a 7- to 10-day schedule to within 28 days of harvest.
	<b>Diseases Suppressed</b> Alternaria blotch ( <i>Alternaria mali</i> ) White rot ( <i>Botryosphaeria dothidea</i> ) Quince rust ( <i>Gymnosporangium clavipes</i> )	13.8 fl oz	For diseases and mites that are only suppressed use the high rate of 13.8 fl oz and make applications on a 7-day interval.  Omega 500F applied as cover sprays on a 7- to 10-day schedule will provide control/suppression of mites, however if applications of Omega 500F are discontinued then the application of a specific miticide may be required.
	<b>Mites Controlled</b> Two-spotted spider mite ( <i>Tetranychus urticae</i> ) European red mite ( <i>Panonychus ulmi</i> )	13.8 fl oz	Applications are based on a tree size requiring a dilute spray of 200 gallons per acre.  <b>Restrictions</b> DO NOT make more than 10 applications or apply more than 8.625 pints of Omega 500F per growing season. DO NOT apply within 28 days of harvest.
	<b>Mites Suppressed</b> Apple rust mite ( <i>Aculus schlectendali</i> )	13.8 fl oz	Restricted Entry Interval (REI) = 2 days (48 hours), for high exposure activities (i.e., hand harvesting/ pruning/pinching/ training).

80613



<b>Brassica (Cole) Leafy Vegetables, Crop Group 5, plus Turnip greens (in Crop Group 2)</b>	Club root <i>Plasmodiophora brassicae</i>	Transplant: 6.45 fl. oz. / 100 gallons  Soil Incorporation: 2.6 pints	<p><b>Application Directions:</b></p> <p><b>Transplant Soil drench:</b> Immediately after transplanting, make a single application at the rate listed here as the transplant water at 3.4 fluid ounces of solution per plant.</p> <p><b>Soil Incorporation:</b> Alternatively, if desired and for soil with low infiltration rates, apply 2.6 pints per acre in a minimum bandwidth of 9 inches along the planting row and incorporate to a soil depth of 6 to 8 inches with a precision incorporator in the same operation. Apply in a water volume of at least 50 gallons per acre. Transplant the seedlings into the treated band. If planting into a bed, a broadcast application can be made prior to forming the bed.</p> <p>Note: This product may delay the start of harvest by up to 8 days, cause some plant stunting, and shorten the harvest period, without adverse effects on the final yield.</p> <p><b>Restrictions</b></p> <p>DO NOT apply more than 3.85 pints per acre per growing season.</p> <p>DO NOT apply within 20 days of harvest on leafy greens such as mustard greens.</p> <p>DO NOT apply within 50 days of harvest on heading vegetables such as cabbage and broccoli.</p> <p>Turnip roots from turnip plants treated with Omega 500F must not be used for human or livestock consumption.</p> <p>Restricted Entry Interval, REI = 2 days, for high exposure activities (i.e., hand harvesting/pruning/pinching/ training).</p>
---	--	---	--

Includes all members of Crop Group 5, Brassica (Cole) Leafy Vegetables: broccoli, Chinese broccoli, broccoli raab (rapini), Brussels sprouts, cabbage, Chinese cabbage (bok choy), Chinese cabbage (napa), Chinese mustard cabbage, cauliflower, cavalo broccolo, collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, and rape greens. Includes the following member of Crop Group 2, Leaves of Root and Tuber Vegetables: turnip greens.

<b>Bushberry, Crop Subgroup 13-07B,</b>	Twig blight and fruit rot <i>Phomopsis vaccinii</i>  Anthracnose (Ripe rot) <i>Colletotrichum acutatum</i> <i>C. gloeosporioides</i>  Botrytis fruit rot <i>Botrytis cinerea</i>	1.25 pints	<p><b>Application Directions:</b></p> <p>Applications for fruit rots should be made on a 7- to 10-day interval, corresponding roughly to applications at green tip, pink tip, early bloom, full bloom, blossom drop and small green fruit to some blue fruit. Use adequate water to provide coverage of foliage, flowers and fruit.</p> <p><b>Restrictions</b></p> <p>DO NOT use more than 7.5 pints per acre per growing season.</p> <p>DO NOT use an adjuvant in the spray mixture with Omega 500F on this crop.</p> <p>DO NOT apply within 30 days of harvest (30-day PHI).</p> <p>Restricted Entry Interval, REI = 3 days, for high exposure activities (i.e., hand harvesting/pruning/pinching/ training).</p>
---	--	------------	---

Includes all members of the Crop Subgroup 13-07B, Bushberry: aronia berry, blueberry (highbush and lowbush), Chilean guava, currant (buffalo, black, red, and native), elderberry, European barberry, gooseberry, highbush cranberry, edible honeysuckle, huckleberry, jostaberry, juneberry, lingonberry, salal, sea buckthorn, and cultivars, varieties, and/or hybrids of these.

90413

<b>Carrot</b>	<p>Southern Blight (<i>Sclerotium rolfsii</i>)</p> <p>Sclerotinia Rot (<i>Sclerotinia sclerotiorum</i>)</p> <p>Alternaria Blight (<i>Alternaria dauci</i>)</p>	1 pint	<p><b>Application Directions:</b> The initial application for control of southern blight and sclerotinia rot should be made approximately 45 days prior to harvest or earlier if disease appears. If required, a second application can be made 14 days after the initial application. Apply in 30 to 50 gallons of water per acre as a directed band spray over the crop.</p> <p>For control of alternaria blight initiate applications when disease conditions are favorable for disease development or when disease symptoms first appear. Repeat applications as needed at a 7-day interval.</p> <p><b>Restrictions</b> DO NOT make more than 4 applications per crop cycle. DO NOT apply within 7 days of harvest (7-day PHI). Restricted Entry Interval (REI) = 2 days (48 hours). DO NOT apply more than 6 pints per growing season.</p>
<b>Ginseng</b>	<p>Rhizoctonia root rot (<i>Rhizoctonia solani</i>)</p> <p>Alternaria blight (<i>Alternaria panax</i>)</p> <p>Botrytis blight (<i>Botrytis cinerea</i>)</p> <p>White mold (<i>Sclerotinia spp.</i>)</p>	1 to 1.5 pints	<p><b>Application Directions:</b> For control of rhizoctonia root rot use 1 pt/A beginning at transplant then continue on a 14-day interval. For control of alternaria blight, botrytis blight, and white mold, use 1 pt/A beginning when the disease first appears or when conditions are favorable for disease development. Repeat applications as needed on a 7- to 14-day interval. Make a uniform application of the fungicide in a minimum of 100 gallons of water per acre. Under conditions favorable for severe disease development, use the 1.5 pint rate.</p> <p><b>Restrictions</b> DO NOT apply more than 6 pints per growing season. DO NOT apply within 30 days of harvest (30-day PHI). Restricted Entry Interval, REI = 5 days, for high exposure activities (i.e., hand harvesting/pruning/pinching/ training).</p>
<b>Lettuce, Head and Leaf</b>	<p>Sclerotinia Drop (<i>Sclerotinia minor</i>, <i>Sclerotinia sclerotiorum</i>.)</p>	1 to 1.5 pints	<p><b>Application Directions:</b> Omega 500F should be applied at 1 to 1.5 pints per acre as either a foliar band or broadcast spray or as a soil drench application at thinning. Use at least 50 gallons of water per acre. Use the higher rate in fields with a history of moderate to severe disease incidence. Omega may be used with all types of lettuce, however, DO NOT apply after thinning as phytotoxicity may occur.</p> <p><b>Restrictions</b> DO NOT apply more than 1.5 pints per growing season. DO NOT use an adjuvant with Omega 500F on this crop. DO NOT apply within 30 days of harvest (30-day PHI). Restricted Entry Interval, REI = 8 days, for medium exposure activities (i.e., irrigation, scouting). Restricted Entry Interval, REI = 12 days, for high exposure activities (i.e., hand harvesting).</p>

100413

<b>Edible-podded Legume Vegetables, (Crop Subgroup 6A, Except Peas)</b>  <b>Succulent Bean, includes Lima Bean (Crop Subgroup 6B, Except Peas)</b>  <b>Dry Beans (Crop Subgroup 6C, Except Peas and Soybeans)</b>	White mold, <i>Sclerotinia sclerotiorum</i>	0.5 to 0.85 pints	<b>Application Directions:</b> For control of white and gray molds, make the first application at 10-30% bloom (i.e. when 10-30% of the plants have at least one (1) open bloom). If needed, a second application may be applied 7 to 10 days later. Use adequate water to provide coverage of foliage and flowers. Under conditions favorable for severe disease development, use the 0.85 pint rate.  <b>Restrictions</b> DO NOT use more than 1.75 pints per acre per growing season. DO NOT apply within 14 days of harvest for edible-podded and succulent beans (14-day PHI). DO NOT apply within 30 days of harvest for dry and Lima beans (30-day PHI). Restricted Entry Interval, REI = 3 days, for high exposure activities (i.e., hand harvesting/pruning/pinching/ training).  OMEGA 500F may be applied through sprinkler system irrigation equipment on beans. See calibration directions preceding this section.
	Gray mold, <i>Botrytis cinerea</i>		

Edible-podded Legume Vegetables Subgroup 6A, except pea includes: Bean *Phaseolus* spp. runner bean, snap bean, wax bean; Bean *Vigna* spp. asparagus bean, Chinese longbean, moth bean, yardlong bean, jackbean, sword bean.

Succulent Shelled Pea and Bean Subgroup 6B, except pea includes: Bean *Phaseolus* spp. lima bean (green), broad bean (succulent); Bean *Vigna* spp. blackeye pea, cowpea, southern pea.

Dried Shelled Pea and Bean (Except Soybean) Subgroup 6C, Except Pea includes dried cultivars of the following beans: Bean *Lupinus* spp. grain lupin, sweet lupin, white lupin, white sweet lupin; Bean *Phaseolus* spp. field bean, kidney bean, lima bean (dry), navy bean, pinto bean, tepary bean; Bean *Vigna* spp. adzuki bean, blackeyed pea, catjang, cowpea, Crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, broad bean.

<b>Lettuce, Head and Leaf</b>	Sclerotinia Drop ( <i>Sclerotinia minor</i> , <i>Sclerotinia sclerotiorum</i> .)	1 to 1.5 pints	<b>Application Directions:</b> Omega 500F should be applied at 1 to 1.5 pints per acre as either a foliar band or broadcast spray or as a soil drench application at thinning. Use at least 50 gallons of water per acre. Use the higher rate in fields with a history of moderate to severe disease incidence. Omega may be used with all types of lettuce, however, DO NOT apply after thinning as phytotoxicity may occur.  <b>Restrictions</b> DO NOT apply more than 1.5 pints per growing season. DO NOT use an adjuvant with Omega 500F on this crop. DO NOT apply within 30 days of harvest (30-day PHI). Restricted Entry Interval, REI = 8 days, for medium exposure activities (i.e., irrigation, scouting). Restricted Entry Interval, REI = 12 days, for high exposure activities (i.e., hand harvesting).
-------------------------------	--	----------------	---

110413

<b>Onion, Bulb</b> <b>Crop Subgroup</b> <b>3-07A,</b>	Botrytis Leaf Blight <i>(Botrytis squamosa)</i> Botrytis Neck Rot <i>(Botrytis allii)</i> Downy Mildew <i>(Peronospora destructor)</i> Purple Blotch <i>(Alternaria porri)</i>	1 pint	<p><b>Application Directions:</b>            Initiate applications when conditions are favorable for disease development or when first disease symptoms appear. Repeat applications on a 7 to 10-day schedule. Use sufficient water to obtain adequate coverage but no less than 5 gallons per acre.</p> <p><b>Restrictions</b>            DO NOT make more than 6 applications per growing cycle.            DO NOT use an adjuvant with Omega 500F on this crop.            DO NOT apply within 7 days of harvest (7-day PHI).            Restricted Entry Interval, REI = 2 days</p> <p>OMEGA 500F may be applied through sprinkler system irrigation equipment on onions. See calibration directions preceding this section.</p>
Includes all members of the Crop Subgroup 3-07A, Onion, Bulb, such as: daylily, bulb; fritillaria, bulb; garlic, bulb; garlic, great-headed, bulb; garlic, serpent, bulb; lily, bulb; onion, bulb; onion, Chinese, bulb; onion, pearl; onion, potato, bulb; shallot, bulb; and cultivars, varieties, and/or hybrids of these.			
<b>Peanuts</b>	Sclerotinia blight <i>(Sclerotinia minor)</i>	1 to 1.5 pints	<p><b>Application Directions:</b>            Apply at 45-70 days after planting or when conditions become conducive to disease development, then make a second application approximately 3-4 weeks later. If disease conditions remain favorable, make a third application approximately 3-4 weeks after the second. If the high rate was used for the first two applications use the low rate for the third application.</p> <p><b>Restrictions</b>            DO NOT use more than 4 pints per acre during any single growing season.            DO NOT apply within 30 days of threshing for harvest.            DO NOT allow livestock to graze in treated areas.            DO NOT feed hay or threshings from treated field to livestock.            DO NOT apply by aerial application equipment.            Restricted Entry Interval, REI = 2 days, for high exposure activities (i.e., hand harvesting/pruning/pinching/ training).</p> <p>OMEGA 500F may be applied through sprinkler system irrigation equipment. Use 1 1/2 pints of product per acre in solid set, portable wheel move, center pivot, motorized lateral move or traveling gun sprinkler irrigation equipment. See calibration directions preceding this section.</p>

120813

<b>Potato</b>	Late blight ( <i>Phytophthora infestans</i> )	5.5 fl. oz.	<b>Application instructions:</b> For late blight and white mold control, begin applications when the plants are 6 to 8 inches tall or when conditions favor disease development. Repeat applications at intervals of 7 to 10 days. When white mold pressure is low to moderate, use 5 1/2 fluid ounces. When conditions favor moderate to high white mold pressure, increase the rate to 8 fluid ounces.  <b>Restrictions</b> DO NOT apply more than 3.5 pints per acre during each growing season. DO NOT apply within 14 days of harvest. Restricted Entry Interval, REI = 4 days, for high exposure activities (i.e., hand harvesting/pruning/pinching/ training).  OMEGA 500F may be applied by aerial application or through sprinkler system irrigation equipment on potatoes. See calibration directions preceding this section.
	White mold ( <i>Sclerotinia sclerotiorum</i> )	5.5 to 8 fl. oz.	

#### **WARRANTY AND LIMITATION OF DAMAGES**

Seller warrants to those persons lawfully acquiring title to this product that at the time of first sale of this product by Seller that this product conformed to its chemical description and was reasonably fit for the purposes stated on the label when used in accordance with Seller's directions under normal conditions of use, and Buyers and users of this product assume the risk of any use contrary to such directions. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, EXCEPT AS PROVIDED ELSEWHERE IN WRITING CONTAINING AN EXPRESS REFERENCE TO THIS WARRANTY AND LIMITATION OF DAMAGES, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OR GUARANTY,

**OF FITNESS OR OF MERCHANTABILITY, AND NO AGENT OF INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY SELLER IS AUTHORIZED TO DO SO.** To the extent consistent with applicable law, in no event shall Seller's liability for any breach of warranty or guaranty exceed the purchase price of the product as to which a claim is made. Buyers and users of this product are responsible for all loss or damage from use or handling of this product which results from conditions beyond the control of Seller, including, but not limited to, incompatibility with other products unless otherwise expressly provided in Directions for Use of this product, weather conditions, cultural practices, moisture conditions or other environmental conditions outside of the ranges that are generally recognized as being conducive to good agricultural and/or horticultural practices.

**ISK Biosciences Corporation**  
7470 Auburn Road, Suite A  
Concord, Ohio 44077 U.S.A.

Omega® 500F is a registered trademark of Ishihara Sangyo Kaisha, LTD

Omega 500F 11232010

130613