

71512-12

05-14-2007

1/12



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505C)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

71512-12

Date of Issuance:

MAY 14 2007

NOTICE OF PESTICIDE:

- Registration
- Reregistration

(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Flazasulfuron 25 WG

Name and Address of Registrant (include ZIP Code):

ISK Biosciences
7470 Auburn Rd.
Concord, OH 44077

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit such data.
2. Within 18 months of the date of this registration, submit one-year storage stability and corrosion characteristics study for review and acceptance.
3. Make the following label changes before you release the product for shipment:
 - a. Revise the EPA Registration Number to read, "EPA Reg. No. 71512-12."

Signature of Approving Official:

Jim Tompkins, Product Manager (25)
Herbicide Branch, Registration Division (7505P)

Date:

5-14-07

2/12

- b. Under "Precautionary Statements";

Simplify this paragraph as the following:

"Harmful, if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse."

- c. "First Aid";

Interchange the sequence between "If in eyes" and "If on skin or clothing."

- d. Under "User Safety Recommendations"

Add the following statement:

"Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry."

- e. Under "Environmental Hazards";

Add the following statements:

- Add "This product is toxic to fish" at the beginning of current Environmental Hazard Statement.
- This product may contaminate water through drift of spray in wind, or drift of soil from treated areas.
- This product has a high potential for runoff for several months or more after application. Poorly drained soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, stream, and springs will reduce the potential for contamination of water form rainfall-runoff. Runoff of this product would be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.
- There is a potential for injury to crops irrigated with run-off water containing flazasulfuron.

- f. Under " Use Precautions";

Change the title to "Application Restrictions".

3/12

Add the following statements:

- The product must be applied by applicators approved by the State to make this type of herbicide application.
- Do not apply flazasulfuron 25WG aerially.
- Do not apply flazasulfuron 25 WG through any irrigation system.
- Do not apply flazasulfuron 25 WG to areas where children can contact treated turf within 48 hours of treatment.

g. Add "Spray Drift Management" section and add the following statements:

- Avoiding spray drift at the application site is the responsibility of the applicator.
- The applicator must be familiar with the effects of temperature inversions.
- Spray is applied with coarse or extremely coarse sprays (based on the ASAE standard).

Add the following spray drift control statements:

- AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR. The interaction of many equipment-and-weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making decisions. Where states have more stringent regulations, they should be observed.
- Do not treat areas where either possible downwind movement into the soil or surface washing may cause contact of Flazasulfuron 25WG herbicide with bentgrass greens and stressed grasses.
- Avoid making applications when spray particles may be carried by air currents to areas where sensitive crops and plants are growing. Do not spray near sensitive plants if windy is gusty, below 2 mph, or in excess of 10 mph and moving in the direction of adjacent areas of sensitive crops or plants. Do not apply during temperature inversions. Always make applications when there is some air movement to determine the direction and distance of possible spray drift. Leave an adequate buffer zone of 100 feet between area to be treated and sensitive plants."
- To avoid injury to desirable plants, equipment used to apply Flazasulfuron 25WG herbicide should be thoroughly cleaned (see PROCEDURE FOR CLEANING SPRAY EQUIPMENT) before reusing to apply any other chemicals."

(1) INFORMATION ON DROPLET SIZE

The best drift management strategy is to apply large droplets and to limit or eliminate small droplets. Applying large droplets reduces drift potential, but will

4/
12

not prevent drift if applications are made improperly or under unfavorable environmental conditions (see sections below).

(2) CONTROLLING DROPLET SIZE

- Volume - Use sufficient volume to form droplets large enough to avoid drift potential.
- Pressure - Pressure and nozzle type and orientation should be carefully managed to avoid formation of fine droplets.
- Number of nozzles - Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Properly designed solid stream nozzles should produce the lowest drift potential. Select nozzles, which do not have a wide discharge profile.

(3) CALIBRATION

Equipment should be calibrated regularly according to the manufacturer's specifications.

(4) WIND

Applications should not be made when wind speed exceeds 10 mph. Use caution when applying in wind speeds less than 2-3 mph because a temperature inversion may be present and wind direction may vary. Many factors, including droplet size and equipment, determine drift potential at any wind speed. The applicator should be familiar with local wind patterns and should monitor wind conditions at the site at time of application.

(5) SENSITIVE AREAS

It is the applicator's responsibility to exercise reasonable prudence when considering the potential for drift into any area, including sensitive areas (e.g. areas where people or nontarget plants may be present, bodies of water, known habitat for threatened or endangered species, etc.).

(6) TEMPERATURE AND HUMIDITY

Low humidity and high temperature increase the evaporation rate of droplets and therefore increase spray drift potential. The applicator should compensate for temperature and humidity.

5/12

(7) TEMPERATURE INVERSIONS

Because of high drift potential, applications should not be made when droplets may reach a temperature inversion layer. *It is the applicator's responsibility to identify the presence of a temperature inversion at the time of application.*

Accurate measurements of temperature, relative humidity, and wind speed help determine if an inversion exists. Local sources of weather information may help identify the presence of temperature inversions.

h. Under "Warranty and Limitations of Damages";

Add "To the fullest extent permitted by law," before "Except as provided elsewhere . . . and no agent of seller is authorized to do so;" "In no event . . . a claim is made;" and "Buyers and users of this product . . . to good agricultural and /or horticultural practices."

Submit one copy of the revised final printed label for the record before you release the product for shipment. If the above conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Enclosure

6/12

Flazasulfuron 25WG

HERBICIDE

A selective herbicide for control of annual and perennial grasses, sedges and broadleaf weeds in Professionally Managed Bermudagrass and Zoysiagrass and for removal of overseeded cool-season turfgrasses from Bermudagrass fairways.

ACTIVE INGREDIENT:	By Wt.
Flazasulfuron*.....	25.0%
OTHER INGREDIENTS:.....	75.0%
Total.....	100.0%

* N-[[[(4,6-dimethoxy-2-pyrimidinyl) amino]carbonyl]-3-(trifluoromethyl)-2-pyridinesulfonamide

Contains 0.25 pounds active ingredient per pound of formulated product.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

SEE SIDE PANEL FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS.

READ ENTIRE LABEL CAREFULLY AND USE ONLY AS DIRECTED.

ACCEPTED
with COMMENTS
in EPA Letter Dated

EPA Registration No. 71512-<>
EPA Establishment No. <>

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

Net Contents:

71512-12
MAY 14 2007

ISK Biosciences Logo
ISK Biosciences Corporation
7470 Auburn Road, Suite A
Concord, Ohio 44077

PRECAUTIONARY STATEMENTS

7/12

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Harmful if inhaled. Avoid breathing dust. Remove contaminated clothing and wash clothing before reuse. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling.

FIRST AID	
If swallowed	<ul style="list-style-type: none">• Call a poison control center or doctor for treatment immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOT LINE NUMBER	
For 24-Hour Medical Emergency Assistance call 1-888-484-7546. For Chemical Emergency, Spill, Leak, Fire or Accident, call CHEMTREC 1-800-424-9300.	

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants, shoes plus socks, protective eyewear and waterproof gloves.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash waters or rinsate.

No bee caution required.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

8/
12

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. Keep children and pets out of treated areas until sprays have dried. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the **Restricted Entry Interval (REI)** of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, waterproof gloves, and shoes plus socks.

Sod and seed farms are within the scope of the Worker Protection Standard.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides, 40 CFR part 170. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Turf grasses on golf courses and other non-residential turf areas such as industrial parks, tank farms, professionally managed college and professional sports fields and commercial lawns are not within the scope of the Worker Protection Standard. Do not enter or allow others to enter the treated area until sprays have dried.

GENERAL INFORMATION

Flazasulfuron 25WG is a 25 percent water dispersible granular formulation of the herbicide flazasulfuron. Flazasulfuron 25WG is a post emergence herbicide with some pre-emergence activity for control of a broad spectrum of weeds as a foliar spray. Flazasulfuron 25WG can be used on golf courses and other non-residential turf areas such as industrial parks, tank farms, sod farms, seed farms, professionally managed college and professional sports fields and commercial lawns for control of cool season grasses and weeds from tolerant grasses.

Weed growth stops within hours after the application, however symptom progress from discoloration or chlorosis to necrosis generally requires from 3 to 6 weeks. Speed of control is generally a function of weather with faster action during warmer weather and actively growing weeds and turf.

Flazasulfuron 25WG controls weeds by inhibiting the acetolactate synthase (ALS) biochemical process. Some weeds may contain naturally occurring populations that are resistant to ALS inhibiting herbicides. Applications of ALS inhibiting herbicides, when used alone, over a period of time may lead to biotypes that are resistant to ALS herbicides. This then leads to a reduction in the level of control obtained through the use of these herbicides. To prevent or delay the build-up of ALS resistant weeds, weed management programs should include the use of appropriate registered herbicides for control of these weeds that have a different mode of action. Applications of herbicides with a different mode of action should be used during the same year or in sequential years.

9/12

USE PRECAUTIONS

- Repeat applications may be made at 2 – 6 weeks after the application for optimum weed control.
- The maximum yearly application rate is 9.0 oz. per acre.
- Use only on bermudagrass or zoysiagrass or severe injury may result.
- Do not apply to new seeded, sodded or sprigged turfgrass.
- Allow at least 6 weeks from the last application to the time of overseeding when applied at 1.5 to 2.25 oz/A. Allow 10 weeks for the 3 oz/A rate.
- Do not use on golf course putting greens.
- Do not apply when grasses are under stress as injury may occur.
- Make applications to actively growing weeds and turf.
- To minimize spray drift to non-target areas, do not spray if winds are above 10 mph, use large droplet size and pressure appropriate for the nozzles to produce medium to large droplets.
- If cool season grasses upslope of sensitive bentgrass greens are to be removed, allow a 100 foot untreated buffer.
- Do not run mowing equipment from treated fairways onto sensitive bentgrass greens when the turf is wet from spray, irrigation or dew. Allow a 100 foot buffer of untreated turf to minimize tracking potential.

MIXING AND LOADING INSTRUCTIONS

Ensure the spray system is clean and free of residues from previous applications. Fill the spray tank 1/2 full with clean water. Ensure the agitation system is operating and sufficient to provide uniform spray mixing during application and until the spray tank has been emptied. Complete filling the spray tank to the desired level.

Prepare no more spray mixture than is needed for the immediate application. Avoid the overnight storage of Flazasulfuron 25WG spray mixtures.

Tank Mixtures

Flazasulfuron 25WG may be tank mixed with most post emergent herbicides recommended for use on bermudagrass and zoysiagrass turf. Read and follow all manufacturer's label recommendations for the companion product. Flazasulfuron 25WG is generally compatible with other insecticides, fungicides, fertilizers and micronutrient products provided sufficient free water is available for dispersion of all the tank mix products. However, the physical compatibility of Flazasulfuron 25WG with tank mix partners should be evaluated before use.

Adjuvant Use Requirements and Precautions

The use of a non-ionic surfactant at 0.25 percent by volume (1 qt/100 gal) provides maximum performance.

Spray Equipment Clean Out:

After spraying Flazasulfuron 25WG and before using sprayer equipment for any other applications, the sprayer must be thoroughly cleaned using the following procedure.

1. Drain tank; thoroughly rinse inside of spray tanks with clean water (rinse about 1 minute per 25 gallons of tank capacity). Loosen and physically remove any visible deposits with a stiff brush.
2. Fill the tank with clean water and add 1 gallon of household ammonia (contains 3% active) for every 100 gallons of water. Circulate the cleaning solution through the tank and hoses for at least 15 minutes. Flush the cleaning solution through the hoses, boom and nozzles (1/4 volume of tank capacity) and then drain the tank.
3. Repeat step 1.
4. Repeat step 2.
5. Remove the nozzles and screen and clean separately in a bucket containing cleaning agent and water.
6. Rinse the tank, boom and hoses with clean water.
7. If only ammonia is used as a cleaner, the rinsate solution from both steps 2 and 4 may be applied back to the crop as recommended on the label. Do not exceed the maximum label use rate. If other cleaners are used, consult the cleaner label for rinsate disposal instructions. If no instructions are given, dispose of the rinsate on site or at an approved waste disposal facility.

(Attention: Do Not use Chlorine bleach with ammonia as a dangerous gas will form).

10/12

APPLICATION INFORMATION

Broadcast: Flazasulfuron 25WG can be applied as a broadcast treatment in a minimum of 20 gallons of water per acre. Higher spray volumes up to 60 gallons of water per acre can be recommended where weed populations are dense or with adverse growing conditions.

Spot Treatments With Backpack Sprayers: Dissolve 0.017 to 0.034 oz of Flazasulfuron 25WG per gallon of water, add 2 teaspoons (1/3 oz.) of a nonionic surfactant and spray mixture at a rate of 2 gallons per 1000 sq. ft.

Aerial Application: Do Not apply Flazasulfuron 25WG aerially.

Chemigation: Do Not apply Flazasulfuron 25WG through any type of irrigation system.

Rainfastness: Rainfall of one inch or greater within 3 hours of application to sensitive weeds and within 6 hours of application to less sensitive or larger weeds may necessitate reapplication of Flazasulfuron 25WG or may result in reduced weed control.

Bermudagrass and Zoysiagrass

Flazasulfuron 25WG is a selective herbicide for some pre-emergence and post-emergence control of broadleaf weeds, grass and sedges in bermudagrass and zoysiagrass. It may be applied to these turf grasses on golf courses and other non-residential turf areas such as industrial parks, tank farms, sod farms, seed farms, professionally managed college and professional sports fields and commercial lawns. Not for use in areas where children can contact treated turf. For best results, apply to actively growing weeds and turf.

Turf Tolerance

Flazasulfuron 25WG has been evaluated and can be used on the following warm season grass cultivars.

Bermudagrass: Common, Midiron, MS Express, MS Supreme, Quickstand, Sahara, Tifsport, Tifway 319, Riviera and Vaymont.

Zoysiagrass: Common, Emerald, El Toro, Meyer

Other warm season grasses such as Argentine Bahiagrass, Carpetgrass, Centipedegrass and St. Augustinegrass are intolerant.

Cool season grasses such as Ryegrass, Fescue, Bentgrass and Kentucky Bluegrass are intolerant.

Removal of Overseeded Cool Season Turf from Bermudagrass

Use 0.5 to 1.5 oz per acre to remove overseeded perennial ryegrass, 1.5 oz per acre to remove Italian (annual) ryegrass and 1.5 to 2.25 oz per acre to remove *Poa trivialis* from established bermudagrass turf. Flazasulfuron 25WG shows a slight rate response with control occurring within 4 - 5 weeks after the application with the highest rate giving better and faster control. Therefore, the bermudagrass turf should be at 50 to 60% greenup at the time of the application for optimal maintenance of a green turf situation.

Rate Recommendation Table For Weed Control

Flazasulfuron 25WG should be applied at rates from 1.5 to 3.0 oz/A for control of a large number of weeds. Broadcast applications should not exceed 0.069 oz per 1000 sq ft per application, with a maximum of three applications per season. Some difficult to control weeds may require multiple applications. For best control apply Flazasulfuron 25WG to grass and sedge weeds prior to initial tillering and when broadleaf weeds have no more than 6 true leaves.

Weed Species Controlled	Product Application Rate Ounces/1000 sq. ft	Product Application Rate Ounces/Acre
Annual bluegrass	0.069	3.0
Barnyardgrass	0.069	3.0
Broadleaf signalgrass	0.069	3.0
Quackgrass	0.069	3.0
Rattail fescue	0.034	1.5
Roughstalk bluegrass	0.069	3.0
Ryegrass, perennial and annual (transition)	0.011 - 0.034	0.5 - 1.5
Smooth crabgrass (< 4 leaf stage of growth)	0.069	3.0
Tall fescue	0.034	1.5
Cocks-comb kyllinga	0.069	3.0
Cylindric sedge	0.069	3.0
False green kyllinga	0.069	3.0
Fragrant kyllinga	0.069	3.0
Green kyllinga	0.069	3.0
Purple nutsedge (2 - 3 applications needed)	0.069	3.0
Rice flatsedge	0.069	3.0
Yellow nutsedge	0.069	3.0
Black medic	0.034	1.5
Bristly mallow	0.034	1.5
Carolina geranium	0.034	1.5
Catsear dandelion	0.034	1.5
Common chickweed	0.034	1.5
Common periwinkle	0.069	3.0
Cutleaf eveningprimrose (partial control)	0.069	3.0
Dandelion	0.069	3.0
Dogfennel	0.052 to 0.069	2.25 to 3.0
Field madder	0.034	1.5
Field pansy	0.034	1.5
Hairy bittercress	0.034	1.5
Henbit	0.034	1.5
Johnny jumpup	0.052	2.25
Large hop clover	0.034	1.5
Lawn burweed	0.034	1.5
Mouse-ear chickweed	0.034	1.5
Parsely-piert	0.034	1.5
Prostrate knotweed	0.052	2.25
Southwest bedstraw	0.034	1.5
Spotted spurge (partial control)	0.069	3.0
Virginia buttonweed (may require repeat apps)	0.069	3.0
Wandering cudweed (partial control)	0.069	3.0
White clover	0.034	1.5
Yellow rocket	0.034	1.5
Yellow woodsorrel	0.069	3.0

Partial control means good activity, but not always at a level considered commercially acceptable. Repeat applications and/or tank mixes can improve levels of control.

12/12

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, fold and roll back bags, clamp and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call CHEMTREC 1-(800) 424-9300.

To confine spill: Cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal

Completely empty container into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and Local authorities, by burning. If burned, stay out of smoke.

WARRANTY AND LIMITATION OF DAMAGES

Seller warrants to those persons lawfully acquiring title to this product that at the time of first sale of this product by Seller that this product conformed to its chemical description and was reasonably fit for the purposes stated on the label when used in accordance with Seller's directions under normal conditions of use, and Buyers and users of this product assume the risk of any use contrary to such directions. **EXCEPT AS PROVIDED ELSEWHERE IN WRITING CONTAINING AN EXPRESS REFERENCE TO THIS WARRANTY AND LIMITATION OF DAMAGES, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OR GUARANTY, INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY, AND NO AGENT OF SELLER IS AUTHORIZED TO DO SO.** In no event shall Seller's liability for any breach of warranty or guaranty exceed the purchase price of the product as to which a claim is made. Buyers and users of this product are responsible for all loss or damage from use or handling of this product which results from conditions beyond the control of Seller, including, but not limited to, incompatibility with other products unless otherwise expressly provided in Directions for Use of this product, weather conditions, cultural practices, moisture conditions or other environmental conditions outside of the ranges that are generally recognized as being conducive to good agricultural and/or horticultural practices.