



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

January 27, 2026

Michele Lussos
Regulatory Consultant
LG Chem Ltd c/o Ag Chem Consulting
12644 Chapel RD
Clifton, VA 20124

Subject: Label Amendment - Registration Review Mitigation for Lambda-Cyhalothrin
Product Name: LAMBDASTAR INSECTICIDE
EPA Registration Number: 71532-20
Case Number: 672650
Application Dates: February 18, 2022

Dear Michele Lussos:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Lambda-Cyhalothrin Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must submit one copy of the final printed labeling before you release the product for

shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Caleb Carr by phone at 202-566-0636, or via email at carr.caleb@epa.gov.

Sincerely,



Julie Javier, Team Leader
Risk Mitigation and Implementation Branch 4
Pesticide Re-Evaluation Division
Office of Pesticide Programs

ENCLOSURE: Stamped label

RESTRICTED USE PESTICIDE
Due to Toxicity to Fish and Aquatic Organisms

For retail sale to and use only to Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

Lambda-Cyhalothrin | GROUP 3A Insecticide

LAMBDASTAR INSECTICIDE

For the Control of a Variety of Insect Pests on Selected Crops

Active Ingredient:

Lambda-cyhalothrin..... 13.1%

Inert Ingredients:.....86.9%

Total 100.0%

Contains petroleum distillates.

Contains 1 lb. of active ingredient per gallon.

LambdaStar Insecticide is an emulsifiable concentrate.

KEEP OUT OF REACH OF CHILDREN

DANGER/PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

[See inside booklet for additional Precautionary Statement, [First Aid,] and Direction for Use.]

EPA Reg. No. 71532-20

EPA Est. No. 5905-AR-01

5905-GA-01

5905-IA-01

44616-MO-01

ACCEPTED

Jan 27, 2026

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under

EPA Reg. No. 71532-20

Net Contents: **Gallons**

Manufactured for:
LG Chem, LTD.

128 YEOUI-DAERO, YEONGDEUNGPO-GU
SEOUL 07336

FIRST AID	
If on skin or clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continuing rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Do not give any liquid to the person. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For 24-hour medical emergency assistance (human or animal) call 1-800-222-1222. For chemical emergency assistance (spill, leak, fire, or accident) call: CHEMTREC 1-800-424-9300.</p>	
<p>Note to Physician – Contains petroleum distillate – vomiting may cause aspiration pneumonia.</p>	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

DANGER-PELIGRO

Corrosive. Causes skin burns. May be fatal if swallowed or inhaled. Causes substantial but temporary eye injury. Do not get in eyes, on skin or clothing. Do not breathe vapor or spray mist. Harmful if absorbed through skin. Wear protective clothing, gloves, eyewear (goggles, face shield, or safety glasses) and respirator as indicated under Personal Protective Equipment. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash clothing before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Skin exposure may also result in a sensation described as a tingling, itching, burning, or prickly feeling. Onset may occur immediately to 4 hours after exposure and may last 2 to 30 hours, without damage. Wash exposed areas once with soap and water. Relief from the skin sensation may be obtained by applying an oil-based cream.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below.

Mixers, loaders, applicators and other handlers must wear:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves, made of barrier laminate, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils or viton \geq 14 mils

- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron when cleaning equipment, mixing, or loading
- A minimum of a NIOSH-approved elastomeric half mask respirator with vapor (OV) cartridges and combination R, or P filters; OR a NIOSH-approved gas mask with OV canisters; OR a NIOSH-approved powered air purifying respirator with OV cartridges and combinations HE filters.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide is extremely toxic to fish and aquatic organisms and toxic to wildlife. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

Non-target Organism Advisory

This product is highly toxic to bees and other pollinating insects exposed to direct treatment or residues in/on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are foraging the treatment area. **Protect pollinating insects by following label directions intended to minimize drift and reduce pesticide risk to these organisms.**

Physical and Chemical Hazards

Combustible liquid. Do not use or store near heat or open flame.

DIRECTIONS FOR USE

Restricted Use Pesticide

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Removable chemical extraction probes (also known as "stingers") used in suction/extraction systems must be rinsed within the pesticide container prior to removal.

This labeling must be in the possession of the user at the time of application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted-entry interval, and notification to workers. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

For hand detasseling or mechanically assisted detasseling of corn (field, pop, and sweet) grown for seed and hand harvesting of sweet corn grown for grain, the restricted-entry interval (REI) is 48 hours

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves, made of barrier laminate, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils or viton \geq 14 mils
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUE

GENERAL INFORMATION

Initial and residual control is contingent upon thorough crop coverage. Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gallons per acre by air or 10 gallons per acre by ground unless otherwise specified in this label. When foliage is dense or pest pressure is high (heavier insect or egg pressure, larger larval stages), use of higher application volumes and/or higher use rates may improve initial and residual control.

For cutworm control, LambdaStar Insecticide may be applied before, during or after planting. For soil incorporated applications, use higher rates for improved control.

Resistance Management

For resistance management, LambdaStar Insecticide contains a Group 3A Insecticide. Any insect population may contain individuals naturally resistant to LambdaStar Insecticide and other Group 3A insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of LambdaStar Insecticide or other Group 3A insecticides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticide from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issue (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.

- Adopt an integrated pest management program for insecticide use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact LG Chem Ltd at www.lgchem.com.

MANDATORY SPRAY DRIFT MANAGEMENT

Aerial Applications:

- Do not release spray at a height greater than 10 feet above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to select nozzle and pressure that deliver medium or coarser droplets according to the most current version of the American Society of Agricultural & Biological Engineers Standard 641 (ASABE S641).
- Do not apply when wind speeds exceed 15mph at the application site. If the wind speed is greater than 10mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- If the windspeed is 10 miles per hour or less, applicators must use $\frac{1}{2}$ swath displacement upwind at the downwind edge of the field. When the windspeed is between 11-15 miles per hour, applicators must use $\frac{3}{4}$ swath displacement upwind at the downwind edge of the field.
- Do not apply during temperature inversion

Airblast Applications:

- Sprays must be directed into the canopy.
- Do not apply when wind speeds exceed 15mph at the application site.
- User must turn off outward pointing nozzles at row ends and when spraying outer row.
- Do not apply during temperature inversions.

Ground Boom Applications:

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to use a Medium or coarser droplet size according to the most current version of the American Society of Agricultural & Biological Engineers Standard 572 (ASAE S572).
- Do not apply when wind speeds exceed 15 mph at the application site.
- Do not apply during temperature inversions.

Boomless Ground Applications:

- Applicators are required to select nozzle and pressure that deliver a Medium or coarser droplet size according to the most current version of the American Society of Agricultural & Biological Engineers Standard 572 (ASAE S572), for all applications.
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversion.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with higher flow rate.
- Pressure – Use the lowest pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle – Use a spray nozzle that is designed for the intended application. Consider using nozzle designed to reduce drift.

Controlling Droplet Size – Aircraft

- Adjust Nozzles – Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with airflow in flight.

Boom Height – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

Release Height - Aircrafts

Higher release heights increase the potential for spray drift.

Shielded Sprayers:

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

Handheld Technology Applications:

Take precautions to minimize spray drift.

Temperature and Humidity:

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

Temperature Inversions:

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or aircraft smoke generator. Smoke that layers ad moves laterally in concentrated cloud (under low wind conditions) indicated an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Wind:

Drift potential generally increases with wind speed. AVOID APPLICATION DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

BUFFER ZONE

VEGETATIVE FILTER STRIPS

Construct and maintain a vegetative filter strip, according to the width specified below, of grass or other permanent vegetation between the field edge and nearby down gradient aquatic habitat (such as, but not limited to, lakes; reservoirs; rivers; streams; marshes or natural ponds; estuaries; and commercial fish farm ponds). Not intended for use on rice.

Only apply products containing LambdaStar Insecticide onto fields where a maintained vegetative filter strip of at **least 25 feet** exists between the field edge and where a down gradient

aquatic habitat exists. This minimum required width of 25 feet may be reduced or removed under the following conditions:

- For Western irrigated agriculture, a maintained vegetative filter strip of at least 10 feet wide is required. Western irrigated agriculture is defined as irrigated farmland in the following states: WA, OR, CA, ID, NV, UT, AZ, MT, WY, CO, NM, and TX (west of I-35).
 - o For Western irrigated agriculture, if a sediment control basin is present, a vegetative filter strip is not required.
- In all other areas, a vegetative filter strip with a minimum width of 25 feet is required, unless the following conditions are met. The vegetative filter strip requirement may be reduced from 25 feet to 15 feet if at least one of the following applies:
 - o The area of application is considered prime farmland (as defined in 7 CFR § 657.5).
 - o Conservation tillage is being implemented on the area of application. Conservation tillage is defined as any system that leaves at least 30% of the soil surface covered by residue after planting. Conservation tillage practices can include mulch-till, no-till, or strip-till.
 - o A functional terrace system is maintained on the area of application.
 - o Water and sediment control basin for the area of application are functional and maintained.
 - o The area of application is less than or equal to 10 acres.

For further guidance on vegetated filter strips, refer to the following publication for information on constructing and maintaining effective buffers: Conservation Buffers to Reduce Pesticide Losses. National Resource Conservation Services.

<https://www.regulations.gov/document?D=EPA-HQ-OPP-2008-0331-0175>

In the State of New York, a 25 foot vegetated, non-cropped buffer strip untraversed by drainage tiles must be maintained between a treated field and a coastal salt marsh or stream that drains into a coastal salt marsh, for both aerial or ground application. For aerial applications, the 25 foot vegetated non-cropped buffer strip for runoff protection would be part of the larger 150 foot buffer strip (or 450 foot buffer strip for ULV application) required for spray drift.

Ground Application

- Do not apply within 25 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).

Ultra Low Volume (ULV) Aerial Application

- Do not apply within 450 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, river, streams, marshes, ponds, estuaries, and commercial fish ponds). Applications made by mosquito control districts and other public health officials are exempt for this requirement.

Non-ULV Aerial Application

- Do not apply within 150 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).

Following best management practices can help reduce risk to terrestrial pollinators. Examples of best management practices includes applying pesticides in the evening and at night when pollinators are not foraging and checking to confirm hive locations before spraying. For additional resources on pollinator best management practices, visit

<https://www.epa.gov/pollinator-protection/find-best-management-practices-protect-pollinators>.

Managed pollinator protection plans are developed by states/tribes to promote communication between growers, landowners, farmers, beekeepers, pesticide users, and other pest management professionals to reduce exposure of bees to pesticides. If available, visit state plans for additional information on how to protect pollinators.

How to Report Bee Kills

It is recommended that users contact both the state lead agency and the U.S. Environmental Protection Agency to report bee kills due to pesticide application. Bee kills can be reported to EPA at beekill@epa.gov. To contact your state lead agency, see the current listing of state pesticide regulatory agencies at the National Pesticide Information Center's website: http://npic.orst.edu/reg/state_agencies.html.

TANK MIX APPLICATION

When tank mixing with any other agricultural product, always add LambdaStar Insecticide last. Fill the tank with one half to two thirds volume of the mixing diluent. Make sure all other products are fully dispersed in the mixing diluent before adding the recommended rate of LambdaStar Insecticide to the tank. Add the remainder of the mixing diluent volume. It is recommended that mixing and spray equipment have continuous agitation for best results. Follow the precautions and limitations of the most restricted product in the tank mixture.

While LambdaStar Insecticide has good flexibility for tank mixing with other agricultural products, a jar test for physical compatibility is recommended for untried mixtures using proper ratios and mixing sequences of all ingredients to be included in the mixture.

LambdaStar Insecticide is an aqueous based formulation. It is recommended that no type of non-emulsifiable oils be used in combination with LambdaStar Insecticide. If adjuvants are used, use only:

- Nonionic Surfactant (NIS) containing at least 75% surface agent, or
- Non-phytotoxic Crop Oil Concentrate (COC) including once refined Vegetable Oil concentrate (VOC), or
- Methylated Sunflower Oils (MSO) containing a minimum of 17% emulsifier.

Adjuvants other than NIS or COC may be used providing the product meets the following criteria:

1. Contains only EPA exempt ingredients.
2. Is non-phytotoxic to the target crop.
3. Is compatible in mixture (may be established through a jar test).
4. Is supported locally for use with LambdaStar Insecticide on the target crop through

proven field trials and through university and extension recommendations.

In addition, the following may be used as diluents:

Crop Oil Concentrate
Methylated Sunflower
Oils Urea-Ammonium
Nitrate

It is recommended that the following not be used in combination with LambdaStar Insecticide as diluents or adjuvants:

Non-emulsifiable
Oils Diesel Fuel
Straight Mineral Oil

CHEMIGATION

Sprinkler Irrigation Application

Apply LambdaStar Insecticide at rates and timing described elsewhere in this label.

As local recommendations differ, consult your local State Extension Service or other local experts for recommendations on adjuvant or diluent types (see TANK MIX APPLICATION) rates and mixing instructions. These recommendations should be proven, through university and extension field trials, to be effective with LambdaStar Insecticide applied by chemigation.

Check the irrigation system to insure uniform application of water to all areas. Thorough coverage of foliage is required for good control. Good agitation in the pesticide supply tank should be maintained prior to and during the entire application period.

Apply by injecting the labeled rate of LambdaStar Insecticide into the irrigation system using a metering device that will introduce a constant flow and by distributing the product to the target area in 0.1-0.2 acre-inch of water. In general, use the least amount of water required for proper distribution and coverage. It is recommended that the product be injected into the main irrigation line ahead of a right angle turn in the line to insure adequate dispersion or mixing in the irrigation water. Once the application is completed, flush the entire irrigation and injection system with clean water before stopping the system.

In addition to the above recommendations, if application is being made during a normal irrigation set of a stationary sprinkler, the labeled rate of LambdaStar Insecticide for the area covered should be injected into the system only during the end of the irrigation set for sufficient time to provide adequate coverage and product distribution.

It is not recommended that LambdaStar Insecticide be applied through an irrigation system connected to a public water system. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves and average of at least 25 individuals daily at least 60 days out of the year

Use Precautions - Sprinkler Irrigation Application

A. Apply this product only through (sprinkler including center pivot, lateral move, end tow, side [wheel] roll, traveler, big gun, solid set, or hand move) irrigation system(s). Do not apply this

product through any other type of irrigation system.

B. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

C. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

D. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

E. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

F. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.

G. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

H. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

I. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

J. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

K. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

L. Any alternatives to the above required safety devices must conform to the list of EPA-approved alternative devices.

M. Do not apply when wind speed favors drift beyond the area intended for treatment or non-uniform distribution of treated water.

N. Do not apply through chemigation systems connected to public water systems.

SPECIFIC USE DIRECTIONS – AGRICULTURAL USES

Crop	Target Pests	Rate		Remarks
		lb. a.i/A	fl. oz./A	
ALAFAFA AND ALFALFA GROWN FOR SEED	Alfalfa Caterpillar	0.015-0.025	1.92 – 3.20	<ul style="list-style-type: none"> • Apply only to fields planted to pure stands of alfalfa. • Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. • Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gallons per acre by air and a minimum of 10 gallons per acre by ground. When foliage is dense and/or pest populations are high 5-10 gallons per acre by air or 20 gallons per acre by ground and higher use rates are recommended. Use higher rates for increased residual control. • Avoid application when bees are actively foraging by applying during the early morning or during the evening hours. Be aware of bee hazard resulting from a cool evening and/or morning dew. It may be advisable to remove bee shelters during and for 2-3 days following application. Do not apply directly into bee shelters. • Do not apply more than 0.03 lb. a.i. (0.24 pt.) per acre per cutting. • Do not apply more than 0.12 lb. a.i. (0.96 pt.) per acre per season. • Do not apply within 1 day of harvest for forage or within 7 days of harvest for hay. • Do not apply more than the maximum rate per acre specified for each pest in the "Rate" column to the left during a single application.
	Army cutworm			
	Cutworm spp.			
	Green Cloverworm			
	Leafhopper spp.			
	Looper spp.			
	Threecornered Alfalfa Hopper			
	Velvetbean Caterpillar			
	Webworm spp.			
	Alfalfa Seed Chalcid (Adult)			
	Alfalfa Weevil			
	Armyworm			
	Bean Leaf Beetle (Adult)			
	Blister Beetle spp.			
	Blue Alfalfa Aphid			
	Clover Leaf Weevil spp.			
	Clover Root Borer (Adult)			
	Clover Root Curculio spp. (Adult)			
	Clover Stem Borer (Adult)			
	Corn Earworm			
	Cowpea Aphid			
	Cowpea Curculio (Adult)			
	Cowpea Weevil (Adult)			
	Cucumber Beetle Spp. (Adult)			
	Egyptian Alfalfa Weevil			
	Fall Armyworm ¹			
	Grape Colaspis (Adult)			
	Grasshopper spp.			
	Green June Beetle (Adult)			
	Green Peach Aphid ³			
	Japanese Beetle (Adult)			
	Meadow Spittlebug			
	Mexican Bean Beetle			
	Pea Aphid			
	Pea Weevil (Adult)			
	Plant Bug spp. Including Lygus spp. ³			
	Spotted Alfalfa Aphid			
	Stink Bug spp.			
	Sweet Clover Weevil (Adult)			
	Thrips spp. ⁴			
	Western Yellow-striped Armyworm			
	Whitefringed Beetle spp. (Adult)			
	Yellow-striped			

¹ Use higher rates for large larvae.

² Suppression only.

³ See resistance statement under GENERAL INFORMATION.

⁴ Does not include Western Flower Thrips.

	Armyworm			
	Beet Armyworm ^{1,3}	0.03	3.84	
	Blotch Leafminer ³			
	Spider Mites ²			

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
CANOLA	Cutworm spp. Armyworm spp. Diamondback Moth Flea Beetle Cabbage Seedpod Weevil Lygus Bug Grasshoppers Looper spp.	0.015-0.03	1.92-3.84	<ul style="list-style-type: none"> • Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold. • Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gallons per acre by air and a minimum of 10 gallons per acre by ground. • Do not apply within 7 days of harvest. • Do not apply more than 0.09 lb. a.i. (0.72 pt.) per acre per year. • Do not apply more than 0.03 lb. a.i. (3.84 fl oz of product) per acre in a single application.
	Cabbage Aphid	0.03	3.84	
CEREAL GRAINS: Corn (At-Plant): Field Corn Popcorn Seed Corn Sweet Corn	Corn Rootworm Larvae (Western, Northern, Southern, Mexican) Cutworm spp. Seedcorn Maggot Seedcorn Beetle Lesser Cornstalk Borer White Grub spp. Wireworm spp. Red Imported Fire Ant ¹	0.005 lb. a.i. per 1,000 ft. of row ²	0.66 fl. oz. per 1,000 ft. of row ²	<ul style="list-style-type: none"> • Banded Applications: Apply at planting as a 5-7 inch T-band sprayed across the open seed furrow between the furrow openers and the press wheels or as a band application behind the press wheel. • In-Furrow Applications: Apply into the seed furrow through spray nozzles or microtubes behind the planter furrow openers and in front of the press wheel. • Apply a minimum of 3 gallons of finished spray per acre. • For hand detasseling or mechanically assisted detasseling of corn (field, pop, and sweet) grown for seed and hand harvesting of sweet corn grow for grain, the restricted-entry interval (REI) is 48 hours. • Do not harvest or graze livestock or cut treated crops for feed within 21 days of at-plant application. • Do not apply more than 0.09 lb. a.i. (0.72 pt.) per acre per crop at-plant. • For field corn, popcorn, and seed corn, do not apply more than 0.12 lb. a.i. (0.96 pt.) per acre per crop from at-plant and foliar applications. • For sweet corn do not apply more than 0.48 lb. a.i. (3.84 pts.) per acre per crop from at-plant and foliar applications. <p>¹ Suppression only.</p>

²Lbs. a.i. and fl. oz./A of LambdaStar Insecticide applied at 0.66 fl. oz./1000 ft. of row for various row spacings:

Row Spacing	40"	38"	36"	34"	32"	30"
Linear Ft./A	13,068	13,756	14,520	15,374	16,335	17,424
Lbs. a.i./A	0.067	0.07	0.075	0.079	0.084	0.09
Fl. oz./A	8.6	9.1	9.6	10.1	10.8	11.5

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
CEREAL GRAINS: Corn (Foliar): Field Corn Popcorn Seed Corn	Cutworm spp. Western Bean Cutworm ¹ Corn Earworm ¹ Green Cloverworm Meadow Spittlebug	0.015-0.025	1.92-3.20	<ul style="list-style-type: none"> • Apply as required by scouting or locally prescribed corn growth stages, usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds or other locally recommended methods. • Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of target location. Apply in a minimum of 2 gallons per acre by air and a minimum of 10 gallons per acre by ground. • For hand detasseling or mechanically assisted detasseling of corn (field, pop, and sweet) grown for seed and hand harvesting of sweet corn grown for grain, the restricted-entry interval (REI) is 48 hours. • For chinch bug control, begin applications when bugs migrate from small grains or grass weeds to small corn. Direct spray to the base of corn plants. Repeat applications at 3-5 day intervals if needed. LambdaStar Insecticide may only suppress heavy infestations and/or subsequent migrations. • For control of adult corn rootworm beetles (<i>Diabrotica</i> species) as part of an aerial-applied corn rootworm control program, use a minimum of 3.84 fl. oz. per acre (0.03 lb. a.i. per acre). • Do not apply within 21 days of harvest. • Do not allow livestock to graze in treated areas or harvest treated corn forage as feed for meat or dairy animals within 1 day after last treatment. Do not feed treated corn fodder or silage to meat or dairy animals within 21 days after last treatment. • Do not apply more than 0.12 lb. a.i. (0.96 pt.) per acre per crop from at-plant and foliar applications.
	Tobacco Budworm ^{1,4} European Corn Borer ¹ Southwestern Corn Borer ¹ Lesser Cornstalk Borer Stalk Borer ¹ Hop Vine Borer ¹ Armyworm ² Fall Armyworm ² Yellow-striped Armyworm ² Webworm spp. Flea Beetle spp. Seedcorn Beetle Western Corn Rootworm Beetle (Adult) Northern Corn Rootworm Beetle (Adult) Southern Corn Rootworm Beetle (Adult) Mexican Corn Rootworm Beetle (Adult) Bean Leaf Beetle Cereal Leaf Beetle Japanese Beetle (Adult) Sap Beetle (Adult) Stink Bug spp. Grasshopper spp. Corn Leaf Aphid ³ Bird Cherry-Oat Aphid ³ English Grain Aphid ³	0.02-0.03	2.56-3.84	
	Beet Armyworm ^{2,4} Chinch Bug Green Bug ^{3,4} Southern Corn Leaf Beetle ³ Mexican Rice Borer ¹	0.03	3.84	

	Rice Stalk Borer ¹ Sugarcane Borer ¹			<ul style="list-style-type: none"> • Do not apply more than 0.06 lb. a.i. (0.48 pt.) per acre after silk initiation with no more than 0.03 lb. a.i. (0.24 pt.) applied per acre per application. • Do not apply more than 0.03 lb. a.i. (0.24 pt.) per acre after corn has reached the milk stage (yellow kernels with milky fluid). • Do not apply more than the maximum rate per acre specified for each pest in the "Rate" column to the left during a single application. <p>¹For control before the larva bores into the plant stalk or ear.</p> <p>² Use higher rates for large larvae.</p> <p>³Suppression only.</p> <p>⁴See resistance statement under GENERAL INFORMATION.</p>
--	---	--	--	--

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
CEREAL GRAINS: Corn (Foliar): Sweet Corn	Corn Earworm	0.02-0.03	2.56-3.84	<ul style="list-style-type: none"> • Apply as required by scouting, or locally prescribed corn growth stages, usually at intervals of 4 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds or other locally recommended methods and should be targeted for control before insects enter the stalk or ear. • Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage and ears (if present). Apply in a minimum of 2 gallons per acre by air and a minimum of 10 gallons per acre by ground. • For hand detasseling or mechanically assisted detasseling of corn (field, pop, and sweet) grown for seed and hand harvesting of sweet corn grow for grain, the restricted-entry interval (REI) is 48 hours. • For control of adult corn rootworm beetles (<i>Diabrotica</i> species) as part of an aerial applied corn rootworm control program, use a minimum of 3.2 fl. oz. per acre (0.025 lb. a.i. per acre). • Do not apply within 1 day of harvest. • Do not allow livestock to graze in treated areas or harvest treated corn forage as feed for meat or dairy
	Fall Armyworm ¹			
	Southern Armyworm ¹			
	Beet Armyworm ^{1,3}			
	Yellow-Striped Armyworm ¹			
	Cutworm spp.			
	Armyworm ¹			
	Western Bean Cutworm			
	Webworm spp.			
	European Corn Borer			
	Southwestern Corn Borer			
	Common Cornstalk Borer			
	Western Corn Rootworm Beetle (Adult)			
	Northern Corn Rootworm Beetle (Adult)			
	Southern Corn Rootworm Beetle (Adult)			
	Mexican Corn Rootworm Beetle (Adult)			
	Japanese Beetle (Adult)			
	Sap Beetle (Adult)			
	Flea Beetle spp.			
	Tarnished Plant Bug			
	Stink Bug spp.			
	Chinch Bug			
	Aster Leafhopper			
	Grasshopper spp.			

	Aphid spp. ^{2,3} Spider Mite spp. ²			animals within 1 day after last treatment. <ul style="list-style-type: none">• Do not feed treated corn fodder or silage to meat or dairy animals within 21 days after last treatment.• Do not apply more than 0.48 lb. a.i. (3.84 pts.) per acre per crop from at-plant and foliar applications.• Do not apply more than 0.03 lb a.i. (3.84 fl oz of product) per acre in a single application.
	Corn Silkfly (Adult) ²	0.03	3.84	¹ Use higher rates for large larvae. ² Suppression only. ³ See resistance statement under GENERAL INFORMATION.

CROP	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
CEREAL GRAINS Rice, Wild Rice	Bird Cherry-Oat Aphid Chinch Bug Fall Armyworm Grasshopper spp. Greenbug Leafhopper spp. Rice Stink Bug Riceworm Rice Water Weevil (Adult) Sharpshooter spp. True Armyworm Yellow Sugarcane Aphid Yellowstriped Armyworm	0.025-0.04	3.20-5.12	<ul style="list-style-type: none">• Mixers/loaders supporting aerial applications to wild rice at a rate of 0.04 lb. a.i. per acre, and treating 1200 acres (or more) per day must wear a minimum of a NIOSH-approved elastomeric half mask respirator with vapor (OV) cartridges and combination R, or P filters; OR a NIOSH-approved gas mask with OV canisters; OR a NIOSH-approved powered air purifying respirator with OV cartridges and combinations HE filters.• Apply as required by scouting. Timing and frequency of application should be based upon insect populations reaching locally determined economic thresholds. Determine the need for repeat applications, usually at intervals of 5-7 days, by scouting.
	European Corn Borer ¹ Mexican Rice Borer ¹ Rice Seed Midge ¹ Rice Stalk Borer ¹ Sugarcane Borer ¹	0.03-0.04	3.84-5.12	<ul style="list-style-type: none">• LambdaStar Insecticide can be safely used when propanil products are being used for weed control.• Apply by air or by ground equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gallons of water (or a total carrier volume) per acre but ensure sufficient volume is used to provide adequate coverage. In addition, adding an emulsifiable crop oil (e.g., 1 pt. per acre) when lower aerial application volumes are used is recommended to help

				<p>improve coverage, reduce evaporation, and improve efficacy. Apply a minimum of 10 gallons per acre by ground.</p> <ul style="list-style-type: none"> For control of rice water weevil in dry seeded rice, make a foliar application as indicated by scouting for the presence of adults and/or feeding scars, usually within a time-frame of 0-5 days after permanent flood establishment. Do not exceed 10 days from starting permanent flood until insecticide application unless scouting indicates weevils have not been previously present. Adults may also be treated at later stages of rice development to reduce overwintering populations. For control of rice water weevil in water seeded rice, make the first foliar application after pinpoint flood as indicated by scouting for the presence of adults and/or feeding scars usually when rice has emerged 0.5 inch above the waterline. Under conditions of prolonged migration into the field, start field scouting for rice water weevil adults and/or feeding scars 3-5 days after the initial treatment and, if needed, apply a second application within 7-10 days of the first application. Adults may also be treated at later stages of rice development to reduce overwintering populations. <u>California:</u> In addition to above directions for control of rice water weevil in water seeded rice, LambdaStar Insecticide may be applied at the 1-3 leaf growth stage, with the majority at the 2 leaf growth stage. Adults are vulnerable on levees and in the water. Larvae are vulnerable while feeding on the leaf prior to entering the soil. Monitor for adults, based upon field history and density of population. Monitor field edges and levee areas for adults. Treat in the following manner: a) spray the inside perimeter of the field, or b) spray the entire field. <u>California:</u> Pre-flood, Pre-plant broadcast soil application for control of rice water weevil in wet-sown rice culture. Uniformly broadcast LambdaStar Insecticide at 3.8-5.1 fl. oz. per acre (0.03 – 0.04 lb. a.i. per acre) as a pre-flood, pre-plant application in wet-sown rice culture. Apply in a minimum of 2 gallons of water (or a total carrier volume) per acre by air or a minimum of 20 gallons of water per acre by ground. For improved efficacy, light
--	--	--	--	--

				<p>incorporation of this product into the upper 1-2 inches of soil following application is recommended – a “roller” may be used for this incorporation. Apply pinpoint flood not more than 5 days after the soil application of this product, or weevil control may be reduced. Scout for feeding scars after plant emergence and apply a second foliar treatment if needed. Do not apply more than 5.1 fl. oz. (0.04 lb. a.i.) per acre under this use pattern.</p> <ul style="list-style-type: none"> • Greenbug is known to have many biotypes. Lambdastar Insecticide may only provide suppression. If satisfactory control is not achieved with the first application of Lambdastar Insecticide, a resistant biotype may be present. Use alternate chemistry for control. • For control of stem borers, scout fields, when rice growth is near panicle differentiation, for early symptoms of damaging populations exhibited as discoloration (orange—tan) around the junction of the leaf sheath and leaf blade which is caused by feeding of young larvae within the sheath. Applications must be made before larvae bore into rice stems. Make the first application at panicle differentiation to 2 inch panicle for partial control. Make the second application at boot to heading for maximum control. All rice varieties are susceptible to stem borer damage, but Cocodrie and Priscilla are particularly susceptible. • Do not release flood water within 7 days of an application. • Do not apply more than 0.12 lb. a.i. (0.96 pt.) per acre per season. • Do not apply more than 0.04 lb. a.i. (0.32 pt.) per acre within 21 to 27 days of harvest. • Do not apply within 21 days of harvest. • Do not use treated rice fields for the aquaculture of edible fish and crustacea. • Do not apply as an ultra-low volume (ULV) spray. • Do not apply more than 0.04 lb a.i. (5.12 fl oz of product) per acre in a single application. <p>¹ For control before the larvae bores into the plant stalk.</p>
--	--	--	--	---

		Rate		
Crop	Target Pests	lb. a.i./A	fl. oz./A	Remarks
CEREAL GRAINS: Sorghum (Grain)	Cutworm spp. Sorghum Midge	0.015-0.02	1.92-2.56	<ul style="list-style-type: none"> • Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
	Armyworm Beet Armyworm ^{1,3} Fall Armyworm ¹ Yellow-striped Armyworm ¹ Corn Earworm Webworm spp. European Corn Borer ² Southwestern Corn Borer ² Lesser Cornstalk Borer ² Flea Beetle spp. Stink Bug spp. Grasshopper spp.	0.02-0.03	2.56-3.84	<ul style="list-style-type: none"> • Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of target location. Apply in a minimum of 2 gallons per acre by air and a minimum of 10 gallons per acre by ground. • For sorghum midge control, begin applications when 25% of the sorghum heads have emerged and are in tip bloom. Repeat applications at 5-day intervals if needed. • For chinch bug control, begin applications when bugs migrate from small grains or grass weeds to small sorghum. Direct spray to the base of sorghum plants. Repeat applications at 3- to 5-day intervals if needed. LambdaStar Insecticide may only suppress heavy infestations and/or subsequent migrations.
	Chinch Bug Mexican Rice Borer ² Rice Stalk Borer ² Sugarcane Borer ²	0.03	3.84	<ul style="list-style-type: none"> • Do not apply within 30 days of harvest. • Do not apply more than 0.08 lb. a.i. (0.64 pt.) per acre per season. • Do not apply more than 0.06 lb. a.i. (0.48 pt.) per acre per season after crop emergence. • Do not apply more than 0.02 lb. a.i. (0.16 pt.) per acre per season once crop is in soft dough stage. • Do not apply more than the maximum rate per acre specified for each pest in the "Rate" column to the left during a single application.
				¹ Use higher rates for large larvae only.
				² For control before the larva bores into the plant stalk.
				³ See resistance statement under GENERAL INFORMATION.

		Rate		
Crop	Target Pests	lb. a.i./A	fl. oz./A	Remarks
CEREAL GRAINS: Barley Buckwheat Oats Rye Wheat Wheat Hay Triticale	Cutworm spp. Army Cutworm	0.015-0.025	1.92-3.20	<ul style="list-style-type: none"> • Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
	Armyworm Fall Armyworm Yellow-striped Armyworm Flea Beetle spp. Cereal Leaf Beetle Stink Bug spp. English Grain Aphid ¹ Russian Wheat Aphid ¹ Bird Cherry-Oat Aphid ¹ Grasshopper spp. Orange Blossom Wheat Midge Hessian Fly ⁴	0.02-0.03	2.56-3.84	<ul style="list-style-type: none"> • Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage. Apply in a minimum of 2 gallons per acre by air and a minimum of 10 gallons per acre by ground. • For chinch bug control, repeat applications at 3- to 5-day intervals if needed. LambdaStar Insecticide may only suppress heavy infestations and/or migrations.
	Grass Sawfly	0.025-0.03	3.20-3.84	<ul style="list-style-type: none"> • Greenbug is known to have many biotypes. LambdaStar Insecticide may provide suppression only. In this situation, a second application using an alternative chemistry may be needed. • Do not apply within 30 days of harvest. • Do not allow livestock to graze in treated areas or harvest treated wheat forage as feed for meat or dairy animals within 7 days after last treatment. Do not feed treated straw to meat or dairy animals within 30 days after last treatment. • Do not apply more than 0.06 lb. a.i. (0.48 pt.) /A per season. • Do not apply more than the maximum rate per acre specified for each pest in the "Rate" column to the left during a single application.
	Chinch Bug Greenbug ^{1,2} Corn Leaf Aphid ² Mite Spp. ²	0.03	3.84	<p>¹ Best control is obtained before insects begin to roll leaves. Once wheat has started to boot, LambdaStar Insecticide may provide suppression only. Higher rates and increased coverage will be necessary.</p> <p>² Suppression only.</p> <p>³ See resistance statement under GENERAL INFORMATION.</p> <p>⁴ Make applications when adults emerge.</p>

		Rate		
Crop	Target Pests	lb. a.i./A	fl. oz./A	Remarks
COLE CROPS: Broccoli Brussels Sprouts Cabbage Cavalo Broccolo Cauliflower Chinese Broccoli (gai lon) Chinese Cabbage (napa) Chinese Mustard Cabbage (gai choy) Kohlrabi	Alfalfa Looper Cabbage Looper Imported Cabbageworm Southern Cabbageworm Cutworm spp. Cabbage Webworm	0.015-0.025	1.92-3.20	<ul style="list-style-type: none"> • Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
	Diamondback Moth ³ Armyworm Beet Armyworm ^{1,3} Fall Armyworm ¹ Yellow-striped Armyworm Corn Earworm Flea Beetle spp. Japanese Beetle (Adult) Vegetable Weevil (Adult) Grasshopper spp. Leafhopper spp. Plant Bug spp. Including Lygus spp ³ Stink Bug spp. Meadow Spittlebug Aphid spp. ^{2,3} Whitefly spp. ^{2,3} Thrips spp. ² Spider Mite spp. ²	0.02-0.03	2.56-3.84	<ul style="list-style-type: none"> • Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gallons per acre by air and a minimum of 10 gallons per acre by ground. • Do not apply as foliar broadcast application using a mechanically pressurized handgun on Brassica (head and stem). • Do not apply within 1 day of harvest. • Do not apply more than 0.24 lb. a.i. (1.92 pts.) per acre per season. <p>¹For control of first and second instar only.</p> <p>²Do not apply more than the maximum rate per acre specified for each pest in the "Rate" column to the left during a single application.</p> <p>³Suppression only.</p> <p>³See resistance statement under GENERAL INFORMATION.</p>

		Rate		
Crop	Target Pests	lb. a.i./A	fl. oz./A	Remarks
COTTON	Cutworm spp. Tobacco Thrips Soybean Thrips	0.015-0.02	1.92-2.56	<ul style="list-style-type: none"> Apply as required by scouting, usually at intervals of 5-7 days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
	Lygus Bug spp. ³ Pink Bollworm Cabbage Looper Cotton Leafperforator Saltmarsh Caterpillar Cotton Leafworm Cotton Fleahopper	0.02-0.03	2.56-3.84	<ul style="list-style-type: none"> Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gallons per acre by air and a minimum of 10 gallons per acre by ground.
	Cotton Bollworm Tobacco Budworm ³ Boll Weevil Fall Armyworm Beet Armyworm ^{1,3} European Corn Borer Brown Stink Bug Green Stink Bug Southern Green Stink Bug Two-spotted Spider Mite ² Cotton Aphid ^{2,3} Bandedwing Whitefly ^{2,3} Sweetpotato Whitefly ^{2,3}	0.025-0.04	3.20-5.12	<ul style="list-style-type: none"> Applications may also be made with equipment adapted and calibrated for ULV sprays. LambdaStar Insecticide may be mixed with once-refined vegetable oil and applied in a minimum of at least 1 qt. of finished spray per acre. Under light bollworm/budworm infestation levels, 0.02 lb. a.i. (0.16 pt.) per acre may be applied in conjunction with intense field monitoring. For boll weevil control spray on a 3- to 5-day schedule. When applied according to label directions for control of cotton bollworm and tobacco budworm, LambdaStar Insecticide also provides ovicidal control of unhatched <i>Heliothis</i> spp. Eggs. Do not apply within 21 days of harvest. Do not graze livestock in treated areas. Do not apply more than 0.2 lb. a.i. (1.6 pts.) per acre per season. Do not make more than a total of 10 synthetic pyrethroid applications (of one product or combination of products) to a cotton crop in one growing season. Do not apply more than the maximum rate per acre specified for each pest in the "Rate" column to the left during a single application.

¹For control of first and second instar only.

²Suppression only.

³See resistance statement under GENERAL INFORMATION.

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
CUCURBIT VEGETABLES	Armyworm spp. ¹ Blister Beetle spp. Cabbage Looper Corn Earworm Cricket spp. Cucumber Beetle spp. (adults) Cutworm spp. Flea Beetle spp. Grasshopper spp. June Beetle spp. Leaffooted Bug Leafhopper spp. Lygus Bug spp. ¹ Melonworm Pickleworm Plant Bug spp. Rindworm spp. Complex Saltmarsh Caterpillar Squash Beetle Squash Bug spp. Squash Vine Borer spp. Stink Bug spp. Thrips spp. ^{1,2} Tobacco Budworm ¹ Webworm spp.	0.02-0.03	2.56-3.84	<ul style="list-style-type: none"> Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of all plant parts. Apply in a minimum of 2 gallons per acre by air and a minimum of 10 gallons per acre by ground. Do not apply as foliar broadcast application using a mechanically pressurized handgun on Cucurbit Vegetables. Use higher application volumes and/or rates when foliage is dense, pest populations are high, larvae are large, weather conditions are adverse and/or as plant size increases. Use higher rates for longer residual. Insects that bore or tunnel into leaves, vines, stems or fruit must be controlled before penetration. Only exposed insects (larvae and/or adults) can be controlled with foliar applications of LambdaStar Insecticide. Do not apply more than 0.18 lb. a.i. (1.44 pts.) per acre per season. Do not apply within 1 day of harvest. Do not apply more than 0.03 lb a.i. (3.84 fl oz of product) per acre in a single application.
Muskmelon (hybrids and/or cultivars of <i>Cucumis melo</i>) — includes: true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon Pumpkin Squash, summer (<i>Cucurbita pepo</i> var. <i>melopepo</i>) — includes: crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini Squash, winter (<i>Cucurbita maxima</i> ; <i>C. moschata</i>) — includes: butternut squash, calabaza, hubbard squash (<i>C. mixta</i> ; <i>C. pepo</i>) — includes: acorn squash, spaghetti squash Watermelon — includes:	Aphid spp. ¹ Leafminer spp. ^{1,3} Spider Mite spp. ³ Whitefly spp. ^{1,3}	0.03	3.84	¹ See resistant statement under GENERAL INFORMATION. ² Does not include Western Flower Thrips. ³ Suppression only.

hybrids and/or varieties of <i>Citrullus lanatus</i>				
---	--	--	--	--

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
FRUITING VEGETABLES: Tomato and Tomatillo Peppers (bell and non-bell) Eggplant Ground Cherry Pepino	Cabbage Looper Cutworm spp. Hornworm spp.	0.015-0.025	1.92-3.20	<ul style="list-style-type: none"> Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gallons per acre by air and a minimum of 10 gallons per acre by ground. Do not apply as foliar broadcast application using a mechanically pressurized handgun on Fruiting Vegetables. Do not apply within 5 days of harvest. Do not apply more than 0.36 lb. a.i. (2.88 pts.) per acre per season. Do not apply more than the maximum rate per acre specified for each pest in the "Rate" column to the left during a single application. <p>¹For control of first and second instar only. ²Suppression only. ³See resistance statement under GENERAL INFORMATION. ⁴For control before the larva bores into the plant stalk or fruit. ⁵Does not include Western Flower Thrips.</p>
	Tomato Fruitworm Tobacco Budworm ³ Tomato Pinworm Beet Armyworm ^{1,3} Southern Armyworm ¹ Yellow-striped Armyworm ¹ Fall Armyworm ¹ European Corn Borer ⁴ Leafminer spp. ² Colorado Potato Beetle ³ Flea Beetle spp. Grasshopper spp. Leafhopper spp. Aphid spp ^{2,3} Whitefly spp ^{2,3} Meadow Spittlebug Stink Bug spp. Plant Bug spp. Stalk Borer ⁴ Blister Beetle spp. Japanese Beetle (Adult) Pepper Weevil (Adult) ² Vegetable Weevil (Adult) Tomato Psyllid ^{2,3} Spider Mite spp. ² Thrips ⁵ Cucumber Beetle spp. (Adult)	0.02-0.03	2.56-3.84	

Crop	Target Pests	Rate		Remarks
		lb.a.i./A	fl.oz./A	
GRASS FORAGE, FODDER AND HAY Pasture and Rangeland Grass, Grass Grown for hay or Silage and Grass Grown for Seed	Army Cutworm Cutworm spp. Essex Skipper Range Caterpillar Striped Grass Looper	0.015-0.025	1.92-3.2	<ul style="list-style-type: none"> Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage. Apply in a minimum of 2 gallons per acre by air and a minimum of 10 gallons per acre by ground.
	Beet Armyworm Billbug spp. ³ Bird Cherry-Oat Aphid ¹ Black Grass Bug Black Turfgrass Beetle (adult) Blue Stem Midge Cereal Leaf Beetle Chinch Bug Crane Fly spp. Cricket spp. English Grain Aphid ¹ Fall Armyworm Flea Beetle spp. Grass Mealybug Grass Sawfly (adult) Grasshopper spp. Green June Beetle (adult) Greenbug ^{1,2} Japanese Beetle (adult) Katydid spp. Leafhopper spp. Mite spp. ³ Russian Wheat Aphid ¹ Southern Armyworm Spittlebug spp. Stink Bug spp. Sugarcane Aphid Thrips spp. Tick spp. True Armyworm Webworm spp. Yellowstriped Armyworm	0.02-0.03	2.56-3.84	<ul style="list-style-type: none"> Use higher application volumes and rates when foliage is dense, pest populations are high, larvae are large and/or weather conditions are adverse. Use higher rates for longer residual. For chinch bug control, LambdaStar Insecticide may only suppress heavy infestations and/or migrations. In this situation, a second application using an alternative chemistry may be needed. Greenbug is known to have many biotypes. LambdaStar Insecticide may provide suppression only. In this situation, a second application using an alternative chemistry may be needed. Pasture and rangeland grass may be used for grazing or cut for forage 0 days after application. Do not cut grass to be dried and harvested for hay until 7 days after the last application. <p>Grass grown for seed: Straw and mature seed (seed screenings) may be used as feed 7 days after the last application. Regrowth of grass grown for seed may be used for grazing, cut for forage or cut to be dried and harvested for hay.</p> <ul style="list-style-type: none"> Do not apply more than 0.03 lb. a.i. (0.24 pt.) per acre per cutting for pastures, rangeland and grasses grown for seed. A minimum re-treatment interval (RTI) of 30 days is indicated for pastures and

				<p>rangeland receiving 0.03 lb. a.i. per acre which have not been cut between applications.</p> <ul style="list-style-type: none"> • Do not apply more than 0.09 lb. a.i. (0.72 pt.) per acre per season. • Do not apply more than the maximum rate per acre specified for each pest in the "Rate" column to the left during a single application. <p>¹Best control is obtained before insects begin to roll leaves.</p> <p>²See resistance statement under GENERAL INFORMATION.</p> <p>³Suppression only.</p>
--	--	--	--	---

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
LEGUME VEGETABLES (BEANS AND PEAS): Edible Podded (only) <i>Canavalia</i> □ <i>ladiata</i> - sword bean	Cutworm spp. Green Cloverworm Imported Cabbageworm Saltmarsh Caterpillar Velvetleaf Caterpillar Mexican Bean Beetle	0.015-0.025	1.92-3.20	<ul style="list-style-type: none"> • Apply as required by scouting. Usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
<i>Canavalia ensiformis</i> – jackbean <i>Glycine max</i> – Soybean (immature seed) Edible Podded, Succulent Shelled or Dried Shelled <i>Phaseolus</i> spp. – includes: field, kidney, lima, navy, pinto, runner, snap, tepary, and wax beans <i>Vigna</i> spp.- includes: adzuki, asparagus, moth, mung, rice, urd and yard long beans, black- eyed pea, catjang, Chinese longbean, cowpea, Crowder pea, and Southern pea <i>Pisum</i> spp. – includes: dwarf, edible-pod, English, field, garden, green, snow and sugar snap peas <i>Cajanus cajan</i> – Pigeon pea Succulent Shelled or Dried Shelled <i>Vicia faba</i> .- broadbean (favabean) Dried Shelled (only) <i>Lupinus</i> spp. – includes: grain, sweet, white and sweet white lupines <i>Cicer arietinum</i> – Chickpea (garbanzo bean) <i>Cyamopsis</i> <i>tetragonoloba</i> – guar <i>Lablab</i> <i>pupureus</i> – Lablab bean (hyacinth bean) <i>Lens</i> <i>esculata</i> –	0.02-0.03	2.56-3.84	<ul style="list-style-type: none"> • Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gallons per acre by air and a minimum of 10 gallons per acre by ground. • Do not apply as foliar broadcast application using a mechanically pressurized handgun on Legume Vegetables. • For edible podded and succulent shelled legume vegetables, do not apply within 7 days of harvest. • For dried shelled legume vegetables, do not apply within 21 days of harvest. • Do not apply more than 0.12 lb. a.i. (0.96 pt.) per acre per season. • For succulent and dried shelled peas and beans, do not graze livestock in treated areas or harvest vines for forage or hay. • Do not apply more than the maximum rate per acre specified for each pest in the "Rate" column to the left during a single application. 	
	Beet Armyworm ^{3,4} Soybean Looper ^{3,4} Lesser Cornstalk Borer ³ Leafminer spp. ^{3,4} Whitefly spp. ^{3,4}	0.03	3.84	<p>¹For control before the larva bores into the plant stalk or pods.</p> <p>²Use higher rates for large larvae.</p> <p>³For suppression only.</p> <p>⁴See resistance statement under GENERAL INFORMATION.</p> <p>⁵Does not include Western Flower Thrips.</p>

Lentils	Spider Mite Spp. ³			
---------	-------------------------------	--	--	--

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
LEGUME VEGETABLES: Soybean	Corn Earworm Velvetbean Caterpillar Green Cloverworm Cabbage Looper Painted Lady (Thistle) Caterpillar Saltmarsh Caterpillar Woollybear Caterpillar Cutworm spp. Bean Leaf Beetle Mexican Bean Beetle Western Corn Rootworm Beetle (Adult) Northern Corn Rootworm Beetle (Adult) Southern Corn Rootworm Beetle (Adult) Mexican Corn Rootworm Beetle (Adult) Three-Cornered Alfalfa Hopper Potato Leafhopper Thrips spp. ⁵ Soybean Aphid ⁴	0.015-0.025	1.92-3.20	<ul style="list-style-type: none"> Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. Do not graze or harvest treated soybean forage, straw, or hay for livestock feed. Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gallons per acre by air and a minimum of 10 gallons per acre by ground. Do not apply as foliar broadcast application using a mechanically pressurized handgun on Legume Vegetables. For control of adult corn rootworm beetles (<i>Diabrotica</i> species) as part of an aerial applied corn rootworm control program, use a minimum of 2.56 fl. oz. per acre (0.02 lb. a.i. per acre). Do not apply within 30 days of harvest. Do not apply more than 0.06 lb. a.i. (0.48 pt.) per acre per season.
	Armyworm ¹ Fall Armyworm ¹ Yellow-striped Armyworm ¹ Tobacco Budworm ³ Webworm spp. European Corn Borer Silverspotted Skipper Japanese Beetle (Adult) Blister Beetle spp. Stink Bug spp. Plant Bug spp. Grasshopper spp.	0.025-0.03	3.20-3.84	<ul style="list-style-type: none"> Do not apply more than the maximum rate per acre specified for each pest in the "Rate" column to the left during a single application. <p>¹ Use higher rates for large larvae. ² Suppression only. ³ See resistance statement under GENERAL INFORMATION. ⁴ Use lower rates for early season applications and/or lighter populations. ⁵ Does not include Western Flower Thrips.</p>
	Beet Armyworm ^{2,3} Soybean Looper ^{2,3} Lesser Cornstalk Borer ² Spider Mite spp. ²	0.03	3.84	

--	--	--	--	--

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
LETTUCE (HEAD AND LEAF)	Alfalfa Looper Cabbage Looper Imported Cabbageworm Cutworm spp. Saltmarsh Caterpillar Green Cloverworm	0.015-0.025	1.92-3.20	<ul style="list-style-type: none"> • Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
	Diamondback Moth ³ Armyworm Beet Armyworm ^{1,3} Fall Armyworm ¹ Southern Armyworm Corn Earworm Tobacco Budworm ³ European Corn Borer Flea Beetle spp. Japanese Beetle (Adult) Vegetable Weevil (Adult) Grasshopper spp. Leafhopper spp. Plant Bug spp. Including Lygus spp. ³ Stink Bug spp. Meadow Spittlebug Aphid spp. ^{2,3} Whitefly spp. ^{2,3} Spider Mite spp. ²	0.02-0.03	2.56-3.84	<ul style="list-style-type: none"> • Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gallons per acre by air and a minimum of 10 gallons per acre by ground. • Do not apply as foliar broadcast application using a mechanically pressurized handgun on Lettuce (head and leaf). • Do not apply within 1 day of harvest. • Do not apply more than 0.3 lb. a.i. (2.4 pts.) per acre per season. • Do not apply more than the maximum rate per acre specified for each pest in the "Rate" column to the left during a single application. <p>'For control of first and second instar only. ²Suppression only. ³See resistance statement under GENERAL INFORMATION.</p>
ONION (BULB) AND GARLIC	Cutworm spp. Seedcorn Maggot (Adult) Onion Maggot (Adult) Leafminer spp. (Adult)	0.015-0.025	1.92-3.20	<ul style="list-style-type: none"> • Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
	Armyworm spp. ¹ Onion Thrips ³ Tobacco Thrips ³ Western Flower Thrips ^{2,3} Flower Thrips ^{2,3} Aphid spp. ² Plant Bug spp. Stink Bug spp.	0.02-0.03	2.56-3.84	<ul style="list-style-type: none"> • Use the higher label rates as thrips population increases and avoid rescue situations. • Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage. Apply in a minimum of 2 gallons per acre by air and a minimum of 10 gallons per acre by ground. • For thrips control by aerial application, the

				<p>addition of 1 % COC v/v, ¼% NIS v/v, or a silicone adjuvant (follow manufacturer's use directions) may enhance the deposition of the spray and increase plant coverage.</p> <ul style="list-style-type: none"> • Do not apply as foliar broadcast application using a mechanically pressurized handgun on Onion (dry bulb) and Garlic. • Do not apply within 14 days of harvest. • Do not apply more than 0.24 lb. a.i. (1.92 pts.) per acre per season. • Do not apply more than the maximum rate per acre specified for each pest in the "Rate" column to the left during a single application. <p>¹For control of the first and second instars only.</p> <p>²Suppression only.</p> <p>³See resistance statement under GENERAL INFORMATION.</p>
--	--	--	--	---

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
PEANUT	Cutworm spp. Green Cloverworm Velvetbean Caterpillar Red-necked Peanut Worm Three-cornered Alfalfa Hopper Potato Leafhopper	0.015-0.025	1.92-3.20	<ul style="list-style-type: none"> • Apply as required by scouting, usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. • Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gallons per acre by air and a minimum of 10 gallons per acre by ground. • Do not apply within 14 days of harvest. • Do not apply more than 0.12 lb. a.i. (0.96 pt.) per acre per season. • Do not apply more than the maximum rate per acre specified for each pest in the "Rate" column to the left during a single application.
	Corn Earworm Fall Armyworm ¹ Bean Leaf Beetle Southern Corn Rootworm (Adult) Vegetable Weevil Whitefringed Beetle (Adult) Stink Bug spp. Tobacco Thrips Grasshopper spp.	0.02-0.03	2.56-3.84	<ul style="list-style-type: none"> • Use higher rates for large larvae. • Suppression only. • See resistance statement under GENERAL INFORMATION.
	Beet Armyworm ^{2,3} Soybean Looper ^{2,3} Lesser Cornstalk Borer ² Spider Mite spp. ² Aphid spp. ²	0.03	3.84	
POME FRUITS:	Leafroller spp. Codling Moth Tufted Apple Budworm Oriental Fruit Moth Lesser Appleworm Green Fruitworm Tent Caterpillar spp. Tentiform Leaf Miner spp. Apple Maggot (Adult) Cherry Fruit Fly spp.	0.02-0.04	2.56-5.12	<ul style="list-style-type: none"> • Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. • Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage or target area. Apply in a minimum of 10 gallons

(Adult) Pear Sawfly Plum Curculio Japanese Beetle Plant Bug spp. Stink Bug spp. Leafhopper spp. Periodical Cicada Apple Aphid Rosy Apple Aphid Pear Psylla ¹ San Jose Scale (fruit infestations only) Orange Tortrix Omnivorous Leafroller Spirea Aphid ¹ Tree Borer spp. Webworm spp.			per acre by air and a minimum of 50 gallons per acre by ground. • Do not apply within 21 days of harvest. • Do not apply more than 0.2 lb. a.i. (1.6 pts.) per acre per year. • Do not apply more than 0.16 lb. a.i. (1.28 pts.) per acre per year post bloom. • Do not apply more than 0.04 lb a.i. (5.12 fl oz of product) per acre in a single application. • Do not apply as a foliar broadcast application using a mechanically pressurized handgun on orchards • Do not apply as a soil, drench, or ground-directed application using a mechanically pressurized handgun on orchards.
¹ Suppression only.			

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
STONE FRUITS: Apricot Sweet and Tart Cherry Nectarine Peach Plum Chickasaw Plum Damson Plum Japanese Plum Plumcot Prune	Leafroller spp. Peach Twig Borer Oriental Fruit Moth Peachtree Borer spp. Green Fruitworm Tent Caterpillar spp. American Plum Borer Cherry Fruit Fly spp. (Adult) Plum Curculio Rose Chafer Japanese Beetle Plant Bug spp. Stink Bug spp. Leafhopper spp. Periodical Cicada Black Cherry Aphid Apple Maggot (Adult) Codling Moth June Beetle Pear Sawfly Thrips spp.	0.02-0.04	2.56-5.12	<ul style="list-style-type: none"> • Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds and IPM recommendations. • Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage or target area. Apply in a minimum of 10 gallons per acre by air and a minimum of 50 gallons per acre by ground. • Do not apply within 14 days of harvest. • Do not apply more than 0.2 lb. a.i. (1.6 pts.) per acre per year. • Do not apply more than 0.16 lb. a.i. (1.28 pts.) per acre per year post bloom. • Do not apply more than 0.04 lb a.i. (5.12 fl oz of product) per acre in a single application. • Do not apply as a foliar broadcast application using a mechanically pressurized handgun on orchards • Do not apply as a soil, drench, or

				ground-directed application using a mechanically pressurized handgun on orchards.
SUGARCANE	Sugarcane Borer ¹ Rice Stalk Borer ¹ Sugarcane Beetle (Adult) ² Yellow Sugarcane Aphid ³ Mexican Rice Borer ¹ Pygmy Mole Cricket Sugarcane Aphid ³ West Indian Cranefly	0.025-0.04	3.20-5.12	<ul style="list-style-type: none"> • Apply as required by scouting, usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold. • Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage or target area. Apply in a minimum of 2 gallons per acre by air and a minimum of 10 gallons per acre by ground. • Do not apply within 21 days of harvest. • Do not apply more than 0.16 lb. a.i. (1.28 pts.) per acre per season. • Do not apply more than 0.04 lb a.i. (5.12 fl oz of product) per acre in a single application. <p>¹For control before the larva bores into the plant stalk. ²Suppression only of beetles active above ground. ³See resistance statement under GENERAL INFORMATION.</p>

		Rate		
Crop	Target Pests	lb. a.i./A	fl. oz./A	Remarks
SUNFLOWER	Sunflower Beetle Cutworm spp.	0.015- 0.025	1.92-3.20	<ul style="list-style-type: none"> • Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. • Apply with ground or air equipment using sufficient water to obtain full coverage of sunflower heads and/or foliage. Apply in a minimum of 2 gallons per acre by air and a minimum of 10 gallons per acre by ground. • Do not apply within 45 days of harvest. • Do not apply more than 0.12 lb. a.i. (0.96 pt.) per acre per season.
	Sunflower Moth Banded Sunflower Moth Fall Armyworm ¹ Woollybear Caterpillar Spotted Cabbage Looper Painted Lady (Thistle) Caterpillar Seed Weevil (Adult) Stem Weevil (Adult) Head-Clipper Weevil (Adult) Japanese Beetle (Adult) Sunflower Maggot	0.02-0.03	2.56-3.84	

	(Adult) Leafhopper spp. Meadow Spittlebug Stink Bug spp. Grasshopper spp.			<ul style="list-style-type: none"> • Do not apply more than 0.09 lb. a.i. (0.72 pt.) per acre per season after bloom initiation. • Do not apply as a ultra-low volume (ULV) spray.
	Beet Armyworm ^{2,3} Spider Mite spp. ²	0.03	3.84	<ul style="list-style-type: none"> • Do not apply more than the maximum rate per acre specified for each pest in the "Rate" column to the left during a single application. <p>¹ Use higher rates for large larvae. ²Suppression only. ³See resistance statement under GENERAL INFORMATION.</p>
TOBACCO	Tobacco Budworm ³ Tobacco Hornworm Cabbage Looper Corn Earworm Salt Marsh Caterpillar Armyworm spp. ¹ Cutworm spp. Webworm spp. Tobacco Flea Beetle (Adult) Cucumber Beetle spp. (Adult) Blister Beetle spp. Vegetable Weevil (Adult) Japanese Beetle (Adult) Grasshopper spp. Tree Cricket spp. Katydid spp. Plant Bug spp. ³ Stinkbug spp. Tobacco Thrips spp. ² Tobacco Aphid spp. ^{2,3} Tobacco Hornworm Potato Tuberworm	0.015-0.03	1.92-3.84	<ul style="list-style-type: none"> • Apply as required by scouting, usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold. • Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage. Apply in a minimum of 2 gallons per acre by air and a minimum of 10 gallons per acre by ground. • Do not apply as foliar broadcast application using a mechanically pressurized handgun on Tobacco. • Do not apply within 40 days of harvest. • Do not apply more than 0.09 lb. a.i. (0.72 pt.) per acre per year. • Do not apply more than 0.03 lb a.i. (3.84 fl oz of product) per acre in a single application. <p>¹For control of first and second instar only. ²Suppression only. ³See resistance statement under GENERAL INFORMATION.</p>

		Rate		
Crop	Target Pests	lb. a.i./A	fl. oz./A	Remarks
TREE NUTS: Almond Beech Nut Brazil Nut Butternut Cashew Chestnut Chinquapin Filbert (Hazelnut) Hickory Nut Macadamia Nut (Bush Nut) Pistachio Walnut, Black Walnut, English (Persian)	Leafroller spp. Navel Orangeworm Codling Moth Filbertworm Peach Twig Borer Walnut Husk Fly spp. (Adult) Ants (excluding Harvester, Pharaoh, Fire and Carpenter) Plant Bug spp. Stink Bug spp. Chinch Bug Leaffooted Bug Walnut Aphid	0.02-0.04	2.56-5.12	<ul style="list-style-type: none"> • Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold. • Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage or target area. Apply in a minimum of 10 gallons per acre by air and a minimum of 50 gallons per acre by ground. • Do not apply within 14 days of harvest. • Do not apply more than 0.16 lb. a.i. (1.28 pts.) per acre per year. • Do not apply more than 0.12 lb. a.i. (0.96 pt.) per acre per year post bloom. • Do not apply more than 0.04 lb a.i. (5.12 fl oz of product) per acre in a single application. • Do not apply as a foliar broadcast application using a mechanically pressurized handgun on orchards • Do not apply as a soil, drench, or ground-directed application using a mechanically pressurized handgun on orchards.
Pecan	Hickory Shuckworm Pecan Casebearer spp. Pecan Weevil Pecan Aphid spp. Pecan Spittlebug Pecan Phylloxera spp. Stink Bug spp.	0.02-0.04	2.56-5.12	

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
TUBEROUS AND CORM VEGETABLES (Potato, Sweet Potato, Yams and Related)	Cutworm spp. Leafhopper spp. Saltmarsh Caterpillar Sweet Potato Hornworm Woolybear Caterpillar spp.	0.015-0.025	1.92-3.20	<ul style="list-style-type: none"> Apply as required by scouting, usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of all above ground plant parts. Apply in a minimum of 2 gallons per acre by air and a minimum of 10 gallons per acre by ground. Do not apply as foliar broadcast application using a mechanically pressurized handgun on Tuberous and Corm Vegetables. Use higher application volumes and/or rates when foliage is dense, pest populations are high, larvae are large, weather conditions are adverse and/or as plant size increases. Use higher rates for longer residual. Insects that bore or tunnel into leaves, vines, stems, tubers or corms must be controlled before penetration. Only exposed insects (larvae and/or adults) can be controlled with foliar applications of LambdaStar Insecticide. Do not apply within 7 days of harvest. Do not apply more than 0.12 lb. a.i. (0.96 pt.) per acre per season. Do not apply more than the maximum rate per acre specified for each pest in the "Rate" column to the left
	Aphid spp. ¹ Armyworm spp. ¹ Blister Beetle spp. Colorado Potato Beetle ¹ Corn Earworm Cricket spp. Cucumber Beetle spp. (adults) European Corn Borer Flea Beetle spp. (adults) Grasshopper spp. Looper spp. ¹ Lygus Bug spp. ¹ Plant Bug spp. Potato Psyllid Potato Tuberworm Stink Bug spp. Sweet Potato Leaf Beetle (adults) Sweet Potato Vine Borer Thrips spp. ^{1, 2} Tortoise Beetle spp. Webworm spp. Weevil spp. (adults)	0.02-0.03	2.56-3.84	
	Leafminer spp. ^{1,3} Whitefly spp. ^{1,3} Spider Mite spp. ³	0.03	3.84	

				<p>during a single application.</p> <p>¹See resistance statement under GENERAL INFORMATION.</p> <p>²Does not include Western Flower Thrips.</p> <p>³Suppression only.</p>
--	--	--	--	--

OTHER USES

		Rate		
Crop	Target Pests	lb. a.i./A	fl. oz./A	Remarks
CONIFER AND DECIDUOUS TREES: Plantations Nurseries	Pine Tip Moth spp. Spruce Budworm Bagworm Tent Caterpillar spp. Leafroller spp. Gypsy Moth Webworm spp. Tussock Moth spp. Pine Sawfly spp. Sawfly spp. Pine Chafer Japanese Beetle May Beetle spp. June Beetle spp. Pine Colaspis Beetle Leaf Beetle spp. Pales Weevil Pine Weevil spp. Pine Conelet Bug Spittlebug spp. Pine Leaf Chermid Balsam Wooly Aphid Balsam Twig Aphid Birch Leafminer Black Pine Weevil Elm Leaf Beetle European Elm Bark Beetle Mealybug spp. ¹ Pine Needle Scale Pine Tortoise Scale Poplar Aphid spp.	0.02-0.04	2.56-5.12	<ul style="list-style-type: none"> • To control exposed foliage, flower, cone, seed, and bark feeding insects, apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. • Apply with ground equipment using sufficient water to obtain full coverage of target site. Apply in a minimum of 2 gallons per acre by air and a minimum of 10 gallons per acre by ground. • Do not apply more than 0.24 lb. a.i. (1.92 pts.) per acre per year. • Do not apply more than 0.04 lb a.i. (5.12 fl oz of product) per acre in a single application. <p style="text-align: center;">¹ Suppression only.</p>
CONIFER AND DECIDUOUS TREES: Seed Orchards	Coneworm spp. Seed Bug spp. Thrips spp.	See Remarks	See Remarks	<ul style="list-style-type: none"> • For high volume sprayers, dilute 5.12 fl. oz. per 100 gallons of water and apply 5-10 gallons of finished spray per tree. • For low volume sprayers, dilute 20 fl. oz. per 100 gallons of water and apply 100 gallons of finished spray per acre. • For aerial applications, apply 15 fl. oz. per acre in a minimum of 10 gallons finished spray per acre. • Do not apply as foliar broadcast application using a mechanically pressurized handgun on orchards. • Do not apply as a soil, drench, or ground-directed application using a mechanically pressurized handgun on orchards. • Do not apply more than 0.5 lb. a.i. (4 pts.) per acre per year.
NON-	See Crop Outlets on this	See Crop	See Crop	<ul style="list-style-type: none"> • Spray non-cropland adjacent to

CROPLAND (Excluding Public Land)	label for target pest and rates.	Outlets	Outlets	<p>agricultural areas to control migratory insects, which may threaten crops.</p> <ul style="list-style-type: none"> • Follow general use directions, rates, and spray recommendations found elsewhere in this label for the adjacent crop outlet and target pests. • Use highest labeled rates for dense/large foliage, high insect populations, and larger larval stages. • Repeat as necessary to maintain control. • Do not exceed 0.2 lb. a.i. (1.6 pts.) per acre per year. • Do not graze livestock in treated areas.
---	----------------------------------	---------	---------	---

Rate Conversion Chart

Lb. A.I. Per Acre	Fl. Oz. Per Acre	Pints Per Acre	Treated Acres Per Gallon
0.015	1.92	0.12	66
0.02	2.56	0.16	50
0.025	3.20	0.20	40
0.03	3.84	0.24	33
0.04	5.12	0.32	25

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill or leak on floor or paved surfaces, soak up with sand, earth, or synthetic absorbent. Remove to chemical waste area.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Disposal:

For Containers equal to or less than 5 Gallons : Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

For Containers greater than 5 Gallons : Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Refillable Container. Refill this container with Lambda-cyhalothrin only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

For Bulk, Mini-Bulk, EZ Handler® and Boomerang Container Disposal

Return container to point of purchase for reuse with seal intact and in salable condition.

Container Precautions

Before refilling RETURNABLE CONTAINERS, inspect thoroughly for damage such as cracks, punctures, bulges, dents, abrasions, and damaged or worn threads on closure devices.

After filling and before transporting, check for leaks.

Do not refill or transport damaged or leaking container.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of LG Chem, LTD., or Seller. To the extent consistent with applicable law all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold LG Chem, LTD., and Seller harmless for any claims relating to such factors.

LG Chem, LTD., warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or LG Chem, LTD., and Buyer and User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW LG CHEM, LTD., MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law in no event shall LG Chem, LTD., or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF LG CHEM, LTD., AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF LG CHEM, LTD., OR SELLER, THE REPLACEMENT OF THE PRODUCT.

LG Chem, LTD., and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of LG Chem, LTD.