

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION PESTICIDES AND TOXIC SUBSTANCES

SEP 2 7 2007

Jane M. Miller Agent for LG Life Sciences, Ltd. C/o BIOLOGIC, Inc. 115 Obtuse Hill Road Brookfield, CT 06804

SUBJECT:

Application for Pesticide Notification - Add Optional Statement; Add Statement

Inadvertently Removed from Export Only Section

MetaStar ST Seed Treatment Fungicide

EPA Reg. No. 71532-22

Application Dated September 7, 2007

Dear Ms. Miller:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Terri Stowe of my staff at 703-305-6117.

Sincerely,

Linda Arrington

Notifications & Minor Formulations Team Leader

(C. Holloman)

Registration Division (7505P) Office of Pesticide Programs

Please read instructions on r	everse before complet	tina form.			Form An	oroved	. OMB No.	2070-006	기 (180. Approval expires 2-28-9
⊕EPA	United States Environmental Protection Agen Washington, DC 20460					Registration Amendment Other		OPP Identifier Number	
	······································	Application	on for Pes	tici	le - Sect	tion	l		
1. Company/Product Number 71532-22				EPA I	Product Man	ager		3. P	roposed Classification
4. Company/Product (Name) MetaStar ST		,	PN 2						Nestricted
5. Name and Address of App LG Life Sciences, Ltd c/o BIOLOGIC, Inc. 1 Brookfield, CT 06804	•		(b) to E)(i), m : PA F		s sim	ilar or iden		n FIFRA Section 3(c)(3) omposition and labeling
			Section						
Amendment - Explain Resubmission in resp	onse to Agency letter	dated			Final printed Agency lett "Me Too" A	er dat Applica	ed ation.	N _C	IV 3, 2007 DTIFICATION SEP 2 7 2007
Explanation: Use addition Final label and notification to is consistent with the provisio confidential statement of form understand that if this notifica subject to enforcement action	add optional directions ns of PR Notice 98-10 a rula of this product. I un tion is not consistent wi	"See back pan and the EPA re nderstand that ith the terms of	el" on front legulations at 40 it is a violation FR Notice 98-	label a O CFR of 18 -10 an	152.46, and U.S.C. Sec. 1 d 40 CFR 152	no oth	er changes l willfully ma	nave been i ke any falsi	made to the labeling or the e statement to EPA. I further
1. Material This Product Will	Be Packaged In:						· · · · · · · · · · · · · · · · · · ·		
		No. per container	Yes No If "Yes" No. per			2. Type of	Metal Plastic Glass Paper Other (Specify)		
3. Location of Net Contents I	nformation ontainer	4. Size(s) Ret	tail Container			5. Lo	cation of La	bel Directi	ons
6. Manner in Which Label is		Lithog Paper Stenci	raph glued led		Other				
			Section	- 11	1				
1. Contact Point (Complete	items directly below fo	or identificatio	n of individua	l to be	contacted,	if nec	essary, to p	rocess this	s application.)
Name Jane M. Miller			Title Agent for LO	3 Life	Sciences, L	₋td.		Telephor (203) 74	ne No. (include Area Code) 0-1200
I certify that the stater I acknowledge that and both under applicable I	y knowlinglly false or i		all attachmen						6. Date Application Received (Stamped)
2. Signature	Ma		3. Title Agent for LG	Life S	ciences, Ltd.		 		
4. Typed Name Jane M. Mitter			5. Date September 7, 2007						

September 7, 2007

BIOLOGIC INC. agribusiness professionals

September 7, 2007

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
US Environmental Protection Agency
One Potomac Yard
2777 S. Crystal Drive
Room S-4900, 4th Floor
Arlington, VA 22202

NOTIFICATION SEP 2 7 2007

Attention:

Ms. Mary Waller (PM #21)

RE: 71532-22, "MetaStar ST"

Final Labeling

Dear Ms. Waller:

As per the EPA Notice of Pesticide Registration dated July 3, 2007, please find enclosed final labeling with the corrections made as outlined in the registration notice. We have also noted the following:

- 1. To the front panel, we have added the optional statement "See back panel and label booklet for additional precautionary information" which is necessary if the product will be labeled with a booklet.
- 2. On the approved label, we had inadvertently left off the sentence "Seed treated in accordance with the export use pattern must be labeled as follows:" on page 7 (under the table Seed Treatment Export Use). THIS SENTENCE MUST BE INCLUDED before the text "FOR EXPORT ONLY NOT FOR DOMESTIC SALES OR USE" so that it makes sense to the reader (user). This is the same text that appears on the cited label.

We have enclosed two (2) copies of the final labeling with the additional text underlined. Please stamp one copy and return to us so that we can use these for state registrations.

Should you have any questions or wish to reach me, please feel free to contact my office at (203) 740-1200.

Sincerely,

Jane M. Miller

Agent for LG Life Sciences, Ltd.

NOTIFICATION SEP 2 7 2007

MetaStar ST Seed Treatment Fungicide

[For use as a seed treatment for control of systemic downy mildew, Pythium seed rot, Pythium damping-off, and early season Phytophthora diseases of certain crops]

Active Ingredient:

Contains 2.65 lbs. Metalaxyl per gallon.

KEEP OUT OF REACH OF CHILDREN WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en datalle. (If you do not understand the label, find someone to explain it to you in detail.)

[See back panel and label booklet for additional precautionary information.]

Product of South Korea Formulated in the USA

LG Life Sciences, Ltd. 20 Yoido-dong Youngdungpo-gu Seoul 150-721, Korea

EPA Reg. No.: 71532-22

EPA Est. No.: 71532-KOR-001

Net Contents: 2.5 gal.

FIRST AID	
If on Skin or	• Take off contaminated clothing.
Clothing:	• Rinse skin immediately with plenty of water for 15-20 minutes.
	• Call a poison control center or doctor for treatment advice.
If in Eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes.
	• Remove contact lenses, if present, after the first 5 minutes, then continue
	rinsing eye.
	• Call a poison control center or doctor for treatment advice.
If Swallowed:	• Call a poison control center or doctor immediately for treatment advice.
	• Have person sip a glass of water if able to swallow.
	• Do not induce vomiting unless told to do so by a poison control center or
	doctor.
	• Do not give anything by mouth to an unconscious person.
Have the product of	container or label with you when calling a poison control center or doctor, or
going for treatmen	t. You may also contact the Poison Control Center at 1-800-222-1222.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals WARNING/AVISO

Causes skin irritation. Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Do not get on skin or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- · Coveralls over short-sleeved shirt and short pants
- · Chemical-resistant gloves such as nitrile, butyl, neoprene, and/or barrier laminate
- · Chemical-resistant footwear plus socks

Also, wear a chemical resistant apron when mixing or loading.

Follow manufacturer's instructions for cleaning/maintaining PPE. If not such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change clothing.

Environmental Hazards

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Groundwater Advisory Statement

This chemical is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Chemical resistant gloves such as barrier laminate or nitrile rubber or neoprene rubber or Viton.
- Chemical resistant footwear plus socks

In California - Only for use in commercial seed treating establishments. Not allowed for "on-farm" use.

NOTE: Seed treated commercially or for sale must be dyed an unnatural color using an EPA approved dye before sale.

Federal law requires that bags containing treated seed shall be labeled with the following information: "This seed has been treated with MetaStar ST fungicide. Do not use for feed, food or oil purposes. Store away from feeds and foodstuffs."

MetaStar ST is a systemic fungicide seed dressing specifically for control of systemic downy mildews, *Pythium* and *Phytophthora* spp. For control of other soilborne diseases, combination of Captan and Vitavax® registered formulations are compatible. Do not use with other seed treatment products unless previous experience assures compatibility.

MetaStar ST may be applied as a water-based slurry with other registered seed treatment insecticides and fungicides through standard slurry or mist-type commercial seed treatment equipment.

NOTE: MetaStar ST is a systemic fungicide having a specific mode of action. MetaStar ST could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted. Therefore, LG Life Sciences cannot assume liability for crop damage resulting from resistant strains of fungi. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible MetaStar ST resistant strains of fungi which may occur. FAILURE TO FOLLOW DIRECTIONS ON THIS LABEL MAY RESULT IN CROP INJURY OR POOR DISEASE CONTROL.

SEED TREATMENT

SEED	INSTRUCTIONS FOR USE
Canola*	For Pythium damping-off control: Apply MetaStar ST as a seed treatment
Mustard Seed*	at the rate of 0.25-0.5 fl. oz. per 100 lbs. of seed.
Rapeseed*	
	Use Restrictions: Do not harvest mustard greens. Rape greens grown and
	harvested from MetaStar ST treated seed must not be used for human and
	feed consumption. Rapeseed grown and harvested from MetaStar ST
	treated seed is only for industrial uses and cannot be used for edible oil or
	any other human/feed consumption.
	*Not registered for use in California.
Carrots*	For Pythium damping-off control: Apply MetaStar ST as a seed treatment
	at the rate of 0.7 fl. oz. per 100 lbs. of seed. *Not registered for use in
	California.
Cotton	For Pythium seed rot and damping-off control: Apply MetaStar ST as a
	seed treatment at the rate of 0.75-1.5 fl. oz. per 100 lbs. of seed.
Soybeans	For Pythium damping-off and early season Phytophthora control: Apply
	MetaStar ST as a seed treatment at the rate of 0.75-1.5 fl. oz. per 100 lbs. of
	seed.
Spinach*	For Pythium damping-off control: Apply MetaStar ST as a seed treatment
,	at the rate of 0.75 fl. oz. per 100 lbs. of seed. Do not apply MetaStar ST to
	spinach which is destined to be grown in the greenhouse. Containers or bags
·	containing MetaStar ST-treated spinach seed must be labeled with this
	restriction. *Not registered for use in California.
Peanuts*	For Pythium seed rot and damping-off control: Apply MetaStar ST as a
	seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed. *Not registered
	for use in California.
Sorghum	For control of <i>Pythium</i> damping-off apply 0.375-0.75 fl. oz. of MetaStar ST

	Final Labeling 9///0/
	per 100 lbs. of seed applied as a seed treatment. For control of systemic downy mildew (races susceptible to chemical control), apply 0.75-1.5 fl. oz. MetaStar ST per 100 lbs. of seed on cultivars which possess sorghum downy mildew resistance. To those cultivars which do not possess resistance to a particular race of sorghum downy mildew in heavily infected areas with recent severe history, apply 1.5-3.0 fl. oz. of MetaStar ST per 100 lbs. of seed as a seed treatment. Only high quality sorghum seed (90% germination and above) should be treated with MetaStar ST .
Sunflowers	For control of systemic downy mildew: Apply MetaStar ST as a seed treatment at the rate of 1.5-3.0 fl. oz. per 100 lbs. of seed.
Beets, Garden Sugarbeets	For <i>Pythium</i> damping-off control: Apply MetaStar ST as a seed treatment at the rate of 0.7 fl. oz. per 100 lbs. of seed.
Forage Legumes	Alfalfa, clover, lespedeza, beans (for forage), soybeans (for forage), soybean hay, peanuts (for forage), peanut hay, peas (for forage), pea vine hay, cowpeas (for forage), cowpea hay, trefoil, vetch and velvet beans (for forage). For <i>Pythium</i> damping-off and early season <i>Phytophthora</i> control: Apply MetaStar ST as a seed treatment at the rate of 0.75-1.5 fl. oz. per 100 lbs. of seed.
Grain Crops	Wheat, barley, rye, oats, millet, milo, corn (field corn, sweet corn and popcorn), buckwheat and triticale. For <i>Pythium</i> damping-off control: Apply MetaStar ST as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed. For systemic downy mildew
	control of sweet corn: Apply MetaStar ST as a seed treatment at the rate of 1.5 fl. oz. per 100 lbs. of seed.
Rice*	For <i>Pythium</i> damping-off control: Apply MetaStar ST as a seed treatment at the rate of 0.75-1.5 fl. oz. per 100 lbs. of seed. If using MetaStar ST treated rice do not use metalaxyl or mefenoxam for control of <i>Pythium</i> in water-seeded rice. *Not registered for use in California.
Seed and Pod Vegetables	Black-eyed peas, chickpeas (Garbanzos), cowpeas, dill, edible soybeans, field beans, field peas, garden peas, green beans, kidney beans, lima beans, lupines, navy beans, okra, peas, pole beans, snap beans, spring beans, wax beans, and lentils.
	For <i>Pythium</i> damping-off control: Apply MetaStar ST as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed. For early season <i>Phytophthora</i> control: Apply MetaStar ST as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed. For control of <i>Pythium</i> damping-off and systemic downy mildew of peas: Apply MetaStar ST as a seed treatment at the rate of 1.5 fl. oz. per 100 lbs. of seed.
Cucumbers*	For <i>Pythium</i> damping-off control: Apply MetaStar ST as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed. *Not registered for use in California.
Turf Grasses	For <i>Pythium</i> damping-off control on golf courses, home lawns reclamation and erosion control projects: Apply MetaStar ST as a seed treatment at the

	rate of 1.5 fl. oz. per 100 lbs. of seed.
Forage Grasses	Grasses grown for hay, grazing or silage corn (fodder or silage), sorghum (hay or silage), and small grain grown for hay, grazing or silage.
	For <i>Pythium</i> damping-off control: Apply MetaStar ST as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed.

SEED TREATMENT-EXPORT USE

For treatment of the following seed for export only to countries requiring treatment with metalaxyl before importation.

metalaxyl before i	
SEED	INSTRUCTIONS FOR USE
Brassica (Cole)	For Pythium damping-off control: Apply MetaStar ST as a seed treatment
Leafy Vegetables	at the rate of 0.75-1.5 fl. oz. per 100 lbs. of seed.
Canola	For Pythium damping-off control: Apply MetaStar ST as a seed treatment
	at the rate of 0.5-1.7 fl. oz. per 100 lbs. of seed (32-110 ml per 100 kg of
	seed).
Corn	For Pythium damping-off control: Apply MetaStar ST as a seed treatment
	at the rate of 0.75-1.5 fl. oz. per 100 lbs. of seed. For systemic downy
	mildew control: Apply MetaStar ST as a seed treatment at 3.0 fl. oz. per 100
	lbs. of seed.
Cucurbit	For Pythium damping-off control: Apply MetaStar ST as a seed treatment
Vegetables	at the rate of 0.75 fl. oz. per 100 lbs. of seed.
Fruiting	For Pythium damping-off control: Apply MetaStar ST as a seed treatment
Vegetables	at the rate of 0.75-1.5 fl. oz. per 100 lbs. of seed.
Leafy Vegetables	For Pythium damping-off control: Apply MetaStar ST as a seed treatment
	at the rate of 0.75-1.5 fl. oz. per 100 lbs. of seed.
Onions	For Pythium damping-off control: Apply MetaStar ST as a seed treatment
	at the rate of 0.75-1.5 fl. oz. per 100 lbs. of seed.
Millet	For Pythium damping-off control: Apply MetaStar ST as a seed treatment
	at the rate of 0.75-1.5 fl. oz. per 100 lbs. of seed. For systemic downy
	mildew control: Apply MetaStar ST as a seed treatment at 2.5 fl. oz. per 100
	lbs. of seed.
Peas	For Pythium damping-off and early season Phytophthora control: Apply
	MetaStar ST as a seed treatment at the rate of 3.5 fl. oz. per 100 lbs. of seed.
Root and Tuber	For Pythium damping-off control: Apply MetaStar ST as a seed treatment
Vegetables	at the rate of 0.75-1.5 fl. oz. per 100 lbs. of seed.
	Do not apply MetaStar ST to potatoes.
Sorghum	For Pythium damping-off control: Apply MetaStar ST as a seed treatment
	at the rate of 0.75-1.5 fl. oz. per 100 lbs. of seed. For systemic downy
	mildew control: Apply MetaStar ST as a seed treatment at 5.25 fl. oz. per
	100 lbs. of seed.
Sunflowers	For systemic downy mildew control: Apply MetaStar ST as a seed
	treatment at 9.0 fl. oz. per 100 lbs. of seed.

Seed treated in accordance with the export use pattern must be labeled as follows: FOR EXPORT ONLY – NOT FOR DOMESTIC SALES OR USE

DIRECTIONS FOR USE AT REDUCED RATES IN COMBINATION WITH OTHER FUNGICIDES

MetaStar ST is a systemic fungicide seed dressing which may be applied in combination with other EPA-registered fungicides to aid in control of *Pythium*. **MetaStar ST** may be applied as a water-based slurry through standard slurry of mist-type commercial seed treatment equipment.

SEED	INSTRUCTIONS FOR USE
Corn (field), Wheat,	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> :
Oats, Barley, Rye	Apply MetaStar ST as a commercial seed treatment at the rate of 0.10-
	0.375 fl. oz. per 100 lbs. of seed. Apply only in combination with EPA
	registered rates of seed treatment fungicides.
Soybeans, Field	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> :
beans, Kidney beans,	Apply MetaStar ST as a commercial seed treatment at the rate of 0.10-
Navy beans, Pinto	0.375 fl. oz. per 100 lbs. of seed. Apply only in combination with EPA
beans, Lentils, Peas	registered rates of broad-spectrum seed treatment fungicides.
Sunflowers	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> :
	Apply MetaStar ST as a commercial seed treatment at the rate of 0.10-
	0.375 fl. oz. per 100 lbs. of seed. Apply only in combination with EPA
	registered rates of broad-spectrum seed treatment fungicides.
Peanuts*	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> :
	Apply MetaStar ST as a commercial seed treatment at the rate of 0.10-
	0.375 fl. oz. per 100 lbs. of seed. Apply only in combination with EPA
	registered rates of broad-spectrum seed treatment fungicides. *Not
	registered for use in California.
Rice*	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> :
	Apply MetaStar ST as a commercial seed treatment at the rate of 0.10-
	0.375 fl. oz. per 100 lbs. of seed. Apply only in combination with EPA
	registered rates of broad-spectrum seed treatment fungicides. *Not
	registered for use in California.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. **Pesticide Storage:** Store at temperatures above 40° F. Crystals may form at lower storage temperatures. If this occurs, place the product in a warm room (68°F or above) and roll or shake the container at frequent intervals until all crystals are dissolved.

For minor spills, leaks, etc. follow all precautions indicated on this label and clean-up immediately. Take special care to avoid contamination of equipment and facilities during clean-up procedures and disposal of wastes.

Pesticide Disposal: Wastes resulting from the use of this product are toxic. Improper disposal of unused pesticide spray mixture, or rinsate is a violation of Federal law. Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures. For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office.

Container Disposal: Do not reuse empty container. Triple rinse (or equivalent) and offer for

recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Returnable/Refillable Containers: Do not rinse container. Do not empty remaining formulated product. Do not break seals. Return intact to point of purchase.

Conditions of Sale and Limitation of Warranty and Liability

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying and using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of LG Life Sciences, Ltd. or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold LG Life Sciences, Ltd. and Seller harmless for any claims relating to such factors.

LG Life Sciences, Ltd. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or LG Life Sciences, Ltd. and Buyer and User assume the risk of any such use. To the extent permitted by law, LG Life Sciences, Ltd. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, LG Life Sciences, Ltd. or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT PERMITTED BY LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF LG LIFE SCIENCES, LTD. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF LG LIFE SCIENCES, LTD. OR SELLER, THE REPLACEMENT OF THE PRODUCT.

LG Life Sciences, Ltd. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty of liability, which may not be modified except by written agreement signed by a duly authorized representative of LG Life Sciences, Ltd.