



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs

Registration Division (7505T)

1200 Pennsylvania Ave., N.W.

Washington, D.C. 20460

NOTICE OF PESTICIDE:

☒ Registration

☐ Reregistration

(under FIFRA, as amended)

EPA Reg. Number:

71532-44

Date of Issuance:

11/6/25

Term of Issuance:

Unconditional

Name of Pesticide Product:

GlamPrime

Name and Address of Registrant (include ZIP Code):

LG Chem Ltd
c/o Ag-Chem Consulting
12644 Chapel Road
Clifton, VA 20124

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

Continues page 2

Signature of Approving Official:

Emily Schmid

Emily Schmid, Product Manager 25
Herbicide Branch, Registration Division (7505T)

Date:

11/6/25

2. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 71532-44."
3. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The record for this product currently contains the following CSF(s):

- Basic CSF dated 09/30/2024

If you have any questions, please contact Margaret Golembiewski at (202)566-0304 or at golembiewski.margaret@epa.gov.

Enclosure

GlamPrime

Herbicide for Glyphosate Resistant Crops. Selective broad-spectrum weed control in glyphosate resistant crops. Non-selective, broad-spectrum weed control for many cropping systems, farmsteads, and Conservation Reserve Program acres.

Active Ingredient:

Glyphosate, N-(phosphonomethyl)-glycine, as isopropylamine salt* 41.0%

Other Ingredients: 59.0%

Total: 100.0%

*Contains 480 grams per liter or 4 pounds per U.S. gallon of the active ingredient glyphosate, in the form of isopropylamine salt. Equivalent to 358 grams per liter or 3 pounds per U.S. gallon of the acid, glyphosate.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID	
If In Eyes	<ul style="list-style-type: none"> - Hold eye open and rinse slowly and gently with water for 15-20 minutes. - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. - Call a poison control center or doctor for treatment advice.
If Swallowed	<ul style="list-style-type: none"> - Call a poison control center or doctor immediately for treatment advice. - Have person sip a glass of water if able to swallow. - Do not induce vomiting unless told to do so by the poison control center or doctor. - Do not give anything by mouth to an unconscious person.
If Inhaled	<ul style="list-style-type: none"> - Move person to fresh air. - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. - Call a poison control center or doctor for further treatment advice.
If on Skin or Clothing	<ul style="list-style-type: none"> - Take off contaminated clothing. - Rinse skin immediately with plenty of water for 15-20 minutes. - Call a poison control center or doctor for treatment advice.
Emergency Numbers For 24-hour medical emergency assistance (human or animal) call 1-800-222-1222. For chemical emergency assistance (spill, leak, fire, accident) call ChemTrec at 1-800-424-9300.	
Have the product label with you when calling a poison control center, or doctor, or going for treatment.	

[Optional referral statements when booklets and container labels are used:]

[See label booklet for [complete] [additional] [First Aid,] [Precautionary Statements], [Directions For Use], and [Storage and Disposal].]

Manufactured [For][By]:

LG Chem Ltd.


128 Yeoui-daero, Yeongdeungpo-gu

Seoul, Korea 07336

EPA Reg. No. 71532-

EPA Est. No.

Net Contents:



PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Causes moderate eye irritation. Avoid contact with eyes or clothing.

Not all products recommended on this label are registered for use in California. Check the registration status of each product in California before using. Read the entire label before using this product. Use only according to label instructions. **AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS (EXCEPT AS SPECIFIED FOR INDIVIDUAL GLYPHOSATE RESISTANT CROPS), DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.**

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Protective eyewear
- Shoes plus socks

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with GlamPrime concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning of equipment or disposing of equipment washwaters or rinsate.

PHYSICAL OR CHEMICAL HAZARDS

Spray solutions of **GlamPrime** should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers. **DO NOT MIX, STORE OR APPLY GLAMPRIE OR SPRAY SOLUTIONS OF GLAMPRIE IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.** **GlamPrime** or spray solutions of **GlamPrime** react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette, or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal Law to use **GlamPrime** in any manner inconsistent with its labeling.

Do not apply **GlamPrime** in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

AGRICULTURAL USE REQUIREMENTS

Use **GlamPrime** only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of **GlamPrime** that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical Resistant Gloves Category A such as butyl rubber, neoprene rubber, natural rubber, or nitrile rubber >14 mils
- Protective eyewear
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Keep people and pets off treated areas until spray solution has dried to prevent transfer of this product onto desirable vegetation.

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, **GlamPrime** is a Group 9 herbicide. Any weed population may contain or develop plants naturally resistant to **GlamPrime** and other Group 9 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of **GlamPrime** or other Group 9 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixture with herbicides from a different group if such use is permitted: where information on resistance in target weed species is available use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.

PRODUCT INFORMATION (How this product works)

Product Description: This product is a post-emergent, systemic herbicide with no soil residual activity. It is non-selective and gives broad-spectrum control of many annual weeds, perennial weeds, woody brush and trees. It is formulated as a water-soluble liquid. It may be applied through most standard industrial or field-type sprayers after dilution and thorough mixing with water or other carriers according to label instructions.

No additional surfactant in the spray solution is needed or recommended. This includes additives containing surfactants, buffering agents or pH adjusting agents when **GlamPrime** is the only pesticide used unless otherwise directed. Ammonium sulfate, drift control additives, or dyes and colorants may be used. See the "MIXING" section of this label for instructions.

Time to Symptoms: This product moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days, but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay development of visual symptoms. Visible effects are a gradual wilting and yellowing of the plant that advances to complete browning of above-ground growth and deterioration of underground plant parts.

Stage of Weeds: Annual weeds are easiest to control when they are small. Best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity. Refer to the “**ANNUAL WEEDS**”, “**PERENNIAL WEEDS**”, and “**WOODY BRUSH AND TREES RATE TABLES**” for recommendations for specific weeds.

Always use the higher rate of this product per acre within the recommended range when weed growth is heavy or dense or weeds are growing in an undisturbed (non-cultivated) area.

Do not treat weeds under poor growing conditions such as drought stress, disease or insect damage, as reduced weed control may result. Reduced results may also occur when treating weeds heavily covered with dust.

Cultural Considerations: Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed, or cut, and have not been allowed to regrow to the recommended stage for treatment.

Rainfastness: Heavy rainfall soon after application may wash this product off of the foliage and a repeat application may be required for adequate control.

Spray Coverage: For best results, spray coverage should be uniform and complete. Do not spray weed foliage to the point of runoff.

Mode of Action: The active ingredient in this product inhibits an enzyme found only in plants and microorganisms that is essential to formation of specific amino acids.

No Soil Activity: Weeds must be emerged at the time of application to be controlled by this product. Weeds germinating from seed after application will not be controlled. Unemerged plants arising from unattached underground rhizomes or rootstocks of perennials will not be affected by the herbicide and will continue to grow.

Biological Degradation: Degradation of this product is primarily a biological process carried out by soil microbes.

Tank Mixing: This product does not provide residual weed control. For subsequent residual weed control, follow a label approved herbicide program. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Mixing this product with herbicides or other materials not recommended on this label may result in reduced performance.

Annual Maximum Use Rate: Except as otherwise specified in a crop section of this label, the combined total of all treatments must not exceed 8 quarts of this product (6 lb of glyphosate) per acre per year. For applications in non-crop sites or in tree, vine, or shrub crops, the combined total of all treatments must not exceed 10.6 quarts of this product (7.95 lb of glyphosate) per acre per year. The maximum use rates stated throughout this product's labeling apply to this product combined with the use of all other herbicides containing glyphosate or sulfosate as the active ingredient, whether applied as mixtures or separately. Calculate the application rates and ensure that the total use of this and other glyphosate or sulfosate containing products does not exceed stated maximum use rate. **NOTE:** Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences.

MIXING

Clean sprayer parts immediately after using this product by thoroughly flushing with water.

NOTE: REDUCED RESULTS MAY OCCUR IF WATER CONTAINING SOIL IS USED, SUCH AS VISIBLY MUDDY WATER OR WATER FROM PONDS AND DITCHES THAT IS NOT CLEAR.

Mixing with Water

This product mixes readily with water. Mix spray solutions of this product as follows: Fill the mixing or spray tank with the required amount of water. Add the specified amount of this product near the end of the filling process and mix well. Use caution to avoid siphoning back into the carrier source. Use approved anti-back-siphoning devices where required by state or local regulations. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, avoid the use of mechanical agitators, terminate by-pass and return lines at the bottom of the tank and, if needed, use an approved anti-foam or defoaming agent.

Tank Mixing Procedure

Mix labeled tank mixtures of this product with water as follows:

1. Place a 20- to 35-mesh screen or wetting basket over filling port.
2. Through the screen, fill the spray tank one-half full with water and start agitation.
3. If ammonium sulfate is used add it slowly through the screen into the tank. Continue agitation. Ensure that dry ammonium sulfate is completely dissolved in the spray tank before adding other products.

4. If a wettable powder is used, make a slurry with the water carrier, and add it SLOWLY through the screen into the tank. Continue agitation.
5. If a flowable formulation is used, premix one part flowable with one part water. Add diluted mixture SLOWLY through the screen into the tank. Continue agitation.
6. If an emulsifiable concentrate formulation is used, premix one part emulsifiable concentrate with two parts water. Add diluted mixture slowly through the screen into the tank. Continue agitation.
7. Continue filling the spray tank with water and add the required amount of this product near the end of the filling process.
8. Add individual formulations to the spray tank as follows: wettable powder, flowable, emulsifiable concentrate, drift control additive, and water soluble liquid.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed. Keep by-pass line on or near the bottom of the tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50 mesh. Always predetermine the compatibility of labeled tank mixtures of this product with water carrier by mixing small proportional quantities in advance.

Refer to the “**Tank Mixing**” section of “**PRODUCT INFORMATION**” for additional precautions.

Mixing for Hand-Held Sprayers

Prepare the desired volume of spray solution by mixing the amount of this product in water as shown in the following table:

Spray Solution						
Volume	Desired Amount of GlamPrime					
	1/2%	1%	1 1/2%	2%	5%	10%
1 gal.	2/3 oz.	1 1/3 oz.	2 oz.	2 2/3 oz.	6 1/2 oz.	13 oz.
25 gals.	1 pt.	1 qt.	1 1/2 qts.	2 qts.	5 qts.	10 qts.
100 gals.	2 qts.	1 gal.	1 1/2 gals.	2 gals.	5 gals.	10 gals.
2 tablespoons = 1 fluid ounce						

For use in knapsack sprayers, it is suggested that the specified amount of this product be mixed with water in a larger container. Fill sprayer with the mixed solution.

Ammonium Sulfate

The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of this product, particularly under hard water conditions, drought conditions or when tank mixed with certain residual herbicides, on annual and perennial weeds. The equivalent rate of ammonium sulfate in a liquid formulation may also be used. Ensure that dry ammonium sulfate is completely dissolved in the spray tank before adding herbicides. Thoroughly rinse the spray system with clean water after use to reduce corrosion.

NOTE: When using ammonium sulfate, apply this product at rates recommended in this label. Lower rates will result in reduced performance.

Colorants or Dyes

Agriculturally approved colorants or marking dyes may be added to this product. Colorants or dyes used in spray solutions of this product may reduce performance, especially at lower rates or dilutions. Use colorants or dyes according to the manufacturer's recommendations.

Drift Control Additives

Drift control additives may be used with all equipment types, except wiper applicators, sponge bars and Controlled Droplet Applicator (CDA) equipment. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label. The use of drift control additives can affect spray coverage, which may result in reduced performance.

APPLICATION EQUIPMENT AND TECHNIQUES

Do not apply this product through any type of irrigation system.

Do not apply when winds are gusty or under any other condition that favors drift.

This product may be applied with the following application equipment:

Aerial - Fixed Wing and Helicopter.

Ground Broadcast Spray - Boom or boomless systems, pull-type sprayer, floaters, pick-up sprayers, spray coupes, and other ground broadcast equipment.

Hand-Held or High-Volume Spray Equipment - Knapsack and backpack sprayers, pump-up pressure sprayers, handguns, handwands, mistblowers*, lances and other hand-held and motorized spray equipment used to direct the spray onto weed foliage.

*This product is not registered in California or Arizona for use in mistblowers.

Selective Equipment - Recirculating sprayers, shielded and hooded sprayers, wiper applicators, and sponge bars.

Injection Systems - Aerial or ground injection sprayers.

Controlled Droplet Applicator (CDA) - Hand-held or boom-mounted applicators which produce a spray consisting of a narrow range of droplet sizes.

APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES. DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION THAT FAVORS DRIFT.

Aerial Equipment

DO NOT APPLY THIS PRODUCT USING AERIAL SPRAY EQUIPMENT EXCEPT UNDER CONDITIONS AS SPECIFIED WITHIN THIS LABEL.

Use the recommended rates of this herbicide in 3 to 15 gallons of water per acre unless otherwise specified on this label. Unless otherwise specified, do not exceed 1 quart per acre. Refer to the individual use area sections of this label for recommended volumes, application rates, and further instructions. This product plus dicamba tank mixtures may not be applied by air in California.

Ensure uniform application - To avoid streaked, uneven or overlapped application, use appropriate marking devices.

AERIAL SPRAY DRIFT MANAGEMENT

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops:

1. The distance of the outermost nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.

Nozzles must always point backward, parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they should be observed.

Importance of Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see the “Wind”, “Temperature and Humidity”, and “Temperature Inversions” sections of this label).

Controlling Droplet Size

- **Volume:** Use high-flow-rate nozzles to apply the highest practical spray volume. Nozzles with the higher rated flows produce larger droplets.
- **Pressure:** Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher-flow-rate nozzles instead of increasing pressure.
- **Number of nozzles:** Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle orientation:** Orienting nozzles so that the spray is released backwards, parallel to the airstream, will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- **Nozzle type:** Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.
- **Boom length:** For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.
- **Application height:** Applications must not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces the exposure of the droplets to evaporation and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller droplets, etc.).

Wind

Drift potential is lowest between wind speeds of 2 to 10 miles per hour. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application must be avoided below 2 miles per hour due to variable wind direction and high inversion potential.

NOTE: Local terrain can influence wind patterns. Every applicator must be familiar with local wind patterns and how they affect drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications must not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sunsets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

The product must only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e. g., when wind is blowing away from the sensitive areas).

Avoid direct application to any body of water.

Aircraft Maintenance

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR IS MOST SUSCEPTIBLE. The maintenance of an organic coating (paint), which meets aerospace specification MIL-C-38413, may prevent corrosion.

FOR AERIAL APPLICATION IN FRESNO COUNTY, CALIFORNIA ONLY

See “**PRODUCT INFORMATION**”, “**MIXING**”, “**APPLICATION EQUIPMENT AND TECHNIQUES**”, and “**SPRAY DRIFT MANAGEMENT**” sections of the label booklet for essential product information prior to making aerial application.

See “**CROPS**” section of the label booklet for specific directions on the use of this product.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS (EXCEPT AS SPECIFIED FOR INDIVIDUAL GLYPHOSATE RESISTANT CROPS), DESIRABLE PLANTS AND TREES BECAUSE SEVERE INJURY IS LIKELY TO RESULT.

FOR FRESNO COUNTY, CA ONLY **From February 15th through March 31st only**

For aerial application outside these dates, refer to the above section of this label.

This section only applies to the area contained inside the following boundaries within Fresno County, California only.

North: Fresno County line
South: Fresno County line
East: State Highway 99
West: Fresno County line

Use Information

Always read and follow the label directions and precautionary statements for all product used in the aerial application.

Observe the following directions to minimize off-site movement during aerial application of **GlamPrime**. Minimization of offsite movement is the responsibility of the grower, Pest Control Advisor, and aerial applicator.

Written Recommendations

A written recommendation **MUST** be submitted by or on behalf of the applicator to the Fresno County Agricultural Commissioner 24 hours prior to application. This written recommendation **MUST** state the proximity of surrounding crops, and that conditions of each manufacturer's applicable product label(s) and this label have been satisfied.

Aerial Applicator Training and Equipment

Aerial application of **GlamPrime** is limited to pilots who have successfully completed a Fresno County Agricultural Commissioner and California Department of Pesticide Regulation approved training program for aerial application of herbicides. All aircraft must be inspected, critiqued in flight, and certified at a Fresno County Agricultural Commissioner approved fly-in. Test and calibrate spray equipment at intervals sufficient to ensure that proper rates of herbicides and adjuvants are being applied during commercial use.

Applicator must document such calibrations and testing. Demonstration of performance at Fresno County Agricultural Commissioner approved "fly-in" constitutes such documentation, or other written records showing calculations and measurements of flight and spray parameters acceptable to the Fresno County Agricultural Commissioner.

Application at Night - Do not apply this product by air earlier than 30 minutes prior to sunrise and/or later than 30 minutes after sunset without prior permission from the Fresno County Agricultural Commissioner.

FOR AERIAL APPLICATIONS IN MISSISSIPPI

The distance of the outermost nozzles on the boom must not exceed 75% of the length of the wingspan or rotor. In many cases, reducing this distance to 65% of the length of the wingspan or rotor will improve drift control without affecting the swath width.

Nozzles must always discharge backward parallel with the air stream and never discharge downwards more than 45 degrees on fixed wing aircraft or forward of the prevailing air flow on rotary winged aircraft. Avoid the use of nozzles with wide-angle discharge. Do not apply this product when winds are in excess of 10 miles per hour.

Do not apply this when there is a low level inversion where fine spray particles could be suspended in still air and move outside the target area when the inversion layer moves. These conditions may occur when wind speeds are less than 2 mph.

Use the following guidelines when applications are made near crops or other desirable vegetation:

1. Do not apply within 100 feet of any desirable vegetation or crops.
2. If wind up to 5 miles per hour is blowing toward desirable vegetation or crops, do not apply within 500 feet upwind of the desirable vegetation or crops.
3. Winds blowing from 5 to 10 miles per hour toward desirable vegetation or crops will likely require buffer zones in excess of 500 feet.

Aerial Application Restrictions:

Aerial application is prohibited in Zone I, south of Highway 8 in the counties listed below, from March 15th through April 30th, except by permit from an authorized employee of the Mississippi Department of Agriculture and Commerce, Bureau of Plant Industry (Ph. 1-888-257-1285).

Aerial application is prohibited in Zone II, north of Highway 8 in the counties listed below, from March 25th through April 30th, except by permit from an authorized employee of the Mississippi Department of Agriculture and Commerce, Bureau of Plant Industry (Ph. 1-888-257-1285).

The Bureau of Plant Industry may at anytime, based on current planting and environmental conditions, modify the above restrictions for either zone or county therein

Zone I: South of Highway 8 in the counties of Bolivar, Sunflower, Leflore, and Grenada plus the entire counties of Carroll, Holmes, Humphreys, Washington, Sharkey, Issaquena, Yazoo, and Warren.

Zone II: North of Highway 8 in the counties of Bolivar, Sunflower, Leflore, and Grenada plus the entire counties of Tallahatchie, Tate, Quitman, Coahoma, Tunica, Panola, and Desoto.

FOR AERIAL APPLICATION IN ARKANSAS ONLY

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See “**PRODUCT INFORMATION**” and “**Mixing**” sections of the label booklet for essential product performance information.

USE DIRECTIONS

Avoid drift. Do not apply into still air where there is a temperature inversion layer low enough for fine spray particles to become suspended and move outside the target area when the inversion layer moves. Do not apply when winds are gusty or under any other condition that favors drift. Drift is likely to cause damage to any vegetation contacted. To prevent injury to adjacent desirable vegetation, appropriate buffer zones must be maintained.

Use the specified rate of this product in 3 to 15 gallons of water per acre.

Use sufficient carrier volume and appropriate equipment set-up to form droplets large enough to avoid drift potential. Coarse droplets in the 300 to 500 (VMD) micron range are recommended.

Applications should typically be made with the nozzle release point at 8 to 15 feet above the top of the target plants unless a greater height is required for aircraft safety.

Ground Broadcast Equipment

Use the specified rates of this product in 3 to 40 gallons of water per acre as a broadcast spray unless otherwise specified. As density of weeds increases, spray volume should be increased within the recommended range to ensure complete coverage. Carefully select proper nozzles to avoid spraying a fine mist. For best results with ground application equipment, use flat spray nozzles. Check for even distribution of spray droplets.

Hand-Held or High-Volume Equipment

Apply to foliage of vegetation to be controlled. For applications made on a spray-to-wet basis, spray coverage should be uniform and complete. Do not spray to the point of runoff. Use coarse sprays only. For recommended rates and timing, refer to the “**ANNUAL WEEDS-HAND-HELD OR HIGH-VOLUME EQUIPMENT**” section of this product label.

Selective Equipment

This product may be applied through recirculating spray systems, shielded applicators, hooded sprayers, wiper applicators or sponge bars, after dilution and thorough mixing with water, to listed weeds growing in any non-crop site specified on this label. In cropping systems, hooded sprayers, shielded sprayers, and wipers may be used in row-middles (in between rows of crop plants) where any dripping or leaking will not contact crop foliage. Such equipment must be capable of preventing all crop contact with herbicide solutions and operated without leakage of spray mists or dripping onto crop. Wipers over-the-top of crops may be used only when specifically recommended in this product's labeling.

AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION.

Contact of the herbicide solution with desirable vegetation may result in damage or destruction. Applicators used above desirable vegetation should be adjusted so that the lowest spray stream or wiper contact point is at least 2 inches above the desirable vegetation. Droplets, mist, foam or splatter of the herbicide solution setting on desirable vegetation may result in discoloration, stunting or destruction.

Applications made above the crops should be made when the weeds are a minimum of 6 inches above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations or when the height of the weeds varies so that not all weeds are contacted. In these instances, repeat treatment may be necessary.

Recirculating Spray System

A recirculating spray system directs the spray solution onto weeds growing above desirable vegetation, while spray solution not intercepted by weeds is collected and returned to the spray tank for reuse.

Shielded and Hooded Applicators

When applied under the conditions described in the following paragraphs for shielded and hooded applications, this product at recommended rates will control those weeds listed in the “**ANNUAL WEEDS RATE TABLE**” and “**PERENNIAL WEEDS RATE TABLE**” sections of this label. A hooded sprayer is a type of shielded applicator where the spray pattern is fully enclosed including top, sides, front and back, thereby shielding the crop from the spray solution. Keep shields on these sprayers adjusted to protect

desirable vegetation. When applying to crops grown on raised beds, ensure that the hood is designed to completely enclose the spray solution. If necessary, extend the front and rear flaps of the hoods to reach the ground in deep furrows.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION.

This equipment must be set up and operated in a manner that avoids bouncing or raising the hoods off the ground in any way. If the hoods are raised, spray particles may escape and come into contact with the crop, causing damage or destruction of the crop. Avoid operation on rough or sloping ground where the spray hoods might be raised off the ground.

Use hoods designed to minimize excessive dripping or run-off down the insides of the hoods. A single, low pressure/ low drift flat-fan nozzle with an 80 to 95 degree spray angle positioned at the top center of the hood is recommended. Spray volume should be 20 to 30 gallons per acre. These procedures will reduce the potential for crop injury:

- The spray hoods must be operated on the ground or skimming across the ground.
- Leave at least an 8-inch untreated strip over the drill row. For example, if the crop row width is 38 inches, the maximum width of the spray hood should be 30 inches.
- Maximum tractor speed: 5 miles per hour to avoid bouncing of the spray hoods.
- Maximum wind speed: 10 miles per hour.
- Use low-drift nozzles that provide uniform coverage within the treated area.

Crop injury may occur when the foliage of treated weeds comes into direct contact with leaves of the crop. Do not apply this product when the leaves of the crop are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution may contact the crop and cause discoloration, stunting or destruction.

Wiper Applicators

When applied under the conditions described in the following paragraphs, this product **CONTROLS** many weeds, including volunteer corn, Texas panicum, common rye, shattercane, sicklepod, spanishneedles, and bristly starbur; and **SUPPRESSES** many weeds including Florida beggarweed, Bermudagrass, hemp dogbane, dogfennel, guineagrass, johnsongrass, milkweed, silverleaf nightshade, redroot pigweed, giant ragweed, smutgrass, sunflower, Canada thistle, musk thistle, vaseygrass, and velvetleaf.

Wiper applicators are devices that physically wipe appropriate amounts of this product directly onto the weed. Equipment must be designed, maintained and operated to prevent the herbicide solution from contacting desirable vegetation. Operate this equipment at ground speeds no greater than 5 miles per hour. Performance may be improved by reducing speed in areas of heavy weed infestations to ensure adequate wiper saturation.

Better results may be obtained if two applications are made in opposite directions. Avoid leakage or dripping onto desirable vegetation. Adjust height of applicator to ensure adequate contact with weeds. Keep wiping surfaces clean. Be aware that, on sloping ground, the herbicide solution may migrate, causing dripping on the lower end and drying of the wicks on the upper end of a wiper applicator.

Do not use wiper equipment when weeds are wet.

Mix only the amount of solution to be used during a 1-day period, as reduced activity may result from use of leftover solutions. Clean wiper parts immediately after using this product by thoroughly flushing with water.

Do not add surfactant to the herbicide solution.

For Rope or Sponge Wick Applicators - Mix 1 gallon of this product with 2 gallons of water to prepare 33 percent solution. Apply this solution to weeds listed above in this section.

For Panel Applicators - Solutions ranging from 33 to 100 percent of this product in water may be used in panel wiper applicators.

Injection Systems

This product may be used in aerial or ground injection spray systems. It may be used as a liquid concentrate or diluted prior to injecting into the spray stream. Do not mix this product with the concentrate of other products when using injection systems.

CDA Equipment

The rate of this product applied per acre by vehicle-mounted Controlled Droplet Application (CDA) equipment must not be less than the amount recommended in this label when applied by conventional broadcast equipment.

For vehicle-mounted CDA equipment, apply 2 to 15 gallons of water per acre. For the control of annual weeds with handheld CDA units, apply a 20 percent solution of this product at a flow rate of 2 fluid ounces per minute and a walking speed of 1.5 miles per hour (1 quart per acre). For the control of perennial weeds, apply a 20 to 40 percent solution of this product at a flow rate of 2 fluid ounces per minute and a walking speed of 0.75 miles per hour (2 to 4 quarts per acre). Controlled droplet application equipment produces a

spray pattern that is not easily visible. Extreme care must be exercised to avoid spray or drift contacting the foliage or any other green tissue of desirable vegetation, as damage or destruction may result.

ANNUAL AND PERENNIAL CROPS (Alphabetical)

NOTE: THIS SECTION GIVES DIRECTIONS THAT APPLY TO ALL LISTED CROPS WITHIN THIS SECTION GROUPED ALPHABETICALLY BELOW. SEE THE INDIVIDUAL CROP CATEGORIES FOR SPECIFIC INSTRUCTIONS, PREHARVEST INTERVALS, AND ADDITIONAL PRECAUTIONS AND RESTRICTIONS.

See the “GLYPHOSATE RESISTANT CROPS” section of this label.

TYPES OF APPLICATIONS: Chemical Fallow, Preplant Fallow Beds, Preplant, Pre-emergence, At-Planting, Hooded Sprayers in Row-Middles, Shielded Sprayers in Row-Middles, Wiper Applications in Row-Middles, and Post-Harvest Treatments.

USE INSTRUCTIONS: Apply this product during fallow intervals preceding planting, prior to planting or transplanting, at planting, or pre-emergent to annual and perennial crops listed in this label, except where specifically limited. For any crop not listed in this label, applications must be made at least 30 days prior to planting. Unless otherwise specified, weed control applications may be made according to the rates listed in the “**ANNUAL WEEDS**”, “**PERENNIAL WEEDS**”, and “**WOODY BRUSH AND TREES RATE TABLES**” in this label. Repeat applications may be made up to a maximum of 8 quarts (6lb of glyphosate) per acre per year.

Post-directed hooded sprayers and wiper equipment capable of preventing all crop contact with herbicide solutions may be used in mulched or unmulched row-middles after crop establishment. Where specifically noted below, wipers may also be used above certain crops to control tall weeds. Refer to the “Selective Equipment” section of this label for essential precautions when using hooded sprayers or wipers to avoid crop injury caused by leakage of spray mists or dripping onto crops. Crop injury is possible with these applications and shall be the sole responsibility of the applicator.

The maximum use rates stated throughout this product's labeling apply to this product combined with the use of all other herbicides containing glyphosate or sulfosate as the active ingredient, whether applied as mixtures or separately. Calculate the application rates and ensure that the total use of this and other glyphosate or sulfosate containing products does not exceed stated maximum use rate.

PRECAUTIONS: Avoid contact of herbicide with foliage, green shoots or stems, bark, exposed roots (including those emerging from plastic mulch), or fruit of crops because severe injury or destruction may result. Broadcast applications made at emergence will result in injury or death to emerged seedlings. Apply before seed germination in coarse sandy soils to further minimize the risk of injury. See “**APPLICATION EQUIPMENT AND TECHNIQUES**” section of this label for additional information. The crop receiving spray in treated area will be killed. Take care to avoid drift or spray outside the target area for the same reason.

RESTRICTIONS: When making pre-emergence and at-planting applications, applications must be made before crop emergence to avoid severe crop injury. Unless otherwise specified in this product's labeling, treatments with selective equipment including wipers and hooded sprayers must be made at least 14 days prior to harvest. Post-harvest or fallow applications must be made at least 30 days prior to planting any non-labeled crop. In crops where spot treatments are allowed, do not treat more than 10 percent of the total field to be harvested. For broadcast post-emergent treatments, do not harvest or feed treated vegetation for 8 weeks following application, unless otherwise specified. Unless otherwise specified, do not apply more than 1.5 quarts (1.13lb of glyphosate) per acre per application for control of annual weeds. Refer to “**ANNUAL WEEDS**” table for rates. Unless otherwise specified, do not apply more than 5 quarts (3.75lb of glyphosate) per acre per application for control of perennial weeds. Refer to the “**PERENNIAL WEEDS**” table for rates. Unless otherwise specified, do not apply more than 16 applications per year. Unless otherwise specified, do not apply more than 5 quarts (3.75lb of glyphosate) per acre per application for control of woody brush and trees. Refer to the “**WOODY BRUSH AND TREES**” table for rates.

Cereal and Grain Crops

LABELED CROPS: Barley, Buckwheat, Millet (pearl, proso), Oats, Rice, Rye, Quinoa, Teff, Teosinte, Triticale, Wheat (all types), and Wild Rice.

RESTRICTIONS: Do not treat rice fields or levees when field contains water. Do not apply more than 5 quarts (3.75lb of glyphosate) of this product per acre per single application, except for preharvest and red rice control applications. Do not apply more than 8 quarts (6lb of glyphosate) of this product per acre per year. When applying at reduced rates, do not apply more than 3 applications per year. Retreatment Interval (RTI): 7 days. Do not harvest or feed vegetation from an area for 8 weeks following broadcast application.

TYPES OF APPLICATIONS: Those listed in the “Annual and Perennial Crops” section plus the following: Red Rice Control Prior to Planting Rice, Spot Treatment (except Rice), Over-the-Top Wiper Applications (Feed Barley and Wheat only), and Preharvest (Feed Barley and Wheat only).

Preplant, Pre-emergence, At-Planting

USE INSTRUCTIONS: This product may be applied before, during or after planting of cereal crops. Applications must be made prior to emergence of the crop.

Red Rice Control Prior to Planting Rice

USE INSTRUCTIONS: Apply 1.5 quarts of this product in 5 to 10 gallons of water per acre. Flush fields prior to application to obtain uniform germination and stand of red rice. Make application when the majority of the red rice plants are in the 2-leaf stage and no more than 4 inches tall. Red rice plants with less than 2 true leaves may be only partially controlled.

PRECAUTIONS: Avoid spraying during low humidity conditions, as reduced control may result.

RESTRICTIONS: Do not treat rice fields or levees when the fields contain floodwater. Do not re-flood treated fields for 8 days following application. Retreatment Interval (RTI): 7 days. Do not flood treated fields for 8 days following application. Do not harvest or feed vegetation from an area for 8 weeks following broadcast application.

Spot Treatment (except Rice)

USE INSTRUCTIONS: This product may be applied as a spot treatment in cereal crops. Apply this product before heading in small grains.

PRECAUTIONS: The crop receiving spray in the treated area will be killed. Take care to avoid drift or spray outside target area for the same reason.

RESTRICTIONS: Do not treat more than 10 percent of the total field area to be harvested. Retreatment Interval (RTI): 7 days. Application must be made a minimum of 14 days prior to harvest.

Over-the-Top Wiper Applications (Feed Barley and Wheat only)

USE INSTRUCTIONS: Wiper applications may be used in feed barley and wheat. To control common rye or cereal rye, apply after the weeds have headed and achieved maximum growth, and when the rye is at least 6 inches above the wheat crop.

RESTRICTIONS: Allow at least 35 days between application and harvest. Do not use roller applicators. Retreatment Interval (RTI): 7 days. When applying at reduced rates, do not apply more than 3 applications per year.

Preharvest (Feed Barley and Wheat only)

USE INSTRUCTIONS: This product provides weed control when applied prior to harvest of wheat or feed barley. For wheat, apply after the hard-dough stage of grain (30 percent or less grain moisture). For feed barley, apply after the haddough stage and when the grain contains 20 percent moisture or less. Stubble may be grazed immediately after harvest. This product may be applied using either aerial or ground spray equipment. For ground applications, apply this product in 10 to 20 gallons of water per acre. For aerial applications, apply this product in 3 to 10 gallons of water per acre.

RESTRICTIONS: Do not apply more than 1 quart of this product (0.75lb glyphosate) per acre. Allow 7 days between application and harvest or grazing. Preharvest application is not recommended for wheat or barley grown for seed, as a reduction in germination or vigor may occur. When applying at reduced rates, do not apply more than 2 applications per year.

Post-Harvest

USE INSTRUCTIONS: This product may be applied after harvest of cereal crops. Higher rates may be required for control of large weeds, which were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

RESTRICTIONS: For any crop not listed on this label, applications must be made at least 30 days prior to planting the next crop. Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation. Retreatment Interval (RTI): 7 days. When applying at reduced rates, do not apply more than 2 applications per year.

Corn

TYPES OF CORN: Field Corn, Seed Corn, Silage Corn, Sweet Corn and Popcorn.

For Glyphosate Resistant Corn, see the Glyphosate Resistant crops section of this label

Preplant, Pre-emergence, At-Planting, Pre-harvest

USE INSTRUCTIONS: This product may be applied alone or in a tank-mixture before, during or after planting corn. Applications must be made prior to emergence of the crop.

TYPES OF APPLICATIONS: Chemical Fallow, Preplant Fallow Beds, Preplant, Pre-emergence, At-planting, Hooded Sprayers in Row-Middles, Shielded Sprayers in Row-Middles, Wiper Applications in Row-Middles, Preharvest, and Post-Harvest Treatments.

TANK MIXTURES: Apply these tank mixtures in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre.

2,4-D

Carfentrazone-ethyl

Atrazine

Dimethenamid

Dimethenamid-P and Atrazine

Rimsulfuron and thifensulfuron-methyl

Flufenacet and Metribuzin	Acetochlor
Isoxaflutole	Linuron
Dicamba	Dicamba and atrazine
Atrazine and S-metolachlor	Pendimethalin
Metalaxyl and Mancozeb	Flumetsulam
Acetochlor and Atrazine	Simazine
Sodium salt of diflufenzopyr and sodium salt of dicamba	Azoystrobin and Propiconazole
Pinoxaden	S-metolachlor

For difficult-to-control annual weeds such as fall panicum, barnyardgrass, crabgrass, shattercane, and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply this product at 2 pints per acre in these tank mixtures. For other labeled annual weeds, apply 1.5 to 2 pints of this product per acre when weeds are less than 6 inches tall, and 2 to 3 pints when weeds are over 6 inches tall. When using nitrogen solutions as the carrier, use rate may need to be increased for acceptable weed control.

RESTRICTIONS: Applications of 2,4-D or dicamba must be made at least 7 days prior to planting corn. For Southern states, do not apply in nitrogen solutions to tough-to-control grasses such as barnyardgrass, fall panicum, broadleaf signalgrass, annual ryegrass, and any perennial weeds. The area covered by this recommendation includes from Route 50 South in Illinois and Indiana and the following states: Alabama, Arkansas, Delaware, Florida, Georgia, Kentucky, Louisiana, Maryland, Mississippi, New Jersey, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia, and West Virginia. Retreatment Interval (RTI): 7 days. When applying at reduced rates, do not apply more than 3 applications per year. Do not harvest or feed vegetation from an area for 8 weeks following broadcast application.

Hooded Sprayers

USE INSTRUCTIONS: This product may be used through hooded sprayers for weed control between the rows of corn. Only hooded sprayers that completely enclose the spray pattern may be used. See additional instruction for the use of hooded sprayers in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

PRECAUTIONS: Contact of this product in any manner to any vegetation to which treatment is not intended may cause damage. Such damage shall be the sole responsibility of the applicator.

RESTRICTIONS: Corn must be at least 12 inches tall, measured without extending leaves. Do not apply more than 1 quart of this product (0.75 lb of glyphosate) per acre for each application and no more than 3 quarts (2.25lb of glyphosate) per acre per year for hooded sprayer applications. Do not apply more than 3 applications per year. Application must be made a minimum of 14 days prior to harvest. Retreatment Interval (RTI): 7 days.

Spot Treatment

USE INSTRUCTIONS: For spot treatments, apply this product prior to silking of corn.

PRECAUTIONS: The crop receiving spray in the treated area will be killed. Take care to avoid drift or spray outside target area for the same reason.

RESTRICTIONS: Do not treat more than 10 percent of the total field area to be harvested. Retreatment Interval (RTI): 7 days. Application must be made a minimum of 14 days prior to harvest. When applying at reduced rates, do not apply more than 3 applications per year.

Preharvest

USE INSTRUCTIONS: Make applications at 35 percent grain moisture or less. Ensure that maximum kernel fill is complete and the corn is physiologically mature (black layer formed). **RESTRICTIONS:** Allow a minimum of 7 days between application and harvest. Not to be used for preharvest application for corn grown for seed, as a reduction in germination or vigor may occur. When applying at reduced rates, do not apply more than 2 applications per year. Retreatment Interval (RTI): 7 days. For ground applications, apply up to 3 quarts of this product per acre. For aerial applications, apply up to 2 quarts of this product per acre.

Post-Harvest

USE INSTRUCTIONS: This product may be applied after harvest of corn. Higher rates may be required for control of large weeds which were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

RESTRICTIONS: Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation. When applying at reduced rates, do not apply more than 2 applications per year. Retreatment Interval (RTI): 7 days. Application of this product must be made a minimum of 30 days prior to planting any crop not listed on this label.

Cotton

TYPES OF APPLICATIONS: Those listed in the "Annual and Perennial Crops" section plus the following: Selective Equipment, Spot Treatment, Pre-harvest. For glyphosate-resistant Cotton, including glyphosate-resistant Flex Cotton, see the glyphosate-resistant section of this label.

Preplant, Pre-emergence, At-Planting

USE INSTRUCTIONS: This product may be applied before, during or after planting cotton. Applications must be made prior to emergence of the crop.

Hooded Sprayer, Selective Equipment

USE INSTRUCTIONS: This product may be applied through hooded sprayers, recirculating sprayers, shielded applicators, or wiper applicators in cotton. Allow at least 7 days between application and harvest. **PRECAUTIONS:** See the "Selective Equipment" part of the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for information on proper use and calibration of this equipment.

Spot Treatment

USE INSTRUCTIONS: For spot treatments, apply this product prior to boll opening of cotton.

PRECAUTIONS: The crop receiving spray in treated area will be killed. Take care to avoid drift or spray outside target area for the same reason.

RESTRICTIONS: Do not treat more than 10 percent of the total field area to be harvested. When applying at reduced rates, do not apply more than 3 applications per year. Retreatment Interval (RTI): 7 days.

Pre-Harvest

USE INSTRUCTIONS: This product provides weed control and cotton regrowth inhibition when applied prior to harvest of cotton. For weed control, apply at rates given in the "ANNUAL WEEDS", "PERENNIAL WEEDS", and "WOODY BRUSH AND TREES RATE TABLES" sections of this label. For cotton regrowth inhibition, apply 1 pint to 2 quarts of this product (0.375- 1.5lb of glyphosate) per acre. Up to 2 quarts of this product (1.5lb of glyphosate) may be applied using either aerial or ground spray equipment. Apply after sufficient bolls have developed to produce the desired yield of cotton. Applications made prior to this time could affect maximum yield potential.

TANK MIXTURES: This product may be tank mixed with S,S,S-Tributyl phosphorotrithioate, Thidiazuron and Diuron, or Ethephon to provide additional enhancement of cotton leaf drop.

RESTRICTIONS: Allow a minimum of 7 days between application and harvest of cotton. Not to be used for preharvest application for cotton grown for seed, as a reduction in germination or vigor may occur. THE USE OF ADDITIVES, OTHER THAN THOSE LISTED ON THIS LABEL, FOR PRE-HARVEST APPLICATION TO COTTON IS PROHIBITED. Retreatment Interval (RTI): 7 days. When applying at reduced rates, do not apply more than 2 applications per year.

Fallow Systems

LABELED CROPS: This product may be applied during the fallow period prior to planting or emergence of any crop on this label.

TYPES OF APPLICATIONS: Chemical Fallow, Preplant Fallow Beds, Aid-to-Tillage.

Chemical Fallow

USE INSTRUCTIONS: This product may be applied during the fallow period prior to planting or emergence of any crop listed on this label. This product may be used as a substitute for tillage to control annual weeds in fallow fields. Also, broadcast or spot treatments will control or suppress many perennial weeds in fallow fields. Ground or aerial application equipment may be used. Tank mixtures with 2,4-D and dicamba may be used. Applications up to 2 quarts (1.5lb of glyphosate) per acre may be made by aerial application in fallow sites where there is sufficient buffer to prevent injury due to drift onto adjacent crops.

PRECAUTIONS: Refer to the specific product labels for crop rotation restrictions and cautionary statements of all products used in tank mixtures. **RESTRICTIONS:** For any crop not listed on this label, applications must be made at least 30 days prior to planting. Do not apply dicamba tank mixtures by air in California. Some crop injury may occur if dicamba is applied within 45 days of planting. When applying at reduced rates, do not apply more than 3 applications per year. Retreatment Interval (RTI): 7 days. Do not harvest or feed vegetation from an area for 8 weeks following broadcast application.

Preplant Fallow Beds

USE INSTRUCTIONS: This product may be applied to fallow beds prior to planting or emergence of any crop listed on this label. For any crop not listed on this label, applications must be made at least 30 days prior to planting. This product will control weeds listed in the "ANNUAL WEEDS", "PERENNIAL WEEDS", and "WOODY BRUSH AND TREES RATE TABLES" sections of this label.

TANK MIXTURES: In addition, 12 fluid ounces of this product plus 2 to 3 fluid ounces of Oxyflurofen per acre will control the following weeds with the maximum height or length indicated: 3 inches-common cheeseweed, chickweed, groundsel; 6 inches-London rocket, shepherd's purse. 16 fluid ounces of this product plus 2 to 3 fluid ounces of Oxyflurofen per acre will control the following weeds with the maximum height or length indicated: 6 inches-common cheeseweed, groundsel, marestail (*Conyza canadensis*), 12 inches-chickweed, London rocket, shepherd's purse.

RESTRICTIONS: When applying at reduced rates, do not apply more than 3 applications per year. Do not harvest or feed vegetation from an area for 8 weeks following broadcast application. Retreatment Interval (RTI): 7 days. Application must be made a minimum of 30 days prior to planting any crop not listed on this label.

Aid-to-Tillage

USE INSTRUCTIONS: This product may be used in conjunction with tillage practices in fallow systems or preplant to labeled crops to control downy brome, cheat, volunteer wheat, tansy mustard and foxtail. Apply 12 fluid ounces of this product in 3 to 10 gallons of water per acre. Make applications before weeds are 6 inches in height. Application must be followed by conventional tillage practices no later than 15 days after treatment and before regrowth occur. Allow at least 1 day after application before tillage.

PRECAUTIONS: Tank mixtures with residual herbicides may result in reduced performance.

RESTRICTIONS: When applying at reduced rates, do not apply more than 3 applications per year. Do not harvest or feed vegetation from an area for 8 weeks following broadcast application. Retreatment Interval (RTI): 7 days. Application must be made a minimum of 30 days prior to planting any crop not listed on this label.

Grain Sorghum (Milo)

TYPES OF APPLICATIONS: Those listed in the “Annual and Perennial Crops” section plus the following: Spot Treatment, Over-the-Top Wiper Applications, Pre-harvest.

Preplant, Pre-emergence, At-Planting

USE INSTRUCTIONS: This product may be applied alone or in tank-mixture before, during or after planting grain sorghum. Applications must be made prior to emergence of the crop.

TANK MIXTURES: Apply these tank mixtures in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre.

Atrazine	Alachlor and Atrazine
Atrazine and S-metolachlor	Alachlor
Acetochlor, Atrazine and terbutylazine	S-metolachlor

For difficult-to-control annual weeds such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply this product at 2 pints (0.75lb of glyphosate) per acre in these tank mixtures. For other labeled annual weeds, apply 1.5 to 2 pints of this product (0.563 – 0.75 lb of glyphosate) per acre when weeds are less than 6 inches tall, and 2 to 3 pints when weeds are over 6 inches tall. When using nitrogen solutions as the carrier, the use rate may need to be increased for acceptable weed control. **RESTRICTIONS:** When applying at reduced rates, do not apply more than 3 applications per year. Retreatment Interval (RTI): 7 days. Do not harvest or feed vegetation from an area for 8 weeks following broadcast application

Spot Treatment, Over-the-Top Wiper Applications

USE INSTRUCTIONS: This product may be applied as a spot treatment in grain sorghum. Make spot treatments before heading of milo. This product may be applied with wiper applicators to control or suppress the weeds listed under “Wiper Applicators” in the “Selective Equipment” section of this label.

PRECAUTIONS: The crop receiving spray in treated area will be killed. Take care to avoid drift or spray outside target area for the same reason.

RESTRICTIONS: For spot treatment, do not treat more than 10 percent of the total field area to be harvested.

For wiper applicators, allow at least 40 days between application and harvest. Do not use roller applicators. Do not feed or graze treated milo fodder. Do not ensile treated vegetation. When applying at reduced rates, do not apply more than 3 applications per year. Application must be made a minimum of 14 days prior to harvest for spot treatment. Retreatment Interval (RTI): 7 days.

Hooded Sprayers

USE INSTRUCTIONS: This product may be used through hooded sprayers for weed control between the rows of milo. Only hooded sprayers that completely enclose the spray pattern may be used. See additional instruction for the use of hooded sprayers in the “APPLICATION EQUIPMENT AND TECHNIQUES” section of this label. Crop injury may occur when the foliage of treated weeds comes into direct contact with leaves of the crop. Do not apply this product when the leaves of the crop are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution may contact the crop and cause discoloration, stunting or destruction.

PRECAUTIONS: Treat before milo sends tillers between the drill rows. If such tillers are contacted with the spray solution, the main plant may be killed. Contact of this product in any manner to any vegetation to which treatment is not intended may cause damage. To the extent consistent with applicable law, such damage shall be the sole responsibility of the applicator.

RESTRICTIONS: Milo must be at least 12 inches tall, measured without extending leaves. Do not graze or feed milo forage or fodder following applications of this product through hooded sprayers. Do not apply more than 1 quart of this product (0.75 lb of glyphosate) per acre per application and no more than 3 quarts (2.25lb of glyphosate) per acre per year for hooded sprayer applications. Do not apply more than 3 applications per year. Retreatment Interval (RTI): 7 days.

Preharvest

USE INSTRUCTIONS: Make applications at 30 percent grain moisture or less.

PRECAUTIONS: As with other herbicides that cause sudden plant death, avoid preharvest applications of this product to milo infected with charcoal rot as lodging can occur

RESTRICTIONS: Do not apply more than 2 quarts of this product (1.5 lb of glyphosate) per acre. Allow a minimum of 7 days between application and harvest of sorghum. Preharvest application is not recommended for sorghum grown for seed, as a reduction in germination or vigor may occur. The use of this product for preharvest grain sorghum (milo) is not registered in California. When applying at reduced rates, do not apply more than 2 applications per year. Retreatment Interval (RTI): 7 days.

Post-Harvest

USE INSTRUCTIONS: This product may be applied after harvest of grain sorghum. Higher rates may be required for control of large weeds which were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

This product may be applied to grain sorghum (milo) stubble following harvest to suppress or control regrowth. Apply 1 quart of this product (0.75lb of glyphosate) per acre for control, or 1.5 pints of this product (0.563lb of glyphosate) per acre for suppression.

RESTRICTIONS: Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation. When applying at reduced rates, do not apply more than 2 applications per year. Retreatment Interval (RTI): 7 days. Application must be made a minimum of 30 days prior to the planting of any crop not listed on this label.

Herbs and Spices

LABELED CROPS: Allspice, Angelica, Star anise, Annatto (seed), Balm, Basil, Borage, Burnet, Camomile, Caper buds, Caraway, Black caraway, Cardamom, Cassia bark, Cassia buds, Catnip, Celery seed, Chervil (dried), Chive, Chinese chive, Cinnamon, Clary, Clove buds, Coriander leaf (cilantro or Chinese parsley), Coriander seed (cilantro), Costmary, Culantro (leaf), Culantro (seed), Cumin, Curry (leaf), Dill (dillweed), Dill (seed), Epazote, Fennel seed (common and Florence), Fenugreek, White ginger flower, Grains of paradise, Horehound, Hyssop, Juniper berry, Lavender, Lemongrass, Lovage (leaf and seed), Mace, Marigold, Marjoram (including oregano), Mexican oregano, Mioga flower, Mustard (seed), Nasturtium, Nutmeg, Parsley (dried), Pennyroyal, Pepper (black and white), Pepper leaves, Peppermint, Perilla, Poppy (seed), Rosemary, Rue, Saffron, Sage, Savory (summer and winter), Spearmint, Stevia leaves, Sweet bay, Tansy, Tarragon, Thyme, Vanilla, Wintergreen, Woodruff, and Wormwood.

TYPES OF APPLICATIONS: Those listed in the “Annual and Perennial Crops” section plus the following: Over-the-Top Wiper Applications (Peppermint and Spearmint only), and Spot Treatments (Peppermint and Spearmint only).

PRECAUTIONS: When applying this product prior to transplanting or direct-seeding crops into plastic mulch, care must be taken to remove residues of this product, which could cause crop injury, from the plastic prior to planting.

RESTRICTIONS: Residues can be removed by a single 0.5-inch application of water, either by natural rainfall or via a sprinkler system. For some crops below, it is recommended to make applications 3 days before transplanting or planting. When applying at reduced rates, do not apply more than 6 applications per year. Do not harvest or feed vegetation from an area for 8 weeks following broadcast application. Retreatment Interval (RTI): 7 days. Application must be made a minimum of 30 days prior to the planting of any crop not listed on this label.

Over-the-Top Wiper Applications, Spot Treatments (Peppermint and Spearmint only)

USE INSTRUCTIONS: This product may be used as a spot treatment or wiper application in spearmint and peppermint. Apply spot treatments on a spray-to-wet basis with hand-held equipment, such as backpack and knapsack sprayers, pump-up pressure sprayers, handguns, hand-wands or any other hand-held or motorized spray equipment used to direct the spray solution to a limited area. In wiper applications, the applicator should be adjusted so that the wiper contact point is at least 2 inches above the crop. Weeds should be a minimum of 6 inches taller than the crop.

PRECAUTIONS: The crop receiving spray in the treated area will be killed. Take care to avoid drift or spray outside the target area for this reason. In wiper applications, contact of the herbicide solution with the crop may result in damage or destruction.

RESTRICTIONS: Allow at least 7 days between application and harvest. Further applications may be made in the same area at 30-day intervals. In spot treatment applications, no more than 10 percent of the total field area to be harvested should be treated at one time. When applying at reduced rates, do not apply more than 6 applications per year.

Oil Seed Crops

LABELED CROPS: Borage, Buffalo gourd (seed), Canola, Crambe, Flax, Jojoba, Lesquerella, Meadowfoam, Mustard (seed), Rape, Safflower, Sesame, and Sunflower.

For glyphosate-resistant canola, see the “**GLYPHOSATE-RESISTANT CROPS**” section of this label. **TYPES OF**

APPLICATIONS: Those listed in the “Annual and Perennial Crops” section.

USE INSTRUCTIONS: This product may be applied before, during or after planting oil seed crops. Broadcast applications must be made prior to emergence of the listed oil seed crops. Wiper applicators or hooded sprayers may be used between the rows once the crop is established.

TANK MIXTURES: For sunflowers, a tank mixture with Prowl may be applied before, during or after planting in conventional tillage systems, into a cover crop, established sod or in previous crop residue.

RESTRICTIONS: Do not apply more than 2 quarts of this product (1.5 lb of glyphosate) per acre on canola. Do not apply more than 1 quart of this product (0.75lb of glyphosate) per acre for sunflowers as a single preplant or pre-emergent application per year. Do not feed or graze sunflower forage following application of this product. Retreatment Interval (RTI): 7 days. Do not feed or graze sunflower forage following application of this product. Application must be made at least 30 days prior to planting any crop not listed on this label.

Soybeans

TYPES OF APPLICATIONS: Those listed in the “Annual and Perennial Crops” section plus the following: Spot Treatment, Preharvest, Selective Equipment.

For glyphosate-resistant soybeans, see the “**GLYPHOSATE-RESISTANT CROPS**” section of this label.

Preplant, Pre-emergence, At-Planting

USE INSTRUCTIONS: This product may be applied alone or in a tank-mixture before, during or after planting soybeans. Applications must be made prior to emergence of the crop.

TANK MIXTURES: Apply these tank mixtures in 10 to 20 gallons of water per acre.

Carfentrazone-ethyl	Sulfentrazone and Chlorimuron Ethyl
Quizalofop-p-ethyl	Clomazone Sulfentrazone
Sulfentrazone S-metolachlor and metribuzin	Difenoconazole
Metribuzin and chlorimuron ethyl	S-metolachlor
Cloransulam-methyl	MSMA
Dimethenamid-P	Alachlor
Fluazifop-P-butyl and fenoxaprop-P-ethyl	Linuron
Fomesafen	Metribuzin
Pendimethalin and Imazaquin	Flumioxazin
Imazaquin and Imazethapyr and Pendimethalin	

This product may be tank-mixed with 2,4-D or 2,4-DB. See the 2,4-D label for intervals between application and planting. For difficult-to-control annual weeds such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply this product at 2 pints per acre in these tank mixtures. For other labeled annual weeds, apply 1.5 to 2 pints of this product (0.563 – 0.75 lb of glyphosate) per acre when weeds are less than 6 inches tall, and 2 to 3 pints (0.75 – 1.13 lb of glyphosate) when weeds are over 6 inches tall.

Spot Treatment

USE INSTRUCTIONS: For spot treatments, apply this product prior to initial pod set in soybeans.

PRECAUTIONS: The crop receiving spray in treated area will be killed. Take care to avoid drift or spray outside target area for the same reason.

RESTRICTIONS: Do not treat more than 10 percent of the total field area to be harvested. Retreatment Interval (RTI): 7 days. When applying at reduced rates, do not apply more than 3 applications per year.

Preharvest

USE INSTRUCTIONS: This product provides weed control when applied prior to harvest of soybeans. Apply at rates given in the “ANNUAL WEEDS”, “PERENNIAL WEEDS”, and “WOODY BRUSH AND TREES RATE TABLES”. This product may be applied using either aerial or ground spray equipment. Apply after pods have set and lost all green color. Care should be taken to avoid excessive seed shatter loss due to ground application equipment.

RESTRICTIONS: Do not apply more than 5 quarts per acre of this product (3.75lb of glyphosate) for preharvest applications. Do not apply more than 2 quarts per acre of this product (1.5lb of glyphosate) by air. Allow a minimum of 7 days between application and harvest of soybeans. Do not graze or harvest treated hay or fodder for livestock feed within 25 days of last preharvest application. (If the application rate is 1 quart (0.75lb of glyphosate) per acre or lower, the grazing restriction is reduced to 14 days after last preharvest application.) Preharvest application is not recommended for soybeans grown for seed, as a reduction in germination or vigor may occur. Retreatment Interval (RTI): 7 days.

Selective Equipment

USE INSTRUCTIONS: This product may be applied through recirculating sprayers, shielded applicators, hooded sprayers, wiper applicators or sponge bars in soybeans. Allow at least 7 days between application and harvest.

PRECAUTIONS: See the “Selective Equipment” part of the “APPLICATION EQUIPMENT AND TECHNIQUES” section of this label for information on proper use and calibration of this equipment.

RESTRICTIONS: When applying at reduced rates, do not apply more than 3 applications per year.

Sugarcane

TYPES OF APPLICATIONS: Those listed in the “Annual and Perennial Crops” section.

Preplant, Pre-emergence, At-Planting

USE INSTRUCTIONS: This product may be applied in or around sugarcane fields or in fields prior to the emergence of plant cane.

RESTRICTIONS: Do not apply to vegetation in or around ditches, canals, or ponds containing water to be used for irrigation. When applying at reduced rates, do not apply more than 3 applications per year. Retreatment Interval (RTI): 7 days.

Spot Treatment

USE INSTRUCTIONS: This product may be applied as a spot treatment in sugarcane. For control of volunteer or diseased sugarcane, make a 1 percent solution of this product in water and spray-to-wet the foliage of vegetation to be controlled. Volunteer or diseased sugarcane should have at least 7 new leaves. **PRECAUTIONS:** Avoid spray contact with healthy cane plants since severe damage or destruction may result. **RESTRICTIONS:** Do not feed or graze treated sugarcane foliage following application. When applying at reduced rates, do not apply more than 3 applications per year. Retreatment Interval (RTI): 7 days.

Fallow Treatments

USE INSTRUCTIONS: This product may be used as a replacement for tillage in fields that are lying fallow between sugarcane crops. This product may also be used to remove the last stubble of ratoon cane. For remove of last stubble of ratoon cane, apply 4 to 5 quarts of this product in 10 to 40 gallons of water per acre to new growth having at least 7 new leaves. Allow 7 or more days after application before tillage. Ground or aerial application equipment may be used. Applications up to 3 quarts per acre may be made by aerial application in fallow sites where there is sufficient buffer to prevent injury due to drift onto adjacent crops. Tank mixtures with 2,4-D and dicamba may be used. **RESTRICTIONS:** Do not apply more than 3 quarts (2.25 lbs of glyphosate) of this product per acre per aerial application. When applying at reduced rates, do not apply more than 3 applications per year. Do not apply to vegetation in or around ditches, canals or ponds containing water to be used for irrigation. Retreatment Interval (RTI): 7 days. Do not harvest or feed vegetation from an area for 8 weeks following broadcast application.

Hooded Sprayers

USE INSTRUCTIONS: This product may be used through hooded sprayers for weed control between the rows of sugarcane. See the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for additional use instructions.

PRECAUTIONS: Droplets, mist, foam, or splatter of the herbicide solution setting on the crop may result in discoloration, stunting or destruction. Such damage shall be the sole responsibility of the applicator.

RESTRICTIONS: Do not allow treated weeds to come into contact with the crop. When applying at reduced rates, do not apply more than 3 applications per year. Retreatment Interval (RTI): 7 days. Do not apply this product when crop leaves are growing in direct contact with weeds to be treated.

Vegetable Crops

NOTE: THIS "VEGETABLE CROPS" SECTION GIVES DIRECTIONS THAT APPLY TO ALL LISTED VEGETABLE CROPS WITHIN SECTION 8-10 GROUPED ALPHABETICALLY BELOW. SEE THE INDIVIDUAL CROP CATEGORIES FOR SPECIFIC INSTRUCTIONS, PREHARVEST INTERVALS, PRECAUTIONS AND RESTRICTIONS. **TYPES OF**

APPLICATIONS: Chemical Fallow, Preplant Fallow Beds, Preplant, Pre-emergence, Prior to Transplanting Vegetables, At-Planting, Hooded Sprayers in Row-Middles, Shielded Sprayers in Row-Middles, Wiper Applications in Row-Middles, and Post-Harvest, Directed Applications (Nonbearing Ginseng), Over-the-Top Wiper Applications (Rutabagas only).

PRECAUTIONS: When applying this product prior to transplanting or direct-seeding crops into plastic mulch, care must be taken to remove residues of this product, which could cause crop injury, from the plastic prior to planting. Residues can be removed by a single 0.5-inch application of water, either by natural rainfall or via a sprinkler system. Care should be taken to ensure that the wash water flushes off the plastic mulch and does not enter transplant holes. Applications made at emergence will result in injury or death to emerged seedlings. Avoid contact of herbicide with foliage, green shoots or stems, bark, exposed roots (including those emerging from plastic mulch), or fruit of crops because severe injury or destruction may result. When making pre-emergence and at-planting applications, applications must be made before crop emergence to avoid severe crop injury. Apply before seed germination in coarse sandy soils to further minimize the risk of injury. In crops with vines, hooded sprayer, shielded sprayer and wiper applications to row-middles should be made prior to vine development otherwise severe injury or destruction may result. See "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for additional information.

RESTRICTIONS: Unless otherwise specified in this product's labeling, treatments with selective equipment including wipers and hooded sprayers must be made at least 14 days prior to harvest. Post-harvest or fallow applications must be made at least 30 days prior to planting any non-labeled crop.

Brassica Vegetables

LABELED CROPS: Broccoli, Chinese broccoli (gai Ion), Broccoli raab (rapini), Brussels sprouts, Cabbage, Chinese cabbage (bok choy), Chinese cabbage (napa), Chinese mustard cabbage (gai choy), Cauliflower, Cavalo broccolo, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, and Rape greens.

RESTRICTIONS: When applying at reduced rates, do not apply more than 6 applications per year. Retreatment Interval (RTI): 7 days. Do not harvest or feed vegetation from an area for 8 weeks following broadcast application. Allow a minimum of 14 days between application and harvest for hooded sprayer, shielded sprayer, and wiper applications. Application must be made a minimum of 30 days prior to the planting of any crop not listed on this label.

Bulb Vegetables

LABELED CROPS: Garlic, Great-headed garlic, Leek, Onion (dry bulb and green), Welsh Onion, and Shallot.

RESTRICTIONS: When applying at reduced rates, do not apply more than 6 applications per year. Do not harvest or feed vegetation from an area for 8 weeks following broadcast application. Allow a minimum of 14 days between application and harvest for hooded sprayer, shielded sprayer, and wiper applications. Retreatment Interval (RTI): 7 days. Make only one preharvest application. Do not make a preharvest application using aerial application equipment. For preharvest application, allow a minimum of 30 days between application and harvest. Do not make a preharvest application on garlic or onion grown for seed.

Cucurbit Vegetables and Fruits

LABELED CROPS: Chayote (fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cucumber, Gherkin, Edible gourd (includes hyotan, cucuzza, hechima, and Chinese okra), Melons (all), Momordica spp. (includes balsam apple, balsam pear, bittermelon, and Chinese cucumber), Muskmelon (includes cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey ball melon, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon), Pumpkin, Summer squash (includes crook-neck squash, scallop squash, straightneck squash, vegetable marrow, and zucchini), Winter squash (includes butternut squash, calabaza, hubbard squash, acorn squash, and spaghetti squash), and Watermelon.

RESTRICTIONS: For Cantaloupe, Casaba melon, Crenshaw melon, Cucumber, Gherkin, Gourds, Honeydew melon, Honey ball melon, Mango melon, Melons (all), Muskmelon, Persian melon, Pumpkin, Squash (summer, winter), and Watermelon, allow at least 3 days between application and planting. Retreatment Interval (RTI): 7 days. When applying at reduced rates, do not apply more than 6 applications per year. Do not harvest or feed vegetation from an area for 8 weeks following broadcast application. Allow a minimum of 14 days between application and harvest for hooded sprayer, shielded sprayer, and wiper applications. Application must be made a minimum of 30 days prior to the planting of any crop not listed on this label.

Leafy Vegetables

LABELED CROPS: Amaranth (Chinese spinach), Arugula (roquette), Beet greens, Cardoon, Celery, Chinese celery, Celtuce, Chaya, Chervil, Edible-leaved chrysanthemum, Garland chrysanthemum, Corn salad, Cress (garden and upland), Dandelion, Dock (sorrel), Dokudami, Endive (escarole), Florence fennel, Gow kee, Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Radicchio (red chicory), Rhubarb, Spinach, New Zealand spinach, Vine spinach, Swiss chard, Watercress (upland), and Water spinach.

RESTRICTIONS: For Watercress, avoid applications within 3 days prior to seeding and during the period between seeding and emergence to minimize the risk of injury. Allow a minimum of 14 days between application and harvest for hooded sprayer, shielded sprayer, and wiper applications. Application must be made a minimum of 30 days prior to the planting of any crop not listed on this label. Retreatment Interval (RTI): 7 days. Do not harvest or feed vegetation from an area for 8 weeks following broadcast application. When applying at reduced rates, do not apply more than 6 applications per year.

Fruiting Vegetables

LABELED CROPS: Eggplant, Groundcherry (*Physalis* spp.), Pepino, Pepper (includes bell pepper, chili pepper, cooking pepper, pimiento, sweet pepper), Tomatillo, and Tomato.

RESTRICTIONS: For Eggplant, Ground cherry, Pepper (all), and Tomatillo, allow at least 3 days between application and planting. For Tomato, hooded or shielded sprayer applications in row-middles are not recommended. When applying at reduced rates, do not apply more than 6 applications per year. Do not harvest or feed vegetation from an area for 8 weeks following broadcast application. Retreatment Interval (RTI): 7 days. Allow a minimum of 14 days between application and harvest for hooded sprayer, shielded sprayer, and wiper applications. Application must be made a minimum of 30 days prior to the planting of any crop not listed on this label.

Legume Vegetables (Succulent or Dried)

LABELED CROPS: Bean (*Lupinus*: includes grain lupin, sweet lupin, white lupin, and white sweet lupin), Bean (*Phaseolus*: includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, and wax bean), Bean (*Vigna*: includes adzuki bean, asparagus bean, black-eyed pea, catjang, Chinese longbean, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, and yardlong bean), Broad bean (fava), Chickpea (garbanzo), Guar, Jackbean, Lablab bean, Lentil, Pea (*Pisum*: includes dwarf pea, edible-podded pea, English pea, field pea, garden pea, green pea, snowpea, and sugar snap pea), Pigeon pea, Soybean (immature seed), and Sword bean.

RESTRICTIONS: When applying at reduced rates, do not apply more than 6 applications per year. Do not harvest or feed vegetation from an area for 8 weeks following broadcast application. Retreatment Interval (RTI): 7 days. Allow a minimum of 14 days between application and harvest for hooded sprayer, shielded sprayer, and wiper applications.

Root and Tuber Vegetables

LABELED CROPS: Arracacha, Arrowroot, Chinese artichoke, Jerusalem artichoke, Beet (garden), Burdock, Canna, Carrot, Cassava (bitter and sweet), Celeriac, Chayote (root), Chervil (turnip-rooted), Chicory, Chufa, Dasheen (taro), Galangal, Ginger, Ginseng, Horseradish, Leren, Kava (turnip-rooted), Parsley (turnip-rooted), Parsnip, Potato, Radish, Oriental radish, Rutabaga, Salsify, Black salsify, Spanish salsify, Skirret, Sweet potato, Tanier, Turmeric, Turnip, Wasabi, Yacon, Yam bean, and True yam.

RESTRICTIONS: When applying at reduced rates, do not apply more than 6 applications per year. Retreatment Interval (RTI): 7 days.

Directed Applications (Non-bearing Ginseng only)

USE INSTRUCTIONS: This product may be used for weed control in established non-bearing ginseng. Applications may be made with boom equipment, CDA, shielded sprayers, hand-held and high volume wands, lances, and orchard guns or with wiper application equipment.

RESTRICTIONS: Direct applications so that there is no contact of this product with the ginseng plant. Applications must be made at least one year prior to harvest. Retreatment Interval (RTI): 7 days.

Over-the-Top Wiper Applications (Rutabagas only)

USE INSTRUCTIONS: Wiper applicators may be used over-the-top of rutabagas.

PRECAUTIONS: Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction.

RESTRICTIONS: Allow at least 14 days between application and harvest of rutabagas. When applying at reduced rates, do not apply more than 6 applications per year. Retreatment Interval (RTI): 7 days.

Miscellaneous Crops

LABELED CROPS: Aloe vera, Asparagus, Bamboo shoots, Globe artichoke, Okra, Peanut (ground nut), Pineapple, Strawberry, and Sugar beet.

TYPES OF APPLICATIONS: Those listed in the “**Annual and Perennial Crops**” section plus the following: Weed Control, Site Preparation, Spot Treatment (Asparagus).

For glyphosate-resistant sugar beets, see the “**GLYPHOSATE-RESISTANT CROPS**” section of this label.

PRECAUTIONS: In crops with vines, hooded sprayer, shielded sprayer and wiper applications to row- middles should be made prior to vine development otherwise severe injury or destruction may result. See “**APPLICATION EQUIPMENT AND TECHNIQUES**” section of this label for additional information.

RESTRICTIONS: Avoid contact of herbicide with foliage, green shoots or stems, bark, exposed roots (including those emerging from plastic mulch), or fruit of crops because severe injury or destruction may result. When making pre-emergence and at-planting applications, applications must be made before crop emergence to avoid severe crop injury. Apply before seed germination in coarse sandy soils to further minimize the risk of injury. Unless otherwise specified in this product's labeling, treatments with selective equipment including wipers and hooded sprayers must be made at least 14 days prior to harvest. Post-harvest or fallow applications must be made at least 30 days prior to planting any non-labeled crop. When applying at reduced rates, do not apply more than 3 applications per year. Retreatment Interval (RTI): 7 days.

Weed Control, Site Preparation

USE INSTRUCTIONS: This product may be applied for weed control or for site preparation prior to planting or transplanting crops listed in this section.

PRECAUTIONS: When applying this product prior to transplanting or direct-seeding crops into plastic mulch, care must be taken to remove residues of this product, which could cause crop injury, from the plastic prior to planting. Residues can be removed by a single 0.5-inch application of water, either by natural rainfall or via a sprinkler system. Care should be taken to ensure that the wash water flushes off the plastic mulch and does not enter transplant holes. Applications made at emergence will result in injury or death to emerged seedlings.

RESTRICTIONS: Do not apply within a week before the first asparagus spears emerge. Do not feed or graze treated pineapple forage following application. Retreatment Interval (RTI): 7 days. Do not harvest or feed vegetation from an area for 8 weeks following application. When applying at reduced rates, do not apply more than 3 applications per year. Allow at least 21 days between residue removal and transplanting.

Spot Treatment (Asparagus)

USE INSTRUCTIONS: This product may be applied immediately after cutting, but prior to the emergence of new spears.

RESTRICTIONS: Do not treat more than 10 percent of the total field area to be harvested. Do not harvest within 5 days of treatment. When applying at reduced rates, do not apply more than 3 applications per year. Retreatment Interval (RTI): 7 days.

Post-Harvest (Asparagus)

USE INSTRUCTIONS: This product may be applied after the last harvest and all spears have been removed. If spears are allowed to regrow, delay application until ferns have developed. Delayed treatments should be applied as a directed or shielded spray in order to avoid contact of the spray with ferns, stems or spears.

PRECAUTIONS: Direct contact of the spray with the asparagus may result in serious crop injury. Select and use recommended types of spray equipment for post-emergence post-harvest applications. A directed spray is any application where the spray pattern is aligned in such a way as to avoid direct contact of the spray with the crop. A shielded spray is any application where a physical barrier is positioned and maintained between the spray and the crop to prevent contact of spray with the crop.

RESTRICTIONS: When applying at reduced rates, do not apply more than 2 applications per year. Retreatment Interval (RTI): 7 days. Do not harvest or feed vegetation from an area for 8 weeks following broadcast application. Application must be made a minimum of 30 days prior to the planting of any crop not listed on this label.

TREE, VINE, AND SHRUB CROPS (Alphabetical)

NOTE: THIS SECTION GIVES DIRECTIONS THAT APPLY TO ALL LISTED TREE, VINE, AND SHRUB CROPS GROUPED ALPHABETICALLY BELOW. SEE THE INDIVIDUAL CROP CATEGORIES FOR SPECIFIC INSTRUCTIONS, PREHARVEST INTERVALS, PRECAUTIONS AND RESTRICTIONS.

TYPES OF APPLICATIONS: Preplant (Site Preparation) Broadcast Sprays, Weed Control, Middles (between rows of trees, vines or bushes), Strips (within rows of trees, vines or bushes), Selective Equipment (shielded sprayers, wiper treatments), Directed Sprays, Spot Treatments, Perennial Grass Suppression, Cut Stump. Applications may be made with boom equipment, CDA equipment, shielded sprayers, hand-held and high-volume wands, lances, orchard guns or with wiper applicator equipment, except as directed.

USE INSTRUCTIONS: This product may be applied in middles (between rows of trees or vines), strips (within rows of trees or vines), and for weed control or perennial grass suppression in established tree fruit and nut groves, orchards, berries, and vineyards. It may also be used for site preparation prior to planting or transplanting these crops. Apply 1 pint to 5 quarts per acre according to the “ANNUAL WEEDS” and “PERENNIAL WEEDS RATE TABLES” sections of this label. Utilize rates at the higher end of the recommended rate range when weeds are stressed, growing in dense populations or are greater than 12 inches tall. Repeat applications may be made up to a maximum of 10.6 quarts (7.95 lb of glyphosate) per acre per year. The maximum use rates stated throughout this product's labeling apply to this product combined with the use of all other herbicides containing glyphosate or sulfosate as the active ingredient, whether applied as mixtures or separately. Calculate the application rates and ensure that the total use of this and other glyphosate or sulfosate

containing products does not exceed stated maximum use rate.

PRECAUTIONS: Extreme care must be exercised to avoid contact of herbicide solution, spray, drift or mist with foliage or green bark of trunk, branches, suckers, fruit or other parts of trees, canes and vines. Avoid applications when recent pruning wounds or other mechanical injury has occurred. Contact of this product with other than matured brown bark can result in serious crop damage or destruction. Only shielded or directed sprayers may be used in crops with potential for crop contact, and then only where there is sufficient clearance. For applications in strips (within rows of trees), only selective equipment (directed sprays, hooded sprayers, shielded applicators, or wipers) should be used to minimize the potential for leakage or drift of herbicide sprays onto crop. For berry crops, hooded or shielded sprayers must be fully enclosed including top, sides, front and back. Only wipers or shielded applicators capable of preventing all contact with crop may be used. See “APPLICATION EQUIPMENT AND TECHNIQUES” section of this label for additional directions and precautions.

RESTRICTIONS: Allow a minimum of 3 days between application and transplanting. Only wipers or shielded applicators capable of preventing all contact with the crop may be used. For berry crops, hooded or shielded sprayers must be fully enclosed including top, sides, front and back. For applications in strips (within rows of trees), only selective equipment (directed sprays, hooded sprayers, shielded applicators, or wipers) may be used in order to minimize the potential for overspray or drift of this product onto the crop. Only shielded or directed sprayers may be used in crops with potential for crop contact, and then only where there is sufficient clearance.

Middles (Between Rows of Trees, Vines or Bushes)

USE INSTRUCTIONS: This product will control or suppress annual and perennial weeds and ground covers growing between the rows of labeled tree and vine crops. If weeds are under drought stress, irrigate prior to application. Reduced control may result if weeds have been mowed prior to application.

TANK MIXTURES: A tank mixture of this product plus Oxyflurofen may be used for annual weeds in middles between rows of citrus crops, tree fruits, tree nuts and vine crops. This mixture is recommended when weeds are stressed or growing in dense populations. 16 to 32 fluid ounces per acre of this product plus 3 to 12 fluid ounces per acre of Oxyflurofen will control annual weeds with a maximum height or diameter of 6 inches, including crabgrass, common groundsel, junglerice, common lambsquarters, redroot pigweed, London rocket, common ryegrass, shepherd's purse, annual sowthistle, filaree (suppression), horseweed/marestail (*Conyza canadensis*), stinging nettle and common purslane (suppression). 16 to 32 fluid ounces per acre of this product plus 3 to 12 fluid ounces per acre of Oxyflurofen will control common cheeseweed (malva) or hairy fleabane (*Conyza bonanensis*) with a maximum height or diameter of 3 inches. **RESTRICTIONS:** Do not apply these tank mixtures in Puerto Rico.

Strips (In Rows of Trees, Vines or Bushes)

TANK MIXTURES: This product may be applied in rows of tree or vine crops in tank mixtures with the following products: Napropamide, Simazine, Diuron, Oxyflurofen, Norflurazon, Bromacil and Diuron, Oryzalin, Pendimethalin.

Do not apply these tank mixtures in Puerto Rico.

Refer to the individual product labels for specific crops, rates, geographic restrictions, and precautionary statements. This product may be tank-mixed with the products listed providing that the product to be tank-mixed is registered for use on this site.

Perennial Grass Suppression

This product will suppress perennial grasses such as bahiagrass, Bermudagrass, tall fescue, orchardgrass, Kentucky bluegrass, and quackgrass that are grown as ground covers in tree and vine crops. For suppression of tall fescue, fine fescue, orchardgrass and quackgrass, apply 8 fluid ounces of this product (0.188lb of glyphosate) in 10 to 20 gallons of water per acre. For suppression of Kentucky bluegrass covers, apply 6 fluid ounces of this product (0.141lb of glyphosate) per acre.

RESTRICTIONS: Do not add ammonium sulfate.

For best results, mow cool season grass covers in the spring to even their height and apply this product 3 to 4 days after mowing.

For suppression of vegetative growth and seedhead inhibition of bahiagrass for approximately 45 days, apply 6 fluid ounces of this product (0.141lb of glyphosate) in 10 to 25 gallons of water per acre. Apply 1 to 2 weeks after full green up or after mowing to a uniform height of 3 to 4 inches. This application must be made prior to seed head emergence.

For suppression up to 120 days, apply 4 fluid ounces of this product per acre, followed by an application of 2 to 4 fluid ounces (0.05-0.094lb of glyphosate) per acre about 45 days later. Make no more than 2 applications per year. For burndown of Bermudagrass, apply 1 to 2 quarts of this product (0.75-1.5lb of glyphosate) in 3 to 20 gallons of water per acre. Use this treatment only if reduction of the Bermudagrass stand can be tolerated. When burndown is required prior to harvest, allow at least 21 days to ensure sufficient time for burndown to occur.

For suppression of Bermudagrass, apply 6 to 16 fluid ounces of this product (0.141 – 0.375lb of glyphosate) per acre east of the Rocky Mountains and 16 fluid ounces of this product per acre west of the Rocky Mountains. Apply in a total spray volume of 3 to 20 gallons per acre, no sooner than 1 to 2 weeks after full green-up. If the Bermudagrass is mowed prior to application, maintain a minimum of 3 inches in height. Sequential applications may be made when regrowth occurs and Bermudagrass injury and stand reduction can be tolerated. East of the Rocky Mountains, rates of 6 to 10 fluid ounces of this product per acre should be used in shaded conditions or where a lesser degree of suppression is desired.

Cut Stump (Tree Crops)

USE INSTRUCTIONS: Cut stump applications of this product may be made during site preparation for site renovation, prior to transplanting tree crops. This product will control regrowth of cut stumps and resprouts of many types of tree species, some of which are listed below.

Citrus Trees: Calamondin, Chironja, Citron, Citrus hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin (Tangerine), Orange (all), Pummelo, Tangelo, and Tangor.

Fruit Trees: Apple, Apricot, Cherry (sweet, sour), Crabapple, Loquat, Mayhaw, Nectarine, Olive, Peach, Pear, Plum/Prune (all), and Quince.

Nut Trees: Almond, Beechnut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (hazelnut), Hickory Nut, Macadamia, Pecan, Pistachio, and Walnut (black, English).

Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut trees or resprouts close to the soil surface. Apply a 50 to 100 percent solution of this product to the freshly cut surface immediately after cutting. Delays in application may result in reduced performance. For best results, applications should be made during periods of active growth and full leaf expansion.

PRECAUTIONS: Some sprouts, stems, or trees may share the same root system. Adjacent trees having a similar age, height and spacing may signal shared roots. Whether grafted or shared, injury is likely to occur to non-treated stems/trees when one or more trees sharing common roots are treated.

RESTRICTIONS: DO NOT MAKE CUT STUMP APPLICATIONS WHEN THE ROOTS OF ADJACENT DESIRABLE TREES MAY BE GRAFTED TO THE ROOTS OF THE CUT STUMP. INJURY RESULTING FROM ROOT GRAFTING MAY OCCUR IN ADJACENT TREES. Do not apply more than 10.67 quarts (7.95 lb of glyphosate) of this product per acre per single application. Do not apply more than 10.67 quarts (7.95 lb of glyphosate) of this product per acre per year.

Berry Crops

LABELED CROPS: Blackberry (including bingleberry, black satin berry, boysenberry, Cherokee blackberry, chesterberry, Cheyenne blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, Himalayaberry, hullberry, juneberry, lavacaberry, lowberry, lucretiaberry, marionberry, nectarberry, olallieberry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, and youngberry), Blueberry, Cranberry, Currant, Elderberry, Gooseberry, Huckleberry, Loganberry, and Salal.

TYPES OF APPLICATIONS: Those listed in the “Tree, Vine, and Shrub Crops” section plus Spot Treatment in Cranberry Production and Post-Harvest Treatments in Cranberry Production.

RESTRICTIONS: To avoid damage, herbicide sprays must not be allowed to contact desirable vegetation, including green shoots, canes, or foliage. Allow a minimum of 30 days between last application and harvest in cranberries. Allow a minimum of 14 days between last application and harvest in other berry crops. Do not make directed sprays within the cranberry bush areas prior to berry harvest. When applying at reduced rates, do not apply more than 10 applications per year for established trees. When applying at reduced rates, do not apply more than 3 applications per year for prior to planting or transplanting stages. Retreatment Interval (RTI): 7 days. Allow a minimum of 14 days between last application and harvest of other berry crops. Allow a minimum of 3 days between application of this product and transplanting. Allow a minimum of 30 days between application and the planting of any crop not listed on the label. Do not apply more than 10.67 quarts (7.95 lb of glyphosate) of this product per acre per year. Do not apply more than 5 quarts (3.75lb of glyphosate) of this product per acre per single application.

Spot Treatment in Cranberry Production

USE INSTRUCTIONS: Spot treatments may be used to control weeds growing in dry ditches (interior and perimeter) of cranberry production areas. Hand-held sprayers or other appropriate application equipment listed under “**APPLICATION EQUIPMENT AND TECHNIQUES**” in this label may be used. Drop water level to remove standing water in ditches prior to application. In hand-held sprayers, use 1 to 2 percent solution of this product. Spray to wet vegetation, not to run-off.

PRECAUTIONS: Use nozzles that emit medium- to large sized droplets to minimize drift in order to avoid crop injury.

RESTRICTIONS: For treatments after draw-down of water in dry ditches, allow 2 or more days after treatment before reintroduction of water to achieve maximum weed control. Apply this product within 1 day after drawdown to ensure application to actively growing weeds. Allow a minimum of 30 days between last application and harvest of cranberries. Do not make applications by air. Do not apply directly to water. Do not apply more than 10.67 quarts (7.95 lbs of glyphosate) of this product per acre per year. Do not apply more than 5 quarts (3.75lb of glyphosate) of this product per acre per single application. When applying at reduced rates, do not apply more than 10 applications per year. Retreatment Interval (RTI): 7 days. Spray adequately to wet the vegetation only; do not spray to the point of run-off. Do not make applications by air. Do not apply directly to water.

Post-Harvest Treatments in Cranberry Production

USE INSTRUCTIONS: Application of this product may be made after the harvest of cranberries to control weeds growing within the field. Best results will be obtained if applications are made to vines that appear dormant (after they have turned red). Hand-held sprayers, wipers, or other appropriate application equipment listed under “**APPLICATION EQUIPMENT AND TECHNIQUES**” in this label may be used. If using hand-held sprayers, use a 0.5 to 1 percent solution of this product. Spray to wet vegetation, not to run-off. If using handheld boom sprayers, apply 2 to 4 quarts of this product (1.5-3lb of glyphosate) per acre.

PRECAUTIONS: Even though vines appear dormant, contact of the herbicide solution with desirable vegetation may result in damage or severe plant injury. Cranberry plants that are directly sprayed may be killed.

RESTRICTIONS: Make applications only after cranberries have been harvested. Do not treat more than 10 percent of the total bog. Allow a minimum of 6 months after last application and next harvest of cranberries. Do not apply this product through the irrigation system. Do not make applications by air. Do not apply directly to water. Retreatment Interval (RTI): 7 days. Do not apply more than 2 applications per year. Do not apply more than 10.67 quarts (7.95 lb of glyphosate) of this product per acre per year. Do not apply more than 5 quarts (3.75lb of glyphosate) of this product per acre per single application.

Citrus

LABELED CROPS: Calamondin, Chironja, Citron, Citrus Hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Orange (all), Pummelo, Satsuma Mandarin, Tangelo (ugli), and Tangor.

TYPES OF APPLICATIONS: Those listed in the “Tree, Vine, and Shrub Crops” section.

USE INSTRUCTIONS: (The directions below pertain to applications in Florida and Texas): For burndown or control of the weeds listed below, apply the recommended rates of this product in 3 to 30 gallons of water per acre. Where weed foliage is dense, use 10 to 30 gallons of water per acre. For goatweed, apply 2 to 3 quarts of this product (1.5-2.25lb of glyphosate) per acre. Apply in 20 to 30 gallons of water per acre when plants are actively growing. Use 2 quarts (1.5lb of glyphosate) per acre when plants are less than 8 inches tall and 3 quarts (2.25lb of glyphosate) per acre when plants are greater than 8 inches tall. If goatweed is greater than 8 inches tall, the addition of Bromacil and Diuron or Diuron may improve control. Refer to the individual product labels for specific crops, rates, geographic restrictions and precautionary statements.

S = Suppression	B = Burndown	C = Control	PC = Partial Control	
Weed Species	GlamPrime RATE PER ACRE			
	1 QT.	2 QTS.	3 QTS.	5 QTS.
Bermudagrass	B	--	PC	C
Guineagrass				
Texas and Florida Ridge	B	C	C	C
Florida Flatwoods	--	B	C	C
Paragrass	B	C	C	C
Torpedograss	B	--	PC	C

RESTRICTIONS: Allow a minimum of 1 day between last application and harvest in citrus crops. For citron groves, apply as directed sprays only. Retreatment Interval (RTI): 7 days. When applying at reduced rates, do not apply more than 3 applications per year for prior to planting or transplanting. When applying at reduced rates, do not apply more than 10 applications per year for established trees. Do not apply more than 10.67 quarts (7.95 lbs of glyphosate) of this product per acre per year. Do not apply more than 5 quarts (3.75lb of glyphosate) of this product per acre per single application

Miscellaneous Tree Food Crops

LABELED CROPS: Cactus (fruit and pads), Palm (heart, leaves), and Palm (oil).

TYPES OF APPLICATIONS: Those listed in the “Tree, Vine, and Shrub Crops” section.

RESTRICTIONS: Do not apply more than 5 quarts (3.75lb of glyphosate) of this product per acre per single application. . Do not apply more than 10.67 quarts (7.95 lbs of glyphosate) of this product per acre per year. When applying at reduced rates, do not apply more than 10 applications per year for established trees. Retreatment Interval (RTI): 7 days. Allow a minimum of 3 days between application and transplanting. Allow a minimum of 30 days between application and the planting of any crop not listed on this label.

Non-Food Tree Crops

LABELED CROPS: Pine, Poplar, Eucalyptus, Christmas trees, Other Non-food Tree Crops.

TYPES OF APPLICATIONS: Those listed in The “Tree, Vine, and Shrub Crops” section.

Directed Sprays, Spot Treatment, Wiper Applications

USE INSTRUCTIONS: This product may be used as a post-directed spray and spot treatment around established poplar, eucalyptus, Christmas trees and other non-food tree crops.

PRECAUTIONS: Care must be exercised to avoid contact of spray, drift or mist with foliage or green bark of established Christmas trees and other pine trees. Desirable plants may be protected from the spray solution by using shields or coverings made of cardboard or other impermeable material.

RESTRICTIONS: THIS PRODUCT IS NOT TO BE USED AS AN OVER-THE-TOP BROADCAST SPRAY IN CHRISTMAS TREES AND OTHER PINE TREES. Do not apply more than 5 quarts (3.75lb of glyphosate) of this product per acre per single application for directed spray and spot treatment. Do not apply more than 10.67 quarts (7.95 lbs of glyphosate) of this product per acre per single application for wiper application. Do not apply more than 10.67 quarts (7.95 lbs of glyphosate) of this product per acre per year. When applying at reduced rates, do not apply more than 10 applications per year for established trees. Retreatment Interval (RTI): 7 days. Do not apply this product as a broadcast application over the top of plantations or tree crops.

Site Preparation

USE INSTRUCTIONS: This product may be used prior to planting nonfood tree crops.

PRECAUTIONS: Precautions should be taken to protect nontarget plants during site preparation applications.

RESTRICTIONS: Do not apply more than 5 quarts (3.75lb of glyphosate) of this product per acre per single application. Do not apply more than 10.67 quarts (7.95 lb of glyphosate) of this product per acre per year. Retreatment Interval (RTI): 7 days. Allow a minimum of 3 days between application and transplanting. Allow a minimum of 30 days between application and the planting of any crop not listed on this label. When applying at reduced rates, do not apply more than 10 applications per year for established trees. When applying at reduced rates, do not apply more than 3 applications per year for prior to planting or transplanting stages.

Pome Fruit

LABELED CROPS: Apple, Crabapple, Loquat, Mayhaw, Pear (including oriental pear), and Quince.

TYPES OF APPLICATIONS: Those listed in the “Tree, Vine, and Shrub Crops” section.

RESTRICTIONS: Allow a minimum of 1 day between last application and harvest in Pome crops. Do not apply more than 5 quarts (3.75lb of glyphosate) of this product per acre per single application. Do not apply more than 10.67 quarts (7.95 lb of glyphosate) of this product per acre per year. Retreatment Interval (RTI): 7 days. When applying at reduced rates, do not apply more than 10 applications per year for established trees. When applying at reduced rates, do not apply more than 3 applications per year for prior to planting or transplanting stages.

Stone Fruit

LABELED CROPS: Apricot, Cherry (sweet, tart), Nectarine, Olive, Peach, Plum/Prune (all types), and Plumcot.

TYPES OF APPLICATIONS: Those listed in the “Tree, Vine, and Shrub Crops” section.

RESTRICTIONS: Allow a minimum of 17 days between last application and harvest in stone fruit crops. For olive groves, apply as directed sprays only. Do not apply more than 5 quarts (3.75lb of glyphosate) of this product per acre per single application. Do not apply more than 10.67 quarts (7.95 lb of glyphosate) of this product per acre per year. Retreatment Interval (RTI): 7 days. When applying at reduced rates, do not apply more than 10 applications per year for established trees. When applying at reduced rates, do not apply more than 3 applications per year for prior to planting or transplanting stages. Remove suckers and low-hanging limbs at least 10 days prior to application.

Restrictions on Application Equipment

For CHERRIES, any application equipment listed in this section may be used in all states. Any application equipment listed in this section may be used in apricots, nectarines, peaches and plums/ prunes growing in Arizona, California, Colorado, Idaho, Kansas, Kentucky, New Jersey, North Dakota, Oklahoma, Oregon, Texas, Utah, and Washington, except for peaches grown in the states specified in the following paragraph. In all other states, use wiper equipment only. For PEACHES, grown in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina, and Tennessee only, apply with a shielded boom sprayer or shielded wiper applicator, which prevents any contact of this product with the foliage or bark of trees. Apply no later than 90 days after first bloom. Applications made after this time may result in severe damage. Remove suckers and low-hanging limbs at least 10 days prior to application. Avoid applications near trees with recent pruning wounds or other mechanical injury. Apply only near trees that have been planted in the orchard for 2 or more years. EXTREME CARE MUST BE TAKEN TO ENSURE NO PART OF THE PEACH TREE IS CONTACTED.

Tree Nuts

LABELED CROPS: Almond, Beechnut, Betelnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Coconut, Filbert (hazelnut), Hickory nut, Macadamia, Pecan, Pine nut, Pistachio, and Walnut (black, English).

TYPES OF APPLICATIONS: Those listed in the “Tree, Vine, and Shrub Crops” section.

RESTRICTIONS: Allow a minimum of 3 days between last application and harvest of tree nuts, except coconut. Allow 14 days between application and harvest in coconut. Do not apply more than 5 quarts (3.75lb of glyphosate) of this product per acre per single application. Do not apply more than 10.67 quarts (7.95 lb of glyphosate) of this product per acre per year. When applying at reduced rates, do not apply more than 10 applications per year for established trees. Retreatment Interval (RTI): 7 days. Allow a minimum of 30 days between application and the planting of any crop not listed on this label.

Tropical and Subtropical Trees and Fruits

LABELED CROPS: Ambarella, Atemoya, Avocado, Banana, Barbados cherry (acerola), Biriba, Blimbi, Breadfruit, Cacao (cocoa) bean, Canistel, Carambola (starfruit), Cherimoya, Coffee, Custard apple, Dates, Durian, Feijoa, Figs, Governor's plum, Guava, Llama, Imbe, Imbu, Jaboticaba, Jackfruit, Longan, Lychee, Mamey apple, Mango, Mangosteen, Marmalade box (genip), Mountain papaya, Papaya, Pawpaw, Plantain, Persimmon, Pomegranate, Pulasan, Rambutan, Rose apple, Sapodilla, Sapote (black, mamey, white), Spanish lime, Soursop, Star apple, Sugar apple, Surinam cherry, Tamarind, Tea, Ti (roots and leaves), and Wax jambu.

TYPES OF APPLICATIONS: Those listed in The “Tree, Vine, and Shrub Crops” section plus Bananacide (Banana only).

RESTRICTIONS: Allow a minimum of 1 day between last application and harvest in banana, guava, papaya, and plantain crops. Allow a minimum of 14 days between last application and harvest for any other tropical or subtropical tree fruit. Allow a minimum of 28 days between last application and harvest in coffee crops. In coffee and banana, delay applications 3 months after transplanting to allow the new coffee or banana plant to become established. Do not apply more than 5 quarts (3.75lb of glyphosate) of this product per acre per single application. Do not apply more than 10.67 quarts (7.95 lb of glyphosate) of this product per acre per year. When applying at reduced rates, do not apply more than 10 applications per year for established trees. Retreatment Interval (RTI): 7 days.

Bananacide (Banana only)

USE INSTRUCTIONS: This product may be used to destroy banana plants infected with the Banana Bunchy Top Virus as well as non-infected banana plants to establish disease free buffers around plantations. Remove all fruit from the plants within the treatment area prior to treatment. Inject 0.04 fluid ounce (1 mL) of this product's concentrate per 2 to 3 inches of pseudostem diameter. Make the injection at least one foot above the ground, except for very small plants, which should be injected vertically into the top. Any subsequent regrowth must also be destroyed. All plants and mats (or units) adjacent (within a 4-foot radius) to a treated mat shall be mechanically destroyed. For control of the Banana Bunchy Top Virus, it is critical that the grower follow a strict control program involving monitoring for diseased plants, spraying to control the aphid vector, and destruction of all infected mats (or units). An infected plant may not show symptoms of the banana bunchy top virus for up to 125 days, therefore it is critical that the entire mat (or unit) containing the diseased plant be destroyed immediately.

RESTRICTIONS: Do not apply more than 0.5 fluid ounce (15 mL) of this product's concentrate per mat (or unit). Remove all fruit from plants and mats (or units) prior to treatment. Do not harvest any fruit or plant materials from treated mats (or units) following injection. Do not allow livestock to consume treated plant materials. Following transplant of new banana plants into treated areas, allow plants to become established for 3 months before applying this product for weed control. Retreatment Interval (RTI): 7 days.

Vine Crops

LABELED CROPS: Grapes (raisin, table, wine), Hops, Kiwi, and Passion fruit.

TYPES OF APPLICATIONS: Those listed in the “Tree, Vine, and Shrub Crops” section.

USE INSTRUCTIONS: Applications must not be made when green shoots, canes or foliage are in the spray zone. In the northeast and Great Lakes regions, applications must be made prior to the end of bloom stage of grapes to avoid injury, or make applications with shielded sprayers or wiper equipment.

RESTRICTIONS: Allow a minimum of 14 days between last application and harvest in vine crops. Do not use selective equipment in kiwi. Do not apply more than 5 quarts (3.75lb of glyphosate) of this product per acre per single application. Do not apply more than 10.67 quarts (7.95 lb of glyphosate) of this product per acre per year. When applying at reduced rates, do not apply more than 10 applications per year for established trees. Retreatment Interval (RTI): 7 days.

PASTURE GRASSES, FORAGE LEGUMES AND RANGELANDS

USE INSTRUCTIONS: This product may be applied to turf or pasture grasses, forage legumes, and rangelands for weed control as directed below. Apply 12 fluid ounces to 5 quarts per acre according to the “ANNUAL WEEDS”, “PERENNIAL WEEDS”, and “WOODY BRUSH AND TREES RATE TABLES” in the product label booklet.

RESTRICTIONS: Follow the specific limitations below with regard to application methods, timing, treatment rates, and post-application intervals. All applications must be made at least 30 days before planting any crop that is not specified for treatment in the label booklet or its supplemental labeling.

Alfalfa, Clover, and Other Forage Legumes

LABELED CROPS: Alfalfa, Clover, Kenaf, Kudzu, Lespedeza, Leucaena, Lupin, Sainfoin, Trefoil, and Velvet bean, Vetch (all types).

TYPES OF APPLICATIONS: Preplant, Pre-emergence, At-Planting, Spot Treatment, Wiper Applications Over-the-Top, Renovation, Preharvest (except Kenaf and Leucaena).

RESTRICTION: Plant Back Interval (PBI): Crops listed on this label may be planted into the treated area at any time; for any other crop, wait at least 30 days between application and planting.

Preplant, Pre-emergence, At-Planting

USE INSTRUCTIONS: This product may be applied before, during or after planting crops listed in this section. Applications must be made prior to emergence of the crop.

RESTRICTIONS: Remove domestic livestock before application.

Preharvest (except Kenaf and Leucaena)

USE INSTRUCTIONS: This product may be used in declining stands or any stand where severe crop injury or destruction is acceptable. This product will control annual and perennial weeds, including quackgrass, when applied prior to crop harvest. Applications may be made at any time of the year. For control of quackgrass, apply in the spring, late summer or fall when quackgrass is actively growing. Treatments for quackgrass must be followed by deep tillage for complete control.

RESTRICTIONS: Make only one application to an existing crop stand per year. The treated crop and weeds can be harvested and fed to livestock according to the intervals below. Application must be made a minimum of 30 days prior to the planting of any crop not listed on this label.

	Maximum Single Application Rate	Minimum Interval between application and harvesting/grazing
Alfalfa	2 quarts per acre (1.5lb of glyphosate)	36 hours
All other labeled legumes above	3 pints per acre (2.25lb of glyphosate)	3 days

This application may destroy an alfalfa stand and may severely injure or destroy other labeled crops such as clover. Preharvest application is not recommended for alfalfa grown for seed, as a reduction in germination or vigor may occur.

Spot Treatment or Wiper Applications Over-the-Top

USE INSTRUCTIONS: This product may be applied as a spot treatment or with wiper applicators. For wipers, see the “Wiper Applicators” in the “Selective Equipment” section of the product label booklet. Applications may be made in the same area at 30-day intervals.

RESTRICTIONS: For spot treatment and wiper applications, apply in areas where the movement of domestic livestock can be controlled. No more than 10 percent of the total field area should be treated at one time. Remove domestic livestock before application and wait 3 days after application before grazing livestock or harvesting. When applying at reduced rates, do not apply more than 3 applications per year. Retreatment Interval (RTI): 30 days.

Renovation

USE INSTRUCTIONS: This product may be applied as a broadcast spray to renovate existing stands of alfalfa, clover, and other labeled forage legumes. If the crop is to be grazed or harvested for feed, use up to 2 quarts (1.5lb of glyphosate) per acre in alfalfa and up to 3 pints (1.13lb of glyphosate) per acre in other labeled legumes. For complete removal of established stands of clover, it may be necessary to use the higher treatment rates listed in the “PERENNIAL WEEDS RATE TABLE” in the label booklet.

RESTRICTIONS: When treatment rates of 2 quarts (1.5lb of glyphosate) per acre for alfalfa or 3 pints (1.13lb of glyphosate) per acre for other forage legumes are used, remove domestic livestock before application and wait 3 days after application before reintroduction. If treatment rates above these levels are necessary, do not graze or harvest treated foliage for livestock feed. Crops listed for treatment in the label booklet may be planted into the treated area at any time; for other crops, wait 30 days between application and planting. Remove domestic livestock before application. Make only one application to an existing crop stand per year. Crops listed in this label may be planted into the treated area at any time; for all other crops, wait 30 days or more between application and planting

Conservation Reserve Program (CRP)

TYPES OF APPLICATIONS: Renovation (Rotating out of CRP), Site Preparation, Post-emergence Weed Control in Dormant CRP Grasses, Wiper Applications Over-the-Top.

Renovation (Rotating out of CRP), Site Preparation

USE INSTRUCTIONS: This product may be used to prepare CRP land for crop production. Refer to Federal, state or local use guides for CRP renovation recommendations. For any crop not listed for treatment in this product's label booklet, applications

must be made at least 30 days prior to planting.

Post-emergence Weed Control in Dormant CRP Grasses, Wiper Applications Over-the-Top USE INSTRUCTIONS: This product may be used to suppress competitive growth and seed production of undesirable vegetation in CRP acres. Such applications may be made with wiper application equipment or as a broadcast or spot treatment to dormant CRP grasses. For selective applications with broadcast spray equipment, apply 12 to 16 fluid ounces of the product (0.28- 0.375 lb of glyphosate) per acre in early spring before desirable CRP grasses, such as crested and tall wheatgrass, break dormancy and initiate green growth. Late fall applications can be made after desirable perennial grasses have reached dormancy.

PRECAUTIONS: Some stunting of CRP perennial grasses will occur if broadcast applications are made when plants are not dormant. No waiting period is required between application and grazing or harvesting for feed.

RESTRICTIONS: Do not apply more than 3 quarts (2.25lb of glyphosate) per acre per year onto CRP grasses. Do not apply more than 3 quarts (2.25lb of glyphosate) of this product per acre per single application onto CRP land. When applying at reduced rates, do not apply more than 3 applications per year. Retreatment Interval (RTI): 7 days. No waiting period is required between application and grazing or harvesting for feed.

Grass Seed or Sod Production

LABELED CROPS: Any grass (*Gramineae family*) except Corn, Sorghum, Sugarcane and those listed in this product's label booklet under “**Cereal and Grain Crops**”.

TYPES OF APPLICATIONS: Preplant, Pre-emergence, At-Planting, Renovation, Site Preparation, Shielded Sprayers, Wiper Applications Over-the-Top, Spot Treatments, Creating Rows in Annual Ryegrass.

Preplant, Pre-emergence, At-Planting, Renovation, Site Preparation

USE INSTRUCTIONS: This product controls most existing vegetation prior to renovating turf or forage grass seed areas or establishing turf grass grown for sod. Make applications before, during or after planting or for renovation. For maximum control of existing vegetation, delay planting to determine if any regrowth from escaped underground plant parts occurs. Where existing vegetation is growing under mowed turfgrass management, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the spray. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm-season grasses, such as Bermudagrass, summer or fall applications provide best control. Broadcast equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested.

RESTRICTIONS: Do not disturb soil or underground plant parts before treatment. Tillage or renovation techniques such as vertical mowing, coring or slicing should be delayed for 7 days after application to allow proper translocation into underground plant parts. If application rates total 3 quarts (2.25lb of glyphosate) per acre or less, no waiting period between treatment and feeding or livestock grazing is required. If the rate is greater than 3 quarts (2.25lb of glyphosate) per acre, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting. For any crop not listed for treatment in this product's label booklet, applications must be made at least 30 days prior to planting. Applications must be made prior to the emergence of the crop to avoid crop injury. Do not disturb soil or underground plant parts before treatment. Retreatment Interval (RTI): 7 days. Plant Back Interval (PBI): Crops listed on this label may be planted into the treated area at any time; for any other crop, wait at least 30 days between application and planting.

Shielded Sprayers

USE INSTRUCTIONS: Apply 1 to 3 quarts of this product (0.75-2.25lb of glyphosate) in 10 to 20 gallons of water per acre to control weeds between grass seed rows. Uniform planting in straight rows aid in shielded sprayer applications. Best results are obtained when the grass seed crop is small enough to easily pass by the protective shields. For additional instructions, see “**Shielded Applicators**” in the “**APPLICATION EQUIPMENT AND TECHNIQUES**” section of this product's label booklet.

PRECAUTIONS: Contact of this product in any manner to any vegetation to which treatment is not intended may cause damage. Such damage shall be the sole responsibility of the applicator.

Wiper Applications Over-the-Top

USE INSTRUCTIONS: Applicators should be adjusted so that the wiper contact point is at least 2 inches above the desirable vegetation. Weeds should be a minimum of 6 inches above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations, or when height of weeds varies so that not all weeds are contacted. In these instances, repeat treatments may be necessary. For additional instructions, see “**Wiper Applicators**” in the “**APPLICATION EQUIPMENT AND TECHNIQUES**” section of this product's label booklet.

PRECAUTION: Contact of the herbicide solution with desirable vegetation may result in damage or destruction.

Spot Treatments

USE INSTRUCTIONS: Use a 1.0 to 2.0 percent solution.

PRECAUTIONS: The crop receiving the spray in the treated area will be killed. Take care to avoid drift or spray outside the

target area for the same reason.

RESTRICTIONS: Apply this product prior to heading of grasses grown for seed. Handheld equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested. Do not treat more than 10 percent of the total field area. Retreatment Interval (RTI): 7 days. If application rates total 3 quarts (2.25lb of glyphosate) per acre or less, no waiting period between treatment and feeding or livestock grazing is required. If the application rate is greater than 3 quarts (2.25lb of glyphosate) per acre, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting.

Creating Rows in Annual Ryegrass

USE INSTRUCTIONS: Use 1 to 2 pints of this product (0.375 – 0.75lb of glyphosate) per acre. Use the higher rate when the ryegrass is greater than 6 inches tall. Best results are obtained when applications are made before the ryegrass reaches 6 inches in height.

RESTRICTIONS: Set nozzle heights to allow the establishment of the desired row spacing while preventing spray droplets, spray fines, or drift to contact the ryegrass plants not treated. Use of low-pressure nozzles, or drop nozzles designed to target the application over a narrow band are recommended. When applying at reduced rates, do not apply more than 8 applications per year. Retreatment Interval (RTI): 7 days.

Grower assumes all responsibility for crop losses from misapplication.

Pastures

LABELED CROPS: Any grass (*Gramineae family*) except Corn, Sorghum, Sugarcane, and those listed in this product's label booklet under “**Cereal and Grain Crops**”. Grasses that may be treated include Bahiagrass, Bermudagrass, Bluegrass, Brome, Fescue, Guinea grass, Kikuyugrass, Orchardgrass, Pangola grass, Ryegrass, Timothy, and Wheatgrass.

TYPES OF APPLICATIONS: Preplant, Pre-emergence, Spot Treatment, Wiper Applications Over-the-Top, Pasture Renovation, Post-emergent Weed Control (Broadcast Treatments).

Preplant, Pre-emergence, Pasture Renovation

USE INSTRUCTIONS: This product may be applied prior to planting or emergence of forage grasses. In addition, this product may be used to control perennial pasture species listed on this label prior to replanting.

RESTRICTIONS: If application rates total 3 quarts (2.25lb of glyphosate) per acre or less, no waiting period between treatment and feeding or livestock grazing is required. If the rate is greater than 3 quarts (2.25lb of glyphosate) per acre, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting. Crops listed for treatment in the label booklet may be planted into the treated area at any time; for other crops, wait 30 days between application and planting. When applying at reduced rates, do not apply more than 4 applications per year. Retreatment Interval (RTI): 7 days.

Spot Treatment, Wiper Applications Over-the-Top

USE INSTRUCTIONS: This product may be applied as a spot treatment or with wiper applicators in pastures. Applications may be made in the same area at 30-day intervals.

RESTRICTIONS: For spot treatments or wiper application methods using rates of 3 quarts (2.25lb of glyphosate) per acre or less, the entire field or any portion of it may be treated. When spot treatments or wiper application are made using rates above 3 quarts (2.25lb of glyphosate) per acre, no more than 10 percent of the total pasture may be treated at any one time. To achieve maximum performance, remove domestic livestock before application and wait 7 days after application before grazing livestock or harvesting.

Post-emergent Weed Control (Broadcast Treatments)

USE INSTRUCTIONS: This product may be used to suppress competitive growth and seed production of annual weeds and undesirable vegetation in pastures. For selective applications with broadcast spray equipment, apply 12 to 16 fluid ounces of this product (0.28- 0.375 lb of glyphosate) per acre in early spring before desirable perennial grasses break dormancy and initiate green growth. Late fall applications can be made after desirable perennial grasses have reached dormancy.

PRECAUTIONS: Some stunting of perennial grasses will occur if broadcast applications are made when plants are not dormant. No waiting period is required between application and grazing or harvesting for feed. Use of higher application rates will cause stand reductions.

RESTRICTIONS: Do not apply more than 3 quarts (2.25lb of glyphosate) per acre per year onto pasture grasses except for renovation uses (see instructions above). If replanting is needed due to severe stand reduction, applications must be made at least 30 days prior to planting any crop not listed for treatment in this product's label booklet.

Rangelands

TYPES OF APPLICATIONS: Post-emergence.

This product will control or suppress many annual weeds growing in perennial cool and warm-season grass rangelands. Preventing viable seed production is key to the successful control and invasion of annual grassy weeds in rangelands. Follow up applications in sequential years should eliminate most of the viable seeds. Grazing of treated areas should be delayed to encourage growth of desirable perennials. Allowing desirable perennials to flower and reseed in the treated area will encourage successful transition.

USE INSTRUCTIONS: Apply 12 to 16 fluid ounces of this product (0.28- 0.375 lb of glyphosate) per acre to control or suppress many weeds, including downy brome, cheatgrass, cereal rye and jointed goatgrass in rangelands. Apply when most brome plants are in early flower and before the plants, including seed heads, turn color. Allowing for secondary weed flushes to occur in the spring following rain events further depletes the seed reserve and encourages perennial grass conversion on weedy sites. Fall applications are possible and recommended where spring moisture is usually limited and fall germination allows for good weed growth. For medusahead, apply 16 fluid ounces of this product (0.375lb of glyphosate) per acre at the 3-leaf stage. Delaying applications beyond this stage will result in reduced or unacceptable control. Controlled burning may be useful in eliminating the thatch layer produced by slow decaying culms prior to application. Allow new growth to occur before spraying after a burn. Repeat applications in subsequent years may be necessary to eliminate the seedbank before reestablishing desirable perennial grasses in medusahead dominated rangelands.

PRECAUTIONS: Slight discoloration of the desirable grasses may occur, but they will regreen and regrow under moist soil conditions as effects of this product wear off.

RESTRICTIONS: Do not use ammonium sulfate when spraying rangeland grasses with this product. No waiting period between treatment and feeding of livestock grazing is required. Do not apply more than 3 quarts (2.25lbs of glyphosate) per acre per year. Retreatment Interval (RTI): 7 days. When applying at reduced rates, do not apply more than 6 applications per year. Do not apply more than 17.07 fluid ounces (0.398 lb of glyphosate) of this product per acre per single application.

GLYPHOSATE RESISTANT CROPS

The following instructions or those separately published on LG Chem Ltd Supplemental labeling include all applications which can be made onto the specified glyphosate resistant crops during the complete cropping season. **DO NOT** combine these instructions with other directions made for crop varieties that do not contain the glyphosate resistant gene, in the “**ANNUAL AND PERENNIAL CROPS (Alphabetical)**” section of this label. For use on glyphosate-resistant crops only. Use on crops listed in this label that do not carry a glyphosate-resistant trait will cause severe injury or plant death. The Glyphosate resistant designation indicates that the crop variety contains a patented gene that provides tolerance to this product. **NOTE: Glyphosate resistant seed, and the method of selectively controlling weeds using glyphosate on a glyphosate resistant crop, are protected under several U.S. Patents, including 5,352,605 and 5,633,435. A license to use glyphosate resistant must be obtained prior to use.** Monsanto retains ownership of the gene and process technologies, and the Purchaser of the seed receives the right to use the licensed genes and technologies subject to the limited use license conditions. Seed containing the glyphosate resistant trait cannot be used for research and demonstration, reverse engineering or in connection with herbicide registration. Property seed containing the glyphosate resistant trait cannot be saved for replanting or transferred to others for replanting. Contact your Authorized Monsanto Co. Retailer for information on obtaining a limited use license.

For ground applications with broadcast equipment, apply this product in 5 to 20 gallons of spray solution per acre. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment use flat spray nozzles. Check for even distribution of spray droplets.

For aerial applications apply this product in 3 to 15 gallons of water per acre. See the “**APPLICATION EQUIPMENT AND TECHNIQUES**” section of this label for procedures to avoid spray drift that may cause injury to any vegetation not intended for treatment. Use of appropriate buffer zones will help prevent injury to adjacent vegetation. For proper stewardship of aerial applications over-the-top of Glyphosate resistant crops, LG Chem Ltd. recommends that growers and applicators read and follow all precautions and procedures.

ATTENTION: AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN THE GLYPHOSATE RESISTANT GENE. See the “**MIXING**” and “**APPLICATION EQUIPMENT AND TECHNIQUES**” sections of this label for additional directions and restrictions on the application of this product.

Tank mixtures with other herbicides, insecticides, fungicides, micronutrients or foliar fertilizers may result in reduced weed control or crop injury and are not to be used for over-the-top applications of this product unless otherwise noted in this product label, supplemental labeling or fact sheets published separately by LG Chem Ltd. Ammonium sulfate may be mixed with this product for applications to Glyphosate resistant crops. Refer to the “**MIXING**” section for use instructions for ammonium sulfate.

Sprayer Preparation: It is important that sprayer and mixing equipment be clean and free of pesticide residue before making applications of this product. Follow the cleaning procedures specified on the label of the product(s) previously used. **THOROUGHLY CLEAN THE SPRAY TANK AND ALL LINES AND FILTERS TO ELIMINATE POTENTIAL CONTAMINATION FROM OTHER HERBICIDES PRIOR TO MIXING AND APPLYING THIS PRODUCT.**

NOTE: The following directions are based on a clean start at-planting by using a burndown application or tillage to control existing weeds before crop emergence. In no-till and stale seedbed systems, a preplant burndown treatment of this product is recommended to control existing weeds prior to crop emergence. Some weeds, such as black nightshade, broadleaf signalgrass, sicklepod, Texas panicum, sandbur, annual morningglory, woolly cupgrass, shattercane, wild proso, millet, burcucumber, and giant ragweed with

multiple germination times or suppressed (stunted) weeds may require a second application of this product for complete control. The second application should be made after some regrowth has occurred and at least 10 days after a previous application of this product.

Corn with the Glyphosate Resistant Gene

TYPES OF APPLICATIONS: Preplant, Pre-emergence, At-Planting, Post-emergence (In-Crop), Spot Treatment, Preharvest, Post-Harvest.

Maximum Allowable Combined Application Quantities Per Season	
Combined total per year for all applications	8 quarts per acre (6lb of glyphosate)
Preplant, Pre-emergence, At-Planting applications	5 quarts per acre (3.75lb of glyphosate)
Total in-crop applications from emergence through the V8 stage or 30 inches	2 quarts per acre (1.5lb of glyphosate)
Maximum preharvest application rate after maximum kernel fill is complete and the crop is physiologically mature (black layer formation) until 7 days before harvest	1 quart per acre (0.75lb of glyphosate)

Preplant, Pre-emergence, At-Planting

USE INSTRUCTIONS: This product may be applied alone or in a tank-mixture before, during or after planting com.

TANK MIXTURES: This product may be tank mixed with Alachlor and Atrazine, Acetochlor, Acetochlor and Atrazine, or Alcahlor at 50 to 100 percent of labeled rate. Refer to the specific product label and observe all precautions and limitations on the label for any pre-emergence herbicide application, including application timing restrictions, soil restrictions, minimum recropping interval and rotational guidelines—the more restrictive requirements apply.

This product may be tank-mixed with the products listed providing that the product to be tank-mixed is registered for use on this site.

NOTE: For maximum weed control, a post-emergence (in-crop) application of this product should be applied following the use of less than labeled rates of the pre-emergence residual products listed above.

Post-emergence (In-Crop)

USE INSTRUCTIONS: This product may be applied post-emergence to glyphosate resistant corn from emergence through the V8 stage (8 leaves with collars) or until corn height reaches 30 inches, whichever comes first. When applied as directed, this product controls labeled annual grass and broadleaf weeds in glyphosate resistant corn. Many perennial grasses and broadleaf weeds will be controlled or suppressed with one or more application of this product. The post-emergent application of 24 to 32 fluid ounces (0.563- 0.75lb of glyphosate) per acre of this product should be made before the weeds reach a height and/or density that the weeds become competitive with the crop, 4 inch tall weeds or less.

This product may be applied alone as a post-emergence in-crop application to provide control of emerged weeds listed on this label. If new flushes of weeds occur, a sequential application of this product at 24 to 32 fluid ounces (0.563- 0.75lb of glyphosate) per acre will control the labeled grasses and broadleaf weeds.

TANK MIXTURES: This product may be applied in tank mixture with Alachlor and Atrazine, Acetochlor, Acetochlor and Atrazine, and Alachlor at 50 to 100 percent of labeled rate. This product may be applied in tank mixture with halosulfuron-methyl; and atrazine at labeled rates. Refer to the specific product label and observe all precautions and limitations on the label for all products used in tank mixtures, including application timing restrictions, soil restrictions minimum recropping interval and rotational guidelines; the more restrictive requirements apply.

Tank-Mix Partner	Maximum Height of Corn For Application
-------------------------	---

Acetochlor Acetochlor and Atrazine,	11 inches
Alachlor and Atrazine* Alachlor*	5 inches
Halosulfuron- methyl	30 inches
Atrazine	12 inches

*Alachlor and Atrazine and Alachlor make sure the product label has a post-emergence application in Texas before using in Texas.

This product may be tank-mixed with the product listed providing that the product to be tank-mixed is registered for use on this site.

PRECAUTIONS: See the “**Glyphosate Resistant CROPS**” section of this label for precautionary instructions for use in glyphosate resistant crops.

RESTRICTIONS: Single in-crop applications of this product are not to exceed 1 quart per acre. Sequential in-crop applications of this product from emergence through the V8 stage or 30 inches must not exceed 2 quarts (1.5lb of glyphosate) per acre per growing season. Allow a minimum of 10 days between in-crop applications of this product. Allow a minimum of 50 days between application of this product and harvest of corn forage.

Preharvest

USE INSTRUCTIONS: In Glyphosate Resistant corn, up to 1 quart (0.75lb of glyphosate) per acre of this product can be applied preharvest. Make applications at 35 percent grain moisture or less. Ensure that maximum kernel fill is complete and the corn is physiologically mature (black layer formed).

RESTRICTIONS: Allow a minimum of 7 days between application and harvest.

Post-Harvest

USE INSTRUCTIONS: This product may be applied after harvest of corn. Higher rates may be required for control of large weeds that were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

RESTRICTIONS: Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

Cotton with the Glyphosate Resistant Gene

TYPES OF APPLICATIONS: Preplant, Pre-emergence, At-Planting, Post-emergence (Over-the-Top), Selective Equipment, Preharvest.

Maximum Allowable Combined Application Quantities Per Season	
Combined total per year for all applications	8 quarts per acre (6lb of glyphosate)
Preplant, Pre-emergence, At-Planting applications	5 quarts per acre (3.75lb of glyphosate)
Total in-crop applications from ground cracking to layby	4 quarts per acre (3lb of glyphosate)
Maximum preharvest application rate	2 quarts per acre (1.5lb of glyphosate)

PRECAUTIONS: See the “**Glyphosate Resistant CROPS**” section of this label for precautionary instructions for use in Glyphosate Resistant crops.

RESTRICTIONS: The combined total application of this product from cotton emergence until harvest must not exceed 6 quarts (4.5lb of glyphosate) per acre. NO MORE THAN TWO OVER-THE-TOP BROADCAST APPLICATIONS MAY BE MADE FROM CROP EMERGENCE THROUGH THE 4-LEAF (NODE) STAGE OF DEVELOPMENT. NO MORE THAN TWO APPLICATIONS SHOULD BE MADE FROM THE 5-LEAF STAGE THROUGH LAYBY. SEQUENTIAL IN-CROP OVER-THE-TOP OR POST-DIRECTED APPLICATIONS OF THIS PRODUCT MUST BE

AT LEAST 10 DAYS APART AND COTTON MUST HAVE AT LEAST TWO NODES OF INCREMENTAL GROWTH BETWEEN APPLICATIONS. ALLOW A MINIMUM OF 7 DAYS BETWEEN APPLICATION AND HARVEST.

Preplant, Pre-emergence, At-Planting

USE INSTRUCTIONS: This product may be applied before, during or after planting cotton.

Post-emergence Over-the-Top

USE INSTRUCTIONS: This product may be applied by aerial or ground application equipment at rates up to 1 quart (0.75lb of glyphosate) per acre per application post-emergence to glyphosate resistant cotton from the ground cracking stage until the 4-leaf (node) stage of development (until the fifth true leaf reaches the size of a quarter). Over-the-top applications made after the 4-leaf (node) stage of development may result in boll loss, delayed maturity and/or yield loss. Salvage Treatment. This treatment may be used after the 4-leaf stage of development and should only be used where weeds threaten to cause the loss of the crop. One quart per acre may be applied either as an over-the-top application or as a post-directed treatments sprayed higher on the cotton plants and over the weeds. NOTE: SALVAGE TREATMENTS WILL RESULT IN SIGNIFICANT BOLL LOSS, DELAYED MATURITY AND/OR YIELD LOSS. NO MORE THAN ONE SALVAGE TREATMENT SHOULD BE USED PER YEAR.

NOTE: For specific rates of application and instructions, refer to the "ANNUAL WEEDS" and "PERENNIAL WEEDS RATE TABLES" in this booklet.

PRECAUTIONS: See the "Glyphosate Resistant CROPS" section of this label for precautionary instructions for use in Glyphosate Resistant crops.

Selective Equipment

USE INSTRUCTIONS: This product may be applied using precision post-directed or hooded sprayers at rates up to 1 quart per acre per application to glyphosate resistant cotton through layby. At this stage, post-directed equipment should be used which directs the spray to the base of the cotton plants. Contact of the spray with cotton leaves should be avoided to the maximum extent possible. To minimize spray onto the leaves of the cotton plants, place nozzles in a low position directing a horizontal spray pattern under the cotton leaves to contact weeds in the row, and maintain low spray pressure (less than 30 psi). For best results, make applications while weeds are small (less than 3 inches).

PRECAUTIONS: See the "Selective Equipment" part of the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for information on proper use and calibration of this equipment.

Preharvest

USE INSTRUCTIONS: This product may be applied for preharvest annual and perennial weed control as a broadcast treatment to glyphosate resistant cotton after 20 percent boll crack. Up to 2 quarts of this product (1.5lb of glyphosate) may be applied using either aerial or ground spray equipment.

TANK MIXTURES: This product may be tank-mixed with S,S,S-Tributyl phosphorothioate, Thidiazuron and Diuron, or Ethepon (2-Chloroethyl phosphonic acid). NOTE: This product will not enhance the performance of these harvest aids when applied to glyphosate resistant cotton.

RESTRICTIONS: Allow a minimum of 7 days between application and harvest of cotton. Do not apply this product to cotton grown for seed, as a reduction in germination or vigor may occur. **THE USE OF ADDITIVES, OTHER THAN THOSE LISTED ON THIS LABEL, FOR PREHARVEST APPLICATION TO COTTON IS PROHIBITED.**

ATTENTION: USE OF THIS PRODUCT IN ACCORDANCE WITH LABEL DIRECTIONS IS EXPECTED TO RESULT IN NORMAL GROWTH OF GLYPHOSATE RESISTANT COTTON, HOWEVER, VARIOUS ENVIRONMENTAL CONDITIONS, AGRONOMIC PRACTICES AND OTHER FACTORS MAKE IT IMPOSSIBLE TO ELIMINATE ALL RISKS ASSOCIATED WITH THIS PRODUCT, EVEN WHEN APPLICATIONS ARE MADE IN CONFORMANCE WITH THE LABEL SPECIFICATIONS. IN SOME CASES, THESE FACTORS CAN RESULT IN BOLL LOSS, DELAYED MATURITY AND/OR YIELD LOSS.

Flex Cotton with Glyphosate Resistant Gene

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS (EXCEPT AS SPECIFIED FOR INDIVIDUAL GLYPHOSATE RESISTANT CROPS), DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See PRODUCT INFORMATION and MIXING, ADDITIVES, and APPLICATION INSTRUCTIONS sections for essential product performance information.

SPECIALLY FORMULATED FOR EXPANDED GLYPHOSATE RESISTANT COTTON USES.

ATTENTION: USE OF THIS PRODUCT IN ACCORDANCE WITH LABEL DIRECTIONS IS EXPECTED RESULT IN NORMAL GROWTH OF GLYPHOSATE RESISTANT COTTON. HOWEVER, DUE TO THE SENSITIVITY OF COTTON FRUITING TO VARIOUS ENVIRONMENTAL CONDITIONS, AGRONOMIC PRACTICES AND OTHER FACTORS, IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS ASSOCIATED WITH THIS PRODUCT, EVEN WHEN APPLICATIONS ARE MADE IN CONFORMANCE WITH THE LABEL SPECIFICATIONS. IN SOME CASES, THESE FACTORS CAN RESULT IN BOLL LOSS, DELAYED MATURITY AND/OR YIELD LOSS.

The use of the over-the-top applications described in this section on other than Glyphosate Resistant cotton will cause crop injury and reduced yields. Drift of this product from applications made to Glyphosate Resistant cotton onto adjacent fields of post 4-leaf (node) Glyphosate Resistant cotton may cause extensive injury including boll loss, delayed maturity and/or yield loss.

NOTE: The instructions provided in this section are specific to, and should only be used with varieties designated as Glyphosate Resistant cotton. **DO NOT** combine the instructions in this section with those in the **GLYPHOSATE RESISTANT COTTON** section.

TYPES OF APPLICATIONS: preplant, At-Planting, Preemergence, Post-emergence, Preharvest.

<u>Maximum Allowable Combined Application Quantities Per Season</u>	
Combined total per year for all applications	8 quarts per acre (6lb of glyphosate)
Calculate the combined rate to be used for all preplant, in-crop and preharvest applications, to ensure that the total does not exceed the maximum allowed rate per acre per year shown above.	
Preplant, At-Planting, Pre-emergence applications	5 quarts per acre (3.75lb of glyphosate)
Total in-crop applications from ground cracking to 60 percent open bolls	6 quarts per acre (4.5lb of glyphosate)
Maximum allowed from 60 percent bolls open to 7 days prior to harvest	2 quarts per acre (1.5lb of glyphosate)

PRECAUTIONS: See the GLYPHATE RESISTANT CROPS section of the (product brand name) label booklet for precautionary instructions for use in GLYPHOSATE RESISTANT crops.

Preplant, Pre-emergence, At-Planting

USE INSTRUCTIONS: This product may be applied before, during or after planting Glyphosate Resistant cotton.

Post-emergence

USE INSTRUCTIONS: When applied in accordance with the label, **GlamPrime** will control labeled annual grasses and broadleaf weeds in Glyphosate Resistant cotton. To maximize yield potential, spray cotton early to eliminate competing weed. Many perennial weeds will be controlled or suppressed with one or more applications of this product. In an initial application of 1 quart (0.75lb of glyphosate) per acre on 1 to 3 inch tall annual grass and broadleaf weeds is recommended. This product may be applied by ground application equipment at rates up to 1.5 quarts (1.13 lb of glyphosate) per acre per application post-emergence to Glyphosate Resistant cotton. In addition to broadcast applications, post-directed equipment may be used to achieve weed coverage.

NOTE: for specific rates of application and instruction, refer to the ANNUAL and PERENNIAL WEEDS RATE TABLES.

RESTRICTIONS: The maximum rate for any single in-crop application of this product is 1.5 quarts (1.13 lb of glyphosate) per acre made using ground application equipment. In-crop application rates above 32 fluid ounces (0.75lb of

glyphosate) per acre made alone or with the addition of other crop chemical products containing surfactant may cause a crop response including leaf speckling of leaf necrosis. Do not exceed a maximum rate of 1 quart (0.75lb of glyphosate) per acre of this product when making applications by air. Between layby and 60 percent open bolls, the maximum combined total rate of this product that may be applied is 2 quarts (1.5lb of glyphosate) per acre. The maximum combined total of all applications made from crop emergence to 60 percent open bolls must not exceed 6 quarts per acre.

Pre-harvest

USE INSTRUCTIONS: This product may be applied for pre-harvest annual and perennial weed control as a broadcast treatment to Glyphosate Resistant cotton after 60 percent boll crack. Up to 52 ounces of this product may be applied using either aerial or ground spray equipment.

NOTE: This product will not enhance the performance of harvest aids when applied to Glyphosate Resistant cotton.

RESTRICTIONS: Allow a minimum of 7 days between application and harvest of Glyphosate Resistant cotton. Do not apply this product over-the-top beyond first bloom to cotton grown for seed.

Soybeans with the Glyphosate Resistance Gene

TYPES OF APPLICATIONS: Preplant, Pre-emergence, At-Planting, Post-emergence (In-Crop), Preharvest, Post - Harvest.

Maximum Allowable Combined Application Quantities Per Season	
Combined total per year for all applications	8 quarts per acre (6lb of glyphosate)
Preplant, Pre-emergence, At-Planting applications	5 quarts per acre (3.75lb of glyphosate)
Total in-crop applications from cracking throughout flowering	3 quarts per acre (2.25lb of glyphosate)
Maximum preharvest application rate	1 quart per acre (0.75lb of glyphosate)

PRECAUTIONS: See the “GLYPHOSATE RESISTANT CROPS” section of this label for precautionary instructions for use in Glyphosate Resistant crops.

Preplant, Pre-emergence, At-Planting

USE INSTRUCTIONS: This product may be applied before, during or after planting soybeans.

Post-emergence (In-Crop)

USE INSTRUCTIONS: When applied as directed, this product will control labeled annual grasses and broadleaf weeds in Glyphosate Resistant soybeans. Applications of this product can be made in Glyphosate Resistant soybeans from emergence (cracking) throughout flowering. Refer to the “ANNUAL WEEDS RATE TABLE” in this label for rate recommendations for specific annual weeds. An initial application of 1 quart per acre on 2- to 8- inch tall weeds is recommended. Weeds will be 2 to 8 inches tall, 2 to 5 weeks after planting. If the initial application is delayed and weeds are larger, apply a higher rate of this product. This product may be used up to 2 quarts per acre in any single in-crop application for control of annual weeds and where heavy weed densities exist. A 1 to 2 quarts (0.75-1.5lb of glyphosate) per acre rate (single or multiple applications) of this product will control or suppress perennial weeds such as: Bermudagrass, Canada thistle, common milkweed, field bindweed, hemp dogbane, horsenettle, maretail, (horseweed), nutsedge, quackgrass, rhizome johnsongrass, redvine, trumpetcreeper, swamp smartweed, and wirestem muhly. For best results, allow perennial weed species to achieve at least 6 inches of growth before spraying with this product.

Under adverse growing conditions such as drought, hail, wind damage or a poor soybean stand that slows or delays canopy closure, a sequential application of this may be necessary to control late flushes of weeds.

IN THE SOUTHERN STATES, A SEQUENTIAL APPLICATION OF THIS PRODUCT WILL BE REQUIRED TO CONTROL NEW FLUSHES OF WEEDS IN THE GLYPHOSATE RESISTANT SOYBEAN CROP. To control giant ragweed, it is recommended that 1 quart per acre of this product be applied when the weed is 8 to 12 inches tall to increase control and possibly avoid the need for a sequential application.

NOTE: The use of this product for in-crop applications over Glyphosate Resistant soybeans is not registered in California.

RESTRICTIONS: The combined total application from crop emergence through harvest must not exceed 3 quarts (2.25lb of glyphosate) per acre. The maximum rate for any single in-crop application is 2 quarts (1.5lb of glyphosate) per acre. The maximum combined total of this product that can be applied during flowering is 2 quarts (1.5lb of glyphosate) per acre.

Preharvest

USE INSTRUCTIONS: This product provides weed control when applied prior to harvest of soybeans. Up to 1 quart per acre of this product can be applied by aerial or ground application.

PRECAUTIONS: Care should be taken to avoid excessive seed shatter loss due to ground application equipment.

RESTRICTIONS: Allow a minimum of 14 days between final application and harvest of soybean grain or feeding of soybean grain, forage or hay.

Post-Harvest

USE INSTRUCTIONS: This product may be applied after harvest of Glyphosate Resistant soybeans. Higher rates may be required for control of large weeds that were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

Canola with the Glyphosate Resistant Gene

See PRODUCT INFORMATION and MIXING sections of this label booklet for essential product performance information.

INFORMATION

USE ONLY ON CANOLA WHICH CONTAINS THE GLYPHOSATE RESISTANT, GENE DO NOT USE THIS PRODUCT ON CANOLA WITH THE GLYPHOSATE RESISTANT GENE PLANTED IN THE FOLLOWING STATES: ALABAMA, DELAWARE, FLORIDA, GEORGIA, KENTUCKY, MARYLAND, NEW JERSEY, NORTH CAROLINA, SOUTH CAROLINA, TENNESSEE, VIRGINIA, AND WEST VIRGINIA.

Applying this product to canola which is not designated as Glyphosate Resistant will result in severe crop injury and yield loss. Avoid contact with foliage, green stems, or fruit of crops, or any desirable plants which do not contain the Glyphosate Resistant gene since severe injury or destruction will result. The Glyphosate Resistant designation indicates the canola contains a patented gene which provides tolerance to this herbicide. Information on Glyphosate Resistant canola may be obtained from your seed supplier.

USE DIRECTIONS

This product will control many troublesome emerged weeds when applied preplant, pre-emergent and/or with over-the-top applications in Glyphosate Resistant canola. Allow a minimum of 60 days between last application and canola harvest.

<u>Maximum Allowable Combined Yearly Rates</u>	
Preplant and Pre-emergence applications	2 quarts/acre (1.5lb of glyphosate)
Total in-crop application from emergence to 6-leaf	1 quart/acre (0.75lb of glyphosate)

For ground applications with broadcast equipment, apply this product in 5 to 20 gallons of spray solution per acre. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment use flat fan nozzles. Check for even distribution of spray droplets.

For aerial applications apply this product in 3 to 15 gallons of water per acre.

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN THE GLYPHOSATE RESISTANT GENE. DO NOT APPLY DURING LOW-LEVEL INVERSION CONDITIONS, WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITIONS WHICH FAVOR DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION

CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Do not allow the herbicide solution to mist, drip, drift, or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of injury occurring from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing or when there are other meteorological conditions that favor spray drift. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist), which are likely to drift. AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations, which dispense spray as fine spray droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure.

There are no rotational crop restrictions following applications of this product.

Sprayer Preparation

It is important that sprayer and mixing equipment be clean and free of pesticide residue before making applications of this product to Glyphosate Resistant canola. Follow the cleaning procedures specified on the label of the product(s) previously used. Canola can be very sensitive to many herbicides at extremely low concentrations and care should be taken to thoroughly clean all equipment prior to use.

Preplant or Pre-emergent Applications

This product may be applied by aerial or ground application equipment prior to planting or emergence of canola. The maximum combined application rate from all preplant and pre-emergent applications should not exceed 2 quarts per acre per season.

NOTE: In no-till and stale seedbed systems, always use a burndown treatment to control existing weeds before canola emerges. Apply a preplant burndown treatment of 16 to 32 fluid ounces (0.375 – 0.75lb of glyphosate) per acre of this product.

Over-the-top Applications

This product may be applied by aerial or ground application equipment post-emergence to Glyphosate Resistant canola from emergence through the 6-leaf stage of development. Applications made during bolting or flowering may result in crop injury and yield loss. To maximize yield potential, make applications early to eliminate competing weeds.

Single Application: Apply 16 to 24 ounces (0.375 – 0.563lb of glyphosate) per acre no later than the 6-leaf stage for the control of annual weeds. Avoid overlapping applications, which may result in temporary yellowing, delaying flowering, and or growth reduction. Similar injury may result when applications of more than 16 ounces per acre are applied after the 4-leaf stage.

Sequential Applications: Apply 16 ounces (0.375 of glyphosate) per acre to 1- to 3-leaf canola followed by a sequential application at a minimum interval of 10 days, but no later than the 6-leaf stage. Sequential applications are recommended for early emerging annual weeds and perennial weeds such as Canada thistle and quackgrass.

This product will control or suppress most perennial weeds. For some perennial weeds, repeat applications may be required to eliminate crop competition throughout the growing season.

No more than two over-the-top broadcast applications may be made from crop emergence through the 6-leaf stage of development and the total in-crop application should not exceed 32 ounces (0.75lb of glyphosate) per acre.

WEED CONTROL DIRECTIONS

For specific rates of application and instructions for control of various annual and perennial weeds, refer to the WEEDS CONTROLLED section of this label booklet.

Tank mixtures with other herbicides, insecticides, or fungicides may result in reduced weed control or crop injury and are not recommended for over-the-top applications of this product.

Some weeds with multiple germination times or suppressed (stunted) weeds may require sequential applications of this product for control. The second application should be made after some regrowth has occurred and at least 10 days after a previous application of this product.

Sugar Beets with the Glyphosate Resistance Gene

TYPES OF APPLICATIONS: Preplant, Pre-emergence, At-Planting, Post-emergence (In-Crop).

Maximum Allowable Combined Application Quantities Per Season	
Combined total per year for all applications	8 quarts per acre (6lb of glyphosate)
Preplant, Pre-emergence, At-Planting applications	5 quarts per acre (3.75lb of glyphosate)
Emergence to 8-leaf stage	2.5 quarts per acre (1.88lb of glyphosate)
Between 8-leaf stage and canopy closure	2 quarts per acre (1.5lb of glyphosate)

PRECAUTIONS: See the “GLYPHOSATE RESISTANCE CROPS” section of this label for precautionary instructions for use in Glyphosate Resistant crops.

RESTRICTIONS: The combined total application from crop emergence through harvest must not exceed 4.5 quarts (3.375lb of glyphosate) per acre. The maximum rate for any single application from crop emergence until the 8-leaf stage is 1.5 quarts (1.13lb of glyphosate) per acre. The maximum rate for any single application between the 8-leaf stage and canopy closure is 1 quart (0.75lb of glyphosate) per acre. Allow a minimum of 30 days between last application and sugar beet harvest.

Preplant, Pre-emergence, At-Planting

USE INSTRUCTIONS: This product may be applied before, during or after planting of Glyphosate Resistant sugar beets.

Post-emergence (In-crop)

USE INSTRUCTIONS: This product may be applied post-emergent over-the-top to Glyphosate Resistant sugar beets from emergence to 30 days prior to harvest. To maximize yield potential spray sugar beets early to eliminate competing weeds. Up to 4 sequential applications of this product may be made with at least 10 days between applications. Refer to the “ANNUAL WEEDS RATE TABLE” in this label for rate recommendations for specific annual weeds. This product will control or suppress, most perennial weeds. For some perennial weeds, repeat applications may be required to eliminate crop competition throughout the growing season.

NON-CROP USES AROUND THE FARMSTEAD

TYPES OF APPLICATIONS: Non-Selective Weed Control, Trim-and-Edge, Greenhouse/Shadehouse, Chemical Mowing, Cut Stumps, Habitat Management.

Weed Control and Trim-and-Edge

USE INSTRUCTIONS: This product may be used to control annual weeds, perennial weeds and woody brush which are found in any part of the farmstead, including building foundations, along and in fences, in dry ditches and canals, along ditchbanks, farm roads, and shelterbelts, prior to landscape plantings and equipment storage areas.

TANK MIXTURES: This product may be tank-mixed with the following products. Refer to these product labels for approved farmstead sites and application rates. For annual weeds, use 1 quart per acre of this product when weeds are less than 6 inches tall, 1.5 quarts per acre when weeds are 6 to 12 inches tall and 2 quarts per acre when weeds are greater than 12 inches tall. For perennial weeds, apply 2 to 5 quarts (1.5-3.75lb of glyphosate) per acre in these tank mixes. For tank-mixtures with these products

through backpack sprayers, handguns or other high-volume spray-to-wet applications, see the “ANNUAL WEED-HAND-HELD OR HIGH-VOLUME EQUIPMENT” section of this label for recommended rates.

Isopropylamine salt of imazapyr	Metsulfuron methyl bromacil and diuron	Pendimethalin ammonium salt of imazapic	oxadiazon imazapyr and
Dicamba/diglycolamine salt of 3,6-dicchloro-o anisic acid	Sulfometuron methyl and metsulfuron methyl	Simazine	Diuron
Mesotrione		Oryzaline	Simazine
Diuron		Chlorsulfuron	Dicamba
Prodiamine		2,4-D	

This product plus dicamba tank mixtures may not be applied by air in California.

Greenhouse/Shadehouse

This product may be used to control weeds in and around greenhouses and shadehouses. Desirable vegetation must not be present during application and air circulation fans must be turned off.

Chemical Mowing

USE INSTRUCTIONS: This product will suppress perennial grasses listed in this section to serve as a substitute for mowing. Use 6 fluid ounces of this product (0.141lb of glyphosate) per acre when treating Kentucky bluegrass. Use 8 fluid ounces of this product (0.188lb of glyphosate) per acre when treating tall fescue, fine fescue, orchardgrass, bahiagrass, or quackgrass covers. Use 16 fluid ounces of this product (0.375lb of glyphosate) per acre when treating Bermudagrass. Use 64 fluid ounces of this product (1.5lb of glyphosate) per acre when treating torpedograss or paragrass. Apply treatments in 10 to 20 gallons of spray solution per acre. Chemical mowing applications may be made along farm ditches and other parts of farmsteads.

PRECAUTIONS: Use only in areas where some temporary injury or discoloration of perennial grasses can be tolerated.

Cut Stumps

TYPES OF APPLICATION: Treating cut stumps in any non-crop site listed on this label.

USE INSTRUCTIONS: This product will control regrowth of cut stumps and resprouts of many types of woody brush and tree species, some of which are listed below. Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut trees or resprouts close to the soil surface. Apply a 50 to 100 percent solution of this product to the freshly cut surface immediately after cutting. Delays in application may result in reduced performance. For best results, applications should be made during periods of active growth and full leaf expansion.

Alder	Oak	Reed, giant	Tan Oak
Eucalyptus	Pepper, Brazilian	Saltcedar	Willow
Madrone	Pine, Austrian	Sweetgum	

PRECAUTIONS: Some sprouts, stems, or trees may share the same root system. Adjacent trees having a similar age, height and spacing may signal shared roots. Whether grafted or shared, injury is likely to occur to non-treated stems/trees when one or more trees sharing common roots are treated.

RESTRICTIONS: Do not make cut stump applications when the roots of desirable woody brush or trees may be grafted to the roots of the cut stump.

Habitat Management

TYPES OF USES: Habitat Restoration and Maintenance, Wildlife Food Plots.

Habitat Restoration and Maintenance

USE INSTRUCTIONS: This product may be used to control exotic and other undesirable vegetation in habitat management areas. Applications can be made to allow recovery of native plant species, prior to planting desirable native species, and for similar broad-spectrum vegetation control requirements in habitat management areas. Spot treatments can be made to selectively remove unwanted plants for habitat maintenance and enhancement.

Wildlife Food Plots

USE INSTRUCTIONS: This product may be used as a site preparation treatment to control annual and perennial weeds prior to planting wildlife food plots. Any wildlife food species may be planted after applying this product, or native species may be allowed to repopulate the area. If tillage is needed to prepare a seedbed, wait 7 days after application before tillage.

ANNUAL WEEDS RATE TABLE (Alphabetically by Species)

WATER CARRIER VOLUMES OF 3 TO 10 GALLONS PER ACRE FOR GROUND APPLICATIONS AND 3 TO 5 GALLONS PER ACRE FOR AERIAL APPLICATIONS ARE RECOMMENDED.

Apply to actively growing annual weeds. Annual weeds are easiest to control when they are small. Older, mature (hardened) annual weed species may require higher rates even if they meet the size requirements. Do not tank mix with soil residual herbicides when using these rates unless otherwise specified.

For weeds that have been mowed, grazed or cut, allow regrowth to occur prior to treatment.

This product may be used up to 48 fluid ounces (1.13lb of glyphosate) per acre where heavy weed densities exist.

ANNUAL WEEDS RATE TABLE RATE (fluid ounces per acre)

WEED SPECIES	Maximum height/length (in inches)				
	16	24	32	40	48
Ammannia, purple	3	6	12	-	18
Annoda, spurred	-	2	3	5	8
Barley	18	18+	-	-	-
Barnyardgrass	-	3	6	7	9
Bassia, fivehook	-	-	6	-	-
Beggarweed, Florida	-	5	8	-	-
Bittercress	12	20	-	-	-
Bluegrass, annual	10	-	-	-	-
Bluegrass, bulbous	6	-	-	-	-
Brome, downy ^{1,2}	6	12	-	-	-
Brome, Japanese	6	12	24	-	-
Browntop panicum	6	8	12	-	24
Buckwheat, wild ³	-	1	2	-	-
Burcucumber	-	6	12	-	18
Buttercup	12	20	-	-	-
Carolina geranium	-	-	4	-	9
Carpetweed	-	6	12	-	-
Cheat ²	6	20	-	-	-
Chervil	20	-	-	-	-
Chickweed	-	12	18	-	-
Cocklebur	12	18	24	-	36
Copperleaf, hophornbeam	-	2	4	-	6
Copperleaf, Virginia	-	2	4	-	6
Coreopsis, plains	-	6	12	-	18
Corn, volunteer	6	12	20	-	-
Corn, speedwell	12	-	-	-	-
Crabgrass	3	6	12	-	-
Crowfootgrass	-	-	6	-	12
Cutleaf evening primrose	-	-	3	-	6

Devils claw (unicorn plant)	-	3	6	-	-
Dwarf dandelion	12	-	-	-	-
Eastern manna grass	8	12	-	-	-

ANNUAL WEEDS RATE TABLE
RATE
(fluid ounces per acre)

WEED SPECIES	Maximum height/length (in inches)				
	16	24	32	40	48
Eclipta	-	4	8	12	-
Fall panicum	4	-	6	-	12
False dandelion	-	20	-	-	-
False flax, smallseed	12	-	-	-	-
Fiddleneck	-	6	12	-	-
Field pennycress	8	12	-	-	-
Filaree	-	-	6	-	12
Fleabane, annual	6	20	-	-	-
Fleabane, hairy (<i>Conyza bonariensis</i>)	-	-	6	-	10
Fleabane, rough	3	6	12	-	-
Florida pusley	-	-	4	-	6
Foxtail, giant, bristly, yellow	6	12	20	-	-
Foxtail, Carolina	10	-	-	-	-
Foxtail, green	12	-	-	-	-
Goatgrass, jointed	-	12	-	-	-
Goosegrass	-	3	6	-	12
Grain sorghum (milo)	6	12	20	-	-
Groundcherry	-	3	6	-	9
Groundsel, common	-	6	10	-	-
Hemp sesbania	-	2	4	6	8
Henbit	-	-	6	-	12
Horseweed/ Mares tail (<i>Conyza canadensis</i>)	-	6	12	-	18
Itchgrass	6	8	12	-	18
Jimsonweed	-	-	12	-	18
Johnsongrass, seedling	6	12	18	-	24
Junglerice	-	3	6	7	9
Knotweed	-	-	6	-	12
Kochia ⁴	-	3	12	-	-
Lambsquarters	-	6	12	-	20
Little barley	6	12	-	-	-
London Rocket	6	-	24	-	-
Mayweed	-	2	6	12	18
Morningglory, annual (<i>Ipomoea</i> spp.)	-	-	3	-	6
Mustard, blue	6	12	18	-	-
Mustard, tansy	6	12	18	-	-
Mustard, tumble	6	12	18	-	-
Mustard, wild	6	12	18	-	-

Nightshade, black	-	4	6	-	12
WEED SPECIES	Maximum height/length (in inches)				
	16	24	32	40	48
Eclipta	-	4	8	12	-
Fall panicum	4	-	6	-	12
False dandelion	-	20	-	-	-
False flax, smallseed	12	-	-	-	-
Fiddleneck	-	6	12	-	-
Field pennycress	8	12	-	-	-
Filaree	-	-	6	-	12
Fleabane, annual	6	20	-	-	-
Fleabane, hairy (<i>Conyza bonariensis</i>)	-	-	6	-	10
Fleabane, rough	3	6	12	-	-
Florida pusley	-	-	4	-	6
Foxtail, giant, bristly, yellow	6	12	20	-	-
Foxtail, Carolina	10	-	-	-	-
Foxtail, green	12	-	-	-	-
Goatgrass, jointed	-	12	-	-	-
Goosegrass	-	3	6	-	12
Grain sorghum (milo)	6	12	20	-	-
Groundcherry	-	3	6	-	9
Groundsel, common	-	6	10	-	-
Hemp sesbania	-	2	4	6	8
Henbit	-	-	6	-	12
Horseweed/ Mares tail (<i>Conyza canadensis</i>)	-	6	12	-	18
Itchgrass	6	8	12	-	18
Jimsonweed	-	-	12	-	18
Johnsongrass, seedling	6	12	18	-	24
Junglerice	-	3	6	7	9
Knotweed	-	-	6	-	12
Kochia ⁴	-	3	12	-	-
Lambsquarters	-	6	12	-	20
Little barley	6	12	-	-	-
London Rocket	6	-	24	-	-
Mayweed	-	2	6	12	18
Morningglory, annual (<i>Ipomoea</i> spp.)	-	-	3	-	6
Mustard, blue	6	12	18	-	-
Mustard, tansy	6	12	18	-	-
Mustard, tumble	6	12	18	-	-
Mustard, wild	6	12	18	-	-
Nightshade, black	-	4	6	-	12

ANNUAL WEEDS RATE TABLE
RATE
(fluid ounces per acre)

WEED SPECIES	Maximum height/length (in inches)				
	16	24	32	40	48
Nightshade, hairy	-	4	6	-	12
Oats	3	6	18	-	-
Pigweed species	-	12	18	24	-
Prickly lettuce	-	6	12	-	-
Purslane	-	-	3	-	6
Ragweed, common	-	6	12	-	18
Ragweed, giant	-	6	12	-	18
Red rice	-	-	4	-	-
Rye, volunteer/cereal ²	6	18	18+	-	-
Ryegrass	-	-	6	-	12
Sandbur, field	6	12	-	-	-
Sandbur, longspine	6	12	-	-	-
Shattercane	6	12	20	-	-
Sicklepod	-	2	4	-	8
Signalgrass, broadleaf	-	3	6	7	9
Smartweed, ladysthumb	-	-	6	-	9
Smartweed, Pennsylvania	-	-	6	-	9
Sowthistle, annual	-	-	6	-	12
Spanishneedles	-	-	6	-	12
Speedwell, purslane	12	-	-	-	-
Sprangletop	6	12	20	-	-
Spurge, prostrate	-	6	12	-	-
Spurge, spotted	-	6	12	-	-
Spurry, umbrella	6	-	-	-	-
Stinkgrass	-	12	-	-	-
Sunflower	12	18	-	-	-
Swinecress	-	5	12	-	-
Teaweed/Prickly sida	-	2	4	-	6
Texas panicum	6	8	12	-	24
Thistle, Russian ⁵	-	6	12	-	-
Velvetleaf	-	-	6	-	12
Virginia pepperweed	-	18	-	-	-
Waterhemp	-	-	6	-	12
Wheat ²	6	12	18	-	-
Wheat (overwintered)	-	6	12	-	18
Wild oats	3	6	18	-	-
Wild proso millet	-	6	12	-	-
Witchgrass	-	12	-	-	-
Woolly cupgrass	-	6	12	-	-
Yellow rocket	-	12	20	-	-

¹ For control of downy brome in no-till systems, use 24 fluid ounces (0.563lb of glyphosate) per acre.

² Performance is better if application is made before this weed reaches the boot stage of growth.

³ Use 24 fluid ounces (0.563lb of glyphosate) per acre of this product to control wild buckwheat in the cotyledon to 2-leaf stage. Use 32 fluid ounces (0.75lb of glyphosate) per acre to control 2- to 4-leaf wild buckwheat. For improved control of wild buckwheat over 2 inches in size, use sequential treatments of 32 fluid ounces (0.75lb of glyphosate) followed by 32 fluid ounces of this product (0.75lb of glyphosate) per acre.

⁴ Do not treat kochia in the button stage.

⁵ Control of Russian Thistle may vary based on environmental conditions and spray coverage. Whenever possible, a tank mixture with 2,4-D as described below may improve control.

Annual Weeds—Rates for 10 to 40 Gallons per Acre

Apply 1 to 2 quarts of this product (0.75-1.5lb of glyphosate) per acre. Use 1 quart per acre if weeds are less than 6 inches tall, 1.5 quarts (1.13lb of glyphosate) per acre if weeds are 6 to 12 inches tall and 2 quarts (1.5lb of glyphosate) per acre if weeds are greater than 12 inches tall. These rates will provide control of weeds listed in the “ANNUAL WEEDS RATE TABLE” when water carrier volume are 10 to 40 gallons per acre for ground applications. Older, mature (hardened) annual weed species may require higher rates even if they meet the size requirements.

Annual Weeds—Tank Mixtures with 2,4-D, Dicamba, or Picloram

12 to 16 fluid ounces of this product (0.28 – 0.375lb of glyphosate) plus 0.25 pound of dicamba or 0.5 pound of 2,4-D or 1 to 2 fluid ounces of Picloram per acre will control the following weeds with the maximum height or length indicated: 6 inches prickly lettuce, marestail/horseweed, morning glory, kochia (dicamba only) wild buckwheat (Picloram Picloram only); 12 inches cocklebur, lambsquarters, pigweed, and Russian thistle (2,4-D only). 16 fluid ounces of this product (0.375 lbs of glyphosate) plus 0.5 pound of 2,4-D per acre will control the following weeds when they are a maximum height or length of 6 inches: common ragweed, giant ragweed, Pennsylvania smartweed, and velvetleaf.

Refer to the specific product labels for crop rotation restrictions and cautionary statements of all products used in tank mixtures. Some crop injury may occur if dicamba or Picloram is applied within 45 days of planting.

DO NOT APPLY DICAMBA TANK MIXTURES BY AIR IN CALIFORNIA.

Annual Weeds—Hand-Held or High-Volume Equipment

For control of weeds listed in the “ANNUAL WEEDS RATE TABLE”, apply a 0.5 percent solution of this product to weeds less than 6 inches in height or runner length. Apply prior to seedhead formation in grass or bud formation in broadleaf weeds. For annual weeds over 6 inches tall, or unless otherwise specified, use a 1 percent solution. For best results, use a 2 percent solution on harder-to-control perennials, such as Bermudagrass, dock, field bindweed, hemp dogbane, milkweed, and Canada thistle. When using application methods that result in less than complete coverage, use a 5 percent solution for annual and perennial weeds and a 5 to 10 percent solution for woody brush and trees.

Annual Weeds—Tank Mixtures with Atrazine for Fallow and Reduced Tillage Systems

For use only in Colorado, Kansas, Nebraska, Oklahoma, Oregon, South Dakota, and Washington. In Oregon and Washington, do not exceed 1 pound of atrazine per acre.

24 to 28 fluid ounces of this product (0.563 – 0.656lb of glyphosate) plus 1 to 2 pounds of atrazine per acre will control the following weeds: Barnyardgrass (requires 28 ounces for control), Downy brome, Green foxtail, Lambsquarters, Prickly lettuce, Tansy mustard, Pigweed, Field sandbur, Stinkgrass, Russian thistle, Volunteer wheat, Witchgrass, and Kochia (add 0.125 pound of dicamba for control).

PERENNIAL WEEDS RATE TABLE (Alphabetically by Species)

Apply to actively growing perennial weeds.

NOTE: If weeds have been mowed or tilled, do not treat until plants have resumed active growth and have reached the recommended stages.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seed. Repeat treatments must be made prior to crop emergence. Unless otherwise stated, allow 7 or more days after application before tillage. Best results are obtained when soil moisture is adequate for active weed growth.

Weed Species	Rate (QT./A)	Water Volume (GPA)	Hand-Held % Solution
--------------	-----------------	-----------------------	-------------------------

Alfalfa	1-2 (0.75-1.5lb of glyphosate)	3-10	2%
----------------	--------------------------------------	------	----

Make applications after the last hay cutting in the fall. Allow alfalfa to regrow to a height of 6 to 8 inches or more prior to treatment. Applications should be followed with deep tillage at least 7 days after treatment, but before soil freeze-up.

Alligatorweed	4 (3lb of glyphosate)	3-20	1.5%
----------------------	--------------------------	------	------

For partial control, apply when most of the plants are in bloom. Repeat applications will be required to maintain control.

Anise (fennel)	—	—	1-2%
-----------------------	---	---	------

Apply as a spray-to-wet treatment. Optimum results are obtained when plants are treated at the bud to full-bloom stage of growth.

Bahiagrass	3-5 (2.25 – 3.75lb of glyphosate)	3-20	2%
-------------------	--------------------------------------	------	----

Apply when most plants have reached the early head stage.

Bentgrass	1.5 (1.13lb of glyphosate)	10-20	2%
------------------	-------------------------------	-------	----

For suppression in grass seed production areas. For ground applications only. Ensure entire crown area has resumed growth prior to a fall application. Bentgrass should have at least 3 inches of growth. Tillage prior to treatment should be avoided. Tillage 7 to 10 days after application is recommended for best results.

Bermudagrass	3-5 (2.25 – 3.75lb of glyphosate)	3-20	2%
---------------------	--------------------------------------	------	----

For control, apply 5 quarts of this product (3.75lb of glyphosate) per acre. For partial control, apply 3 quarts per acre. Treat when Bermudagrass is actively growing and seedheads are present. Retreatment may be necessary to maintain control.

Bermudagrass water (knotgrass)	1-1.5 (0.75-1.13lb of glyphosate)	5-10	2%
---------------------------------------	--------------------------------------	------	----

Apply 1.5 quarts of this product (1.13lb of glyphosate) in 5 to 10 gallons of water per acre. Apply when water Bermudagrass is 12 to 18 inches in length. Allow 7 or more days before tilling, flushing or flooding the field.

Fall applications only: Apply 1 quart of this product (0.75lb of glyphosate) in 5 to 10 gallons of water per acre. Fallow fields should be tilled prior to application. Apply prior to frost on water Bermudagrass that is 12 to 18 inches in length. This product is not registered in California for use on water Bermudagrass.

Bindweed, field	0.5-5 (0.375 – 3.75lb of glyphosate)	3-20	2%
------------------------	---	------	----

Do not treat when weeds are under drought stress as good soil moisture is necessary for active growth.

For control, apply 4 to 5 quarts of this product (3 – 3.75lb of glyphosate) per acre west of the Mississippi River and 3 to 4 quarts east of the Mississippi River. Apply when the weeds are at or beyond full bloom. For best results, apply in late summer or fall. Fall treatments must be applied before a killing frost.

Also for control, apply 2 quarts of this product (1.5lb of glyphosate) plus 0.5 pound of dicamba in 10 to 20 gallons of water per acre. Do not apply by air.

For suppression on irrigated agricultural land, apply 1 to 2 quarts of this product (0.75 – 1.25lb of glyphosate) plus 1 pound of 2,4-D in 10 to 20 gallons of water per acre with ground equipment only. Applications should be made following harvest or in fall fallow ground when the bindweed is actively growing and the majority of runners are 12 inches or more in length. The use of at least one irrigation will promote active bindweed growth.

For suppression, apply 16 fluid ounces of this product (0.375lb of glyphosate) plus 0.5 pound of 2,4-D in 3 to 10 gallons of water per acre for ground applications and 3 to 5 gallons of water per acre for aerial applications. Apply by air in fallow and reduced tillage systems only. Applications should be delayed until maximum emergence has occurred and when vines are between 6 to 18 inches in length.

In California only, apply 1 to 5 quarts of this product (0.75 -3.75lb of glyphosate) per acre. Actual rate needed for suppression or control will vary within this range depending on local conditions. For suppression on irrigated land where annual tillage is performed, apply 1 quart of this product (0.75lb of glyphosate) in 3 to 10 gallons of water per acre. Apply to bindweed that has reached a length of 12 inches or greater. Allow maximum weed emergence and runner growth. Allow 3 or more days after application before tillage.

Bluegrass, Kentucky	1-2	3-40	2%
	(0.75-1.5lb of glyphosate)		

Apply 2 quarts of this product (1.5lb of glyphosate) in 10 to 40 gallons of water per acre when most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 1 to 1.5 quarts of this product (0.75 – 1.13lb of glyphosate) in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height.

Blueweed, Texas	3-5	3-40	2%
	(2.25 – 3.75lb of glyphosate)		

Apply 4 to 5 quarts of this product (3 – 3.75lb of glyphosate) per acre west of the Mississippi River and 3 to 4 quarts (2.25 - 3lb of glyphosate) per acre east of the Mississippi River. Apply when plants are at or beyond full bloom. New leaf development indicates active growth. For best results, apply in late summer or fall. Fall treatments must be applied before a killing frost.

Brackenfern	3-4	3-40	1-
	(2.25- 3lb of glyphosate)		

1.5% Apply to fully expanded fronds that are at least 18 inches long.

Bromegrass, smooth	1-2	3-40	2%
	(0.75-1.5lb of glyphosate)		

Apply 2 quarts of this product (1.5lb of glyphosate) in 10 to 40 gallons of water per acre when most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 1 to 1.5 quarts of this product (0.75 – 1.13lb of glyphosate) in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height.

Bursage, woolly-leaf	—	3-20	2%
-----------------------------	---	------	----

For control, apply 2 quarts of this product (1.5lb of glyphosate) plus 0.5 pound of dicamba per acre. For partial control, apply 1 quart of this product plus 0.5 pound of dicamba per acre. Apply when plants are producing new active growth which has been initiated by moisture for at least 2 weeks and when plants are at or beyond flowering.

Canarygrass, reed	2-3	3-40	2%
	(1.5-2.25lb of glyphosate)		

For best results, apply when most plants have reached the boot-to-head stage of growth.

Cattail	3-5	3-40	2%
	(2.25-3.75lb of glyphosate)		

Apply when most plants have reached the early head stage.

Clover; red, white	3-5	3-20	2%
	(2.25-3.75lb of glyphosate)		

Apply when most plants have reached the early bud stage. Also for control, apply 16 to 32 fluid ounces of this product (0.375 -0.75lb of glyphosate) plus 0.5 to 1 pound of 2,4-D in 3 to 10 gallons of water per acre.

Cogongrass	3-5	10-40	2%
	(2.25-3.75lb of glyphosate)		

Apply when cogongrass is at least 18 inches tall in late summer or fall. Due to uneven stages of growth and the dense nature of vegetation preventing good spray coverage, repeat treatments may be necessary to maintain control.

Dallisgrass	3-5	3-20	2%
	(2.25-3.75lb of glyphosate)		

Apply when most plants have reached the early head stage.

Dandelion	3-5	3-40	2%
	(2.25-3.75lb of glyphosate)		

Apply when most plants have reached the early bud stage of growth. Also for control, apply 16 fluid ounces of this product (0.375lb of glyphosate) plus 0.5 pound of 2,4-D in 3 to 10 gallons of water per acre.

Dock, curly	3-5	3-40	2%
	(2.25-3.75lb of glyphosate)		

Apply when most plants have reached the early bud stage of growth. Also for control, apply 16 to 32 fluid ounces of this product (0.375 -0.75lb of glyphosate) plus 0.5 to 1 pound of 2,4-D in 3 to 10 gallons of water per acre.

Dogbane, hemp	4	3-40	2%
	(3lb of glyphosate)		

Apply when most plants have reached the late bud to flower stage of growth. Following crop harvest or mowing, allow weeds to regrow to a mature stage prior to treatment. For best results, apply in late summer or fall.

For suppression, apply 16 fluid ounces of this product (0.375lb of glyphosate plus 0.5 pound of 2,4-D in 3 to 10 gallons of water per acre for ground applications and 3 to 5 gallons of water per acre for aerial applications. Delay applications until maximum emergence of dogbane has occurred.

Fescue, (except tall)	3-5	3-20	2%
	(2.25-3.75lb of glyphosate)		

Apply when most plants have reached the early head stage.

Fescue, tall	1-3	3-40	2%
---------------------	-----	------	----

(0.75-2.25lb of glyphosate)

Apply 3 quarts of this product (2.25lb of glyphosate) per acre when most plants have reached boot-to-early seed head stage of development.

Fall applications only: Apply 1 quart of this product (0.75lb of glyphosate) in 3 to 10 gallons of water per acre. Apply to fescue in the fall when plants have 6 to 12 inches of new growth. A sequential application of 1 pint per acre of this product will improve long-term control and control seedlings germinating after fall treatments or the following spring.

Guineagrass	2-3	3-40	1%
--------------------	-----	------	----

(1.5-2.25lb of glyphosate)

Apply when most plants have reached at least the 7-leaf stage of growth. Ensure thorough coverage when using hand-held equipment. In Texas and ridge of Florida, use 2 quarts (1.5lb of glyphosate) for control. In the flatwoods region of Florida, 3 quarts (2.25lb of glyphosate) is required for control.

Horsenettle	3-5	3-20	2%
--------------------	-----	------	----

(2.25-3.75lb of glyphosate)

Apply when most plants have reached the early bud stage.

Horseradish	4	3-40	2%
--------------------	---	------	----

(3lb of glyphosate)

Apply when most plants have reached the late bud to flower stage of growth. For best results, apply in late summer or fall.

Iceplant	—	—	1.5-2%
-----------------	---	---	--------

Iceplant should be at or beyond the early bud stage of growth. Thorough coverage is necessary for best control.

Jerusalem artichoke	3-5	3-20	2%
----------------------------	-----	------	----

(2.25-3.75lb of glyphosate)

Apply when most plants are in the early bud stage.

Johnsongrass	0.5-3	3-40	1%
---------------------	-------	------	----

(0.375 -2.25lb of glyphosate)

In annual cropping systems apply 1 to 2 quarts of this product (0.75 -1.5lb of glyphosate) per acre. Apply 1 quart of this product in 3 to 10 gallons of water per acre. Use 2 quarts of this product (1.5lb of glyphosate) when applying 10 to 40 gallons of water per acre. In non- crop, or area where annual tillage (no-till) is not practiced, apply 2 to 3 quarts of this product (1.5-2.25lb of glyphosate) in 10 to 40 gallons of water per acre.

For best results, apply when most plants have reached the boot-to-head stage of growth or in the fall prior to frost. Allow 7 or more days after application before tillage. Do not tank mix with residual herbicides when using 1 quart of this product per acre.

For burndown of Johnsongrass, apply 1 pint of this product (0.375lb of glyphosate) in 3 to 10 gallons of water per acre before the plants reach a height of 12 inches. For this use, allow at least 3 days after treatment before tillage.

Spot treatment (partial control or suppression) - Apply a 1 percent solution of this product when Johnsongrass is 12 to 18 inches in height. Coverage should be uniform and complete.

Kikuyugrass	2-3 (1.5 – 2.25lb of glyphosate)	3-40	2%
--------------------	-------------------------------------	------	----

Spray when most kikuyugrass is at least 8 inches in height (3- or 4-leaf stage of growth). Allow 3 or more days after application before tillage.

Knapweed	4 (3lb of glyphosate)	3-40	2%
-----------------	--------------------------	------	----

Apply when most plants have reached the late bud to flower stage of growth. For best results, apply in late summer or fall.

Lantana	—	—	1-1.25%
----------------	---	---	---------

Apply at or beyond the bloom stage of growth. Use the higher application rate for plants that have reached the woody stage of growth.

Lespedeza	3-5 (2.25-3.75lb of glyphosate)	3-20	2%
------------------	------------------------------------	------	----

Apply when most plants have reached the early bud stage.

Milkweed, common	3 (2.25lb of glyphosate)	3-40	2%
-------------------------	-----------------------------	------	----

Apply when most plants have reached the late bud to flower stage of growth.

Muhly, wirestem	1-2 (0.75-1.5 lb of glyphosate)	3-40	2%
------------------------	------------------------------------	------	----

Use 1 quart of this product (0.75lb of glyphosate) in 3 to 10 gallons of water per acre. Use 2 quarts of this product (1.5lb of glyphosate) when applying 10 to 40 gallons of water per acre or in pasture, sod, or non-crop areas. Spray when the wirestem muhly is 8 inches or more in height. Do not till between harvest and fall applications or in the fall or spring prior to spring applications. Allow 3 or more days after application before tillage.

Mullein, common	3-5 (2.25-3.75lb of glyphosate)	3-20	2%
------------------------	------------------------------------	------	----

Apply when most are in early bud stage.

Napiergrass	3-5 (2.25 – 3.75lb of glyphosate)	3-20	2%
--------------------	--------------------------------------	------	----

Apply when most plants are early head stage.

Nightshade silverleaf	2 (1.5lb of glyphosate)	3-10	2%
------------------------------	----------------------------	------	----

Applications should be made when at least 60 percent of the plants have berries. Fall treatments must be applied before a killing frost.

Nutsedge; Purple or yellow	0.5-3 (0.375 -2.25lb of glyphosate)	3-40	1-2%
-----------------------------------	--	------	------

Apply 3 quarts of this product (2.25lb of glyphosate) per acre or apply a 1 to 2 percent solution for control of nutsedge plants and immature nutlets attached to treated plants. Treat when plants are in flower or when new nutlets can be found at rhizome tips. Nutlets that have not germinated will not be controlled and may germinate following treatment. Repeat treatments will be required for long-term control of ungerminated tubers.

Sequential applications: 1 to 2 quarts of this product (0.75-1.5lb of glyphosate) in 3 to 10 gallons of water per acre will also provide control. Make applications when a majority of the plants are in the 3- to 5-leaf stage (less than 6 inches tall). Repeat this application, as necessary, when newly emerging plants reach the 3- to 5-leaf stage. Subsequent applications will be necessary for long-term control.

For partial control of existing plants, apply 1 pint to 2 quarts of this product (0.375 -1.5lb of glyphosate) in 3 to 40 gallons of water per acre. Treat when plants have 3 to 5 leaves and most are less than 6 inches tall. Repeat treatments will be required to control subsequent emerging plants or regrowth of existing plants.

Orchardgrass	1-2	3-40	2%
	(0.75-1.5lb of glyphosate)		

Apply 2 quarts of this product (1.5lb of glyphosate) in 10 to 40 gallons of water per acre when most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 1 to 5 quarts of this product (0.75-3.75 lb of glyphosate) in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height.

Orchardgrass sods going to no-till corn: Apply 1 to 1.5 quarts of this product (0.75 -1.13lb of glyphosate) in 3 to 10 gallons of water per acre. Apply to orchardgrass that is a minimum of 12 inches tall for spring application and 6 inches tall for fall applications. Allow at least 3 days following application before planting. A sequential application of atrazine will be necessary for optimum results.

Pampasgrass	—	—	1.5-2%
--------------------	---	---	--------

Pampasgrass should be at or beyond the boot stage of growth. Thorough coverage is necessary for best control.

Paragrass	3-5	3-20	2%
	(2.25-3.75lb of glyphosate)		

Apply when most plants are in the early head stage.

Phragmites	3-5	10-40	1-2%
	(2.25-3.75lb of glyphosate)		

For partial control and best results, treat during late summer or fall when plants are actively growing and in full bloom. Treatment before or after this stage may lead to reduced control. Due to the dense nature of the vegetation, which may prevent good spray coverage or uneven stages of growth, repeat treatments may be necessary to maintain control. Visual control systems will be slow to develop.

Poison hemlock	—	—	1-2%
-----------------------	---	---	------

Apply as a spray-to-wet treatment. Optimum results are obtained when plants are treated at the bud to full-bloom stage of growth.

Pokeweed, common	1.0	3-40	2%
	(0.75lb of glyphosate)		

Apply to actively growing plants up to 24 inches tall.

Quackgrass	1-3	3-40	2%
	(0.75-2.25lb of glyphosate)		

In annual cropping systems, or in pastures and sods followed by deep tillage: Apply 1 quart of this product (0.75lb of glyphosate) in 3 to 10 gallons of water per acre. For 10 to 40 gallons of water per acre, apply 2 quarts of this product (1.5lb of glyphosate). Do not tank mix with residual herbicides when using the 1-quart rate. Spray when quackgrass is 6 to 8 inches in height. Do not till between harvest and fall applications or in fall or spring prior to spring application. Allow 3 or more days after application before tillage. In pastures or sods, use a midboard plow for best results. In pastures, sods or non-crop areas where deep tillage does not follow application: Apply 2 to 3 quarts of this product (1.5 - 2.25lb of glyphosate) in 10 to 40 gallons of water per acre when the quackgrass is greater than 8 inches tall.

Redvine	0.75-2	5-10	2%
	(0.563- 1.5lb of glyphosate)		

For suppression, apply 24 fluid ounces of this product (0.563lb of glyphosate) per acre at each of two applications 7 to 14 days apart or a single application of 2 quarts (1.5lb of glyphosate) per acre. Apply recommended rates in 5 to 10 gallons of water per acre. Apply in late September or early October to plants that are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least 1 week before a killing frost.

Reed, giant	—	—	2%
--------------------	---	---	----

Best results are obtained when applications are made in late summer to fall.

Ryegrass, perennial	1-3	3-40	1%
	(0.75-2.25lb of glyphosate)		

In annual cropping systems apply 1 to 2 quarts of this product (0.75-1.5 lb of glyphosate) per acre. Apply 1 quart of this product (0.75 lb of glyphosate) in 3 to 10 gallons of water per acre. Use 2 quarts of this product (1.5lb of glyphosate) when applying 10 to 40 gallons of water per acre. In non- crop, or areas when annual tillage (no-till) is not practiced, apply 2 to 3 quarts of this product (1.5-2.25lb of glyphosate) in 10 to 40 gallons of water per acre.

For best results, apply when most plants have reached the boot-to-head stage of growth or in the fall prior to frost. Do not tank-mix with residual herbicides when using 1 quart of this product per acre.

Smartweed, swamp	3-5	3-40	2%
	(2.25-3.75lb of glyphosate)		

Apply when most plants have reached the early bud stage of growth. Also for control, apply 16 fluid ounces of this product (0.375lb of glyphosate) plus 0.5 pound of 2,4-D in 3 to 10 gallons of water per acre in the late summer or fall.

Sowthistle, perennial	2-3	3-40	2%
	(1.5-2.25lb of glyphosate)		

Apply when most plants are at or beyond the bud stage of growth. After harvest, mowing or tillage in the late summer or fall, allow at least 4 weeks for initiation of active growth and rosette development prior to the application of this product. Fall treatments must be applied before a killing frost. Allow 3 or more days after application before tillage.

Spurge, leafy	—	3-10	2%
----------------------	---	------	----

For suppression, apply 16 fluid ounces of this product (0.375lb of glyphosate) plus 0.5 pound of 2,4-D in 3 to 10 gallons of water per acre in the late summer or fall. If mowing has occurred prior to treatment, apply when most of the plants are 12 inches tall. Allow 7 or more days after application before tillage.

Starthistle, yellow	2	10-40	2%
	(1.5lb of glyphosate)		

Best results are obtained when applications are made during the rosette, bolting and early flower stages.

Sweet potato, wild	—	—	2%
---------------------------	---	---	----

For partial control, apply to plants that are at or beyond the bloom stage of growth. Repeat applications may be required.

Thistle, artichoke	—	—	2%
---------------------------	---	---	----

For partial control, apply to plants that are at or beyond the bloom stage of growth. Repeat applications may be required.

Thistle, Canada	2-3 (1.5- 2.25lb of glyphosate)	3-40	2%
------------------------	------------------------------------	------	----

Apply when most plants are at or beyond the bud stage of growth. After harvest, mowing or tillage in the late summer or fall, allow at least 4 weeks for initiation of active growth and rosette development prior to the application of this product. Fall treatments must be applied before a killing frost. Allow 3 or more days after application before tillage.

For suppression in the spring, apply 1 quart of this product (0.75lb of glyphosate), or 1 pint of this product (0.375lb of glyphosate) plus 0.5 pound of 2,4-D, in 3 to 10 gallons of water per acre. Allow rosette regrowth to a minimum of 6 inches in diameter before treating. Applications can be made as long as leaves are still green and plants are actively growing at the time of application. Allow 3 or more days after application before tillage.

Timothy	2-3 (1.5-2.25lb of glyphosate)	3-40	2%
----------------	-----------------------------------	------	----

For best results, apply when most plants have reached the boot-to-head stage of growth.

Torpedograss	4-5 (3-3.75lb of glyphosate)	3-40	2%
---------------------	---------------------------------	------	----

For partial control, apply when most plants are at or beyond the seedhead stage of growth. Repeat applications will be required to maintain control. Fall treatments must be applied before frost.

Trumpet creeper	2 (1.5lb of glyphosate)	5-10	2%
------------------------	----------------------------	------	----

For partial control, apply in late September or October, to plants that are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least 1 week before a killing frost.

Vaseygrass	3-5 (2.25-3.75lb of glyphosate)	3-20	2%
-------------------	------------------------------------	------	----

Apply when most plants are in the early head stage.

Velvetgrass	3-5 (2.25-3.75lb of glyphosate)	3-20	2%
--------------------	------------------------------------	------	----

Apply when most plants are in the early head stage.

Wheatgrass, western	2-3 (2.25-3.75lb of glyphosate)	3-40	2%
----------------------------	------------------------------------	------	----

For best results, apply when most plants have reached the boot-to-head stage of growth.

WOODY BRUSH AND TREES RATE TABLE (Alphabetically by Species)

Apply this product after full leaf expansion, unless otherwise directed. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are

obtained when application is made in late summer or fall after fruit formation. In arid areas, best results are obtained when applications are made in the spring to early summer when brush species are at high moisture content and are flowering.

Unless otherwise directed, apply broadcast treatments in 3 to 40 gallons of water per acre. Ensure thorough coverage when using hand-held equipment. Symptoms may not appear prior to frost or senescence with fall treatments. Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatments are made following a frost.

*PC=Partial Control; C=Control

Weed species	Rate QT./A	Hand Held % solution	Comments*
Alder	3-4 (2.25-3lb of glyphosate)	1-1.5%	C
Ash	2-5 (2.25-3.75lb of glyphosate)	1-2%	PC
Aspen, quaking	2-3 (1.5-2.25lb of glyphosate)	1-1.5%	C
Bearmat (Bearclover)	2-5 (1.5-3.75lb of glyphosate)	1-2%	PC
Beech	2-5 (1.5-3.75lb of glyphosate)	1-2%	PC
Birch	2-5 (1.5-3.75lb of glyphosate)	1-1.5%	C
Blackberry	3-4 (2.25-3lb of glyphosate)	1-1.5%	C

Make applications after plants have reached full leaf maturity. Best results are obtained when applications are made in late summer or fall. Applications may also be made after leaf drop and until a killing frost or as long as stems are green. After berries have set or dropped in late fall, blackberry can be controlled by applying a 0.75 percent solution of this product. For control of blackberries after leaf drop and until killing frost or as long as stems are green, apply 3 to 4 quarts of this product in 10 to 40 gallons of water per acre.

Blackgum	2-5 (1.5-3.75lb of glyphosate)	1-2%	C
Bracken	2-5 (1.5-3.75lb of glyphosate)	1-2%	C
Broom; French, Scotch	—	1.5-2%	C
Buckwheat, California	—	1-2%	PC

Thorough coverage of foliage is necessary for best results.

Cascara	2-5 (1.5-3.75lb of glyphosate)	1-2%	PC
Catsclaw	—	1-1.5%	PC
Ceanothus	2-5 (1.5-3.75lb of glyphosate)	1-2%	PC
Chamise	—	1%	C

Thorough coverage of foliage is necessary for best results.

Cherry; bitter, black, pin	2-3 (1.5-2.25lb of glyphosate)	1-1.5%	C
---------------------------------------	-----------------------------------	--------	---

Coyote brush	—	1.5-2%	C
---------------------	---	--------	---

Apply when at least 50 percent of the new leaves are fully developed.

Dogwood	2-5 (1.5-3.75lb of glyphosate)	1-2%	PC
Elderberry	2-3 (1.5-2.25lb of glyphosate)	1-1.5%	C
Elm	2-5 (1.5-3.75lb of glyphosate)	1-2%	PC
Eucalyptus	—	2%	C

For control of eucalyptus resprouts, apply when resprouts are 6 to 12 feet tall. Ensure complete coverage. Avoid application to drought-stressed plants.

Florida holly (Brazilian Peppertree)	2-5 (1.5-3.75lb of glyphosate)	1-2%	PC
Gorse	2-5 (1.5-3.75lb of glyphosate)	1-2%	PC
Hasardia	—	1-2%	PC

Thorough coverage of foliage is necessary for best results.

Hawthorn	2-3 (1.5-2.25lb of glyphosate)	1-1.5%	C
Hazel	2-3 (1.5-2.25lb of glyphosate)	1-1.5%	C
Hickory	2-5 (1.5-3.75lb of glyphosate)	1-2%	PC
Honeysuckle	3-4 (2.25-3lb of glyphosate)	1-1.5%	C
Hornbeam, American	2-5 (1.5-3.75lb of glyphosate)	1-2%	PC
Kudzu	4-5 (3-3.75lb of glyphosate)	2%	C

Repeat applications may be required to maintain control.

Locust, black	2-4 (1.5-3lb of glyphosate)	1-2%	PC
Madrone resprouts	—	2%	PC

Apply to resprouts that are 3 to 6 feet tall. Best results are obtained with spring/early summer treatments.

Manzanita	2-5 (1.5-3.75lb of glyphosate)	1-2%	PC
Maple, red	2-4 (1.5-3lb of glyphosate)	1-1.5%	C

Apply a 1 to 1.5 percent solution when at least 50 percent of the new leaves are fully developed. For partial control, apply 2 to 4 quarts of this product per acre.

Maple, sugar	—	1-1.5%	C
---------------------	---	--------	---

Apply when at least 50 percent of the new leaves are fully develop.

Monkey Flower	—	1-2%	PC
----------------------	---	------	----

Thorough coverage of foliage is necessary for best result.

Oak; black, white	2-4 (1.5-3lb of glyphosate)	1-2%	PC
Oak, post	3-4 (2.25-3lb of glyphosate)v	1-1.5%	C
Oak, northern	—	1-1.5%	C

Apply when at least 50 percent of the new pin leaves are fully developed.

Oak, southern red	2-3 (1.5-2.25lb of glyphosate)	1-1.5%	C
Persimmon	2-5 (1.5-3.75lb of glyphosate)	1-2%	C
Pine	2-5 (1.5-3.75lb of glyphosate)	1-2%	C
Poison ivy/Poison oak	4-5 (3-3.75lb of glyphosate)	2%	C

Repeat applications may be required to maintain control. Fall treatments must be applied before leaves lose green color.

Poplar, yellow	2-5 (1.5-3.75lb of glyphosate)	1-2%	PC
Redbud, eastern	2-5 (1.5-3.75lb of glyphosate)	1-2%	C
Rose, multiflora	2 (1.5lb of glyphosate)	1%	C

Treatments should be made prior to deterioration by leaf-eating insects.

Russian olive	2-5 (1.5-3.75lb of glyphosate)	1-2%	PC
Sage, black	—	1%	C

Thorough coverage of foliage is necessary for best results.

Sage, white	2-5 (1.5-3.75lb of glyphosate)	1-2%	PC
Sage brush, California	—	1%	C

Thorough coverage of foliage is necessary for best results.

Salmonberry	2-3 (1.5-2.25lb of glyphosate)	1-1.5%	PC
Saltcedar	2-5 (1.5-3.75lb of glyphosate)	1-2%	C
Sassafras	2-5 (1.5-3.75lb of glyphosate)	1-2%	PC

Sourwood	2-5 (1.5-3.75lb of glyphosate)	1-2%	PC
Sumac; poison, smooth, winged	2-4 (1.5-3lb of glyphosate)	1-2%	PC
Sweetgum	2-3 (1.5-2.25lb of glyphosate)	1-1.5%	C
Swordfern	2-5 (1.5-3.75lb of glyphosate)	1-2%	PC
Tallowtree, Chinese	—	1%	C

Thorough coverage of foliage is necessary for best results.

Tan oak resprouts	—	2%	PC
--------------------------	---	----	----

Apply to resprouts that are less than 3 to 6 feet tall. Best results are obtained with fall applications.

Thimbleberry	2-3 (1.5-2.25lb of glyphosate)	1-1.5%	C
Tobacco, tree	—	1-2%	PC
Trumpet creeper	2-3 (1.5-2.25lb of glyphosate)	1-1.5%	C
Vine maple	2-5 (1.5-3.75lb of glyphosate)	1-2%	PC
Virginia creeper	2-5 (1.5-3.75lb of glyphosate)	1-2%	C
Waxmyrtle, southern	2-5 (1.5-3.75lb of glyphosate)	1-2%	PC
Willow	3-4 (2.25-3lb of glyphosate)	1-1.5%	C

INDUSTRIAL, RECREATIONAL AND PUBLIC AREAS

When applied as directed for “NON-CROP USES”, under conditions described, **GlamPrime** controls annual and perennial weeds listed on this label growing in areas including:

Airports	Parks
Apartment complexes	Parking areas
Around Farm, Ranch, Commercial or Industrial Structures	Petroleum & Other Tank Farms
Around Ornamental Gardens	Pumping Installations
Around Ornamental Trees & Shrubs	Pipeline, Power, Telephone & Utility Rights-of-Way
Ditch Banks	Preplant to Turf & Ornamental Plants
Driveways & Ramps	Railroads
Dry Ditches & Canals Fences & Fencerows	Recreational & Residential areas
Golf Courses	Schools
Gravel or Ground Bark Mulches	Sidewalks
Habitat Restoration & Management Areas	Sod or turf seed farms
Highways & Roadsides (including aprons, medians & guardrails)	Storage areas
Industrial Plant Sites	Uncropped Farmstead Areas
Lanes, Trails & Access Roads	Utility Substations
Lumberyards	Vacant Lots & Wastelands
Manufacturing sites	Warehouse areas
Office complexes	
Other public areas	

For specific rates of application and instructions for control of particular annual weeds, perennial weeds, woody brush and trees, see the “WEEDS CONTROLLED” section of this label. These applications may be made to large affected areas or as spot treatments. For use in small areas, see alternative instructions below under “Small Area Treatment with Hand-held Sprayers”. **GlamPrime** is a nonselective herbicide that is diluted and applied to the foliage of actively growing weeds as a spot or broadcast application. It is absorbed by the leaves and moves throughout the stem and roots to control the entire plant. Visible symptoms may require a week or more to appear, with burndown usually occurring in 2 to 4 weeks. Symptoms are a gradual wilting and yellowing of the sprayed plant followed by deterioration of both shoots and roots. **GlamPrime** has no herbicidal activity in the soil and will not wash or leach to affect nearby vegetation. Any ornamental species may be planted in treated areas 7 days or more after application. For most effective results, delay mowing, clipping, tilling, planting or sodding of treated areas for at least 7 days after application. This allows time for GlamPrime to move within the plant.

Unless the “Agricultural Use Requirements” on this label are observed, the following restrictions apply: Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

GLAMPRIIME CAN INJURE OR DESTROY ALL VEGETATION CONTACTED. WHEN USED AS A SPOT TREATMENT IN LAWNS, ALL VEGETATION CONTACTED WILL BE DAMAGED. AVOID SPRAY DRIFT CONTACT WITH DESIRABLE LAWN GRASSES, FLOWERS, VEGETABLES, SHRUBS OR TREES. DO NOT CONTACT GREEN BARK OF TREES OR SHRUBS. IF DESIRABLE VEGETATION IS CONTACTED, WASH IMMEDIATELY WITH WATER.

Depending on the type of non-crop application, **GlamPrime** may be applied with boom equipment, high-volume spray equipment and hand-held sprayers as described in the respective portions of the “APPLICATION EQUIPMENT and TECHNIQUES” section of the label. Additionally, the product may be applied with recirculating sprayers, shielded applicators, or wiper applicators in any non-crop site specified on this label. See the “Selective Equipment” part of “APPLICATION EQUIPMENT and TECHNIQUES” section of this label for information on proper use and calibration of this equipment.

Small Area Treatment With Hand-held Sprayers - Add 3 to 6 fluid ounces of **GlamPrime** (0.0703- 0.141lb of glyphosate) to 1 gallon of clean water. Use the lower rate for many grasses and annual weeds. Use the higher specified rate for control of perennials and brush. Use pump-up sprayer, backpack sprayer or other sprayer suitable for small areas. Adjust equipment

to deliver a coarse spray pattern. USE OF HOSE-END SPRAYERS OR SPRINKLER-TYPE DEVICES MAY RESULT IN POOR AND/OR ERRATIC RESULTS.

TANK MIXTURES FOR INDUSTRIAL SITES AND FORESTRY SITE PREPARATIONS

Use on industrial sites including airports, industrial plants, lumberyards, petroleum tank farms, pumping stations, railroads, roadsides, or storage areas where bare ground is desired.

This tank mixture may also be used as a site preparation treatment for sites to be planted to jack pine, loblolly pine, red pine, slash pine and Virginia pine. When applied as directed for "NON-CROP USES" under the conditions described, **GlamPrime** plus sulfometuron-methyl and metsulfuron methyl provides control of annual weeds listed in the "WEEDS CONTROLLED" section of the label for **GlamPrime** and sulfometuron-methyl and metsulfuron methyl, and control or partial control of the perennial weeds listed below.

Apply 1 to 2 quarts of **GlamPrime** (0.75-1.5lb of glyphosate) with 2 to 4 ounces of sulfometuron-methyl and metsulfuron methyl in 10 to 40 gallons of spray solution per acre as a broadcast spray to actively growing weeds. This mixture may be applied by aerial equipment in site prep operations. When applied by air, use the specified rates in 5 to 15 gallons of spray solution per acre. **GlamPrime** plus sulfometuron-methyl and metsulfuron methyl tank mixtures may not be applied by air in California.

For control of annual weeds, use the lower rates of these products.

For control of the listed perennial weeds, use the higher rates of both products. For partial control, use the lower rates.

Bahiagrass (<i>Paspalum notatum</i>)	Dogfennel (<i>Eupatorium capillifolium</i>)	Quackgrass (<i>Agropyron repens</i>)
Bermudagrass* (<i>Cynodon dactylon</i>)	Fescue, tall (<i>Festuca arundinaceus</i>)	Trumpetcreeper* (<i>Campsis radicans</i>)
Broomsedge (<i>Andropogon virginicus</i>)	Johnsongrass** (<i>Sorghum halepense</i>)	Vaseygrass (<i>Paspalum urvillei</i>)
Dock, Curly (<i>Rumex crispus</i>)	Poorjoe** (<i>Diodia teres</i>)	Vervain, blue (<i>Verbena hastate</i>)

*Suppression at the higher rates only.

**Control at the lower rates.

Read and carefully observe the precautionary statements and all other information appearing on the labels of all herbicides used.

TANK MIXTURES NON-CROP SITES

When applied as a tank mixture, **GlamPrime** provides control of the emerged annual weeds and partial control of the emerged perennial weeds listed in this label. When applied as a tank mixture, the following residual herbicides will provide pre-emergence control of the weeds listed in the individual product labels.

GlamPrime plus Diuron

GlamPrime plus Bromacil and Diuron*

GlamPrime plus Oxadiazon

GlamPrime plus Simazine*

GlamPrime plus Oryzalin*

See the "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" section of this label before preparing these tank mixtures.

Read and carefully observe the label claims, precautionary statements, specified rates and all other information on the labels of all products used in these tank mixtures. Use according to the most restrictive label directions for each product in the mixture.

CONTROL OF EMERGED WEEDS

NOTE: For backpack sprayer and handgun applications, see the "HAND-HELD AND HIGH VOLUME EQUIPMENT" section for specified rates.

Annual Weeds - Apply 1 quart per acre of **GlamPrime** (0.75lb of glyphosate) in these tank mixtures when weeds are less than 6 inches tall and 1.5 quarts per acre when weeds are more than 6 inches tall.

Perennial Weeds - For partial control of perennial weeds using these tank mixtures, apply 2 to 5 quarts per acre of **GlamPrime** (1.5-3.75lb of glyphosate). Follow the Directions in the "WEEDS CONTROLLED" section of this label for stage of growth and rate of application for specific perennial weeds.

PRE-EMERGENCE WEED CONTROL

For pre-emergence weed control, refer to the individual product labels for specific non-crop sites, rates, carrier volumes and precautionary statements. Mix only the quantity of spray solution which can be used during the same day. Do not allow these tank mixtures to stand overnight as this may result in reduced weed control.

FARMSTEAD WEED CONTROL

When applied as directed for "NON-CROP USES", under conditions described, **GlamPrime** controls undesirable vegetation listed on this label around areas including farmstead building foundations, along and in fences, shelterbelts and for nonselective farmstead weed control. For specific rates of application and instructions for control of various annual and perennial weeds, see the "WEEDS CONTROLLED" section of this label.

FARM DITCHES

GlamPrime will suppress perennial grasses along farm ditches. Apply **GlamPrime** at a rate of 6 to 8 fluid ounces per acre. Use 8 fluid ounces (0.188lb of glyphosate) per acre when treating tall (coarse) fescue, fine fescue, orchardgrass or quackgrass covers. For best suppression of these species, add ammonium sulfate at a rate of 1.7 pounds per 10 gallons of spray solution. Use 6 fluid ounces (0.141 lb of glyphosate) per acre without ammonium sulfate when treating Kentucky bluegrass. Apply treatments in 10 to 20 gallons of spray solution per acre to actively growing perennial grass covers. For best spray distribution and coverage, use flat fan nozzles. Where broadleaf weed control or suppression is desired, tank mix **GlamPrime** with an appropriate, labeled broadleaf weed herbicide.

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

PESTICIDE STORAGE: STORE ABOVE 10°F (-12°C) TO KEEP PRODUCT FROM CRYSTALLIZING. Crystals will settle to the bottom. If allowed to crystallize, place in a warm room 68°F (20°C) for several days to redissolve and shake or roll to mix well before using.

PESTICIDE DISPOSAL: Wastes resulting from the use of **GlamPrime** that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures. Emptied container retains vapor and product residue. Observe all label safeguards until container is destroyed.

[CONTAINER (EQUAL TO OR LESS THAN 5 GALLONS):] Do not reuse container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available.

[NONREFILLABLE CONTAINER (GREATER THAN 5 GALLONS):] Do not reuse container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling, if available.

[RETURNABLE / REFILLABLE CONTAINERS:] After use, return the container to the point of purchase or designated locations. Refill this container with glyphosate only. DO NOT REUSE THIS CONTAINER FOR ANY OTHER PURPOSE. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

LIMIT OF WARRANTY AND LIABILITY

LG Chem Ltd warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those

Directions under the conditions described therein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

To the extent consistent with applicable law, buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

This Company does not warrant any product reformulated or repackaged from this product except in accordance with this Company's stewardship requirements and with express written permission from this Company.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Upon opening and using this product, buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY, which may not be varied by any verbal or written agreement. If terms are not acceptable, return at once unopened.