



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Biopesticides and Pollution Prevention Division (7511P)  
1200 Pennsylvania Ave., N.W.  
Washington, D.C. 20460

EPA Reg. Number:

71637-2

Date of Issuance:

6/12/2020

NOTICE OF PESTICIDE:

Registration  
 Reregistration  
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Trico

Name and Address of Registrant (include ZIP Code):

KWIZDA AGRO GMBH  
Universitätsring  
A-1010 Vienna  
Austria

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Biopesticides and Pollution Prevention Division prior to use of the label in commerce. In any correspondence on this product, always refer to the above EPA Registration Number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA or the Act).

Registration is in no way to be construed as an endorsement or recommendation of this product by the U.S. Environmental Protection Agency (EPA). In order to protect health and the environment, the Administrator, on his or her motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under the Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration or registration review of your product when the EPA requires all registrants of similar products to submit such data.
2. Submit storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) data as these data requirements are not satisfied. A one-year study is required to satisfy these data requirements. You have 18 months from the date of this registration to provide these data to the EPA.

Signature of Approving Official:

Andrew Bryceland, Team Leader  
Biochemical Pesticides Branch  
Biopesticides and Pollution Prevention Division (7511P)  
Office of Pesticide Programs

Date:

6/12/2020

3. Make the following labeling change before you release this product for shipment:
  - Revise the EPA Registration Number to read, “EPA Reg. No. 71637-2.”
4. Submit one (1) copy of the final printed labeling for the record before you release this product for shipment.

Should you wish to add/retain a reference to your company’s website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA-approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6. A stamped copy of the labeling is enclosed for your records. Please also note that the record for this product currently contains the following acceptable Confidential Statement of Formula (CSF):

- Basic CSF dated 05/28/2020

Any CSFs other than those listed above are superseded.

If you have any questions, please contact Alex Horansky of my team by phone at (703) 347-0128 or via email at [Horansky.alex@epa.gov](mailto:Horansky.alex@epa.gov).

Sincerely,



Andrew Bryceland, Team Leader  
Biochemical Pesticides Branch  
Biopesticides and Pollution  
Prevention Division (7511P)  
Office of Pesticide Programs

Enclosure

**ACCEPTED**

06/12/2020

Under the Federal Insecticide, Fungicide  
and Rodenticide Act as amended, for the  
pesticide registered under  
EPA Reg. No. 71637-2

[Bracketed information is optional text]

**Master Label**

**Sublabel A: Commercial Use on Trees and Shrubs**

**Sublabel B: Residential Use**

**Sublabel C: Agricultural Use**

**TRICO®**

**REPELLENT FOR PREVENTING BROWSING DAMAGE CAUSED  
BY DEER, [RABBITS], ELK, AND MOOSE**

ACTIVE INGREDIENT:

Sheep fat.....6.4%  
OTHER INGREDIENTS.....93.6%  
TOTAL .....100.0%

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See [back] [bottom] [and] [side] [panel/panels] for First Aid, additional precautionary  
statements and Directions for Use

Net Contents:  
Lot/batch No.:

EPA Reg. No.  
EPA Est. No.

Kwizda Agro GmbH  
Universitätsring 6  
A-1010, Vienna  
Austria

[Made in USA]

[TRICO® is a registered trademark of KWIZDA Agro]

[Bracketed information is optional text]  
**Sublabel A: Commercial Use on Trees and Shrubs**

**TRICO®**

**REPELLENT FOR PREVENTING BROWSING, RUBBING AND  
BARK STRIPPING DAMAGE CAUSED BY DEER, [RABBITS], ELK,  
AND MOOSE**

ACTIVE INGREDIENT:

Sheep fat.....6.4%  
OTHER INGREDIENTS.....93.6%  
TOTAL .....100.0%

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See [back] [bottom] [and] [side] [panel/panels] for First Aid, additional precautionary  
statements and Directions for Use

Net Contents: 1 pint, 1 quart, ½ gallon, 1 gallon, 5 gallons, 55 gallons, 275 gallons  
Lot/batch No.:

EPA Reg. No.  
EPA Est. No.

Kwizda Agro GmbH  
Universitätsring 6  
A-1010, Vienna  
Austria

<b>FIRST AID</b>	
IF IN EYES:	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
IF ON SKIN:	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
IF INHALED:	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
IF SWALLOWED:	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>HOTLINE</b>	
<p>Have the product container or label with you when calling a poison control center, doctor, or going for treatment. For emergencies, call the poison control center 1-800-222-1222. For non-emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 (NPIC Web site: <a href="http://www.npic.orst.edu">www.npic.orst.edu</a>), 24 hours a day, seven days a week.</p>	

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION.** Causes moderate eye irritation. Harmful if swallowed or absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Avoid contact with eyes, skin and clothing, and avoid breathing spray mist. Remove and wash contaminated clothing before reuse. Wear protective gloves, long sleeved shirt and long pants, shoes, and socks. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Applicators must wear long sleeved shirt, long pants, shoes, and socks. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

## ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

## PRODUCT INFORMATION

TRICO® is a scent and taste repellent that protects against deer, [rabbits], elk and moose that browse, rub or debark seedlings, trees and shrubs.

For commercial application, TRICO® is a ready-to-use liquid concentrate used directly without any mixing or dilution. For protection against deer browsing, TRICO® is applied to the terminal shoots of stems or trunks of trees or shrubs – for protection against rubbing or debarking.

TRICO® provides effective control for up to 6 months [8 months] and ensures that young plantations have undisturbed growth.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

### Non-Agricultural Use

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

- Keep children and pets out of the treated areas until sprays have dried.
- Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

### Information for Use:

- Apply only at temperatures above 35° F and below 85° F.
- Avoid application in direct sunlight on hot days to avoid damage to leaves — for best results, apply in early morning or late afternoon.
- Before large-scale treatments or application in combination with other products, test product on a small area of that plant type to avoid possible plant damage.
- Protect product from freezing. Freezing makes TRICO® unusable.
- Do not apply other products one week before and one week after application of TRICO®.

Shake the container thoroughly before use. For 55 gallon and 275 gallon drums/containers, stir container contents until homogenous.

Apply only under dry conditions. Drying period depends on weather conditions. Under low humidity conditions, drying is expected to take a maximum of 6 hours. Once dried, TRICO® provides excellent rain resistance.

## APPLICATION INSTRUCTIONS

To control deer, [rabbits], elk, or moose browsing in winter and summer:

Application equipment: Use a hand-held, knapsack or garden sprayer with flat fan or cone nozzle.

TRICO® is sprayed directly on the terminal sprouts and tops of young trees and shrubs which are particularly exposed to damage by browsing of deer, [rabbits], elk, or moose.

Apply 1.1-2.2 gal/acre of undiluted TRICO depending on planting density. For best control, use 0.13 fluid ounces/plant. [1 gallon covers approximately 1,000 plants]. Make only one application in autumn/winter to deter winter browsing.

Newly grown shoots may be protected by a spring application to prevent spring/summer deer, [rabbits], elk, or moose browsing. For best control, use 0.13 fl. oz./plant. [1 gallon covers approximately 1,000 plants]. Make one application only in spring to deter spring/summer browsing. Make no more than two applications of 2.2 gal/acre per year (0.13 fl. oz./plant per year).

To control rubbing:

Rubbing damage by deer, elk, or moose mainly occurs in the summer/fall. Apply 2.2 gal/acre (282 fl. oz./acre) or 0.17-0.68 fl. oz./trunk of undiluted TRICO to the stems or trunks of coniferous and deciduous trees. Adjust quantity to tree size and apply on the threatened parts. Start from the top to the middle of the tree. Make only one application.

To control bark stripping:

Debarking of deciduous and coniferous trees by deer can occur in mostly winter and spring depending on the weather conditions. One application of TRICO is enough to provide sufficient protection against debarking damage for one growing season. Apply 2.2 gal/acre (282 fl. oz./acre) or 3.4 fl. oz./trunk of undiluted TRICO to the vulnerable part of the trunk (approximately 3-4.5 feet above the ground or snow level). Adjust application timing to local conditions. Apply before deer damage might happen. Make only one application.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store only in original container, in a cool, dry area. Keep away from children and pets. Protect from heat and freezing. Store between 35°F and 85°F. From date of manufacture, product can be stored for 24 months. Keep container tightly closed to prevent spillage and evaporation. If spillage occurs, dilute with water and flush away.

**PESTICIDE DISPOSAL:** In case of leak or spill, use absorbent materials to contain liquids and dispose of as waste.

**CONTAINER HANDLING:** Nonrefillable container. Do not refill or reuse this container. Offer for recycling, if available. Clean container promptly after emptying.

Container sizes of 5 gallons or less: Triple rinse as follows: Empty the remaining contents into the application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Container sizes of greater than 5 gallons: Triple rinse as follows: Empty the remaining contents into the application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

After 24 months' storage has been exceeded, do not use and dispose of product in accordance with the instructions above.

Note: Seller warrants that this product complies with the specifications expressed in this label. To the extent consistent with applicable law, Seller makes no other warranties, and disclaims all other warranties, express or implied, including, but not limited to, warranties of merchantability and fitness for the intended purpose. To the extent consistent with applicable law, Seller's total liability or default, breach or failure under this label shall not exceed the amount of the purchase price. To the extent consistent with applicable law, Seller shall have no liability for consequential, special or incidental damages.

The Directions for Use of this product are believed to be adequate and must be followed carefully. It is impossible to eliminate all risks associated with use of this product, however. Ineffectiveness or other unintended consequences may result based on factors beyond the control of the registrant, including but not limited to weather conditions or the manner of use or application.



[Optional Marketing/Advertising Claims]

- Deer, elk and moose repellent
- Prevents Deer, Elk or Moose Damage
- Protects Against Deer, Elk and Moose Damage
- Effective for up to 6 (8) months
- Lasts up to 6 (8) months\*
- Repels (deer) up to 6 (8) months\*
- (Protects) (Effective) up to 6 (8) months\*
- Residual repellency up to 6 (8) months\*
- (Single) (One) (1) Application (lasts) (protects) (repels) (deer) up to 6 (8) months\*
- Weather-Resistant Formula Provides Effective Damage Control
- Weather-Resistant Formula
- Get Long-Lasting (Results) (Effectiveness) (Protection) with (a Single) (One) (1) Application
- Repels by Scent and Taste
- Both Scent and Taste Barrier
- Starts working upon application.
- Water-resistant once dried
- Hardly recognizable smell to humans.
- Dries odorless to humans.
- Effective in all seasons
- Apply year-round
- Proven track record in Europe
- The European market leader
- The product of choice in Sweden
- Apply less than competing products.
- One gallon covers (protects) up to 1,000 plants.\*

\*When applied as directed.

[Bracketed information is optional text]  
**Sublabel B: Residential Use**

**TRICO®**

**REPELLENT FOR PREVENTING BROWSING DAMAGE CAUSED  
BY DEER, [RABBITS], ELK, AND MOOSE**

ACTIVE INGREDIENT:

Sheep fat.....6.4%  
OTHER INGREDIENTS.....93.6%  
TOTAL .....100.0%

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See [back] [bottom] [and] [side] [panel/panels] for First Aid, additional precautionary  
statements and Directions for Use

Net Contents: 1 pint, 1 quart, ½ gallon, 1 gallon, 2.5 gallons  
Lot/batch No.:

EPA Reg. No.  
EPA Est. No.

Kwizda Agro GmbH  
Universitätsring 6  
A-1010, Vienna  
Austria

<b>FIRST AID</b>	
IF IN EYES:	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
IF ON SKIN:	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
IF INHALED:	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
IF SWALLOWED	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>HOTLINE</b>	
<p>Have the product container or label with you when calling a poison control center, doctor, or going for treatment. For emergencies, call the poison control center 1-800-222-1222. For non-emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 (NPIC Web site: <a href="http://www.npic.orst.edu">www.npic.orst.edu</a>), 24 hours a day, seven days a week.</p>	

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION.** Causes moderate eye irritation. Harmful if swallowed or absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Avoid contact with eyes, skin and clothing, and avoid breathing spray mist. Remove and wash contaminated clothing before reuse. Wear protective gloves, long sleeved shirt and long pants, shoes, and socks. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

**ENVIRONMENTAL HAZARDS**

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help

avoid run off to water bodies or drainage systems.

## **PRODUCT INFORMATION**

TRICO® protects against deer, [rabbits], elk and moose that feed on and damage flowers, ornamentals, shrubs, trees as well as vegetable gardens and fruit-bearing trees.

Hardly recognizable to humans, but highly repellent to [rabbits], deer, elk, and moose, it provides – apart from a taste barrier – a very strong scent barrier.

Its proven efficacy has made TRICO® a product of choice [market leader] [in Europe]. Furthermore, with a minimum spray interval of 28 days during periods of high plant growth, it provides superior convenience for the user.

TRICO® provides effective control for up to 4 months, depending on animal pressure and season.

## **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

### **Mixing and Dilution Instructions**

Shake the container thoroughly before use. Use 1 part of TRICO® to 9 parts of water. [for 16 fl. oz. container size only]

Contents of 16 fl. oz. (1/8 gallon) bottle make 160 fl. oz. (1 1/4 gallons) of spray when mixed with 144 fl. oz. (1 1/8 gallons) of water. For heavy browsing pressure use this amount for 2,550 sq. ft. For light to moderate browsing pressure, treat up to 5,100 sq. ft.

[for 32 fl. oz. container size only]

Contents of 32 fl. oz. (1/4 gallon) bottle make 320 fl. oz. (2 1/2 gallons) of spray when mixed with 288 fl. oz. (2 1/4 gallons) of water. For heavy browsing pressure use this amount for 5,100 sq. ft. For light to moderate browsing pressure, treat up to 10,200 sq. ft.

[for 64 fl. oz. container size only]

Contents of 64 fl. oz. (1/2 gallon) bottle make 640 fl. oz. (5 gallons) of spray when mixed with 576 fl. oz. (4 1/2 gallons) of water. For heavy browsing pressure use this amount for 10,200 sq. ft. For light to moderate browsing pressure, treat up to 20,400 sq. ft.

[for 1 gallon container size only]

Contents of 128 fl. oz. (1 gallon) bottle make 1280 fl. oz. (10 gallons) of spray when mixed with 1152 fl. oz. (9 gallons) of water. For heavy browsing pressure use this amount for 20,400 sq. ft. For light to moderate browsing pressure, treat up to 40,800 sq. ft.

[for 2.5 gallon container size only]

Contents of 320 fl. oz. (2.5 gallon) bottle make 3200 fl. oz. (25 gallons) of spray when

mixed with 2880 fl. oz. (22½ gallons) of water. For heavy browsing pressure use this amount for 51,000 sq. ft. For light to moderate browsing pressure, treat up to 102,000 sq. ft.

## APPLICATION INSTRUCTIONS

[For best results]

Spray dry plant leaves, stems, flowers, tree bark and branches that need protection until run-off occurs or plants look wet. For extra protection on low growing plants, spray soil around plant base as well.

Spray treatment: For vegetables and fruit targeted for human consumption, spray on stems and leaves. [To avoid influencing [fresh] [fruit] [produce] [taste] [odor] [fragrance] [smell] do not apply directly to fruit [produce] [within 14 days] [prior to harvest]. [To avoid spray layer on [fresh] [fruit] [produce] do not apply directly to fruit [produce] [within 14 days] prior to harvest].

[Perimeter treatment]: Create a repelling barrier by applying to the surrounding area. Make treatment at least 6 feet wide. Wider applications may be more effective.

Drying period depends on weather conditions. Under low humidity conditions, drying is expected to take a maximum of 6 hours. Once dried, TRICO® provides excellent rain resistance.

Use a hand-held, knapsack or garden sprayer with a flat fan or cone nozzle.

Information for Use:

- Apply only at temperatures between 35° F and 85° F.
- Avoid application in direct sunlight on hot days to avoid damage to leaves. For best results, apply in early morning or late afternoon.
- Before large-scale treatments or application in combination with other products, test product on a small area of that plant type to avoid possible plant damage. Use with caution on open blooms and sensitive varieties such as impatiens, fuchsia, and begonias.
- Do not apply more than 16 fl. oz. of product per 2550 sq. ft.
- Do not exceed six applications per year.

Foliar Application:

[for 16 fl. oz. container size only]

<b>Degree of Browning</b>	<b>Application Rate</b>	<b>Timing of Application Spring/Summer</b>	<b>Timing of Application Autumn/Winter</b>	<b>Area Covered</b>
Heavy Browning	16 fluid ounces concentrate (mix 1:9 with water)	Up to 4 applications	1-2 applications	2,550 sq. ft.
Light to Moderate Browning	16 fluid ounces concentrate (mix 1:9 with water)	Up to 4 applications	1-2 applications	5,100 sq. ft.

[for 32 fl. oz. container size only]

<b>Degree of Browning</b>	<b>Application Rate</b>	<b>Timing of Application Spring/Summer</b>	<b>Timing of Application Autumn/Winter</b>	<b>Area Covered</b>
Heavy Browning	32 fluid ounces concentrate (mix 1:9 with water)	Up to 4 applications	1-2 applications	5,100 sq. ft.
Light to Moderate Browning	32 fluid ounces concentrate (mix 1:9 with water)	Up to 4 applications	1-2 applications	10,200 sq. ft.

[for 64 fl. oz. container size only]

<b>Degree of Browning</b>	<b>Application Rate</b>	<b>Timing of Application Spring/Summer</b>	<b>Timing of Application Autumn/Winter</b>	<b>Area Covered</b>
Heavy Browning	64 fluid ounces concentrate (mix 1:9 with water)	Up to 4 applications	1-2 applications	10,200 sq. ft.
Light to Moderate Browning	64 fluid ounces concentrate (mix 1:9 with water)	Up to 4 applications	1-2 applications	20,400 sq. ft.

[for 1 Gallon container size only]

<b>Degree of Browning</b>	<b>Application Rate</b>	<b>Timing of Application Spring/Summer</b>	<b>Timing of Application Autumn/Winter</b>	<b>Area Covered</b>
Heavy Browning	1 Gallon concentrate (mix 1:9 with water)	Up to 4 applications	1-2 applications	20,400 sq. ft.
Light to Moderate Browning	1 Gallon concentrate (mix 1:9 with water)	Up to 4 applications	1-2 applications	40,800 sq. ft.

[for 2.5 Gallon container size only]

<b>Degree of Browning</b>	<b>Application Rate</b>	<b>Timing of Application Spring/Summer</b>	<b>Timing of Application Autumn/Winter</b>	<b>Area Covered</b>
Heavy Browning	2.5 Gallons concentrate (mix 1:9 with water)	Up to 4 applications	1-2 applications	51,000 sq. ft.
Light to Moderate Browning	2.5 Gallons concentrate (mix 1:9 with water)	Up to 4 applications	1-2 applications	102,000 sq. ft.

In case of fresh deer, elk, moose [or rabbit] damage or rapid plant growth, re-apply after 28 days.

[Tested varieties]

TRICO® has been tested on the following ornamental plants, on which compatibility has occurred at the labeled application rate:

begonia, black pine, common hawthorn, daisy, English dogwood, English hawthorn, European hornbeam, European silver fir, field larkspur, field maple forsythia geranium,, Himalayan crane's bill, Japanese fantail willow, laburnum, lilac, northern white cedar, Norway spruce, oak, Oriental spruce, rose, sage, violet, and weigela.

#### **STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store only in original container, in a cool, dry area. Keep away from children and pets. Protect from heat and freezing. Store between 35°F and 85°F. From date of manufacture, product can be stored for 24 months. Keep container tightly closed to prevent spillage and evaporation. If spillage occurs, dilute with water and flush away.

**CONTAINER HANDLING AND DISPOSAL:** Nonrefillable container. Do not refill or reuse this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. After 24 months' storage has been exceeded, do not use and dispose of product in accordance with the instructions above.

Note: Seller warrants that this product complies with the specifications expressed in this label. To the extent consistent with applicable law, Seller makes no other warranties, and disclaims all other warranties, express or implied, including, but not limited to, warranties of merchantability and fitness for the intended purpose. To the extent consistent with applicable law, Seller's total liability or default, breach or failure under this label shall not

exceed the amount of the purchase price. To the extent consistent with applicable law, Seller shall have no liability for consequential, special or incidental damages.

The Directions for Use of this product are believed to be adequate and must be followed carefully. It is impossible to eliminate all risks associated with use of this product, however. Ineffectiveness or other unintended consequences may result based on factors beyond the control of the registrant, including but not limited to weather conditions or the manner of use or application.



[Optional Marketing/Advertising Claims]

- Deer, elk and moose repellent
- Prevents Deer, Elk or Moose Damage
- Protects against deer, elk or moose that feed on and damage flowers, ornamentals, shrubs, trees, vegetables and fruit-bearing trees
- Protects Against Deer, Elk and Moose Damage
- Deer & rabbit repellent
- Protects against deer & rabbits
- Protects flowers, ornamentals, shrubs, trees, vegetable gardens, and fruit-bearing trees
- (For) Use on flowers, ornamentals, shrubs, trees, vegetable gardens. and fruit trees
- Half as many applications when compared to other repellent products
- Effective for a minimum of 28 days
- Lasts up to 4 months\*
- Repels (deer) up to 4 months\*
- (Protects) (Effective) up to 4 months\*
- Residual repellency up to 4 months\*
- (Single) (One) (1) Application (lasts) (protects) (repels) (deer) up to 4 months\*
- Protects your flower garden and ornamentals for up to 4 months\*
- Weather-Resistant Formula Provides Effective Damage Control
- Weather-Resistant Formula
- Get Long-Lasting (Results) (Effectiveness) (Protection) with (a Single) (One) (1) Application
- [New] [exclusive] Technology
- Starts to repel immediately
- Dual action
- Repels by Scent and Taste
- Both Scent and Taste Barrier
- Starts working upon application.
- Water-resistant once dried
- Does not cause a visible spray layer on the plants and ornamentals.
- Hardly recognizable smell to humans.
- Dries odorless to humans.
- Does not disturb pets
- Effective in all seasons
- Apply year-round
- Proven track record in Europe
- The European market leader
- Apply less than competing products.
- Covers (protects) up to 5,100 sq. ft. (16 fluid ounces)\*\*

- Covers (protects) up to 10,200 sq. ft. (32 fluid ounces)\*\*
- Covers (protects) up to 20,400 sq. ft. (64 fluid ounces)\*\*
- Covers (protects) up to 40,800 sq. ft. (1 gallon)\*\*
- Covers (protects) up to 102,000 sq. ft. (2.5 gallons)\*\*
- New! [Claim valid for 6 months from date of first stamped "accepted" label only]

\*When applied as directed

[or]

\*Reapply TRICO® at 28-day intervals.

\*\*Under moderate browsing pressure.

[Bracketed information is optional text]  
**Sublabel C: Agricultural Use**

**TRICO®**

**REPELLENT FOR PREVENTING DEER, [RABBITS], ELK, AND  
MOOSE DAMAGE**

ACTIVE INGREDIENT:

Sheep fat.....6.4%  
OTHER INGREDIENTS.....93.6%  
TOTAL .....100.0%

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See [back] [bottom] [and] [side] [panel/panels] for First Aid, additional precautionary  
statements and Directions for Use

Net Contents: 1.5 gallons, 2.5 gallons, 5 gallons, 55 gallons, 275 gallons  
Lot/batch No.:

EPA Reg. No.  
EPA Est. No.

Kwizda Agro GmbH  
Universitätsring 6  
A-1010, Vienna  
Austria

<b>FIRST AID</b>	
IF IN EYES:	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
IF ON SKIN:	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
IF INHALED:	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
IF SWALLOWED	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>HOTLINE</b>	
<p>Have the product container or label with you when calling a poison control center, doctor, or going for treatment. For emergencies, call the poison control center 1-800-222-1222. For non-emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 (NPIC Web site: <a href="http://www.npic.orst.edu">www.npic.orst.edu</a>), 24 hours a day, seven days a week.</p>	

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION.** Causes moderate eye irritation. Harmful if swallowed or absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contact with eyes, skin and clothing, and avoid breathing spray mist. Remove and wash contaminated clothing before reuse. Wear protective gloves, long sleeved shirt and long pants, shoes, and socks. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

**ENVIRONMENTAL HAZARDS**

For terrestrial uses: Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

## PRODUCT INFORMATION

TRICO® protects field crops, vegetable plants, vineyards, orchards, strawberry plants, and blueberry plants by preventing deer, elk, [rabbits], and moose from browsing.

Hardly recognizable to humans, but highly repellent to [rabbits], deer, elk, and moose, it provides – apart from a taste barrier – a very strong scent barrier.

TRICO® provides effective control of deer damage to vulnerable, deer attractive agricultural plants.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your State and Tribe, consult the State/Tribal agency responsible for pesticide regulation.

### AGRICULTURAL USE

Use this product only in accordance with its labelling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water) is: long sleeved shirt, long pants, gloves, shoes, and socks.

### APPLICATION INSTRUCTIONS

[For best results]

Shake the container thoroughly before diluting. For 55 gallon and 275 gallon drums/containers, stir contents until homogenous. Use a hand-held, knapsack or garden sprayer with a flat fan or cone nozzle, or farm spraying equipment.

Apply only under dry conditions. Drying period depends on weather conditions. Once dried, TRICO® provides excellent rain resistance.

**Information for Use:**

- Apply only at temperatures between 35° F and 85° F.
- Avoid application in on hot days to avoid damage to the leaves. For best results, apply in early morning or late afternoon.
- Before large-scale treatments or application in combination with other products, test product on a small area of that plant type to avoid possible plant damage.
- Protect product from freezing. Freezing makes TRICO® unusable.
- Do not apply other products one week before and one week after application of TRICO®, unless compatibility has been confirmed. Be aware, sheep fat may increase the effect of other pesticides.
- Do not apply more than 16 fl. oz. of product per 2550 sq. ft.
- Do not exceed six applications per year.

**Field Crops:**

For use on field crops, such as Corn, Soy, Oilseed Rape, Canola, Sunflower, Cotton, Sweet corn, and Hops. [Make application to the whole field or through a perimeter treatment at least 70 feet wide. Wider applications may be more effective.] Use application rates as shown below.

<b>Plant</b>	<b>Application Rate</b>	<b>Number of Applications</b>	<b>Application Timing</b>
Corn, Soy, Oilseed Rape, Canola, Sunflower, Cotton, Sweet corn	1.65 gal/acre in 80–120 gal of water	Up to 4 applications per year	Minimum retreatment interval is 7 days
Hop	1.1-1.65 gal/acre in 20 gal of water	Up to 3 applications per year	Minimum retreatment interval is 14 days

## Vegetables - All

For all vegetables, spray on plants or make a perimeter treatment creating a repelling barrier. [To avoid influencing [fresh] [fruit] [produce] [taste] [odor] [fragrance] [smell] do not apply directly to fruit [produce] [within 14 days] [prior to harvest]. [To avoid spray layer on [fresh] [fruit] [produce] do not apply directly to fruit [produce] [within 14 days] prior to harvest].

Plant	Application Rate	Number of Applications	Application Timing
All	1.65 gal/acre in 20–120 gal of water	Up to 4 applications per year	Minimum retreatment interval is 7 days

## Vineyards, Orchards and Berries:

For vineyards, orchards, strawberries and blueberries, spray to plants. [For berries, treat the entire field or through a perimeter treatment of 70 feet creating a repelling barrier]. [To avoid influencing [fresh] [fruit] [produce] [taste] [odor] [fragrance] [smell] do not apply directly to fruit [produce] [within 14 days] [prior to harvest]. [To avoid spray layer on [fresh] [fruit] [produce] do not apply directly to fruit [produce] [within 14 days] prior to harvest].

Plant	Application Rate	Number of Applications	Application Timing
Grapes (new plantations)	1.1-1.65 gal/acre in 20 gal of water	Up to 2 applications per year	Minimum retreatment interval is 28 days
Grapes (established plantations)	1.1-1.65 gal/acre in 20 gal of water	Up to 2 applications per year	
Orchards	1.1-1.65 gal/acre in 20 gal of water	Up to 2 applications per year	

Strawberries, Blueberries (high and low bush)	1.1-1.65 gal/acre in 20 gal of water	Up to 2 applications per year	
--	--	-------------------------------------	--

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store only in original container, in a cool, dry area. Keep away from children and pets. Protect from heat and freezing. Store between 35°F and 85°F. From date of manufacture, product can be stored for 24 months. Keep container tightly closed to prevent spillage and evaporation. If spillage occurs, dilute with water and flush away.

**PESTICIDE DISPOSAL:** In case of leak or spill, use absorbent materials to contain liquids and dispose of as waste.

**CONTAINER HANDLING:** Nonrefillable container. Do not refill or reuse this container. Offer for recycling, if available. Clean container promptly after emptying.

Container sizes of 5 gallons or less: Triple rinse as follows: Empty the remaining contents into the application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Container sizes of greater than 5 gallons: Triple rinse as follows: Empty the remaining contents into the application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. After 24 months' storage has been exceeded, do not use and dispose of product in accordance with the instructions above.

Note: Seller warrants that this product complies with the specifications expressed in this label. To the extent consistent with applicable law, Seller makes no other warranties, and disclaims all other warranties, express or implied, including, but not limited to, warranties of merchantability and fitness for the intended purpose. To the extent consistent with applicable law, Seller's total liability or default, breach or failure under this label shall not exceed the amount of the purchase price. To the extent consistent with applicable law, Seller shall have no liability for consequential, special or incidental damages.

The Directions for Use of this product are believed to be adequate and must be followed carefully. It is impossible to eliminate all risks associated with use of this product, however. Ineffectiveness or other unintended consequences may result based on factors



beyond the control of the registrant, including but not limited to weather conditions or the manner of use or application.

[Optional Marketing/Advertising Claims]

- Deer, elk and moose repellent
- Prevents Deer, Elk or Moose Damage
- Protects Against Deer, Elk and Moose Damage
- Effective for up to 6 (8) months
- Lasts up to 6 (8) months\*
- Repels (deer) up to 6 (8) months\*
- (Protects) (Effective) up to 6 (8) months\*
- Residual repellency up to 6 (8) months\*
- (Single) (One) (1) Application (lasts) (protects) (repels) (deer) up to 6 (8) months\*
- Weather-Resistant Formula Provides Effective Damage Control
- Weather-Resistant Formula
- Get Long-Lasting (Results) (Effectiveness) (Protection) with (a Single) (One) (1) Application
- Repels by Scent and Taste
- Both Scent and Taste Barrier
- Starts working upon application.
- Water-resistant once dried
- Hardly recognizable smell to humans.
- Dries odorless to humans.
- Effective in all seasons
- Apply year-round
- Proven track record in Europe
- The European market leader
- Apply less than competing products.

\*When applied as directed