

71711-1

6/10/2010

Page 1 of 9



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

Shannon Yanocha
Registration Specialist
Nichino America, Inc.
4550 New Linden Hill Rd., Suite 501
Wilmington, DE 19808

JUN 10 2010

Subject: Moncut® 70WP Fungicide
EPA Reg. No. 71711-1
Your amendment dated May 11, 2010
EPA Decision Number 433994

Dear Ms Yanocha:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act as amended is acceptable.

One copy of the label stamped "Accepted" is enclosed for your records. This label supersedes all labels previously accepted for this product. Please submit one copy of the final printed label before the product is released for shipment.

If you have any questions, please contact Robert Westin by phone at (703) 305-5721 or via email at westin.robert@epa.gov.

Sincerely,

Lisa Jones FOR

Erik Kraft
Acting PM, Team 20
Fungicide Branch
Registration Division (7504P)

Enclosure

289

NICHINO
AMERICA

ACCEPTED

JUN 10 2010

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 71711-1

Moncut® 70WP FUNGICIDE

For Use on Peanuts*, Potatoes and Rice*

*Not for use in California.

Not for use in Nassau and Suffolk counties in New York State.

ACTIVE INGREDIENT:	Percent by Weight
FLUTOLANIL: N-[3-(1-methylethoxy) phenyl]-2-(trifluoromethyl) benzamide	70.0%
OTHER INGREDIENTS:	30.0%
TOTAL	100.0%

EPA Reg. No. 71711-1

EPA Est. No. 67545-AZ-1

KEEP OUT OF REACH OF CHILDREN
CAUTION

FIRST AID	
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-348-5832 for emergency medical treatment information. In case of fire or spills, information may be obtained by calling 1-800-424-9300.	

NICHINO AMERICA, INC.
4550 New Linden Hill Rd., Suite 501
Wilmington, DE 19808
1-888-740-7700

NET CONTENTS: _____

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals
CAUTION

Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride
- Shoes plus socks

If you want more options for chemical-resistant materials, follow the instructions for category A on an EPA chemical-resistance category selection chart.

USER SAFETY REQUIREMENTS

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- remove clothing/PPE immediately if pesticide gets inside, then wash thoroughly and put on clean clothing.
- remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This product is toxic to shrimp. For terrestrial uses, do not apply to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling MUST be in the possession of the user at the time of pesticide application.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency in your State responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride
- Shoes plus socks

GENERAL INFORMATION

MONCUT® 70WP is a systemic fungicide for control of Southern stem rot (white mold) *Sclerotium rolfsii*, and the limb/pod rot complex, *Rhizoctonia solani*, in peanuts*, for control of sheath blight, *Rhizoctonia solani*, in rice*, and for control of black scurf, *Rhizoctonia solani*, in potatoes.

*Not for use in California

MIXING INSTRUCTIONS

- Buffer the water in the spray tank.
- Fill the spray tank $\frac{3}{4}$ full. Turn on spray tank agitation. Pour required amount of this material on the surface of water in the spray tank.
- Add balance of water to spray tank.
- Keep agitation running during filling and spraying operations.
- Do not apply with adjuvants. Do not use oil as a carrier or add other additives to the finished spray.

USE PRECAUTIONS

Rotational Crop Restrictions:

Do not plant rotational crops, other than those listed in the table below for 365 days following the last application of MONCUT 70WP FUNGICIDE.

Crop	Rotational/Plantback Interval
Cotton Peanuts* Potatoes Rice* Soybeans Sugar beets	0 days following application
Wheat	30 days following application
Leafy vegetables (such as lettuce, spinach or celery) Small grain crops other than wheat (such as barley, rye or oats)	150 days following application
Corn (such as field, sweet or popcorn) Sorghum	240 days following application
All other crops	365 days following application

(*not for use on peanuts and rice in California)

APPLICATION DIRECTIONS

PEANUTS (Not for use in California)

Apply MONCUT 70WP FUNGICIDE as a broadcast foliar application using spray equipment commonly used to apply leaf spot fungicide treatments. In fields with a history of high disease, or under prolonged conditions conducive to disease infection and development, higher labeled rates (1.43 lb./A or more) should be used for the most effective control. In such situations, sequential applications will provide more effective control than a single application. MONCUT 70WP FUNGICIDE may be applied in tankmix with leaf spot treatments that fall within the specified timing for MONCUT 70WP FUNGICIDE application.

MONCUT 70WP FUNGICIDE may be applied to peanuts through chemigation. (See application techniques below).

Ground Application: By ground, apply specified rate in 10-40 gallons of water per acre.

Aerial Application: By air, apply specified rate in 5-10 gallons of water per acre.

Crop	Disease	Rate Per Acre (lb./a)	Application Directions
Peanuts	Southern stem rot (white mold, <i>Sclerotium rolfsii</i>); limb/pod rot complex (<i>Rhizoctonia solani</i>)	1.43 - 2.86	One application program: Apply 1.43 - 2.86 lbs. MONCUT 70WP FUNGICIDE per acre about 50-70 days after planting. Use the higher rates in fields known to have high disease incidence.
		0.71 - 1.43	Two application program: Apply 0.71 - 1.43 lbs. MONCUT 70WP FUNGICIDE per acre. Apply 50-70 days after planting and repeat with a second application of 0.71 - 1.43 lbs. MONCUT 70WP FUNGICIDE per acre about 30 days later. Use the higher rate in fields known to have high disease incidence.
		0.36 - 0.71	Four application program: Apply 0.36 - 0.71 lbs. MONCUT 70WP FUNGICIDE per acre. Begin when conditions are favorable for soilborne disease development, generally 50-70 days after planting. Repeat application at 10 to 14 day intervals for a total of four applications.
<ul style="list-style-type: none"> • May be applied at the specified rates in a tank mix for leaf spot treatments. Apply no more than a combined total of 2.86 lbs. MONCUT 70WP FUNGICIDE per acre to peanuts as a broadcast application or through chemigation during a single growing season. • Do not apply later than 40 days prior to digging of peanuts. 			

In-Furrow Application:

Use Rate And Method Of Application (to control *Rhizoctonia*)

Apply 1.1 lbs of MONCUT 70WP FUNGICIDE per acre (see referenced chart) as an in-furrow spray by directing spray uniformly over the seed, bottom and walls of the seed furrow and soil that is used to cover the seed in a 4-8 inch band prior to covering with soil. Use a minimum of 3 gallons of spray volume per acre. Use highest rate when disease conditions are expected to be severe, or if field has a history of *Rhizoctonia* infestation.

**In-Furrow Application Rates
1.1 lb. rate per acre**

Row Spacing	Ounces Product/ 1000 Row Ft.
40" Row	1.31 oz.
38" Row	1.24 oz.
36" Row	1.18 oz.
32" Row	1.11 oz.
30" Row	1.04 oz.

RICE (Not for use in California)

Apply MONCUT 70WP FUNGICIDE aerially in 5-10 gallons of water per acre, using one of the following schedules:

Crop	Disease	Rate Per Acre (lb./a)	Application Directions
Rice	Sheath blight (<i>Rhizoctonia solani</i>)	0.5 – 0.71	Apply 0.5 - 0.71 lbs. MONCUT 70WP FUNGICIDE per acre at first internode elongation, and follow with a second application at the same rate 10 to 14 days later. Use the higher rate where disease pressure is expected to be heavy.
		0.5 – 1.0	In fields where close scouting is practiced to detect and monitor sheath blight, apply MONCUT 70WP FUNGICIDE when 5 -10% of the tillers of a susceptible variety or 10-15% of a moderately susceptible variety have sheath blight lesions. Apply MONCUT 70WP FUNGICIDE at 0.5 - 1.0 lbs. per acre. Continue scouting and reapply if disease begins to move up the rice stem again.
<ul style="list-style-type: none"> • Do not apply more than 1.43 lbs. of MONCUT 70WP FUNGICIDE per acre per year. • Do not apply later than 30 days prior to rice harvest, or beyond 75% heading development stage, whichever occurs first. • This pesticide is toxic to shrimp. Do not apply this product within three miles of any estuarine/marine waterways or watershed. • Flooded fields may be used for aquaculture of crayfish only following rice harvest. 			

POTATOES

(In-furrow Use Only)

Do not apply this product through any type of irrigation system on potatoes.

Crop	Disease	Rate Per Acre (lb./a)	Application Directions
Potatoes	Black scurf (<i>rhizoctonia solani</i>)	0.71 – 1.1 (see reference chart below)	Apply 0.71 - 1.1 lbs. Moncut 70WP FUNGICIDE per acre (see reference chart) as an in-furrow spray by directing spray uniformly around and

	Powdery Scab <i>(Spongospora subterranea)</i> suppression only		over the seed piece in a 4 to 8 inch band prior to covering with soil. Use a minimum of 3 gallons of spray volume per acre. Use the higher rate where disease pressure is expected to be severe, or if field has a history of <i>Rhizoctonia</i> infestation.
--	---	--	---

IN-FURROW APPLICATION RATES

Row Spacing	0.71 lb/acre Ounces Product/ 1000 Row Ft.	1.1 lb/acre Ounces Product/ 1000 Row Ft.
40" Row	0.87 oz	1.31 oz
38" Row	0.83 oz	1.24 oz
36" Row	0.79 oz	1.18 oz
34" Row	0.74 oz	1.11 oz
32" Row	0.70 oz	1.04 oz

CHEMIGATION APPLICATION:

1. Determine the size of the area to be treated.
2. Determine the time required to apply ¼ to ½ inch of water over the area to be treated when the system and injection equipment are operated at normal pressures as recommended by the equipment manufacturer.
3. Using water, determine the injection pump output when operated at normal line pressure.
4. Determine the amount of Moncut 70WP FUNGICIDE required to treat the area covered by the irrigation system.
5. Add the required amount of Moncut 70WP FUNGICIDE to the solution tank with sufficient water to meet the injection time requirements.
6. Make certain the system is fully charged with water before starting injection of the Moncut 70WP FUNGICIDE solution. Time the injection to last at least as long as it takes to bring the system to full pressure.
7. Maintain constant solution tank agitation during the entire injection period.
8. Stop injection equipment after treatment is completed. Continue to operate the system until the Moncut 70WP FUNGICIDE solution has cleared the last sprinkler head. (Also see APPLICATION AND CALIBRATION TECHNIQUES FOR SPRINKLER IRRIGATION section below.

APPLICATION AND CALIBRATION TECHNIQUES FOR SPRINKLER IRRIGATION:

Apply this product only through center pivot, motorized lateral move, traveling gun, solid set, or portable (wheel move, side roll, end tow, or hand move) irrigation system(s). Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.

Do not apply this product through irrigation systems connected to a public water system. 'Public water system' means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days per year.

Controls for both irrigation water and pesticide injection systems must be functionally interlocked, so as to automatically terminate pesticide injection when the irrigation water pump motor stops. A person knowledgeable of the irrigation system and responsible for its operation shall be present so as to discontinue pesticide injection and make necessary adjustments, should the need arise.

The irrigation water pipeline must be fitted with a functional, automatic, quick-closing check valve to prevent the flow of treated irrigation water back toward the water source. The pipeline must also be fitted with a vacuum relief valve and low pressure drain, located between the irrigation water pump and the check valve, to prevent back-siphoning of treated irrigation water into the water source.

Always inject this product into irrigation water after it discharges from the irrigation pump and after it passes through the check valve. Never inject pesticides into the intake line on the suction side of the pump.

Pesticide injection equipment must be fitted with a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump. Interlock this valve to the power system, so as to prevent fluid from being withdrawn from the chemical supply tank when the irrigation system is either automatically or manually turned off.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur. Do not apply when wind speed favors drift beyond the area intended for treatment.

This product may be used through two basic types of sprinkler irrigation systems as outlined in Sections A and B below. Determine which type of system is in place, then refer to the appropriate directions provided for each type.

A. Center Pivot, Motorized Lateral Move and Traveling Gun Irrigation Equipment

For injection of pesticides, these continuously moving systems must use a positive displacement injection pump, of either diaphragm or piston type, constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock and capable of injection at pressures approximately 2–3 times those encountered within the irrigation water line. Venturi applicator units cannot be used on these systems. Thoroughly mix recommended amount of this product for acreage to be covered into same amount of water used during calibration and inject into system continuously for one revolution or run. Mixture in the chemical supply tank must be continuously agitated during the injection run. Shut off injection equipment after one revolution or run, but continue to operate irrigation system until this product has been cleared from last sprinkler head.

B. Solid Set and Portable (Wheel Move, Side Roll, End Tow, or Hand Move) Irrigation Equipment

With stationary systems, an effectively designed in-line venturi applicator unit is preferred which is constructed of materials that are compatible with pesticides; however, a positive-displacement pump can also be used. Determine acreage covered by sprinkler. Fill tank of injection equipment with water and adjust flow to use contents over a thirty to forty-five minute period. Mix desired amount of this product for acreage to be covered with water so that the total mixture of this product plus water in the injection tank is equal to the quantity of water used during calibration, and operate entire system at normal pressures recommended by the manufacturer of injection equipment used, for amount of time established during calibration. Agitation is recommended. This product can be injected at the beginning or end of the irrigation cycle or as a separate application. Stop injection equipment after treatment is completed and continue to operate irrigation system until this product has been cleared from last sprinkler head.

ENDANGERED SPECIES**Restrictions in the State of Arkansas:**

The use of MONCUT 70WP FUNGICIDE (flutolanil) on rice is restricted to protect the endangered fat pocketbook pearly mussel (*Potamilus capax*) and its habitat. Use is prohibited in the following areas of Arkansas:

Mississippi County: Within the basin that drains directly into the Right Hand Chute of Little River, south of Big Lake National Refuge.

Poinsett County: Between Crowley's Ridge and the levee east of the Right Hand Chute of Little River and the St. Francis Floodway. Use is also prohibited west of Rt. 140 and north of Rt. 63 at the SIPHON near Marked Tree. Except that the prohibited area does not include the area bounded by Arkansas Highway 373 on the west, Highway 63 on the east and Highway 14 on the south.

Cross, St. Francis, and Lee Counties: Between Crowley's Ridge and the levee east of the Right Hand Chute of Little River and the St. Francis Floodway as far south as the confluence of L'Anguille River (Lee County).

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

IMPORTANT: READ BEFORE USE

By using this product, user or buyer accepts the following conditions, warranty, disclaimer of warranties, and limitations of liability.

CONDITIONS: The directions for use of this product are believed to be accurate and must be followed carefully. However, because of extreme weather and soil conditions, use methods and other factors beyond the control of Nichino America, Inc. (NAI), it is impossible for NAI to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. To the extent consistent with applicable law, all such risks are assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THERE ARE NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, WHICH EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of NAI is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, NAI disclaims any liability whatsoever for incidental or consequential damages, including, but not limited to, liability arising out of breach of contract, express or implied warranty (including warranties of merchantability and fitness for a particular purpose), tort, negligence, strict liability, or otherwise.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT THE ELECTION OF NICHINO AMERICA, THE REPLACEMENT OF PRODUCT.

© Nichino America, Inc. 2010

Moncut is a registered trademark of Nichino America, Inc.

Label Code 040510