



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

SEP 7 2004

Marie A. Maks, Manager, Regulatory Affairs
Nichino America, Inc.
Linden Park, Suite 501
4550 New Linden Hill Road
Wilmington, DE 19808

SUBJECT: Label amendment to revise Section 3 Applaud Master label in accordance with April 1, 2004 Agency letter.
Product Registration No: 71711-15
Product Name: Applaud® 70 WP Insect Growth Regulator
Date of submission: September 3, 2004


Dear Ms. Maks,

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records. If you have any questions, you may contact Richard J. Gebken, at (703) 305-6701.

Sincerely,

Richard Gebken
Acting Product Manager 10
Insecticide Branch
Registration Division (7505C)

enclosure
072771 1-00015 8707021 wjd



Nichino America, Inc.

2/12

APPLAUD® 70WP Insect Growth Regulator

FOR: Almonds, Beans, Citrus, Cotton, Cucurbita (including cucumbers, melons, pumpkins, squash), Grapes, Lettuce, Longan, Lychee, Peas, Peanuts, Pumpkins, Spanish Lima, Snap Beans, Tomatoes and Onions/Onion (Greenhouse and Outdoor)

ACTIVE INGREDIENT:

Buprofezin, 2-(4-tert-butylphenyl)-3-hydroxy-6-phenylpyridin-1,3,5-triazolin-4-yl-ene

OTHER INGREDIENTS:	70.00%
TOTAL	100.00%

Contains 0.70 lbs. buprofezin per pound of product

Marketing Brands

Applied - Almonds, Beans, Citrus, Grapes, Longan, Lychee, Peas, Peanuts, Pumpkins, Spanish Lima
Counter - Cotton, Cucurbita (including cucumbers, melons, pumpkins, squash), Lettuce, Snap Beans, Tomatoes
Talus - Greenhouse Tomatoes and Outdoor and Greenhouse Onions/Onions
Centaur - Almonds, Citrus, Grapes

EPA Reg. No. 71711-15
EPA Est. No. 67548-AZ-1

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID	
If Inhaled	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-30 minutes.• Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none">• Hold eye open and flush slowly and gently with water for 15-30 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue flushing eye.• Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none">• Call poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER	

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-368-0832 for emergency medical treatment information. In case of fire or spill, information may be obtained by calling 1-800-624-6300.

NOTE TO PHYSICIAN: There is no specific antidote. All treatment should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.

For product information use Internet website www.nichino.net

Net Contents: _____

Active Ingredient Made in Japan; Formulated and Packaged in U.S.A.

Nichino America, Inc.
4550 New Linden Hill Road, Suite 501
Wilmington, DE 19808
www.nichino.net

ACCEPTED
SEP 7 2004

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, the pesticide registered under EPA Reg. No. 71711-15

APPLAUD

APPLAUD

3/12

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals
CAUTION

Harmful if swallowed. Wear thoroughly with soap and water after handling. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Harmful if inhaled. Avoid breathing spray mist. Remove contaminated clothing and wash before reuse.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washing, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should: Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

ENGINEERING CONTROL STATEMENT

When handlers use closed systems, enclosed cabs, or enclosures in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (d) (4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

ENVIRONMENTAL HAZARDS

For immediate use, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

IMPORTANT — Read the entire Directions for Use and Disclaimer of Warranties before using the product.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with the labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with the labeling and with the Worker Protection Standard, 40 CFR Part 170. The Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for labeling, distribution, notification, and emergency assistance. It also contains specific instructions and warnings pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in the box only apply to uses of the product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over long-sleeved shirt and long pants
- Waterproof gloves
- Socks and chemical-resistant footwear

NONAGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, or greenhouses. For other uses, including golf courses and other nonagricultural uses, do not enter treated areas without protective clothing until sprays have dried.

4/12

GENERAL INFORMATION

APPLAUD® 70WP insect growth regulator contains the active ingredient, buprofezin. It is effective against the nymph stages of whitefly, scales, mealybugs, planthoppers, and leafhoppers by inhibiting chitin biosynthesis, suppressing oviposition of adults, and reducing viability of eggs. Treated susceptible pests may remain alive on the plant for 3-7 days, but feeding damage during this time is typically very low. APPLAUD 70WP is not disruptive to beneficial insects and triles.

Good coverage is essential. Apply by ground or air in sufficient water volume. Orient nozzles to assure good coverage. Use of higher volume of water will assure better coverage, especially under adverse conditions such as hot, dry weather, and/or a dense canopy. The entire field should be treated.

Apply when economic infestations occur based on local information.

APPLAUD is not for sale, sale into, distribution, and or use in Nassau and Suffolk counties of New York state.

COURIER is not for sale, sale into, distribution, and or use in Nassau and Suffolk counties of New York state.

TALUS is not for sale, sale into, distribution, and or use in Nassau and Suffolk counties of New York state.

CENTAUR is not for sale, sale into, distribution, and or use in Nassau and Suffolk counties of New York state.

INSECTS CONTROLLED

Whiteflies: Silverleaf whitefly; Sweet potato whitefly; Greenhouse whitefly; Ash whitefly

Mealybugs: Grape mealybug; Vine mealybug; Citrus mealybug; Cornstock mealybug; Longtailed mealybug; Mexican mealybug; Oblique mealybug

Leafhoppers and Planthoppers: Eastern grape leafhopper; Glassy-winged sharpshooter; Variegated leafhopper; Western grape leafhopper; Cherry leafhopper; Potato leafhopper; Brown planthopper

Scales:

Soft Scales: Citricole scale; Flammole scale; Black scale; Brown soft scale; False oleander scale; Hemispherical scale; Indian wax scale and other wax scales; Tesselated scale; White peach scale

Armored Scales: Bofedual scale; Cactus scale; California red scale; Coconut scale; Fern scale; Florida red scale; Oystershell scale; San Jose scale

Margaroid Scale: Cottony cushion scale

MIXING, LOADING, AND HANDLING INSTRUCTIONS

USE PRECAUTIONS

DO NOT use APPLAUD 70WP water-soluble bags in a tankmix with products that contain boron, chromium, or other micronutrients.

GENERAL HANDLING INSTRUCTIONS

APPLAUD 70WP insect growth regulator is sealed in water-soluble bags. Do not open the foil-lined outer packaging until time of use. Do not use a knife or other sharp object to open the foil-lined outer packaging. Once the outer protective packaging has been opened, the water-soluble bag inside can be dropped into the spray tank at the appropriate time.

SPECIFIC MIXING INSTRUCTIONS: To ensure proper mixing of the soluble bags, follow the instructions below:

- APPLAUD 70WP ALONE:** Begin by filling the spray tank with clean water to prime the system (about 6 to 8 inches for most sprayers). Turn off the fill valve and drop the recommended number of soluble bags into the tank, distributing them across the bottom but not covering the circulation and outlet holes. After the bags have soaked for 2 minutes, start the agitation and agitate thoroughly for 2 minutes. With the agitation on, continue to fill the spray tank to the desired level. Agitate during spraying. If spraying must be stopped before emptying the sprayer, resume agitation before spraying the remainder of the load.
- APPLAUD 70WP TANK MIXTURES:** Follow all use directions as listed above under APPLAUD 70WP ALONE with the following exception: after the APPLAUD 70WP bags are thoroughly soaked and the water is added to fill the tank 1/2 full, add the recommended amount of wettable powder, soluble powder, flowable, emulsifiable concentrate, or soluble liquid product, while maintaining agitation. If a spray additive is needed, add a liquid fertilizer (except fertilizers containing boron, chromium or other micronutrients: see USE PRECAUTIONS), surfactant, or crop oil concentrate to the spray tank. Then continue adding water to the tank to achieve the desired level, while maintaining agitation.
- APPLAUD 70WP TANK MIXTURES WITH PRODUCTS IN WATER-SOLUBLE PACKAGING:** Follow all use directions as listed above under APPLAUD 70WP ALONE with the following exception: add the desired number of other water-soluble bags to the tank (if allowed by their label instructions) at the same time the APPLAUD 70WP bags are added to the spray tank.

5/12

HANDLING PRECAUTIONS FOR WATER-SOLUBLE BAGS

Do not allow the foil-lined outer packaging or portions thereof to fall into spray tank as they are not water-soluble and may clog the sprayer.

- Do not handle the soluble bag with wet gloves, as the bag will begin to dissolve on contact with even a small amount of water.
- Do not open the soluble bag or attempt to remove the APPLAUD 70WP from it. The product is not designed to be remeasured or subdivided; the entire contents of the water-soluble bag must be used.

Refer to the charts below for the proper application rate of APPLAUD 70WP insect growth regulator.

RESISTANCE MANAGEMENT

APPLAUD 70 WP belongs to a class of chemicals not previously used as pesticides and not known to be cross-resistant to other classes of insecticides. However, insect pests are known to develop resistance to products used repeatedly. Because resistance development cannot be predicted, the use of this product should conform to resistance management strategies established for the crop and use area. Such strategies may include rotating with insecticides with different modes of action. Consult your local pest control advisor or extension office for details. If resistance to this product develops in your area, it may not provide adequate control. If you experience difficulty with control, and resistance is a likely cause, consult your local state university horticultural specialist or local agricultural authorities for the best alternative method of control. To preserve the usefulness of APPLAUD 70WP, do not make more than two consecutive applications. Prior to subsequent applications, use an alternative chemistry with a different mode of action. If a different crop in an adjacent field has been treated with buprofezin against whitefly within the past 28 days, do not apply APPLAUD 70WP as the first whitefly application. Always consult your local crop advisor for the most appropriate control decision for your area. Resistance management strategies advise against applying rates lower than those recommended on the label.

(For marketing brand, Talus, add: if another insect growth regulator with the mode of action as a chitin synthesis inhibitor (i.e. cyromazine, diflubenzuron, and novaluron) has been used in the greenhouse within 28 days, do not apply Talus as the first whitefly application.)

6/12

APPLICATION RATE CHART FOR APPLAUD 70WP INSECT GROWTH REGULATOR

Crop	Pest	Rate	DOSAGE RATES
			lb Formulated Product
Almonds ¹	San Jose scale	1.5-2.0 lb a/A	2.14-2.86 (36 to 46 oz)
Bananas ²	Whiteflies, leafhoppers, mealybugs, scales	0.31 lb a/A	0.44 (7 oz)
Citrus ³	California red scale Cibicola scale Cottony cushion scale Glassy winged sharpshooter	1.5-2.0 lb a/A	2.14-2.86 (36 to 46 oz)
Cotton ⁴	Whiteflies	0.25 - 0.36 lb a/A	0.36 - 0.60 (6 to 8 oz)
Cucurbits ⁵ (including cucumbers, melons, pumpkin, squash)	Whiteflies	0.25-0.36 lb a/A	0.36 - 0.54 (6 to 9 oz)
Grapes ⁶	Leafhoppers Mealybugs	0.36-0.60 lb a/A	0.54 - 0.71 (9 to 12 oz)
Lettuce ⁷	Whiteflies	0.25-0.36 lb a/A	0.36 - 0.54 (6 to 9 oz)
Loquat, Lychee, Pomegranate, Rambutan, and Spanish Lime ⁸	Scales	1.56 lb a/A	2.25 (36oz)
Snap Bean ⁹	Whiteflies	0.36 lb a/A	0.54 (9 oz)
Tomatoes ¹⁰	Whiteflies Mealybugs Leafhoppers	0.25-0.36 lb a/A 0.36 lb a/A 0.36 lb a/A	0.36 - 0.54 (6 to 9 oz) 0.54 (9 oz) 0.54 (9 oz)
Ornamental plants in greenhouses; both and shadehouses; nurseries; landscape ornamentals; ground covers; field- and container-grown ornamentals; nonbearing fruit and nut trees and vines in nurseries; Christmas trees ¹¹	Whiteflies Mealybugs Leafhoppers Scales	0.25 lb a/100 gal 0.60 lb a/100 gal 0.60 lb a/100 gal 0.60 lb a/100 gal	0.36 lb (6 oz.) 0.71 lb (12 oz.) 0.71 lb (12 oz.) 0.66 lb (14 oz.)

¹ **Almonds**
 Treatment should be applied when peak crawler emergence occurs. Consult local or State agricultural authorities for details. Good coverage is essential. Apply by ground application using a minimum of 400 gallons of water per acre. Apply no more than one application per season. Do not apply within 60 days of harvest.

² **Bananas**
 Treatment should be applied when population level reaches economic threshold. Consult local or State agricultural authorities for details. Good coverage is essential. Apply by ground in a minimum of 15 gallons of water per acre. Use of a higher volume of water will assure better coverage, especially under adverse conditions, such as hot, dry weather and/or a dense canopy. Apply no more than four applications per season. Allow at least 14 days between applications. Do not apply within one day of harvest.

³ **Citrus**
 Treatment should be made when peak crawler emergence occurs. Consult local and State agricultural authorities for details. Good coverage is essential. Apply by ground application using 100 to 400 gallons of water per acre. Apply no more than two applications per season. Allow at least 30 days between applications. Do not apply within 60 days of harvest.

7/12

- 4 **Cotton**
Do not apply within 21 days of harvest in California and 14 days of harvest outside California.
Treatment should be applied when population level reaches economic threshold. Consult local and State agricultural authorities for details.
Good coverage is essential. With ground application, use 10 to 50 gallons of water per acre. For aerial application, use a minimum of 5 gallons per acre. Orient nozzles to assure good coverage. Use of a higher volume of water will assure better coverage, especially under adverse conditions, such as hot, dry weather and/or a dense canopy. Use the higher application rate and spray volume when treating a rapidly increasing insect population. The entire field should be treated.
Apply no more than two applications per season. Allow at least 28 days between applications.
See the RESISTANCE MANAGEMENT section.
- 5 **Cucurbits [including cucumbers; melons (cantaloupe, honeydew melons, watermelons, muskmelons); pumpkins; squash]**
Allow at least 7 days between applications to cucurbit crops.
Do not apply within 7 days of harvest in cucumbers.
Do not apply within 10 days of harvest in California and 7 days outside California in melons, pumpkins, and squash.
Do not make more than 2 applications per crop cycle. Do not make more than 4 applications per year.
- 6 **Grapes**
Treatment should be applied when population level reaches economic threshold. Consult local and State agricultural authorities for details.
Good coverage is essential. Apply by ground application using 50 to 200 gallons of water per acre, depending on the size of the grapes. Orient nozzles to assure good coverage. Use of a higher volume of water will assure better coverage, especially under adverse conditions, such as hot, dry weather and/or a dense canopy.
Apply no more than two applications per crop cycle. Allow at least 14 days between applications.
Do not apply within 30 days of harvest.
- 7 **Lettuce (head and leaf)**
Do not apply within 10 days of harvest in California and 7 days outside California.
Allow at least 7 days between applications to lettuce.
Do not make more than 2 applications per crop cycle. Do not make more than 4 applications per year.
- 8 **Longan, Lychee, Pulasan, Rambutan, and Spanish Lime**
Apply by ground in a minimum of 75 gal/acre spray volume.
Allow at least 14 days between applications.
Do not make more than 2 applications per year.
Do not apply within 21 days of harvest.
Do not exceed 72 oz/A (3.16 lb ai/A) per year.
- 9 **Snap Bean**
Allow at least 14 days between applications.
Do not exceed 0.76 lb ai/A per crop cycle.
Do not plant rotational crops other than those listed on this label for 120 days after the last application of Applaud.
Do not apply within 14 days of harvest.
- 10 **Tomatoes**
Treatment should be applied when population level reaches economic threshold. Consult local and State agricultural authorities for details.
Good coverage is essential. Apply in a minimum finished spray volume of 5 gallons per acre by air or 20 gallons per acre by ground to accommodate for the size of the tomato plants. Orient nozzles to assure better coverage. Use of a higher volume of water will assure better coverage, especially under adverse conditions, such as hot, dry weather and/or a dense canopy. The entire field should be treated.
Apply no more than two applications per growing cycle. Allow at least 28 days between applications.
Do not apply within 7 days of harvest.
See the RESISTANCE MANAGEMENT section.

*For marketing brand Talus, delete this section and replace with: Good coverage is essential. For greenhouse tomatoes, apply by ground on 2 acres as a minimum with 20 gallons water per acre. Apply in sufficient water to obtain complete coverage of all plant parts. Applications may be made with high volume, low volume or ultra low volume (thermal and non-thermal foggers, misters, etc.) ground equipment only. Follow the spray equipment manufacturer's directions to determine the amount of spray solution required to obtain thorough coverage. Consult the spray equipment manufacturer's operator's manual, spray nozzle catalogs and/or your crop advisor for more information.

8/12

11 Ornamentals

- Apply the specified dosage as a foliar spray in sufficient water for complete, uniform coverage, including stems and underside of leaves. Spray to the point of runoff.
- For whitefly or leafhopper control: Make first application as soon as adult insects begin to appear.
For mealybug control: Make first application as soon as insect activity is observed.
- For scale control: Make first application when crawlers are emerging.
- If additional insecticide applications are required for control, use another class of chemistry or a different insect Growth Regulator (IGR) with a different mode of action before making subsequent applications of buprofezin.
- Make no more than 2 applications of buprofezin per crop per season.
- Consult local or state agricultural authorities for details concerning economic thresholds for each target pest.

USE PRECAUTIONS

Apply in a minimum finished spray volume of 5 gallons per acre by air or 20 gallons per acre by ground, unless otherwise specified in Directions for Use.

Crop Rotational Restrictions: All crops listed on this label may be planted at any time. Small grain crops and leafy vegetables may be planted 30 days following application. All other crops may be planted 60 days following application.

Do not apply this product through any type of irrigation system.

For marketing brand Takas in the greenhouse, add: Apply in a minimum of 20 gallons per acre. Be sure to apply in sufficient water to obtain complete coverage of all plant parts. Applications may be made with high volume, low volume, or ultra low volume (thermal and non-thermal foggers, misters, etc.) ground equipment. Follow the spray equipment manufacturer's directions to determine the amount of spray solution required to obtain thorough coverage. Consult the spray equipment manufacturer's operator's manual, spray nozzle catalogs and/or your crop advisor for more information.

9/12

Application Chart for Determining Dosage Rate of Applaud 70WP and Talus

Rate lb a/A	lb Formulated/A	Number of 1.66 lb Packages (containing 3 X 0.36 lb water- soluble bags)	Number of Acres Treated
0.25	0.36		2.8
			8.4
			14.0
		1	3.0
0.31	0.44	3	9.0
		5	15.0
			2.3
			6.8
0.35	0.50		11.4
		1	2.5
		3	7.4
		5	12.3
0.38	0.54		2.0
			6.0
			10.0
		1	2.2
0.50	0.71	3	6.5
		5	10.8
			1.8
			5.5
1.5	2.14		9.2
		1	2.0
		3	6.0
		5	10.0
2.0	2.86		1.4
			4.2
			7.0
		1	1.5
		3	4.6
		5	7.6
			0.5
			1.4
			2.3
		1	0.5
		3	1.5
		5	2.5
			0.4
			1.1
			1.8

10/12

Application Chart for Determining Dosage Rate of Courier

Rate lb ai/A	lb Formulated/A	Number of 1 lb Water-Soluble Bags (within 5 lb outer packaging)	Number of Acres Treated
0.25	0.36	1	2.8
		3	8.4
		5	14.0
0.31	0.44		3.0
			9.0
			15.0
0.35	0.50	1	2.3
		3	6.8
		5	11.4
0.38	0.54		2.5
			7.4
			12.3
0.50	0.71	1	2.0
		3	6.0
		5	10.0
1.5	2.14		2.2
			6.5
			10.8
2.0	2.86	1	1.8
		3	5.5
		5	9.2
0.50	0.71		2.9
			6.0
			10.0
0.50	0.71	1	1.4
		3	4.2
		5	7.0
1.5	2.14		1.5
			4.5
			7.6
1.5	2.14	1	0.6
		3	1.4
		5	2.3
2.0	2.86		0.6
			1.6
			2.5
2.0	2.86	1	0.4
		3	1.1
		5	1.8

11/12

Application Charts for Determining Dosage Rate of Talus

Application Chart for Talus Using 2 oz Water Soluble Bags		
Rate ai/100 gal	oz formulated/100 gal	Number of 2 oz Water Soluble Bags
0.25	6 oz	3
0.50	12 oz	6
0.60	14 oz	7

Application Chart for Talus Using 0.6oz Water Soluble Bags		
Rate ai/100 gal	oz formulated/100 gal	Number of 0.6 oz Water Soluble Bags ¹
0.25	0.36 oz	10 bags in 100 gal 5 bags in 50 gal 1 bag in 10 gal
0.50	0.71 oz	20 bags in 100 gal 10 bags in 50 gal 2 bags in 10 gal
0.60	0.86 oz	24 bags in 100 gal 12 bags in 50 gal 3 bags in 10 gal 1 bag in 4 gal

¹ 0.6 oz water soluble bags (packaged 24 x 0.6 oz water soluble bags per 0.9 lb container) in water

12/1/2

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in original container, unopened, in a cool, dry place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Remove inner water-soluble bag from outer packaging and place inner bag into application equipment. Then dispose of empty outer packaging in a sanitary landfill, or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

IMPORTANT: READ BEFORE USE

By using this product, user or buyer accepts the following conditions, warranty, disclaimer of warranties, and limitations of liability.

CONDITIONS: The directions for use of this product are believed to be accurate and should be followed carefully. However, because of extreme weather and soil conditions, use methods and other factors beyond the control of Nichino America, Inc. (NAI), it is impossible for NAI to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. All such risks are assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: THERE ARE NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, WHICH EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of NAI is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. NAI disclaims any liability whatsoever for incidental or consequential damages, including, but not limited to, liability arising out of breach of contract, express or implied warranty (including warranties of merchantability and fitness for a particular purpose), tort, negligence, strict liability, or otherwise.

LIMITATIONS OF LIABILITY: THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT THE ELECTION OF NICHINO AMERICA, THE REPLACEMENT OF PRODUCT.

090304

Nichino America, Inc.
4650 New Linden Hill Rd., Suite 501
Wilmington, DE 19808

©2004, Nichino America, Inc.

APPLAUD®, COURIER®, TALUS® AND CENTAUR™ are trademarks of Nichino America, Inc.