07/08/2010



### UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

JUL - 8 2010

Nichino America, Inc. c/o Ms. Marie Maks 4550 New Linden Hill Road, Suite 501 Wilmington, DE 19808

Subject:

Supplemental Label for Use on Citrus by Air in Florida Fujimite 5EC Miticide/Insecticide Label EPA Reg. No. 71711-19 Your Letter Dated February 25, 2010 and Your Submission Dated April 9, 2010 Decision No: 430897

Dear Ms. Maks:

The Agency has reviewed your rebuttal regarding the adjuvants prohibition statement for all crops on the Fujimite 5EC Miticide/Insecticide label. The following is a summary of conclusions from the Agency's Science Advisory Council for Chemistry (ChemSAC) in the Health Effects Division, in the memorandum dated April 22, 2010:

- based on submitted residue trials conducted with citrus using non-ionic surfactants (NIS), residues from these trials demonstrate that the addition of the NIS adjuvant does not have a significant effect on residues levels in citrus harvested at a 14-day pre-harvest interval (PHI);
- ChemSAC is willing to translate these data to other crops that have PHIs of at least 14 days or more for applications made with NIS adjuvants, only; and,
- Additional data will be required should the you wish to pursue removal of the adjuvants use prohibition on all registered crops with PHIs less than 14 days, and for other classes of adjuvants.

On the basis of the conclusions in the Agency's memorandum responding to your rebuttal to the adjuvants prohibition, the labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. Two (2) copies of the finished labeling must be submitted prior to releasing the product for shipment. A stamped copy of the label is enclosed for your records.

At your next label printing, or within eighteen (18) months, whichever comes first, you must incorporate the supplemental labeling into the main product labeling. If you have any questions regarding these actions, please contact Rosanna Louie at (703) 308-0037.

Sincerely,

Mark Suarez Product Manager 13, Insecticide Branch Registration Division

Enclosures:

- Fenpyroximate. Response to Health effects Division (HED) Request for Field Trial Data to Support use of Adjuvants. April 22, 2010
- Fujimite 5EC Miticide/Insecticide Supplemental Label for Use on Citrus, Stamped Accepted

## SUPPLEMENTAL LABEL For Use in Florida Only

# FUJIMITE<sup>®</sup> 5EC Miticide/Insecticide For Use on Citrus

#### EPA Reg. No. 71711-19

This supplemental label expires on December 31, 2011 and must not be used or distributed after this date.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling and the EPA approved container label must be in the possession of the user at the time of application.

New application information appears on this supplemental labeling that do not appear on the Section 3 container label.

#### **GENERAL USE RESTRICTIONS**

- Do not apply more than 4.0 pints per acre per growing season.
- Do not make more than 2 applications per growing season.
- Allow 14 days between applications.
- Do not apply within 14 days of harvest.
- Use of non-ionic surfactants are permitted for use on citrus.
- Do not apply to citrus nurseries or citrus in greenhouses.
- Do not apply through any type of irrigation system.
- NOTICE: Before using this product, read the First Aid, Precautionary Statements, Conditions of Sale and Warranty, and complete Directions for Use found on the container labeling. All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.

| Сгор  | Pest  | Use Rate Per Acre                               | Spray Volume   | Application<br>Restrictions   |
|---|---|---|--|---|
| Citrus Crop Group:<br>Calamondin<br>Citrus citron<br>Citrus hybrids   | Asian citrus psyllid <sup>1</sup><br>4.0 pints<br>Citrus rust mite                  | Minimum 10<br>gallons water<br>per acre by air. | <ul> <li>For aerial<br/>applications, do not<br/>apply within 150<br/>feet of any aquatic</li> </ul>                     |   |
| (includes chironja, tangelo, tangor)<br>Grapefruit<br>Kumquat<br>Lemon<br>Lime<br>Mandarin (tangerine)<br>O'range, sour<br>Orange, sweet<br>Pummelo | Citrus Leafminer<br>Citrus Thrips<br>(suppression only)<br>Leafhoppers<br>Mealybugs | 2.0 to 4.0 pints                                | Minimum 100 <sup>2</sup><br>gallons water<br>per acre using<br>ground or<br>airblast (air<br>assist) spray<br>equipment. | area.<br>• For ground<br>applications, do not<br>apply within 75 feet<br>of any aquatic area. |
| Satsuma mandarin<br>White sapote (C <i>asimiroa</i> spp.)   |   | ,<br>I  |  |   |

<sup>2</sup> The use of higher water volumes is recommended for thorough coverage.


©2010 Nichino America, Inc. Fujimite® is a trademark of Nichino America, Inc.

070810



2/3

NICHINO AMERICA, INC. 4550 New Linden Hill Road Wilmington, DE 19808 (888) 740-7700



ΑССЕРТЕД

JUL - 8 2010

Fungicide, and Rodenticide Act,

Under the Federal Insecticide,