

71711-19

12/4/2012

1/32



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON D C 20460

CHEMICAL SAFETY
AND POLLUTION PREVENTION

OFFICE OF

December 04, 2012

Marie Maks
Nichino America, Inc
4550 New Linden Hill Road, Suite 501
Wilmington, DE 19808

Subject Application for Pesticide Amendment
 Fujimate® 5EC Miticide/Insecticide
 EPA Reg No 71711-19
 Your Submission Dated February 12, 2011
 Decision No 450032 and 451915

Dear Ms Maks

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. Two (2) copies of the finished labeling must be submitted prior to releasing the product for shipment. A stamped copy of the label is enclosed for your records.

If you have any questions contact Driss Benmhend at 703-308-9525 or by e-mail at Benmhend.driss@epa.gov

Sincerely,

Mark Suarez
Product Manager 13, Insecticide Branch
Registration Division (7504P)

Enclosure *Label stamped Accepted*

Nichino America, Inc.

2/32

GROUP 21A INSECTICIDE

FUJIMITE® 5EC miticide/insecticide

ACTIVE INGREDIENT

Fenpyroximate *tert*-butyl(E)- α -(1,3-dimethyl-5-phenoxy-pyrazol-4-ylmethyleneaminoxy)-*p*-toluate

5.0%

OTHER INGREDIENTS*

95.0%

TOTAL

100.0%

Contains 0.4 lb active ingredient per U.S. gallon
Contains petroleum distillates

EPA Reg No 71711-19

EPA Est No 67545-AZ-1

Marketing Brands

PORTAL® miticide/insecticide

Ortus Super miticide/insecticide

**KEEP OUT OF REACH OF CHILDREN
WARNING - AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle
(If you do not understand the label, find someone to explain it to you in detail)

FIRST AID	
If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
If swallowed	<p>Immediately call a poison control center or doctor.</p> <ul style="list-style-type: none"> Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-348-5832 for emergency medical treatment information. In case of fire or spills, information may be obtained by calling 1-800-424-9300.	
NOTE TO PHYSICIAN Contains petroleum distillates. Vomiting may cause aspiration pneumonia. Probable mucosal damage may contraindicate the use of gastric lavage.	

NET CONTENTS

Active Ingredient Made in Japan Formulated and Packaged in U.S.A. for

NICHINO AMERICA INC
4550 New Linden Hill Road
Wilmington, DE 19808
1-888-740-7700

**ACCEPTED
DEC 04 2012**

Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended, for the
pesticide registered under

EPA Reg No 71711-19

FUJIMITE 5EC 71711-19
D 38 041411 15 Page 1 of 19

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals

WARNING - AVISO

Causes substantial but temporary eye injury Harmful if inhaled or swallowed Do not get in eyes or on clothing Avoid contact with skin and breathing spray mist

Wear long-sleeved shirt and long pants socks shoes and chemical resistant gloves (such as barrier laminate or viton) Wear protective eyewear (safety glasses goggles or face shield)

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear

- Long-sleeved shirt and long pants
- Chemical resistant gloves (such as barrier laminate or viton)
- Protective eyewear (such as safety glasses goggles or face shield)
- Shoes plus socks
- For overhead exposure wear chemical resistant headgear

When mixing and loading wear a chemical-resistant apron When cleaning equipment wear a chemical-resistant apron

Follow manufacturer s instructions for cleaning/maintaining PPE If no such instructions for washables use detergent and hot water Keep and wash PPE separately from other laundry When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40CFR 170 240(d)(4-6)] the handler PPE requirements may be reduced or modified as specified in the WPS

User Safety Recommendations
Users should
Wash thoroughly with soap and water after handling and before eating drinking chewing gum or using tobacco
Remove and wash contaminated clothing before reuse

ENVIRONMENTAL HAZARDS

This pesticide is very highly toxic to fish and aquatic invertebrates For terrestrial uses do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas Do not contaminate water when cleaning equipment or disposing of equipment washwaters

This product may impact surface water quality due to runoff of rain water This is especially true for poorly draining soils and soils with shallow ground water This product is classified as having medium to high potential for reaching both surface water and aquatic sediment via runoff for several weeks after application A level well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds streams and springs will reduce the potential loading of this chemical from runoff water and sediment Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours Sound erosion control practices will reduce this product s potential to reach aquatic sediment via runoff

ENDANGERED SPECIES RESTRICTIONS

This product may pose a hazard to endangered aquatic species Follow all use directions

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame

Minimum Honey Bee Toxicity

Applications to all crops may be made at anytime Fenpyroximate is practically nontoxic to bees and wasps when used according to this label

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

Do not apply this product in a way that will contact workers or other persons either directly or through drift Only protected handlers may be in the area during application For any requirements specific to your State or Tribe consult the agency in your State responsible for pesticide regulation

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170 This Standard contains requirements for the protection of agricultural workers on farms forests nurseries and greenhouses and handlers of agricultural pesticides It contains requirements for training decontamination notification and emergency assistance It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants soil or water is

- Coveralls worn over short-sleeved shirt and short pants
- Socks
- Chemical-resistant footwear
- Protective eyewear (such as safety glasses goggles or face shield)
- Chemical-resistant gloves (such as barrier laminate or viton)

GENERAL INFORMATION

FUJIMITE® 5EC miticide/insecticide is used for the control of leafhoppers mealybugs mites psylla psyllids and whiteflies **FUJIMITE 5EC miticide/insecticide** stops mite feeding immediately after application **FUJIMITE 5EC miticide/insecticide** controls all motile stages of mites by inhibiting cellular respiration in the mitochondrion of cells which results in rapid cessation of all biological activities including feeding and reproduction Mortality of mites can be observed within 3-7 days after intoxication

FUJIMITE 5EC miticide/insecticide works primarily through contact action **so thorough spray coverage is necessary** Mix with sufficient water and apply as a foliar spray to obtain uniform coverage Dense foliage or excessive growth will often prevent adequate coverage adjust spray volumes accordingly Treat plants when pests are immature or at a susceptible stage and populations are building before crop damage occurs

Target Species	
Apple rust mite (suppression only)	Pacific spider mite
Asian citrus psyllid	Six spotted mite
Banks grass mite	Pear psylla
Broad mite	Pear rust mite
Carmine mite	Pecan leaf scorch mite
Citrus bud mite	Potato leafhopper
Citricola scale	Tomato (Potato) psyllid
Citrus leafminer (suppression only)	Powdery Mildew (suppression only)
Citrus red mite	Strawberry spider mite
Citrus rust mite	Texas citrus mite

Citrus thrips (suppression only)	Tomato russet mite
Cyclamen mite	Two-spotted spider mite
European red mite	Variegated leafhopper
Glassy-winged sharpshooter (suppression only)	White apple leafhopper
Grape leafhopper	Whiteflies
McDaniel mite	Willamette spider mite
Mealybug species	Persea mite
Mint bud mite	Avocado Brown mite

APPLICATION DIRECTIONS

- Applications should be made immediately after the spray solution is prepared
- Apply with properly calibrated spray equipment
- Apply by ground or air using the recommended water spray volume found in the Directions for Use section of this label
- For aerial equipment use larger droplet size (greater than 200 microns)
- Thorough spray coverage is essential for mite and insect control
- For best results apply when pest populations are beginning to build before reaching economic thresholds Consult your local agricultural advisor or state cooperative extension service for further information

USE OF ADJUVANTS

When thorough coverage is a concern it is recommended that a spray adjuvant be used to maximize uniformity of coverage and performance of FujiMite 5EC Use a non-ionic activator type wetting spreading or penetrating adjuvant or horticultural spray oil adjuvant Do not use a dormant oil or binder or sticker type adjuvant Non ionic adjuvants (NIS) should contain at least 75% surfactant Crop oil concentrates (COC) methylated seed or vegetable oils (MSO) organosilicone products (OS) or blends of these adjuvants should contain at least 15% emulsifier/surfactant Check compatibility of any adjuvant used with FujiMite 5EC before using Follow the Directions for Use on each adjuvant product label for rates of use and use restrictions

APPLICATION RESTRICTIONS

- For all other use patterns do not apply within 75 feet of fish-bearing waters
- Use by air on citrus is limited to the states of Florida and Texas
- For aerial use on citrus in Florida do not apply within 150 feet of all aquatic areas
- Do not use products with the same mode of action in consecutive applications
- Do not plant rotational crops other than those listed on this label for 30 days following the last application of this product
- Do not apply through any type of irrigation system
- FujiMite and Portal Brands Do not use in greenhouse structures
- Do not apply by Alternate Row Middle (ARM) spray method

RESISTANCE MANAGEMENT

Repeated use of the same chemistry has been shown to result in the buildup of resistant strains of mites or other insects Do not use **FUJIMITE 5EC miticide/insecticide** in successive miticide applications Rotate the use of **FUJIMITE 5EC Miticide/insecticide** with alternate mode of action insecticides

- Miticides must be rotated with alternate products for resistance management Do not use products with the same mode of action [METI 2 inhibitor/IRAC Group 21A] in consecutive applications

Consult your local crop advisor for the most appropriate alternative products Resistance management strategies recommend that you DO NOT apply rates lower than recommended on the label

MIXING DIRECTIONS

FUJIMITE 5EC Miticide/Insecticide Alone Shake well before using. Begin with clean equipment. Fill spray tank with $\frac{3}{4}$ of the amount of water needed for the intended application and then turn on agitation. Pour recommended amount of product on the surface of water in the spray tank. Add the balance of the water to the spray tank with agitation running. Keep agitation running during filling and spraying operations. If spraying must be stopped before emptying the sprayer, resume agitation before spraying the remainder of the load.

FUJIMITE 5EC Miticide/Insecticide Tank Mixtures Shake well before using. Read and follow all label directions for each tank mix product prior to any tank mixing with **FUJIMITE 5EC miticide/insecticide**. This product can be mixed with other registered pesticides for use on labeled crops or sites in accordance with the most restrictive use directions and precautions. Follow all use directions as listed above under **FUJIMITE 5EC Miticide/Insecticide Alone** with the following exception: after the **FUJIMITE 5EC miticide/insecticide** is thoroughly mixed and the tank is $\frac{3}{4}$ full, add the recommended amount of wettable powder, soluble powder, flowable, emulsifiable concentrate, or soluble liquid product **while maintaining agitation**. Then continue adding water to the tank to achieve the desired level while maintaining agitation.

If you have no experience with the combination you are considering, you should conduct a test to determine physical compatibility. To determine physical compatibility, add the recommended proportions of each chemical with the same proportion of water as will be present in the chemical supply tank into a suitable container, mix thoroughly, and allow to stand for five minutes. If the combination remains mixed or can be readily re-mixed, the mixture is considered physically compatible.

SPRAY DRIFT REDUCTION MANAGEMENT

Avoid spray drift to all other crops and nontarget areas. Do not apply when weather conditions may cause drift. Do not allow this product to drift onto nontarget areas. Drift may result in illegal residues or injury to adjacent crops and vegetation. To avoid spray drift, **DO NOT** apply aerially when wind speed is greater than 10 mph or during periods of temperature inversions. Use of larger droplet size will also reduce spray drift.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR

The interaction of equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making decisions.

Droplet size, boom height, and wind speed are the primary factors determining drift. The specific application conditions required for the use of this product are described below.

Controlling Droplet Size – General Techniques

Volume

Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure

Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.**

Nozzle Type

Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

**Controlling Droplet Size – Aircraft
Number of Nozzles**

Use the minimum number of nozzles with the highest flow rate that provide uniform coverage

Nozzle Orientation

Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.

Nozzle Type

Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

Boom Height and Length – Ground and Aircraft

Boom Height (ground) Setting the boom at the lowest labeled height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.

Boom Height (aircraft) Application more than 10 feet above the canopy increases the potential for spray drift.

Boom Length (aircraft) The minimum boom length should not exceed $\frac{3}{4}$ of the wing length. Using shorter booms decreases drift potential. For helicopters, the minimum boom length should not exceed $\frac{9}{10}$ of the rotary blade to prevent droplets from entering the rotor vortices.

Wind

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given wind speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. **AVOID GUSTY OR WINDLESS CONDITIONS.** **Note:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. This cloud can move in unpredictable directions due to the light and variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, nontarget crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

Shielded Sprayers

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with the uniform deposition of the product.

Air Assisted (Air Blast) Field Crop Sprayers

Air assisted field crop sprayers carry droplets to the target via a downward directed air stream. Some may reduce the potential for drift, but if a sprayer is unsuitable for the application and/or set up improperly, high drift potential can result. It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application, is configured properly, and that drift is not occurring.

Note Air assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration. Consult the application equipment section of this label to determine if use of an air assisted sprayer is recommended.

Air Assisted (Air Blast) Tree And Vine Sprayers

Air assisted tree and vine sprayers carry droplets into the canopy of trees and vines via a radially or laterally directed air stream. These sprayers are not suitable for applying herbicides. In addition to the general drift management practices already described, the following specific practices will further reduce the potential for drift:

- Adjust the deflectors and aiming devices so that spray is only directed into the canopy
 - Block off upward pointed nozzles when there is no overhanging canopy
 - Use only enough air volume to penetrate the canopy and provide good coverage
- Do not allow spray to go beyond the edge of the cultivated area. Spray the outside row only from outside the planting.

APPLICATION RATE CHART FOR
FUJIMITE 5EC MITICIDE/INSECTICIDE

Avocado, black sapote, canistel, mamey sapote, mango, papaya, sapodilla, and star apple		
Pest	Rate/Acre	Use Directions
Mites (see Target Species) Whiteflies	1.0 to 2.0 pints	<p>USE RESTRICTIONS</p> <ul style="list-style-type: none"> • Apply by ground using a minimum of 95 gallons of water per acre or by air using a minimum of 50 gallons of water per acre • Do not apply more than 4.0 pints per acre per growing season • Do not make more than 2 applications per growing season • Allow 14 days between applications • Pre harvest Interval (PHI) 1 day • Do not apply through any type of irrigation system

CITRUS FRUITS (Crop Group 10-10) Australian desert lime, Australian finger lime, Australian round lime, Brown River finger lime, calamondin, citron, citrus hybrids, grapefruit, Japanese summer grapefruit, kumquat, lemon, lime, Mediterranean mandarin, mount white lime, New Guinea wild lime, orange, sour, orange, sweet, pummelo, Russell River lime, satsuma mandarin, sweet lime, tachibana orange, Tahiti lime, tangelo, tangerine (mandarin orange, clementine), tangor, trifoliate orange, unqi fruit, cultivars, varieties, and/or hybrids of these		
Pest	Rate/Acre	Use Directions
Citrus rust mite	4.0 pints	USE RESTRICTIONS All geographies <ul style="list-style-type: none"> • Apply by ground using a minimum of 100 gallons of water per acre. For full size trees use a minimum of 200 gallons of water per acre. • Apply by air using a minimum of 10 gallons of water per acre. • For aerial use in Florida do not apply within 150 feet of all aquatic areas. • Do not apply more than 4.0 pints per acre per growing season. • Do not make more than 2 applications per growing season. • Allow 14 days between applications. • Pre harvest Interval (PHI) 14 days. • Do not apply to citrus nurseries or citrus in greenhouses. • Do not apply through any type of irrigation system. West of the Mississippi River <ul style="list-style-type: none"> • Do not apply more than 8.0 pints per acre per growing season. • Do not apply by air except in Texas. • Apply by air in a minimum of 10 gallons of water in Texas. • In California for control of citricola scale apply by ground using 500 gallons of water.
Asian citrus psyllid ¹ Citrus thrips (suppression only) Citrus leafminer (suppression only) Leafhoppers Mealybugs Other Mites (see Target Species)	2.0 to 4.0 pints	
¹ For best results use for control of adults and nymphs present at time of application when newly expanding foliage flush is present		

COTTON		
Pest	Rate/Acre	Use Directions
Mites (see Target Species)	Early season¹ (when cotton is less than 10 inches in height) 0.4 to 1.0 pints ²	USE RESTRICTIONS <ul style="list-style-type: none"> • Apply by ground using a minimum of 10 gallons of water per acre or by air using a minimum of 3 gallons of water per acre. As canopy density increases use of higher water volume will assure better coverage. • Do not apply more than 2.0 pints per acre per growing season. • Do not make more than 2 applications per growing season. • Allow 14 days between applications. • Pre-harvest Interval (PHI) 14 days. • Do not apply through any type of irrigation system.
	Mid season (when cotton is more than 10 inches in height) 1.0 pint by ground 1.0 to 2.0 pints by air	
Whiteflies	2.0 pints	

¹ For early season use when cotton is less than 10 inches in height. FUJIMITE 5EC Miticide/insecticide may also be applied as a directed spray using ground spray equipment.

² When applying by ground equipment apply as a directed spray for best results.

CUCUMBERS		
Pest	Rate/Acre	Use Directions
Mites (see Target Species) Tomato/Potato Psyllid Whiteflies	1.0 to 2.0 pints	USE RESTRICTIONS <ul style="list-style-type: none"> • Apply by ground using a minimum of 40 gallons of water per acre or by air using a minimum of 10 gallons of water per acre. • Do not apply more than 4.0 pints per acre per crop cycle. • Do not make more than 2 applications per crop cycle. • Allow 14 days between applications. • Pre harvest Interval (PHI) 1 day. • Do not apply through any type of irrigation system.

FIELD CORN, POP CORN, SILAGE CORN, SEED CORN (limited to States of California, Colorado, Hawaii, Kansas, New Mexico, Oklahoma, and Texas)		
Pest	Rate/Acre	Use Directions
Mites (see Target Species)	1.0 to 2.0 pints	<p>USE RESTRICTIONS</p> <ul style="list-style-type: none"> • Apply by ground application using a minimum of 10 gallons of water per acre or by air using a minimum of 5 gallons of water per acre • Do not apply more than 4.0 pints per acre per crop cycle • Do not make more than 2 applications per crop cycle • Allow 14 days between applications • Pre harvest Interval (PHI) 14 days for forage silage stover and grain • Do not apply through any type of irrigation system

FRUITING VEGETABLES (Crop Group 8-10)		
African eggplant, bush tomato, bell pepper, cocona, currant tomato, eggplant, garden huckleberry, goji berry, groundcherry, martynia, naranjilla, okra, pea eggplant, pepino, nonbell pepper, roselle, scarlet eggplant, sunberry, tomatillo, tomato, tree tomato, cultivars, varieties, and/or hybrids of these		
Pest	Rate/Acre	Use Directions
Mites (see Target Species) Tomato/Potato Psyllid Whiteflies	2.0 pints	<p>USE RESTRICTIONS</p> <ul style="list-style-type: none"> • Apply by ground using a minimum of 20 gallons of water per acre or by air using a minimum of 5 gallons of water per acre • Do not apply more than 4.0 pints per acre per crop cycle • Do not make more than 2 applications per crop cycle • Allow 14 days between applications • Pre-harvest Interval (PHI) 1 day • Do not apply through any type of irrigation system

GRAPES		
Raisin Table Wine		
Pest	Rate/ Acre	Use Directions
Mites (see Target Species) Mealybugs Powdery Mildew (suppression)	2.0 pints	USE RESTRICTIONS <ul style="list-style-type: none"> • Apply by ground using a minimum of 50 gallons of water per acre • For vines with a heavy canopy or in high pressure situations higher water volumes are recommended. If lower water volume amounts are used tractor speed must be reduced to ensure complete coverage • Do not apply by air • Do not apply more than 2.0 pints per acre per growing season • Do not make more than 2 applications per growing season • Allow 14 days between applications • Pre-harvest Interval (PHI) 14 days • Do not apply through any type of irrigation system
Willamette spider mite	1.5 to 2.0 pints	
Leafhoppers	1.0 to 2.0 pints ¹	
¹ Use higher rate for dense foliage. Best control of leafhoppers is achieved by applications when majority of the population is in an immature development stage.		

HOPS*		
Pest	Rate/ Acre	Use Directions
Mites (see Target Species)	2.0 to 3.0 pints	USE RESTRICTIONS <ul style="list-style-type: none"> • Apply by ground using a minimum of 100 gallons of water per acre • Do not apply more than 3.0 pints of product per acre per growing season • Do not apply by air • Allow 14 days between applications • Pre-harvest Interval (PHI) 15 days • Do not apply through any type of irrigation system RECOMMENDATIONS <ul style="list-style-type: none"> • For best results apply before mite populations exceed 5 mites per leaf

***NOTE** Leaf yellowing may occur when FUJIMITE 5EC miticide/ insecticide is combined with spray oil in excess of 1% of the spray volume. If this symptom occurs, it is usually more pronounced on newly expanding leaves. This symptom may occur in plants under stress and is worsened by certain conditions including the following

- High Temperatures (air temperatures exceeding 90°F at the time of application or within a few days after application)
- Wet soil conditions and high humidity (rainy, misty, or foggy weather within a few days after application)
- Storm damage (including hail and wind)

LOW GROWING BERRY (Crop Subgroup 13-07G)* EXCEPT CRANBERRIES bearberry, bilberry, blueberry, lowbush, cloudberry, lingonberry, muntries, partridgeberry, strawberry, cultivars, varieties, and/or hybrids of these		
Pest	Rate/Acre	Use Directions
Mites (see Target Species) Whiteflies	1.0 to 2.0 pints	USE RESTRICTIONS <ul style="list-style-type: none"> • Apply by ground application using a minimum of 25 gallons of water per acre • Do not apply by air • Do not apply more than 4.0 pints per acre per crop cycle • Do not make more than 2 applications per crop cycle • Allow 14 days between applications • Pre harvest Interval (PHI) 1 day • Do not apply through any type of irrigation system
<p>*NOTE Temporary pinking of immature green berries may be observed after a Fujimite application on certain strawberry varieties. This effect is transient and does not affect fruit sizing, color or quality. Avoid puddling of spray solution on plastic mulch as this can potentially result in underside scarring of fruit in direct contact with the plastic.</p>		

MELONS (Crop Subgroup 9A) muskmelon, including hybrids and/or varieties of <i>Cucumis melo</i> (including true cantaloupe, cantaloupe, casaba, Santa Claus melon, crenshaw melon, honeydew melon, honey balls, Persian melon, golden pershaw melon, mango melon, pineapple melon, snake melon), citron melon, and watermelon, including hybrids and/or varieties of (<i>Citrullus spp</i>)		
Pest	Rate/Acre	Use Directions
Mites (see Target Species) Whiteflies	2.0 pints	USE RESTRICTIONS <ul style="list-style-type: none"> • Apply by ground application using a minimum of 20 gallons of water per acre • Do not apply by air • Do not apply more than 4.0 pints per acre per crop cycle • Do not make more than 2 applications per crop cycle • Allow 14 days between applications • Pre harvest Interval (PHI) 3 days • Do not apply through any type of irrigation system

MINT (peppermint, spearmint)		
Pest	Rate/Acre	Use Directions
Mites (see Target Species)	1.0 to 2.0 pints	<p>USE RESTRICTIONS</p> <ul style="list-style-type: none"> • Apply by ground using a minimum of 25 gallons of water per acre • Do not apply by air • Do not apply more than 4.0 pints per acre per growing season • Do not make more than 2 applications per growing season • Allow 7 days between applications • Pre-harvest Interval (PHI) 1 day • Do not apply through any type of irrigation system

NONBEARING DECIDUOUS FRUIT, TREE NUT, AND VINES		
Pest	Rate/Acre	Use Directions
Leafhoppers Mealybugs Mites (see Target Species)	1.0 to 2.0 pints	<p>USE RESTRICTIONS</p> <ul style="list-style-type: none"> • Apply by ground using a minimum of 75 gallons of water per acre • Do not apply by air • Do not apply more than 2.0 pints per acre per growing season • Do not make more than 2 applications per growing season • Allow 14 days between applications • Do not harvest edible crops for 12 months following application unless the crop is listed on the label • Do not apply to citrus nurseries or citrus in greenhouses • Do not apply through any type of irrigation system

NONBEARING DECIDUOUS FRUIT, TREE NUT, AND VINES		
Pest	Rate/Acre	Use Directions
Leafhoppers Mealybugs Mites (see Target Species)	2.0 pints	<p>USE RESTRICTIONS</p> <ul style="list-style-type: none"> • Apply by ground using a minimum of 75 gallons of water per acre • Do not apply by air • Do not apply more than 2.0 pints per acre per growing season • Do not make more than 1 application per growing season • Do not harvest edible crops for 12 months following application unless the crop is listed on the label • Do not apply to citrus nurseries or citrus in greenhouses • Do not apply through any type of irrigation system

Ornamentals		
Pest	Rate/Acre	Use Directions
Leafhoppers Mealybugs Mites (see Target Species)	1.0 to 2.0 pints	<p>USE RESTRICTIONS</p> <ul style="list-style-type: none"> • Apply by ground using a minimum of 50 gallons of water per acre • Do not apply by air • Do not apply more than 2.0 pints per acre per growing season • Allow 14 days between applications • Do not apply to citrus nurseries or citrus in greenhouses • Do not apply through any type of irrigation system

POME FRUITS (Crop Group 11 10) Apple, azarole, crabapple, loquat, mayhaw, medlar, pear, pear, Asian, quince, quince, Chinese, quince, Japanese, tejocote, cultivars, varieties, and/or hybrids of these		
Pest	Rate/Acre	Use Directions
Leafhoppers Mealybugs Mites (see Target Species)	1.0-2.0 pints	<p>USE RESTRICTIONS</p> <ul style="list-style-type: none"> • Apply by ground using a minimum of 100 gallons of water per acre • Do not apply by air • Do not apply more than 2.0 pints per acre per growing season • Do not make more than 2 applications per growing season • Allow 14 days between applications • Pre-harvest Interval (PHI) 14 days • Do not apply through any type of irrigation system • Do not apply by Alternate Row Middle (ARM) spray method
Leafhoppers Mealybugs Mites (see Target Species) Pear psylla	2.0 pints	<p>USE RESTRICTIONS</p> <ul style="list-style-type: none"> • Apply by ground using a minimum of 100 gallons of water per acre • Do not apply by air • Do not apply more than 2.0 pints per acre per growing season • Do not make more than 1 application per growing season • Pre harvest Interval (PHI) 14 days • Do not apply through any type of irrigation system • Do not apply by Alternate Row Middle (ARM) spray method

Snap Beans		
Pest	Rate/Acre	Use Directions
Mites (see Target Species) Whiteflies	2.0 pints	USE RESTRICTIONS <ul style="list-style-type: none"> • Apply by ground using a minimum of 30 gallons of water per acre or by air using a minimum of 5 gallons of water per acre • Do not apply more than 4.0 pints per acre per crop cycle • Do not make more than 2 applications per crop cycle • Allow 14 days between applications • Pre harvest Interval (PHI) 1 day • Do not apply through any type of irrigation system

ALMONDS AND PISTACHIOS (West of the Mississippi River)		
Pest	Rate/Acre	Use Directions
Mites (see Target Species)	1.5 to 4.0 pints	USE RESTRICTIONS WEST OF THE MISSISSIPPI RIVER <ul style="list-style-type: none"> • Apply by ground using a minimum of 100 gallons of water per acre • Do not apply by air • Do not apply more than 8.0 pints per acre per growing season • Do not make more than 2 applications per growing season • Allow 14 days between applications • Pre harvest Interval (PHI) 14 days • Do not apply through any type of irrigation system

TREE NUTS (Crop Group 14) (West of the Mississippi River) (excluding almonds and pistachios) Beech nut, Brazil nut, butternut, cashew, chestnut, chinquapin, filbert (hazelnut), hickory nut, macadamia nut, pecan, walnut, black and English		
Pest	Rate/Acre	Use Directions
Mites (see Target Species)	1.5 to 4.0 pints	USE RESTRICTIONS WEST OF THE MISSISSIPPI RIVER <ul style="list-style-type: none"> • Apply by ground using a minimum of 100 gallons of water per acre • Do not apply by air • Do not apply more than 4.0 pints per acre per growing season • Do not make more than 2 applications per growing season • Allow 14 days between applications • Pre-harvest Interval (PHI) 14 days • Do not apply through any type of irrigation system

TREE NUTS (Crop Group 14) and Pistachio Almond, beech nut, Brazil nut, butternut, cashew, chestnut, chinquapin, filbert (hazelnut), hickory nut, macadamia nut, pecan, pistachio, walnut, black and English		
Pest	Rate/Acre	Use Directions
Mites (see Target Species)	2.0 pints	USE RESTRICTIONS ALL GEOGRAPHIES <ul style="list-style-type: none"> • Apply by ground using a minimum of 100 gallons of water per acre • Do not apply by air • Do not apply more than 2.0 pints per acre per growing season • Do not make more than 1 application per growing season • Pre-harvest Interval (PHI) 14 days • Do not apply through any type of irrigation system

GREENHOUSE CUCUMBERS		
Pest	Rate/Acre	Use Directions
Mites (see Target Species) (Tomato) Potato Psyllid Whiteflies	1.0 to 2.0 pints	USE RESTRICTIONS <ul style="list-style-type: none"> • Do not use products with the same mode of action in consecutive applications • Apply by ground using a minimum of 40 gallons of water per acre • Do not apply more than 4.0 pints per acre crop cycle • Do not make more than 2 applications per crop cycle • Do not make more than 4 applications per year • Allow 14 days between applications • Pre harvest Interval (PHI) 1 day • Do not apply through any type of irrigation system • Do not apply in Ultra Low Volume Equipment RECOMMENDATIONS <ul style="list-style-type: none"> • Apply in sufficient water to obtain complete coverage of all plant parts. Applications may be made with high volume or low volume ground equipment only. Follow the spray equipment manufacturer's directions to determine the amount of spray solution required to obtain thorough coverage. Consult the spray equipment manufacturer's operator's manual, spray nozzle catalogs and/or your crop advisor for more information.

GREENHOUSE TOMATOES		
Pest	Rate/Acre	Use Directions
Mites (see Target Species) (Tomato) Potato Psyllid Whiteflies	2.0 pints	<p>USE RESTRICTIONS</p> <ul style="list-style-type: none"> • Do not use products with the same mode of action in consecutive applications • Apply by ground using a minimum of 100 gallons of water per acre • Do not apply more than 4.0 pints per acre per crop cycle • Do not make more than 2 applications per growing season • Do not make more than 4 applications per year • Allow 14 days between applications • Pre-harvest Interval (PHI) 1 day • Do not apply through any type of irrigation system • Do not apply in Ultra Low Volume Equipment <p>RECOMMENDATIONS</p> <ul style="list-style-type: none"> • Apply in sufficient water to obtain complete coverage of all plant parts. Applications may be made with high volume or low volume ground equipment only. Follow the spray equipment manufacturer's directions to determine the amount of spray solution required to obtain thorough coverage. Consult the spray equipment manufacturer's operator's manual, spray nozzle catalogs and/or your crop advisor for more information.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal

PESTICIDE STORAGE Store in a cool dry place

PESTICIDE DISPOSAL Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility

CONTAINER HANDLING

(Non-refillable container equal to or less than 5 gallons)
 Nonrefillable container DO NOT reuse or refill this container Triple rinse (or equivalent) promptly after emptying Triple rinse as follows Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip Fill the container ¼ full with water and recap Shake for 10 seconds Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal Drain for 10 seconds after the flow begins to drip Repeat this procedure two more times Then offer for recycling if available or reconditioning or puncture and dispose of in a sanitary landfill or by incineration or if allowed by State or local authorities by burning If burned stay out of smoke

(Refillable container greater than 5 gallons)
 Refillable container Refill this container with pesticide only Do not reuse this container for any other purpose Cleaning the container before final disposal is the responsibility of the person disposing of the container Cleaning before refilling is the responsibility of the refiller To clean the container before final disposal empty the remaining contents from this container into application equipment or mix tank Fill the container about 10 percent full with water Agitate vigorously or recirculate water with the pump for 2 minutes Pour or pump rinsate into application equipment or rinsate collection system Repeat this rinsing procedure two more times

Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or if allowed by state and local authorities by burning If burned stay out of smoke

IMPORTANT READ BEFORE USE

By using this product user or buyer accepts the following conditions warranty disclaimer of warranties and limitations of liability

CONDITIONS The directions for use of this product are believed to be accurate and must be followed carefully However because of extreme weather and soil conditions use methods and other factors beyond the control of Nichino America Inc (NAI) it is impossible for NAI to eliminate all risks associated with the use of this product As a result crop injury or ineffectiveness is always possible To the extent consistent with applicable law all such risks are assumed by the user or buyer

DISCLAIMER OF WARRANTIES TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THERE ARE NO WARRANTIES EXPRESS OR IMPLIED OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE WHICH EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL No agent of NAI is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein To the extent consistent with applicable law NAI disclaims any liability whatsoever for incidental or consequential damages including but not limited to liability arising out of breach of contract express or implied warranty (including warranties of merchantability and fitness for a particular purpose) tort negligence strict liability or otherwise

LIMITATIONS OF LIABILITY TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT WARRANTY TORT NEGLIGENCE STRICT LIABILITY OR OTHERWISE SHALL NOT EXCEED THE PURCHASE PRICE PAID OR AT THE ELECTION OF NICHINO AMERICA THE REPLACEMENT OF PRODUCT

©2012 Nichino America Inc

FUJIMITE® and PORTAL® are trademarks of Nichino America Inc

100212

21/32

SUPPLEMENTAL LABEL

GROUP 21A INSECTICIDE

Portal® Miticide/Insecticide
EPA Reg No 71711-19

For Use on Fruiting Vegetables

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling
This labeling and the EPA approved container label must be in the possession of the user at the time of application

New use directions appear on this supplemental labeling that supersede the Section 3 container label

NOTICE Before using this product read the First Aid Precautionary Statements Conditions of Sale and Warranty and complete Directions for Use found on the container labeling All applicable directions restrictions and precautions on the EPA registered label are to be followed

APPLICATION RATE CHART

FRUITING VEGETABLES (Crop Group 8-10)		
African eggplant bush tomato bell pepper cocona currant tomato eggplant garden huckleberry goji berry groundcherry martynia naranjilla okra pea eggplant pepino nonbell pepper roselle scarlet eggplant sunberry tomatillo tomato tree tomato cultivars varieties and/or hybrids of these		
Pest	Rate/Acre	Use Directions
Mites (see Target Species) Tomato/Potato Psyllid Whiteflies	2.0 pints	USE RESTRICTIONS <ul style="list-style-type: none"> Do not apply within 75 feet of fish-bearing waters Apply by ground application using a minimum of 20 gallons of water per acre or by air using a minimum of 5 gallons of water per acre Do not apply more than 4.0 pints per acre per crop cycle Do not make more than 2 applications per crop cycle Allow 14 days between applications Pre-harvest Interval (PHI) 1 day Do not apply through any type of irrigation system

©2012 Nichino America Inc
Portal® is a trademark of Nichino America Inc

NICHINO AMERICA
4550 New Linden Hill Road
Suite 501
Wilmington DE 19808
888 740 7700

**ACCEPTED
DEC 04 2012**
Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended, for the
pesticide registered under

EPA Reg No 71711-19

22/32

SUPPLEMENTAL LABEL

GROUP 21A INSECTICIDE

FujiMite® 5EC Miticide/Insecticide
EPA Reg No 71711-19

For Use on Fruiting Vegetables

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling
This labeling and the EPA approved container label must be in the possession of the user at the time of application

New use directions appear on this supplemental labeling that supersede the Section 3 container label

NOTICE Before using this product read the First Aid Precautionary Statements Conditions of Sale and Warranty and complete Directions for Use found on the container labeling All applicable directions restrictions and precautions on the EPA registered label are to be followed

APPLICATION RATE CHART

FRUITING VEGETABLES (Crop Group 8 10) African eggplant bush tomato bell pepper cocona currant tomato eggplant garden huckleberry goji berry groundcherry martynia naranjilla okra pea eggplant pepino nonbell pepper roselle scarlet eggplant sunberry tomatillo tomato tree tomato cultivars varieties and/or hybrids of these		
Pest	Rate/Acre	Use Directions
Mites (see Target Species) Tomato/Potato Psyllid Whiteflies	2 0 pints	USE RESTRICTIONS <ul style="list-style-type: none"> • Do not apply within 75 feet of fish-bearing waters • Apply by ground application using a minimum of 20 gallons of water per acre or by air using a minimum of 5 gallons of water per acre • Do not apply more than 4 0 pints per acre per crop cycle • Do not make more than 2 applications per crop cycle • Allow 14 days between applications • Pre harvest Interval (PHI) 1 day • Do not apply through any type of irrigation system

©2012 Nichino America Inc
FujiMite® is a trademark of Nichino America Inc

NICHINO AMERICA
4550 New Linden Hill Road
Suite 501
Wilmington DE 19808
888 740 7700

**ACCEPTED
DEC 0 4 2012**

**Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended, for the
pesticide registered under**

EPA Reg No 71711-19

23/32

SUPPLEMENTAL LABEL

GROUP 21A INSECTICIDE

Portal® Miticide/Insecticide
EPA Reg No 71711-19

For Use on Citrus

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling
This labeling and the EPA approved container label must be in the possession of the user at the time of application

New use directions appear on this supplemental labeling that supersede the Section 3 container label

NOTICE Before using this product read the First Aid Precautionary Statements Conditions of Sale and Warranty and complete Directions for Use found on the container labeling All applicable directions restrictions and precautions on the EPA registered label are to be followed

APPLICATION RATE CHART

CITRUS FRUITS (Crop Group 10-10) Australian desert lime Australian finger lime Australian round lime Brown River finger lime calamondin chironja-citron citrus hybrids grapefruit Japanese summer grapefruit kumquat lemon lime Mediterranean mandarin mount white lime New Guinea wild lime orange sour orange sweet pummelo Russell River lime satsuma mandarin sweet lime tachibana orange Tahiti lime tangelo tangerine (mandarin) tangor trifoliate orange unqi fruit cultivars varieties and/or hybrids of these		
Pest	Rate/Acre	Use Directions
Citrus rust mite	4 0 pints	USE RESTRICTIONS
Asian citrus psyllid ¹	2 0 to 4 0 pints	<ul style="list-style-type: none"> • Apply by ground using a minimum of 100 gallons of water per acre For full size trees use a minimum of 200 gallons of water per acre • Use by air is limited to Florida and Texas • Apply by air using a minimum of 10 gallons of water per acre • Do not apply within 75 feet of fish-bearing waters For aerial use in Florida do not apply within 150 feet of all aquatic areas • Do not apply more than 4 0 pints per acre per growing season • Do not make more than 2 applications per growing season • Allow 14 days between applications • Pre-harvest Interval (PHI) 14 days • Do not apply to citrus nurseries or citrus in greenhouses • Do not apply through any type of irrigation system
Citrus thrips		
Citrus leafminer (suppression only)		
Leafhoppers		
Mealybugs		
Other Mites		
(see Target Species)		

¹For best results use for control of adults and nymphs present at time of application when newly expanding foliage flush is present

©2012 Nichino America, Inc
Portal® is a trademark of Nichino America, Inc

ACCEPTED
DEC 04 2012

NICHINO AMERICA
4550 New Linden Hill Road
Suite 501
Wilmington DE 19808
888 740 7700

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under

EPA Reg No 71711-19

Expiration date 11/15/15

24/32

SUPPLEMENTAL LABEL

GROUP 21A INSECTICIDE

Portal® Miticide/Insecticide
EPA Reg No 71711-19

For Use on Cucumbers

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling
This labeling and the EPA approved container label must be in the possession of the user at the time of application

New use directions appear on this supplemental labeling that supersede the Section 3 container label

NOTICE Before using this product read the First Aid Precautionary Statements Conditions of Sale and Warranty and complete Directions for Use found on the container labeling All applicable directions restrictions and precautions on the EPA registered label are to be followed

APPLICATION RATE CHART

Cucumbers		
Pest	Rate / Acre	Use Directions
Mites (see Target species) Whiteflies	2.0 pints	USE RESTRICTIONS <ul style="list-style-type: none"> Do not apply within 75 feet of fish bearing waters Apply by ground application using a minimum of 40 gallons of water per acre and by air using a minimum of 10 gallons of water per acre Do not apply more than 4.0 pints per acre per crop cycle Do not make more than 2 applications per crop cycle Allow 14 days between applications Pre harvest Interval (PHI) 1 day Do not apply through any type of irrigation system

©2012 Nichino America Inc
Portal® is a trademark of Nichino America Inc

NICHINO
AMERICA

4550 New Linden Hill Road
Suite 501
Wilmington DE 19808
888 740 7700

ACCEPTED

DEC 04 2012

**Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended, for the
pesticide registered under**

EPA Reg No 71711-19

25/
32

SUPPLEMENTAL LABEL

GROUP 21A INSECTICIDE

FujiMite® 5EC Miticide/Insecticide
EPA Reg No 71711-19

For Use on Cucumbers

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling
This labeling and the EPA approved container label must be in the possession of the user at the time of application

New use directions appear on this supplemental labeling that supersede the Section 3 container label

NOTICE Before using this product read the First Aid Precautionary Statements Conditions of Sale and Warranty and complete Directions for Use found on the container labeling All applicable directions restrictions and precautions on the EPA registered label are to be followed

APPLICATION RATE CHART

Cucumbers		
Pest	Rate/Acre	Use Directions
Mites (see Target Species) Tomato/Potato Psyllid Whiteflies	2.0 pints	USE RESTRICTIONS <ul style="list-style-type: none"> • Do not apply within 75 feet of fish bearing waters • Apply by ground application using a minimum of 40 gallons of water per acre or by air using a minimum of 10 gallons of water per acre • Do not apply more than 4.0 pints per acre per crop cycle • Do not make more than 2 applications per crop cycle • Allow 14 days between applications • Pre-harvest Interval (PHI) 1 day • Do not apply through any type of irrigation system

©2012 Nichino America Inc
FujiMite® is a trademark of Nichino America Inc

NICHINO AMERICA
4550 New Linden Hill Road
Suite 501
Wilmington DE 19808
888 740 7700

ACCEPTED
DEC 04 2012
Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended, for the
pesticide registered under

EPA Reg No 71711-19

26/
32

SUPPLEMENTAL LABEL

GROUP 21A INSECTICIDE

FujiMite® 5EC Miticide/Insecticide
EPA Reg No 71711-19

For Use on Field Corn, Silage Corn, Pop Corn, and Corn Grown for Seed (limited to States of California and Hawaii)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling and the EPA approved container label must be in the possession of the user at the time of application.

New use directions appear on this supplemental labeling that supersede the Section 3 container label.

NOTICE Before using this product read the First Aid Precautionary Statements Conditions of Sale and Warranty and complete Directions for Use found on the container labeling All applicable directions restrictions and precautions on the EPA registered label are to be followed.

APPLICATION RATE CHART

FIELD CORN, POP CORN, SILAGE CORN, SEED CORN (limited to States of California and Hawaii)		
Pest	Rate / Acre	Use Directions
Mites (see Target Species)	1.5 to 2.0 pints	USE RESTRICTIONS <ul style="list-style-type: none"> Do not apply within 75 feet of fish bearing waters Apply by ground application using a minimum of 10 gallons of water per acre or by air using a minimum of 5 gallons per acre Do not apply more than 4.0 pints per acre per crop cycle Do not make more than 2 applications per crop cycle Allow 14 days between applications Pre harvest Interval (PHI) 14 days for forage silage stover and grain Do not apply through any type of irrigation system

©2012 Nichino America Inc
FujiMite® is a trademark of Nichino America Inc

ACCEPTED
DEC 04 2012

Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended, for the
pesticide registered under

**NICHINO
AMERICA**
4550 New Linden Hill Road
Suite 501
Wilmington DE 19808
888 740 7700

EPA Reg No 7174-19

27/32

SUPPLEMENTAL LABEL

GROUP 21A INSECTICIDE

Portal® Miticide/Insecticide
EPA Reg No 71711-19

For Use on Field Corn, Silage Corn, Pop Corn, and Corn Grown for Seed (limited to States of Colorado, Kansas, New Mexico, Oklahoma, and Texas)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling and the EPA approved container label must be in the possession of the user at the time of application.

New use directions appear on this supplemental labeling that supersede the Section 3 container label.

NOTICE Before using this product read the First Aid Precautionary Statements Conditions of Sale and Warranty and complete Directions for Use found on the container labeling All applicable directions restrictions and precautions on the EPA registered label are to be followed.

APPLICATION RATE CHART

FIELD CORN, POP CORN, SILAGE CORN, SEED CORN (limited to States of Colorado, Kansas, New Mexico, Oklahoma, and Texas)		
Pest	Rate / Acre	Use Directions
Mites (see Target Species)	1.5 to 2.0 pints	USE RESTRICTIONS <ul style="list-style-type: none"> Do not apply within 75 feet of fish bearing waters Apply by ground application using a minimum of 10 gallons of water per acre or by air using a minimum of 5 gallons per acre Do not apply more than 4.0 pints per acre per crop cycle Do not make more than 2 applications per crop cycle Allow 14 days between applications Pre-harvest Interval (PHI) 14 days for forage silage stover and grain Do not apply through any type of irrigation system

©2012 Nichino America Inc
Portal® is a trademark of Nichino America Inc

ACCEPTED
DEC 04 2012
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under

NICHINO AMERICA
4550 New Linden Hill Road
Suite 501
Wilmington DE 19808
888 740 7700

EPA Reg No 71711-19

28/32

SUPPLEMENTAL LABEL

GROUP 21A INSECTICIDE

FujiMite® 5EC Miticide/Insecticide
EPA Reg No 71711-19

For Use on Snap Beans

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling and the EPA approved container label must be in the possession of the user at the time of application.

New use directions appear on this supplemental labeling that supersede the Section 3 container label.

NOTICE Before using this product read the First Aid Precautionary Statements Conditions of Sale and Warranty and complete Directions for Use found on the container labeling. All applicable directions restrictions and precautions on the EPA registered label are to be followed.

APPLICATION RATE CHART

Snap Beans		
Pest	Rate / Acre	Use Directions
Mites (see Target species) Whiteflies	2.0 pints	USE RESTRICTIONS <ul style="list-style-type: none"> Do not apply within 75 feet of fish-bearing waters Apply by ground application using a minimum of 30 gallons of water per acre and by air using a minimum of 5 gallons of water per acre Do not apply more than 4.0 pints per acre per crop cycle Do not make more than 2 applications per crop cycle Allow 14 days between applications Pre harvest Interval (PHI) 1 day Do not apply through any type of irrigation system

©2012 Nichino America Inc.
FujiMite® is a trademark of Nichino America Inc.

NICHINO AMERICA
4550 New Linden Hill Road
Suite 501
Wilmington DE 19808
888 740 7700

**ACCEPTED
DEC 04 2012**
Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended, for the
pesticide registered under

EPA Reg No 71711-19

29/
132

SUPPLEMENTAL LABEL

GROUP 21A INSECTICIDE

Portal® Miticide/Insecticide
EPA Reg No 71711-19

For Use on Snap Beans

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling and the EPA approved container label must be in the possession of the user at the time of application.

New use directions appear on this supplemental labeling that supersede the Section 3 container label.

NOTICE Before using this product read the First Aid Precautionary Statements Conditions of Sale and Warranty and complete Directions for Use found on the container labeling. All applicable directions restrictions and precautions on the EPA registered label are to be followed.

APPLICATION RATE CHART

Snap Beans		
Pest	Rate / Acre	Use Directions
Mites (see Target species) Whiteflies	2.0 pints	USE RESTRICTIONS <ul style="list-style-type: none"> Do not apply within 75 feet of fish bearing waters Apply by ground application using a minimum of 30 gallons of water per acre and by air using a minimum of 5 gallons of water per acre Do not apply more than 4.0 pints per acre per crop cycle Do not make more than 2 applications per crop cycle Allow 14 days between applications Pre harvest Interval (PHI) 1 day Do not apply through any type of irrigation system

©2012 Nichino America Inc
Portal® is a trademark of Nichino America Inc

ACCEPTED
DEC 04 2012
Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended, for the
pesticide registered under

NICHINO
AMERICA
4550 New Linden Hill Road
Suite 501
Wilmington DE 19808
888 740 7700

EPA Reg No 71711-19

30/
32

SUPPLEMENTAL LABEL

GROUP 21A INSECTICIDE

Portal® Miticide/Insecticide
EPA Reg No 71711-19

For use on avocado, black sapote, canistel, mamey sapote, mango, papaya, sapodilla, star apple

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling and the EPA approved container label must be in the possession of the user at the time of application.

New use directions appear on this supplemental labeling that supersede the Section 3 container label.

NOTICE Before using this product read the First Aid Precautionary Statements Conditions of Sale and Warranty and complete Directions for Use found on the container labeling All applicable directions restrictions and precautions on the EPA registered label are to be followed

APPLICATION RATE CHART

AVOCADO, BLACK SAPOTE, CANISTEL, MAMEY SAPOTE, MANGO, PAPAYA, SAPODILLA, STAR APPLE		
Pest	Rate / Acre	Use Directions
Mites (see Target species) Whiteflies	2.0 pints	USE RESTRICTIONS <ul style="list-style-type: none"> Do not apply within 75 feet of fish-bearing waters Apply by ground application using a minimum of 95 gallons of water per acre and by air using a minimum of 50 gallons of water per acre Do not apply more than 4.0 pints per acre per growing season Do not make more than 2 applications per growing season Allow 14 days between applications Pre-Harvest Interval (PHI) 1 day Do not apply through any type of irrigation system

©2012 Nichino America Inc
Portal® is a trademark of Nichino America Inc

ACCEPTED
DEC 04 2012
Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended, for the
pesticide registered under

NICHINO
AMERICA
4550 New Linden Hill Road
Suite 501
Wilmington DE 19808
888 740 7700

EPA Reg No 71711-19
D 30 042111 5

31/
32

SUPPLEMENTAL LABEL

GROUP 21A INSECTICIDE

FujiMite® 5EC Miticide/Insecticide
EPA Reg No 71711-19

For use on avocado, black sapote, canistel, mamey sapote, mango, papaya, sapodilla, star apple

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling and the EPA approved container label must be in the possession of the user at the time of application.

New use directions appear on this supplemental labeling that supersede the Section 3 container label.

NOTICE Before using this product read the First Aid Precautionary Statements Conditions of Sale and Warranty and complete Directions for Use found on the container labeling All applicable directions restrictions and precautions on the EPA registered label are to be followed

APPLICATION RATE CHART

AVOCADO, BLACK SAPOTE CANISTEL MAMEY SAPOTE MANGO PAPAYA, SAPODILLA STAR APPLE		
Pest	Rate / Acre	Use Directions
Mites (see Target species) Whiteflies	2.0 pints	USE RESTRICTIONS <ul style="list-style-type: none"> Do not apply within 75 feet of fish-bearing waters Apply by ground application using a minimum of 95 gallons of water per acre and by air using a minimum of 50 gallons of water per acre Do not apply more than 4.0 pints per acre per growing season Do not make more than 2 applications per growing season Allow 14 days between applications Pre Harvest Interval (PHI) 1 day Do not apply through any type of irrigation system

©2012 Nichino America Inc
FujiMite® is a trademark of Nichino America Inc

NICHINO AMERICA
4550 New Linden Hill Road
Suite 501
Wilmington DE 19808
888 740 7700

ACCEPTED
DEC 04 2012

Under the Federal Insecticide, Fungicide and Rodenticide Act, as amended for the pesticide registered under

EPA Reg No 71711-19
D 29 042111 5

32/32

SUPPLEMENTAL LABEL

GROUP 21A INSECTICIDE

FujiMite® 5EC Miticide/Insecticide
EPA Reg No 71711-19

For Use on Citrus

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling
This labeling and the EPA approved container label must be in the possession of the user at the time of application

New use directions appear on this supplemental labeling that supersede the Section 3 container label

NOTICE Before using this product read the First Aid Precautionary Statements Conditions of Sale and Warranty and complete Directions for Use found on the container labeling All applicable directions restrictions and precautions on the EPA registered label are to be followed

APPLICATION RATE CHART

CITRUS FRUITS (Crop Group 10-10) Australian desert lime Australian finger-lime Australian round lime Brown River finger lime calamondin citron citrus hybrids grapefruit Japanese summer grapefruit kumquat lemon lime Mediterranean mandarin mount white lime New Guinea wild lime orange sour orange sweet pummelo Russell River lime satsuma mandarin sweet lime tachibana orange Tahiti lime tangelo tangerine (mandarin) tangor trifoliate orange uniq fruit cultivars varieties and/or hybrids of these		
Pest	Rate/Acre	Use Directions
Citrus rust mite	4 0 pints	USE RESTRICTIONS • Apply by ground using a minimum of 100 gallons of water per acre For full size trees use a minimum of 200 gallons of water per acre • Do not apply within 75 feet of fish-bearing waters • Do not apply more than 4 0 pints per acre per growing season • Do not make more than 2 applications per growing season • Allow 14 days between applications • Pre-harvest Interval (PHI) 14 days • Do not apply to citrus nurseries or citrus in greenhouses • Do not apply through any type of irrigation system
Asian citrus psyllid ¹ Citrus thrips (suppression only) Citrus leafminer (suppression only) Leafhoppers Mealybugs Other Mites (see Target Species)	2 0 to 4 0 pints	
¹ For best results use for control of adults and nymphs present at time of application when newly expanding foliage flush is present		

©2012 Nichino America Inc
FujiMite® is a trademark of Nichino America Inc

NICHINO AMERICA
4550 New Linden Hill Road
Suite 501
Wilmington DE 19808
888 740 7700

**ACCEPTED
DEC 04 2012**

Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended, for the
pesticide registered under

EPA Reg No 71711-19