

71711-20

07-10-2009

1/15



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

JUL 10 2009

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Ms. Marie A. Maks, Manager, Regulatory Affairs
Nichino America, Inc.
Linden Park, Suite 501
4550 New Lindon Hill Road
Wilmington, DE 19808

Subject: New Uses on Brassica, Coffee, and Pomegranate

EPA Reg. Nos.: Buprofezin Technical (EPA Reg. No. 71711-16), Buprofezin 40SC Insect Growth Regulator (EPA Reg. No. 71711-20), Applaud 70 DF Growth Regulator (EPA Reg. No. 71711-21)

Decisions: 396787, 391754, 396789, 391757, 391758


The labeling referred to above, submitted in connection with registration under the Federal insecticide, Fungicide, and Rodenticide Act (FIFRA), is acceptable subject to the comments listed below.

Nichino must conduct and submit the following four studies with the timeframe listed below as a condition of registration:

- a. Acute Neurotoxicity study (OPPTS Guideline No. 870.6200(a)) within 1 year of the date of this letter;
- b. Subchronic Neurotoxicity study (OPPTS Guideline No. 870-6200(b)) within 18 months of the date of this letter;
- c. Immunotoxicity study (OPPTS Guideline No. 870.7800) within 18 month of the date of this letter; and
- d. Developmental Thyroid study (Special Study) within 2 years of the date of this letter.

If you have any questions/comments, please contact Melody Banks on (703) 305-5413.

Sincerely,



Kimberly Nesci
Product Manager 11
Insecticide Branch
Registration Division (7507P)

3/15

GROUP 16 INSECTICIDE

Nichino America, Inc.

BUPROFEZIN 40SC Insect Growth Regulator

Marketing Brand: Courier 40SC, Talus 40SC

ACTIVE INGREDIENT:

Buprofezin: 2-tert-butylimino-3-isopropyl-5-phenyl-1,3,5-thiadiazinan-4-one 40.00%

OTHER INGREDIENTS: 60.00%

TOTAL 100.00%

Contains 3.6 lbs. buprofezin per U.S. gallon

EPA Reg. No. 71711-20

EPA Est. No.: 67545-AZ-1

KEEP OUT OF REACH OF CHILDREN WARNING-AVISO


Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain to you in detail.)

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
If on skin	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-348-5832 for emergency medical treatment information. In case of fire or spills, information may be obtained by calling 1-800-424-9300.	

NOTE TO PHYSICIAN: There is no specific antidote. All treatment should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.

Active Ingredient Made in China; Formulated and Packaged in U.S.A. for

Nichino America, Inc.
4550 New Linden Hill Road, Suite 501
Wilmington, DE 19808



4
15

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING - AVISO

Causes substantial but temporary eye injury. Avoid contact with skin or clothing. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Wear long-sleeved shirt and long pants, socks, shoes, and chemical-resistant gloves.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant (such as nitrile or butyl) gloves
- Protective eyewear
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should: Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

IMPORTANT – Read the entire **Directions for Use** and **Disclaimer of Warranties** before using this product.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls over long-sleeved shirt and long pants
- chemical-resistant (such as nitrile or butyl) gloves
- protective eyewear
- shoes plus socks

NONAGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, or greenhouses. For other uses, including golf courses and other nonagricultural uses, do not enter treated areas without protective clothing until sprays have dried.

GENERAL INFORMATION

BUPROFEZIN 40SC insect growth regulator contains the active ingredient, buprofezin. It is effective against the nymph stages of whitefly, scales, mealybugs, planthoppers, and leafhoppers by inhibiting chitin biosynthesis, suppressing oviposition of adults, and reducing viability of eggs. Treated susceptible pests may remain alive on the plant for 3-7 days, but feeding damage during this time is typically very low. BUPROFEZIN 40SC is not disruptive to beneficial insects and mites.

BUPROFEZIN 40SC is not for sale, sale into, distribution, and or use in Nassau and Suffolk counties of New York state.

INSECTS CONTROLLED

Whiteflies: Silverleaf whitefly; Sweet potato whitefly; Greenhouse whitefly; Ash whitefly; Banded wing whitefly

Mealybugs: Grape mealybug; Vine mealybug; Citrus mealybug; Comstock mealybug; Longtailed mealybug; Mexican mealybug; Obscure mealybug; Striped mealybug; Gill's mealybug

Leafhoppers and Planthoppers: Eastern grape leafhopper; Glassy-winged sharpshooter; Variegated leafhopper; Western grape leafhopper; Cherry leafhopper; Potato leafhopper; Brown planthopper

Scales:

Soft Scales: Citricola scale; Barnacle scale; Black scale; Brown soft scale; False oleander scale; Hemispherical scale; Indian wax scale and other wax scales; Tessellated scale; White peach scale

Armored Scales: Boisduval scale; Cactus scale; California red scale; Coconut scale; Fern scale; Florida red scale; Oystershell scale; San Jose scale

Margarodid Scale: Cottony cushion scale

RESISTANCE MANAGEMENT

BUPROFEZIN 40SC is classified by IRAC in Group 16 – chitin synthesis inhibitors, and is not known to be cross-resistant to other classes of insecticides. However, insect pests are known to develop resistance to products used repeatedly. Because resistance development cannot be predicted, the use of this product should conform to resistance management strategies established for the crop and use area. Such strategies may include rotating with insecticides with different modes of action. Consult your local pest control advisor or extension office for details. If resistance to this product develops in your area, it may not provide adequate control. If you experience difficulty with control, and resistance is a likely cause, consult your local state university horticultural specialist or local agricultural authorities for the best alternative method of control. To preserve the usefulness of BUPROFEZIN 40SC, do not make more than two consecutive applications. Prior to subsequent applications, use an alternative chemistry with a different mode of action. Always consult your local crop advisor for the most appropriate control decision for your area. Resistance management strategies advise against applying rates lower than those recommended on the label.

For greenhouses: If another insect growth regulator with the mode of action as a chitin synthesis inhibitor (i.e. cyromazine, diflubenzuron, and novaluron) has been used in the greenhouse within 28 days, do not apply BUPROFEZIN 40SC as the first whitefly application.

MIXING DIRECTIONS

Shake well before using. Read and follow all label directions for each tank mix product prior to any tank mixing with **BUPROFEZIN 40SC**. This product can be mixed with other registered pesticides for use on labeled crops or sites, in accordance with the most restrictive use directions and precautions. No labeled dose rate should be exceeded. **BUPROFEZIN 40SC** cannot be mixed with any product containing a label prohibition against such mixing.

BUPROFEZIN 40SC ALONE: Begin with clean equipment. Fill spray tank with $\frac{3}{4}$ of the amount of water needed for the intended application and then turn on agitation. Pour recommended amount of product on the surface of water in the spray tank. Add the balance of the water to the spray tank with agitation running. Keep agitation running during filling and spraying operations. If spraying must be stopped before emptying the sprayer, resume agitation before spraying the remainder of the load.

BUPROFEZIN 40SC TANK MIXTURES: Follow all use directions as listed above under **BUPROFEZIN 40SC ALONE** with the following exception: after the **BUPROFEZIN 40SC** is thoroughly mixed and the tank is $\frac{3}{4}$ full, add the recommended amount of wettable powder, soluble powder, flowable, emulsifiable concentrate, or soluble liquid product, **while maintaining agitation**. Then continue adding water to the tank to achieve the desired level, while maintaining agitation.

BUPROFEZIN 40SC TANK MIXTURES WITH PRODUCTS IN WATER-SOLUBLE PACKAGING: Follow all use directions as listed above under **BUPROFEZIN 40SC ALONE** with the following exception: add the desired number of water-soluble bags to the tank (if

6/15

allowed by their label instructions) at the same time the **BUPROFEZIN 40SC** is added. **Note:** If using products in water soluble packaging, do not tank mix with products that contain boron, chromium, or other micronutrients.

BUPROFEZIN 40SC is physically and biologically compatible with many registered pesticides, fertilizers or micronutrients. Contact your supplier for advice when considering mixing **BUPROFEZIN 40SC** with other pesticides, fertilizers or micronutrients. For further information, contact your local Nichino America Representative. If you have no experience with the combination you are considering, you should conduct a test to determine physical compatibility. To determine physical compatibility, add the recommended proportions of each chemical with the same proportion of water, as will be present in the chemical supply tank, into a suitable container, mix thoroughly and allow to stand for five minutes. If the combination remains mixed, or can be readily re-mixed, the mixture is considered physically compatible.

APPLICATION DIRECTIONS

Applications should be made immediately after the spray solution is prepared. Thorough spray coverage is essential for effective control. Applications may be made with high or low volume spray equipment that provides thorough coverage of the plant. Apply with properly calibrated spray equipment. For best results, apply when pest populations are beginning to build, before reaching economic thresholds. Consult your local agricultural advisor, state cooperative extension service, or regional Nichino America representative for recommendations.

USE RESTRICTIONS

- Do not apply this product through any type of irrigation system.
- Crop Rotational Restrictions:**

CROP	PLANTBACK TIMING
All crops registered for use with buprofezin	0 days following application
Leafy vegetables not registered for use with buprofezin	30 days following application
Root and tuber vegetables	60 days following application
All other crops	365 days following application

USE PRECAUTIONS

- Apply in a minimum finished spray volume of 5 gallons per acre by air or 20 gallons per acre by ground, unless otherwise specified in Directions for Use.

APPLICATION RATE CHART FOR BUPROFEZIN 40SC INSECT GROWTH REGULATOR

Crop	Pest	Formulated Product/A	Directions for Use
Head and Stem Brassica Subgroup 5A: Broccoli Brussels sprouts Cabbage Cauliflower Chinese broccoli Chinese cabbage (Napa) Chinese mustard cabbage Cavalo broccolo Kohlrabi	Leafhoppers Planthoppers Whiteflies	9.0 to 13.6 fl oz	USE RESTRICTIONS <ul style="list-style-type: none"> Do not make more than 2 applications per crop cycle. Do not exceed 27.2 fl oz per crop cycle. Allow at least 7 days between applications. Do not make more than 4 applications per year Do not apply within 1 day of harvest.
Cotton	Whiteflies	9.0 to 12.5 fl oz	USE RESTRICTIONS <ul style="list-style-type: none"> Do not make more than 2 applications per growing season. Allow at least 28 days between applications. With ground application, use 10 to 50 gallons of water per acre. For aerial application, use a minimum of 5 gallons of water per acre. Do not apply within 21 days of harvest in California and 14 days outside California. RECOMMENDATIONS <ul style="list-style-type: none"> Treatment should be applied when population level reaches economic threshold. Consult local and state agricultural authorities for details. Good coverage is essential. Orient nozzles to assure good

			<p>coverage.</p> <ul style="list-style-type: none"> • Use of a higher volume of water will assure better coverage, especially under adverse conditions, such as hot, dry weather and/or a dense canopy. Use the higher application rate and spray volume when treating a rapidly increasing insect population. The entire field should be treated.
Cucurbit Crop Group 9: Cucumbers; Melons Cantaloupe Honeydew melons Watermelons Muskmelons) Pumpkins; Squash	Leafhoppers Planthoppers	13.6 fl oz	USE RESTRICTIONS <ul style="list-style-type: none"> • Do not make more than 2 applications per crop cycle. • Allow at least 7 days between applications to cucurbits. • Do not make more than 4 applications per year. • Do not apply within 10 days of harvest in California and 7 days outside California in melons, pumpkins, and squash. • Do not apply within 7 days of harvest in cucumbers.
	Whiteflies	9.0 to 13.6 fl oz	
Leafy Vegetable Crop Group 4: Celery Endive Head lettuce Leaf lettuce Parsley Spinach	Leafhoppers Planthoppers Whiteflies	9.0 to 13.6 fl oz	USE RESTRICTIONS <ul style="list-style-type: none"> • Do not make more than 2 applications per crop cycle. • Allow at least 7 days between applications. • Do not make more than 4 applications per year. • Do not apply within 7 days of harvest.
Snap Beans	Leafhoppers Planthoppers Whiteflies	9.0 to 13.6 fl oz	USE RESTRICTIONS <ul style="list-style-type: none"> • Do not exceed 27.2 fl oz per crop cycle. • Allow at least 14 days between applications. • Do not apply within 14 days of harvest.
Low-Growing Berry Subgroup 13-07G: Strawberry Bearberry Bilberry Blueberry, lowbush Cloudberry Cranberry Lingonberry Muntries Partridgeberry	Leafhoppers Mealybugs Planthoppers Whiteflies	9.0 to 12.0 fl oz	USE RESTRICTIONS <ul style="list-style-type: none"> • With ground application, use at least 80 gallons of water per acre. For aerial application, use at least 5 gallons of water per acre. • Do not make more than 2 applications per crop cycle. • Allow at least 10 days between applications. • Do not apply within 3 days of harvest. RECOMMENDATIONS <ul style="list-style-type: none"> • Good coverage is essential.

Fruiting Vegetable Crop Group 8: Bell pepper Chili pepper Cooking pepper Pimento Eggplant Groundcherry Okra Pepino Sweet pepper Tomatillo Tomato	Leafhoppers Mealybugs Planthoppers	13.6 fl oz	USE RESTRICTIONS <ul style="list-style-type: none"> • Apply no more than two applications per crop cycle. • Allow at least 5 days between applications. • Do not apply within 1 day of harvest. • For Greenhouse tomatoes, apply by ground on 2 acre minimum with 20 gallons of water per acre. RECOMMENDATIONS <ul style="list-style-type: none"> • Treatment should be applied when population level reaches economic threshold. Consult local and state agricultural authorities for details. • Good coverage is essential. Orient nozzles to assure better coverage. Use of a higher volume of water will assure better
--	--	------------	---

8
15

	Whiteflies	9.0 to 13.6 fl oz	<p>coverage, especially under adverse conditions, such as hot, dry weather and/or a dense canopy. The entire field should be treated.</p> <ul style="list-style-type: none"> • For Greenhouse tomatoes, apply in sufficient water to obtain complete coverage of all plant parts. Applications may be made with high volume, low volume or ultra low volume (thermal and non-thermal foggers, misters, etc.) ground equipment only. Follow the spray equipment manufacturer's directions to determine the amount of spray solution required to obtain thorough coverage. Consult the spray equipment manufacturer's operator's manual, spray nozzle catalogs and/or your crop advisor for more information.
<p>Ornamental plants in greenhouses; lath and shadehouses; nurseries; landscape ornamentals; ground covers; field- and container-grown ornamentals; non-bearing fruit and nut trees in nurseries; Christmas trees</p>	Leafhoppers Mealybugs Planthoppers	18.0 fl oz/100 gal	<p>USE RESTRICTIONS</p> <ul style="list-style-type: none"> • Make no more than 2 applications of BUPROFEZIN 40SC per year. <p>RECOMMENDATIONS</p> <ul style="list-style-type: none"> • Apply the specified dosage as a foliar spray in sufficient water for complete, uniform coverage, including stems and underside of leaves. Spray to the point of runoff. • Leafhopper, Planthopper, Whitefly: Make first application as soon as adult insects begin to appear. • Mealybug: Make first application as soon as insect activity is observed. • Scale: Make first application when crawlers are emerging. • If additional insecticide applications are required for control, use another class of chemistry or a different Insect Growth Regulator (IGR) with a different mode of action before making a second application of BUPROFEZIN 40SC. • Consult local or state agricultural authorities for details concerning economic thresholds for each target pest.
	Scales	21.5 fl oz/100 gal	
	Whiteflies	9.0 fl oz/100 gal	

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container, unopened, in a cool, dry place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. DO NOT reuse or refill this container. Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

9
15

IMPORTANT: READ BEFORE USE

By using this product, user or buyer accepts the following conditions, warranty, disclaimer of warranties, and limitations of liability.

CONDITIONS: The directions for use of this product are believed to be accurate and must be followed carefully. However, because of extreme weather and soil conditions, use methods and other factors beyond the control of Nichino America, Inc. (NAI), it is impossible for NAI to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. To the extent consistent with applicable law, all such risks are assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THERE ARE NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, WHICH EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of NAI is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, NAI disclaims any liability whatsoever for incidental or consequential damages, including, but not limited to, liability arising out of breach of contract, express or implied warranty (including warranties of merchantability and fitness for a particular purpose), tort, negligence, strict liability, or otherwise.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT THE ELECTION OF NICHINO AMERICA, THE REPLACEMENT OF PRODUCT.

©2009. Nichino America, Inc.

070809

15/10

SUPPLEMENTAL LABEL

NICHINO AMERICA, INC.

GROUP	16	INSECTICIDE
-------	----	-------------

COURIER® 40SC Insect Growth Regulator For Use on Cucurbit Crop Group 9

EPA Reg. No. 71711-20

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling and the EPA approved container label must be in the possession of the user at the time of application.

New crops appear on this supplemental labeling that do not appear on the Section 3 container label.

NOTICE: Before using this product, read the First Aid, Precautionary Statements, Conditions of Sale and Warranty, and complete Directions for Use found on the container labeling. All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.

Crop	Pest	Formulated Product/A	Directions for Use
Cucurbit Crop Group 9: Cucumbers Melons Cantaloupe Honeydew melons Muskmelons Watermelons Pumpkins Squash	Leafhoppers Planthoppers	13.6 fl oz	USE RESTRICTIONS <ul style="list-style-type: none"> ▪ Do not make more than 2 applications per crop cycle. ▪ Allow at least 7 days between applications to cucurbits. ▪ Do not make more than 4 applications per year. ▪ Do not apply within 10 days of harvest in California and 7 days outside California in melons, pumpkins and squash. ▪ Do not apply within 7 days of harvest in cucumbers.
	Whiteflies	9.0 to 13.6 fl oz	

USE RESTRICTIONS:

Crop Rotational Restrictions

CROP	PLANTBACK TIMING
All crops registered for use with buprofezin	0 days following application
Leafy vegetables not registered for use with buprofezin	30 days following application
Root and tuber vegetables	60 days following application
All other crops	365 days following application

©2009 Nichino America, Inc.
Courier® is a trademark of Nichino America, Inc.

070809

NICHINO AMERICA, INC.
4550 New Linden Hill Road, Suite 501
Wilmington, DE 19808
(302) 636-9001

11
15

SUPPLEMENTAL LABEL

NICHINO AMERICA, INC.

GROUP 16 INSECTICIDE

COURIER® 40SC Insect Growth Regulator For Use on Leafy Vegetable Crop Group 4

EPA Reg. No. 71711-20

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling and the EPA approved container label must be in the possession of the user at the time of application.

New crops appear on this supplemental labeling that do not appear on the Section 3 container label. This supplemental label supersedes any previously approved supplemental label for use on leafy vegetables.

NOTICE: Before using this product, read the First Aid, Precautionary Statements, Conditions of Sale and Warranty, and complete Directions for Use found on the container labeling. All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.

Crop	Pest	Formulated Product/A	Directions for Use
Leafy Vegetable Crop Group 4: Celery Endive Head lettuce Leaf lettuce Parsley Spinach	Leafhoppers Planthoppers Whiteflies	9.0 to 13.6 fl oz	USE RESTRICTIONS <ul style="list-style-type: none"> ▪ Do not make more than 2 applications per crop cycle. ▪ Allow at least 7 days between applications. ▪ Do not make more than 4 applications per year. ▪ Do not apply within 7 days of harvest.

USE RESTRICTIONS:

Crop Rotational Restrictions

CROP	PLANTBACK TIMING
All crops registered for use with buprofezin	0 days following application
Leafy vegetables not registered for use with buprofezin	30 days following application
Root and tuber vegetables	60 days following application
All other crops	365 days following application

©2009 Nichino America, Inc.
Courier® is a trademark of Nichino America, Inc.

070809

NICHINO AMERICA, INC.
4550 New Linden Hill Road, Suite 501
Wilmington, DE 19808
(302) 636-9001

12
15

SUPPLEMENTAL LABEL

NICHINO AMERICA, INC.

GROUP 16 INSECTICIDE

**COURIER® 40SC Insect Growth Regulator
For Use on Snap Beans**

EPA Reg. No. 71711-20

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling and the EPA approved container label must be in the possession of the user at the time of application.

New crops appear on this supplemental labeling that do not appear on the Section 3 container label.

NOTICE: Before using this product, read the First Aid, Precautionary Statements, Conditions of Sale and Warranty, and complete Directions for Use found on the container labeling. All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.

Crop	Pest	Formulated Product/A	Directions for Use
Snap Beans	Leafhoppers Planthoppers Whiteflies	9.0 to 13.6 fl oz	USE RESTRICTIONS <ul style="list-style-type: none"> ▪ Do not exceed 27.2 fl oz per crop cycle. ▪ Allow at least 14 days between applications. ▪ Do not apply within 14 days of harvest.

USE RESTRICTIONS:

Crop Rotational Restrictions

CROP	PLANTBACK TIMING
All crops registered for use with buprofezin	0 days following application
Leafy vegetables not registered for use with buprofezin	30 days following application
Root and tuber vegetables	60 days following application
All other crops	365 days following application

©2009 Nichino America, Inc.
Courier® is a trademark of Nichino America, Inc.

070809

NICHINO AMERICA, INC.
4550 New Linden Hill Road, Suite 501
Wilmington, DE 19808
(302) 636-9001

13
15

SUPPLEMENTAL LABEL

NICHINO AMERICA, INC.

GROUP 16 INSECTICIDE

COURIER® 40SC Insect Growth Regulator For Use on Head & Stem Brassica Subgroup 5A

EPA Reg. No. 71711-20

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling and the EPA approved container label must be in the possession of the user at the time of application.

New crops appear on this supplemental labeling that do not appear on the Section 3 container label.

NOTICE: Before using this product, read the First Aid, Precautionary Statements, Conditions of Sale and Warranty, and complete Directions for Use found on the container labeling. All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.

Crop	Pest	Formulated Product/A	Directions for Use
Head & Stem Brassica Subgroup 5A: Broccoli Brussel sprouts Cabbage Cauliflower Cavalo broccolo Chinese broccoli Chinese cabbage (Napa) Chinese mustard cabbage Kohlrabi	Leafhoppers Planthoppers Whiteflies	9.0 to 13.6 fl oz	USE RESTRICTIONS <ul style="list-style-type: none"> ▪ Do not make more than 2 applications per crop cycle. ▪ Do not exceed 27.2 fl oz per crop cycle. ▪ Allow at least 7 days between applications. ▪ Do not make more than 4 applications per year. ▪ Do not apply within 1 day of harvest.

USE RESTRICTIONS:

Crop Rotational Restrictions

CROP	PLANTBACK TIMING
All crops registered for use with buprofezin	0 days following application
Leafy vegetables not registered for use with buprofezin	30 days following application
Root and tuber vegetables	60 days following application
All other crops	365 days following application

©2009 Nichino America, Inc.
Courier® is a trademark of Nichino America, Inc.
070809

NICHINO AMERICA, INC.
4550 New Linden Hill Road, Suite 501
Wilmington, DE 19808
(302) 636-9001

14
15

SUPPLEMENTAL LABEL

NICHINO AMERICA, INC.

GROUP	16	INSECTICIDE
-------	----	-------------

COURIER® 40SC Insect Growth Regulator For Use on Low-Growing Berry Subgroup 13-07G

EPA Reg. No. 71711-20

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling and the EPA approved container label must be in the possession of the user at the time of application.

New crops appear on this supplemental labeling that do not appear on the Section 3 container label. This supplemental label supersedes any previously approved supplemental label for use on low-growing berry subgroup 13-07G.

NOTICE: Before using this product, read the First Aid, Precautionary Statements, Conditions of Sale and Warranty, and complete Directions for Use found on the container labeling. All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.

Crop	Pest	Formulated Product/A	Directions for Use
Low-Growing Berry Subgroup 13-07G: Strawberry Bearberry Bilberry Blueberry, lowbush Cloudberry Cranberry Ligonberry Muntries Partridgeberry	Leafhoppers Mealybugs Planthoppers Whiteflies	9.0 to 12.0 fl oz	USE RESTRICTIONS <ul style="list-style-type: none"> With ground application, use at least 80 gallons of water per acre. For aerial application, use at least 5 gallons of water per acre. Do not make more than 2 applications per crop cycle. Allow at least 10 days between applications. Do not apply within 3 days of harvest. RECOMMENDATIONS <ul style="list-style-type: none"> Good coverage is essential

USE RESTRICTIONS:

Crop Rotational Restrictions

CROP	PLANTBACK TIMING
All crops registered for use with buprofezin	0 days following application
Leafy vegetables not registered for use with buprofezin	30 days following application
Root and tuber vegetables	60 days following application
All other crops	365 days following application

©2009 Nichino America, Inc.
Courier® is a trademark of Nichino America, Inc.

070809

NICHINO AMERICA, INC.
 4550 New Linden Hill Road, Suite 501
 Wilmington, DE 19808
 (302) 636-9001

15
15

SUPPLEMENTAL LABEL

NICHINO AMERICA, INC.

GROUP	16	INSECTICIDE
-------	----	-------------

**COURIER® 40SC Insect Growth Regulator
For Use on Fruiting Vegetable Crop Group 8**

EPA Reg. No. 71711-20

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling and the EPA approved container label must be in the possession of the user at the time of application.

New crops appear on this supplemental labeling that do not appear on the Section 3 container label. This supplemental label supersedes any previously approved supplemental label for use on fruiting vegetables.

NOTICE: Before using this product, read the First Aid, Precautionary Statements, Conditions of Sale and Warranty, and complete Directions for Use found on the container labeling. All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.

Crop	Pest	Formulated Product/A	Directions for Use
Fruiting Vegetable Crop Group 8: Bell pepper Chili pepper Cooking pepper Pimento Eggplant Groundcherry Okra Pepino Sweet pepper Tomatillo Tomato	Leafhoppers Mealybugs Planthoppers	13.6 fl oz	USE RESTRICTIONS <ul style="list-style-type: none"> Apply no more than 2 applications per crop cycle. Allow at least 5 days between applications. Do not apply within 1 day of harvest. RECOMMENDATIONS <ul style="list-style-type: none"> Treatment should be applied when population level reaches economic threshold. Consult local and state agricultural authorities for details. Good coverage is essential. Orient nozzles to assure better coverage. Use of a higher volume of water will assure better coverage, especially under adverse conditions, such as hot, dry weather and/or a dense canopy. The entire field should be treated.
	Whiteflies	9.0 to 13.6 fl oz	

USE RESTRICTIONS:

Crop Rotational Restrictions

CROP	PLANTBACK TIMING
All crops registered for use with buprofezin	0 days following application
Leafy vegetables not registered for use with buprofezin	30 days following application
Root and tuber vegetables	60 days following application
All other crops	365 days following application

©2009 Nichino America, Inc.
Courier® is a trademark of Nichino America, Inc.

070809

NICHINO AMERICA, INC.
4550 New Linden Hill Road, Suite 501
Wilmington, DE 19808
(302) 636-9001