



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505T)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

Date of Issuance:

71711-72

1/15/26

NOTICE OF PESTICIDE:

Registration

Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

NAI-9071 SC Fungicide

Name and Address of Registrant (include ZIP Code):

Nichino America, Inc.
4550 Linden Hill Rd.
Wilmington, DE 19808

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

Continues page 2

Signature of Approving Official:

Alexandra Boukedes
Digitally signed by
Alexandra Boukedes
Date: 2026.01.15
11:17:05 -05'00'

Date:

1/15/26

for Nathan Mellor, Branch Chief
Fungicide-Herbicide Branch
Registration Division (7505P)

2. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 71711-72."
3. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The record for this product currently contains the following CSFs:

- Basic CSF dated 07/09/2024

If you have any questions, please contact Ernest Kraka at kraka.ernest@epa.gov.

Enclosure



FLUTOLANIL	GROUP	7	FUNGICIDE
THIOPHANATE-METHYL	GROUP	1	FUNGICIDE

NAI-9071 SC Fungicide

ACTIVE INGREDIENTS:

Flutolanil: Benzamide, N-[3-(1-methylethoxy)phenyl]-2-(trifluoromethyl)-..... **14.8%**

Thiophanate-methyl: (Dimethyl [1,2-phenylene)-bis(iminocarbonothioyl)]bis[carbamate])... **24.4%**

OTHER INGREDIENTS: **60.8%**

TOTAL **100.00%**

Contains 1.42 lbs. Flutolanil and 2.34 lbs. Thiophanate-methyl as active ingredients per U.S. gallon.

EPA Reg. No. 71711-XX

EPA Est. No.

[Alternate Brand Names: Annoint™ Fungicide]

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

If swallowed	<ul style="list-style-type: none">Call a poison control center or doctor immediately for treatment advice.Have person sip a glass of water if able to swallow.Do not induce vomiting unless told to by a poison control center or doctor.Do not give anything to an unconscious person.
If on skin	<ul style="list-style-type: none">Take off contaminated clothing.Rinse skin immediately with plenty of water for 15-20 minutes.Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none">Hold eye open and rinse slowly and gently with water for 15-20 minutes.Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-348-5832 for emergency medical treatment information. In case of fire or spills, information may be obtained by calling 1-800-424-9300.

See [attached, inside] booklet for First Aid, Precautionary Statements, and Directions for Use

NET CONTENTS: _____

[Manufactured in ____] [formulated in ____] [and] [packaged in ____] for

Nichino America Inc.

4550 Linden Hill Road, Suite 501

Wilmington, DE 19808

888-740-7700

A C C E P T E D

Jan 15, 2026

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No.

71711-72

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, loaders and applicators must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves, made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton® ≥ 14 mils.
- Chemical-resistant apron for mixers, loaders and other handlers exposed to the concentrate

Handlers using the dust collection system must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves, made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton® ≥ 14 mils (except pilots, groundboom applicators, and airblast applicators)

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **Do not** reuse them.

User Safety Requirements

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic organisms. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. **Do not** apply directly to water, areas where surface water is present, or to intertidal areas below the mean high water mark. **Do not** contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. **Do not** enter or allow worker entry into treated areas during the restricted-entry interval (REI). The REI is 12 hours except as listed in the application rate tables below.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water is:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, polyvinyl chloride (PVC) \geq 14 mils, or Viton® \geq 14 mils.
- Chemical resistant footwear plus socks
- Chemical resistant headgear for overhead exposure

USE INFORMATION

NAI-9071 SC Fungicide is a coformulation containing two fungicides effective for the suppression of black scurf and *Rhizoctonia* stem canker (*Rhizoctonia solani*), *Fusarium* dry rot (*Fusarium sambucinum*; *F. coeruleum*), and silver scurf (*Helminthosporium solani*) on potato seed-pieces. NAI-9071 SC Fungicide contains 14.8% flutolanil and 24.4% thiophanate-methyl in a suspension concentrate formulation. This pesticide will not control bacterial diseases or other diseases not claimed on this label for varieties known to be susceptible to bacterial soft rot. Companion fungicides known to be effective against this organism should be used. Consult your extension potato pathologist for variety and treatment information.

NAI-9071 SC Fungicide is to be used as an integral part of a potato disease management strategy. This strategy includes the use of high-quality certified seed and the incorporation of cultural practices such as: crop rotation, optimal harvest time to minimize disease development, and proper handling of tubers to minimize bruising and improve storage quality. Sound strategy should also include sanitation of equipment and storage areas to reduce disease inoculum and control of storage area temperature, humidity, and ventilation. Consult your local agricultural extension agent for more information on disease management practice.

ROTATIONAL CROP RESTRICTIONS

Crop/Crop Group	Plantback Timing
Brassica (cole) leafy vegetables (crop group 5) Cotton Peanuts Potatoes Rice Soybeans Turnip greens	0 days following application
Wheat	30 days following application
Leafy vegetables (such as lettuce, spinach, or celery) Small grain crops, other than wheat (such as barley, rye, or oats)	150 days following application
Corn (such as field, sweet, or popcorn) Sorghum	240 days following application
All other crops	365 days following application

RESISTANCE MANAGEMENT

For resistance management, please note, the flutolanil component of NAI-9071 SC Fungicide belongs to the succinate dehydrogenase inhibitor class (FRAC Group 7). The thiophanate-methyl component of NAI-9071 SC Fungicide belongs to the thiabendazole and relatives (electrophiles) class (FRAC Group 1). Any fungal population may contain individuals naturally resistant to NAI-9071 SC Fungicide and other Group 7 or Group 1 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Nichino America, Inc. encourages responsible product stewardship to ensure effective long-term control of the fungal diseases on this label. Appropriate resistance-management strategies should be followed.

Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.

MIXING DIRECTIONS

NAI-9071 SC Fungicide Alone: Fill spray tank with $\frac{3}{4}$ of the amount of water needed for the intended application and then turn on agitation. Pour specified amount of product on the surface of the water in the spray tank. Add the balance of the water to the spray tank with agitation running. Keep agitation running during filling and spraying operations. If spraying must be stopped before emptying the sprayer, resume agitation before spraying the remainder of the load. Mix only as much spray solution as can be sprayed within four hours. Storage and use of the previous day's spray mix may result in reduced activity.

NAI-9071 SC Fungicide in Tank Mixtures: It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Begin with clean equipment. Fill spray tank with $\frac{3}{4}$ of the amount of water needed for the intended application and turn on agitation. If using a buffering agent, add after filling the tank with $\frac{3}{4}$ amount of water. **Do not** use oil as carrier or include other additives in the finished spray. Add the recommended amount of tankmix products in the following order while maintaining agitation:

- 1) products in water-soluble packets
- 2) wettable powders
- 3) water-dispersible granulars and/or soluble powders
- 4) flowable liquids (including NAI-9071 SC Fungicide)
- 5) emulsifiable concentrates
- 6) adjuvants and/or oils
- 7) remaining amount of water to achieve the desired level

Always follow the labeled mixing instructions of any partner products. Keep agitation running during filling and spraying operations. If spraying must be stopped before emptying the sprayer, resume agitation before spraying the remainder of the load. Mix only as much spray solution as can be sprayed within four hours. Storage and use of the previous day's spray mix may result in reduced activity.

APPLICATION RESTRICTIONS

Seeds/seed-pieces that have been treated with this product that are then packaged or bagged for future use must contain the following labeling on the outside of the seed/seed-piece package or bag:

- Treated Seed/Seed Pieces - **Do Not** Use for Food, Feed, or Oil Purposes. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting.
- After the seeds/seed-pieces have been planted, **do not** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Exception: Once the seeds/seed-pieces are planted in soil or other planting media, the Worker Protection Standard allows workers to enter the treated area without restriction if there will be no worker contact with the soil/media subsurface.

NAI-9071 SC FUNGICIDE APPLICATION RATE CHART

Potato (seed-piece treatment only)		
Disease	Rate	Directions for Use
Seed-borne black scurf*	1.0 fl oz (0.011 lb Flutolanil and 0.018 lb Thiophanate-methyl) /100 lbs of cut seed- pieces	• Apply to the seed-pieces immediately after cutting. Use dispensing equipment that will give thorough coverage to the cut seed-piece surfaces.
Seed-borne Fusarium dry rot (<i>Fusarium sambucinum</i> ; <i>Fusarium coeruleum</i>)*		• Thoroughly clean and sanitize cutting equipment including knives, belts, rollers, tables, trucks, and planting equipment before planting seed-pieces.
Seed-borne Rhizoctonia stem canker (<i>Rhizoctonia solani</i>)*		
Seed-borne silver scurf (<i>Helminthosporium solani</i>)*		
*suppression		

SEED BAG LABEL REQUIREMENTS

- This seed has been treated with NAI-9071 SC Fungicide, a fungicide containing Flutolanil and Thiophanate-methyl.
- **Do not** use treated seed for feed, food, or oil purposes.

The U.S. Environmental Protection Agency requires the following statements on containers containing seed treated with NAI-9071 SC Fungicide:

- Store treated seed away from food and feedstuffs.
- **Do not** allow children, pets, or livestock to have access to treated seeds.
- Wear long pants, long-sleeved shirt and protective gloves when handling treated seed.
- Treated seeds exposed on the soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting (such as in row ends).
- Dispose of all excess treated seed by burying seed away from bodies of water.
- **Do not** contaminate bodies of water when disposing of planting equipment wash water.
- Dispose of seed packaging or containers in accordance with local requirements.

- Excess treated seed may be used for ethanol production if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.

Note: This product does not contain dye and is not covered by an appropriate tolerance, tolerance exemption, or other clearance under the Federal Food, Drug and Cosmetic Act. To comply with 40 CFR 153.155, therefore, all seed treated commercially with this product must be colored with an EPA-approved dye or colorant of a suitable color to prevent accidental use as food for man or feed for animals.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container, and keep tightly closed when not in use. Store in a cool, dry place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. **Do not** reuse or refill this container. Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill or by incineration, or other methods allowed by state and local authorities.

IMPORTANT: READ BEFORE USE

By using this product, user or buyer accepts the following conditions, warranty, disclaimer of warranties, and limitations of liability.

CONDITIONS: The directions for use of this product are believed to be accurate and must be followed carefully. However, because of extreme weather and soil conditions, use methods, and other factors beyond the control of Nichino America, Inc. (NAI), it is impossible for NAI to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. To the extent consistent with applicable law, all such risks are assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THERE ARE NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, WHICH EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of NAI is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, NAI disclaims any liability whatsoever for incidental or consequential damages, including, but not limited to, liability arising out of breach of contract, express or implied warranty (including warranties of merchantability and fitness for a particular purpose), tort, negligence, strict liability, or otherwise.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES, OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT THE ELECTION OF NICHINO AMERICA, INC., THE REPLACEMENT OF PRODUCT.

©2025 Nichino America, Inc. Annoint is a trademark of Nichino America, Inc. Viton is a registered trademark of The Chemours Company,

APPENDIX

Text that may be included on the container label or promotional information supporting this product: Potato Seed-Piece Treatment for Suppression of Seedling Diseases