

71908-1

11/25/2008

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

1/79

NOV 25 2008

E.I.D. Parry India, Ltd.
 c/o Ms. Karen Pither
 Pither Consulting
 5813 N. Kensington Street
 Kansas City, MO 64119

Subject: Company Name: E.I.D. Parry India, Ltd.
 Product Name: Neemzal T/S1 2%EC Insecticide
 EPA Registration No. 71908-1
 Label Amendment to add new package designs (water soluble packaging) under the new product name "Eco Garden Botanical Insecticide, Miticide and Nematicide."
 Application Dated 12/22/07


Dear Ms. Pither

The amendment referred to above, submitted in connection with registration under FIFRA section 3(c)(5), is acceptable provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Submit five (5) copies of your final printed labeling before you release the product for shipment. Final printed labeling means the label or labeling of the product when distributed or sold. Clearly legible reproductions or photo reductions will be accepted for unusual labels, such as those silk-screened directly onto glass or metal containers or large bags or drum labels.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions. If you have any questions contact Andrew Bryceland, of my staff at (703)305-6928 or by e-mail at: bryceland.andrew@epa.gov. A stamped copy of the label is enclosed for your records.

Sincerely,



Linda A. Hollis, Chief
 Biochemical Pesticides Branch
 Biopesticides and Pollution

CONCURRENCE: Prevention Division (7511P)

SYMBOL	►	Enclosure	<i>Bryceland</i>					
SURNAME	►		<i>11/25/08</i>					
DATE	►							

ACCEPTED

NOV 25 2000

MASTER LABEL

Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended, for
the pesticide registered under
EPA Reg. No. 71908-1

2/79

NeemAzal® T/S 1.2 EC

ACTIVE INGREDIENT:

Azadirachtin	% By Wt.
..... 1.2%	
OTHER INGREDIENTS	98.8%
..... TOTAL 100.00%	

Contains 0.0987 lb. azadirachtin per gallon.

KEEP OUT OF REACH OF CHILDREN
CAUTION

Sublabel 1: NeemAzal® T/S 1.2 EC BOTANICAL - AZADIRACHTIN-BASED BOTANICAL, ANTIPEST, REPELANT AND INSECT GROWTH REGULATOR, FOR USE ON GREENHOUSE AND OUTDOOR FOOD CROPS, ORNAMENTAL FLOWERS, TREES, SHRUBS AND PLANTS.

Sublabel 2: NeemAzal® T/S 1.2 EC BOTANICAL INSECTICIDE, REPELANT, ANTIPEST AND INSECT GROWTH REGULATOR (IGR) BOTANICAL PRODUCT FOR CONTROL OF INSECTS ON INDOOR AND OUTDOOR TREES, SHRUBS, FLOWERS, FRUIT AND NUT TREES, GARDEN VEGETABLES AND PLANTS. INDOOR AND OUTDOOR VEGETABLES, ORNAMENTAL FLOWERS, TREES, TURFGRASS, SHRUBS AND PLANTS, INCLUDING PLANTS GROWN IN CONTAINERS, INTERIORSCAPES, HOME AND GARDEN USE.

Sublabel 3: AZAMAX™ (Alternate Product Name) BOTANICAL INSECTICIDE, MITICIDE AND NEMATICIDE, REPELANT, ANTIPEST AND INSECT GROWTH REGULATOR (IGR) FOR CONTROL OF INSECTS ON FRUIT AND NUT TREES, INDOOR AND OUTDOOR VEGETABLES, ORNAMENTAL FLOWERS, TREES, SHRUBS AND PLANTS INCLUDING PLANTS GROWN IN CONTAINERS, INTERIORSCAPES AND GARDEN USE.

Sublabel 4: ECOGARDEN™ (Alternate Product Name) BOTANICAL INSECTICIDE, MITICIDE AND NEMATICIDE REPELANT, ANTIPEST AND INSECT GROWTH REGULATOR (IGR) FOR CONTROL OF INSECTS ON FRUIT AND NUT TREES, INDOOR AND OUTDOOR VEGETABLES, ORNAMENTAL FLOWERS, TREES, SHRUBS AND PLANTS INCLUDING PLANTS GROWN IN CONTAINERS, INTERIORSCAPES AND GARDEN USE.

Sublabel 5: ECOGARDEN™ (Alternate Product Name) BOTANICAL INSECTICIDE, MITICIDE AND NEMATICIDE SINGLE DOSE DELIVERY CARTRIDGE FOR CONTROL OF INSECTS ON FRUIT AND NUT TREES, INDOOR AND OUTDOOR VEGETABLES, ORNAMENTAL FLOWERS, TREES, SHRUBS AND PLANTS INCLUDING PLANTS GROWN IN CONTAINERS, INTERIORSCAPES AND GARDEN USE.

Sublabel 6: ECOGARDEN™ (Alternate Product Name) BOTANICAL INSECTICIDE, MITICIDE AND NEMATICIDE SINGLE DOSE DELIVERY SACHET FOR CONTROL OF INSECTS ON FRUIT AND NUT TREES, INDOOR AND OUTDOOR VEGETABLES, ORNAMENTAL FLOWERS, TREES, SHRUBS AND PLANTS INCLUDING PLANTS GROWN IN CONTAINERS, INTERIORSCAPES AND GARDEN USE.

Sublabel 7: ECOGARDEN™ (Alternate Product Name) BOTANICAL INSECTICIDE, MITICIDE AND NEMATICIDE SINGLE DOSE DELIVERY VIAL FOR CONTROL OF INSECTS ON FRUIT AND NUT TREES, INDOOR AND OUTDOOR VEGETABLES, ORNAMENTAL FLOWERS, TREES, SHRUBS AND PLANTS INCLUDING PLANTS GROWN IN CONTAINERS, INTERIORSCAPES AND GARDEN USE.

Sublabel 8: ECOGARDEN™ (Alternate Product Name) BOTANICAL INSECTICIDE, MITICIDE AND NEMATICIDE WATER SOLUBLE SACHET FOR CONTROL OF INSECTS ON FRUIT AND NUT TREES, INDOOR AND OUTDOOR VEGETABLES, ORNAMENTAL FLOWERS, TREES, SHRUBS AND PLANTS INCLUDING PLANTS GROWN IN CONTAINERS, INTERIORSCAPES AND GARDEN USE.

Sublabel 9: ECOGARDEN™ (Alternate Product Name) BOTANICAL INSECTICIDE, MITICIDE AND NEMATICIDE WATER SOLUBLE CAPSULE FOR CONTROL OF INSECTS ON FRUIT AND NUT TREES, INDOOR AND OUTDOOR VEGETABLES, ORNAMENTAL FLOWERS, TREES, SHRUBS AND PLANTS INCLUDING PLANTS GROWN IN CONTAINERS, INTERIORSCAPES AND GARDEN USE.

See Directions for Use for a Complete List of Insects Controlled.

NET CONTENTS _____ GALLONS

EPA Reg. No. 71908-1

EPA Est. No. 71908-IND-001
USA Phone Number: (916) 973-1351

Label Revision Dated 7-1-08

EID PARRY (INDIA) LIMITED
Bioproducts Division
234 NSC Bose Road
Chennai (Madras) 600 001 India

Sublabel 1

NeemAzal[®] T/S 1.2 EC

AZADIRACHTIN-BASED BOTANICAL, ANTI FEEDANT, REPELLANT AND INSECT GROWTH REGULATOR

FOR USE ON GREENHOUSE AND OUTDOOR FOOD CROPS,
ORNAMENTAL FLOWERS, TREES, SHRUBS AND PLANTS

FOR ORGANIC PRODUCTION

ACTIVE INGREDIENT:

Azadirachtin	% By Wt.
.....	1.2%
OTHER INGREDIENTS	98.8%
.....	TOTAL 100.00%

Contains 0.0987 lb. (44.8 g) azadirachtin per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call poison control center or doctor for further treatment advice.
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.
If swallowed	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

CALL THE POISON CONTROL HOTLINE 24 HOURS A DAY AT 1-888-478-0798

READ ALL DIRECTIONS BEFORE USING THIS PRODUCT

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution: Harmful if swallowed, absorbed through skin or if inhaled. Causes moderate eye irritation. Avoid breathing vapor. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash before reuse. Wear chemical resistant gloves.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long sleeved shirt and long pants
- Chemical resistant gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates.

For Terrestrial Uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwater or rinsate.

USER SAFETY RECOMMENDATIONS

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

NET CONTENTS _____ GALLONS

EID PARRY (INDIA) LIMITED
Bioproducts Division
234 NSC Bose Road
Chennai (Madras) 600 001 India

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the state or tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS).

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical resistant gloves
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in the box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

PRODUCT MODE OF ACTION

NEEMAZAL® T/S 1.2 EC controls target pests on contact or by ingestion. The product acts on pests by way of repellence, antifeedance and interference with the molting process. The buyer or user is reminded that the degree of efficacy of the product is largely dependent on weather conditions, intensity of pest population, area of application, type of pest, and physical stages of pests and crops.

GENERAL INFORMATION

Read all directions before using this product.

Apply **NeemAzal® T/S 1.2 EC** as directed to any food or non-food crop up to and including the day of harvest, at a maximum rate of 3½ pints/Acre. Refer to the Use Site Section for a complete listing of crops.

MIXING

Shake well before using. Add required amount of **NeemAzal® T/S 1.2 EC** to a clean spray tank with at least one-half of the water to be sprayed. Constant agitation is required, particularly with tank mixes. Agitate the mixture thoroughly and then fill the tank with remaining water and continue agitation. Thorough mixing is necessary for uniform coverage. Non-uniform mixing can cause crop injury or can result in lowered effectiveness. For tank mixes, add other components to the tank containing the **NeemAzal® T/S 1.2 EC** spray mixture and agitate thoroughly. If tank mixture is allowed to sit, agitation is necessary prior to application. Adjusting the spray mixture pH between 5.5 and 7 will provide optimal performance. Always use this product promptly after mixing with water and do not let tank mix sit for any extended period.

COMPATIBILITY: **NeemAzal® T/S 1.2 EC** has been found to be compatible with most commonly used pesticides and fertilizers. To avoid problems, conduct a compatibility test before using this product in a tank mix with other pesticides or with fertilizers. To test for compatibility, mix a small amount of each product, in the appropriate proportions, in a small jar.

PHYTOTOXICITY: **NeemAzal® T/S 1.2 EC** has been evaluated for phytotoxicity on a wide range of crops and ornamentals. However, since testing on all varieties of all crops and ornamentals is not feasible, testing a small portion of the area to be treated for phytotoxicity is recommended before treating the entire area. Further, all possible combinations or sequences of pesticide sprays, including other fertilizers, surfactants, adjuvants and other pesticides, have not been tested, thus testing for phytotoxicity of spray mixtures is recommended. It is further recommended that spray equipment used to apply

NeemAzal® T/S 1.2 EC be thoroughly cleaned before use. The addition of spray adjuvants enhances control in some crops under ideal conditions. Addition of certain adjuvants may cause phytotoxicity therefore, the addition of crop oils and other adjuvants should be thoroughly tested before using. The addition of crop oils to spray mixtures is NOT recommended on ornamental crops. Captain, Bordeaux mixtures, and highly alkaline products cause unacceptable phytotoxicity and/or reduced effectiveness on target pests. Avoid tank mix combinations of **NeemAzal® T/S 1.2 EC** plus compounds known to be incompatible with oil-based formulations to prevent phytotoxicity. "Waxy bloom" on certain crops and ornamental plants is reduced after a **NeemAzal® T/S 1.2 EC** application.

APPLICATION INSTRUCTIONS

For optimal performance spray product as soon as possible when pests are expected or when pests first appear. For foliar applications, apply **NeemAzal® T/S 1.2 EC** in sufficient spray volume and with adequate spray pressure to ensure complete and thorough coverage of all plant surfaces including both the top and bottom of leaves. Avoid excessive runoff. Best results are obtained following 2-3 applications made at 7-10 day intervals. When pest pressure is heavy or plant canopy is dense, use higher rates and increase spray frequency. Spraying in the morning or evening hours will provide the best results. Repeat application if rain occurs within two to three hours of spraying.

SPRAY DIRECTIONS

Apply **NeemAzal® T/S 1.2 EC** as a foliar spray or a drench to soil or non-soil media to control insects. When needed, soil drenches can also be used to control soil-borne pests, including soil-borne larvae of foliar insect pests. When applying as a drench, avoid excessive leaching. **NeemAzal® T/S 1.2 EC** can also be applied through sub-surface soil treatment equipment. Always follow equipment manufacturer's use directions. **NeemAzal® T/S 1.2 EC** may be applied using any powered or manual pesticide application equipment which includes, but is not restricted to, high volume, low volume, ultra-low volume, electrostatic, fogging and chemigation. Follow the original manufacturer's instructions when using these types of equipment.

DRENCH AND ORNAMENTAL SPRAY DIRECTIONS FOR LABELED PLANTS GROWN IN GREENHOUSES, SHADECLOTHS AND NURSERIES

When used as a soil drench, apply one pint of finished spray for each gallon of soil in the pot. For most pests apply 18-21 oz. **NeemAzal® T/S 1.2 EC**

per 100 gallons of water. For treatment of harder to control pests, such as Dipteran leafminers, use up to 27 ounces per 100 gallons of water. Do not exceed 57 oz of **NeemAzal® T/S 1.2 EC** per acre per application.

RATES

Use **NeemAzal® T/S 1.2 EC** at 1-2 pints/acre for most pest and crop conditions. Under extremely heavy pest pressure up to 3 1/2 pints may be used. Do not use less than 5 oz. per acre of **NeemAzal® T/S 1.2 EC** alone. When tank mixed with other insecticidal products, the rate of **NeemAzal® T/S 1.2 EC** may be reduced by 1/2, but not less than 4 oz per acre. Use up to 2.6 oz/1000 square feet for manure and compost treatments.

GENERAL CHEMIGATION INSTRUCTIONS

Apply this product only through low pressure, drip (trickle) or sprinkler (center pivot, lateral move, end tow, side (wheel) roll, traveler, solid set or hand move) irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers, or other experts. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Dilute NeemAzal® T/S 1.2 EC with water before introduction into the system. Use the diluted mixture within 8 hours. Do not apply in irrigation water if the pH exceeds 7.0. The optimum pH range for application is 5.5 - 6.5. If needed, the pH of the irrigation water can be adjusted by use of a suitable buffering agent. Agitation is necessary. Apply at the rate indicated in APPLICATION INSTRUCTIONS using sufficient water to achieve an even distribution.

For Chemigation Systems

Connected to Public Water Systems

- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection.
- The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favors drift beyond the area intended for treatment.

In addition, all directions and requirements specified for Sprinkler Irrigation Systems must be followed.

Sprinkler Irrigation Systems

The system must contain a functional check valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Center pivot, motorized lateral move, or traveling gun types of equipment: Inject into the system for one revolution or run. Shut off injection equipment after one revolution or run, but continue to operate irrigation system until **NeemAzal® T/S 1.2 EC** has been cleared from the last sprinkler head. Do not use end guns. The system should be run at maximum speed for a foliar application.

Wheel move, side roll, end tow, solid set, or hand move types of equipment: Adjust equipment to inject **NeemAzal® T/S 1.2 EC** over a 30-60 minute period. Shut off injection equipment. Continue to operate irrigation system until **NeemAzal® T/S 1.2 EC** has been cleared from the last sprinkler head. **NeemAzal® T/S 1.2 EC** can be injected at the end of the irrigation cycle or as a separate application. Do not use end guns.

NeemAzal® T/S 1.2 EC must be premixed in a supply tank with water and other appropriate tank-mix chemicals. Agitation is necessary at all times.

Attention must be exercised in irrigation waters with a pH greater than 7. If the irrigation cycle will last longer than 8 hours and the NeemAzal® T/S 1.2 EC is premixed in the supply tank, the tank mix must be buffered to a pH of 6 or lower. Please contact your Company sales representative should this situation apply. Application is to be made in sufficient water and of sufficient duration to apply the appropriate rate evenly over the entire treated area.

No field runoff can be permitted during chemigation.

USE SITES

AGRICULTURAL USE SITES – Use **NeemAzal® T/S 1.2 EC** on agricultural use sites including, but not limited to, the following:

BERRIES GROUP, such as: Blackberry, Blueberry, Currant, Elderberry, Gooseberry, Huckleberry, Loganberry, Raspberry (black and red) Note: For Strawberries – see miscellaneous.

BULB VEGETABLES, such as: Garlic, Leek, Onion (dry bulb, green and Welch), Shallot

CEREAL GRAINS and GRAINS GROUP, such as: Barley, Buckwheat, Corn, Millet (pearl and Proso), Oats, Popcorn, Rice, Rye, Sorghum (milo), Teosinte, Triticale, Wheat, Wild rice

CITRUS FRUITS, such as: Calamondin, Citrus citron, Citrus hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Orange (sour and sweet), Pummelo, Satsuma mandarin, White Sapote, Uniq Fruit

COTTON AND TOBACCO

CUCURBIT VEGETABLES, such as: Chayote, Chinese waxgourd, Citron melon, Cucumber, Gherkin, Gourd (edible), Muskmelon, Pumpkin, Squash (summer and winter), Watermelon

FORAGE CROPS, including but not limited to: Alfalfa, Clover, Trevoil or Vetch.

FRUITING VEGETABLES, such as: Eggplant, Groundcherry, Pepino, Pepper (including bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), Tomatillo, Tomato

HERBS AND SPICES GROUP, such as: Allspice, Angelica, Anise (anise seed and star), Annatto (seed), Balm (lemon balm), Basil, Borage, Burnet, Camomile, Caper buds, Caraway, Caraway (black), Cardamom, Cassia bark, Cassia buds, Catnip, Celery seed, Chervil (dried), Chive, Chinese Chive, Cinnamon, Clary, Clove buds, Coriander (cilantro or Chinese parsley – leaf), Coriander (cilantro-seed), Costmary, Culantro (leaf and seed), Cumin, Curry (leaf), Dill (dillweed and seed), Fennel (common, Florence), Fenugreek, Grains of paradise, Horehound, Hyssop, Juniper berry, Lavender, Lemongrass, Lovage (leaf and seed), Mace, Marigold, Marjoram, Mustard (seed), Nasturtium, Nutmeg, Parsley (dried), Pennyroyal, Pepper (black and white), Poppy (seed), Rosemary, Rue, Saffron, Sage, Savory (summer and winter), Sweet bay (bay leaf), Tansy, Tarragon, Thyme, Vanilla, Wintergreen, Woodruff, Wormwood

LEGUME VEGETABLES (Succulent or Dried), such as: Bean, Broad Bean, Chickpea, Guar, Jackbean, Lablab bean, Lentil, Pea, Pigeon Pea, Soybean, Sword bean

LEAFY AND BRASSICA (COLE) VEGETABLES, such as: Amaranth, Arugula, Broccoli, Broccoli raab (rapini), Brussels Sprouts, Cabbage, Cauliflower, Cardoon, Cavallo broccolo, Celery, Chinese Broccoli (gai lon), Chinese Cabbage (bok choy, Napa), Chinese mustard Cabbage (gai choy), Chinese Celery, Celtsce, Chervil, Chrysanthemum (edible-leaved, Garland), Collards, Corn salad, Cress (garden, upland), Dandelion, Dock (sorrel), Endive (escarole), Fennel (florence), Kale, Kohlrabi, Lettuce (head and leaf), Mizuna, Mustard Greens, Mustard Spinach, Orach, Parsley, Purslane (garden, winter), Radicchio (red chicory), Rape Greens, Rhubarb, Spinach, Spinach (New Zealand, vine), Swiss Chard, Turnip Greens

MISCELLANEOUS, such as: Asparagus, Avocado, Banana, Coffee, Cocoa, Cranberry, Fig, Globe artichoke, Grape, Hops, Kiwifruit, Mango, Mushroom, Okra, Olives, Papaya, Pawpaw, Peanut, Persimmon, Pineapple, Pomegranate, Strawberry, Tea, Water chestnut, Watercress, and all other food crops

POME FRUITS GROUP, such as: Apple, Crabapple, Loquat, Mayhaw, Quince, Oriental Pear, or Pear (Comice varieties: DO NOT apply more than 24 fl oz/A. DO NOT apply after pink stage of flowering; test small areas of other varieties of pears for plant safety prior to full scale useage.)

ROOT AND TUBER VEGETABLE GROUP, such as: Arracacha, Arrowroot, Artichoke (Jerusalem, Chinese), Beet (garden, sugar), Burdock (edible), Canna (edible), Carrot, Cassava (bitter and sweet), Celeriac (celery root), Chayote (root), Chervil, (turnip-rooted), Chicory, Chufa, Dasheen (taro), Ginger, Ginseng, Horseradish, Leren, Oriental Radish (daikon), Parsley (turnip-rooted), Parsnip, Potato, Radish, Rutabaga, Salsify (oyster plant, black, Spanish), Skirret, Sweet potato, Tanier, Turmeric, Turnip, Yam bean (jicama, manioc pea), Yam (true)

STONE FRUIT GROUP, such as: Apricot, Cherry (sweet and tart), Nectarine, Peach, Plum (Chickasaw, Damson, Japanese), Plumcot, Prune

TREE AND NUT GROUP, such as: Almond, Beech nut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (hazelnut, Hickory nut, Macadamia nut (bush nut), Pecan, Walnut (black and English), Pistachios

TROPICAL FRUITS, such as: Papaya, Black Sapote, Canistel, Mamey Sapote, Mango, Sapodilla, Star Apple, Guava, Feijoa, Jaboticaba, Wax Jambu, Star Fruit, Passion Fruit, Acerola, Lychee, Longan, Spanish Lime, Rambutan, Pulasan, Sugar Apple, Atemoya, Custard Apple, Cherimoya, Ilama, Soursop, and Biriba.

ORNAMENTAL-USE SITES – **NeemAzal® T/S 1.2 EC** may be used on Ornamental Use sites including, but not limited to, the following:

ORNAMENTAL SHRUBS AND PLANTS, such as: Amaranthus, Aster, Azalea, Ferns, Fuschia, Caladium, Carnation, Chrysanthemum, Dahlia, Daisy, Lilies, Ivy, Ficus, Gardenia, Impatiens, Iris, Jasmine, Lilac, Marigold, Philodendron, Poinsettia, Rose, Zinnia

ORNAMENTAL TREES, such as: Ash, Birch, Cedar, Cypress, Dogwood, Fir, Elm, Juniper, Maple, Oak, Pine, Spruce

CHRISTMAS TREES AND CHRISTMAS TREE PLANTATIONS

NON-CROP USE SITES – Use **NeemAzal® T/S 1.2 EC** on non-crop use sites including, but not limited to, the following:

UNCULTIVATED AGRICULTURAL AREAS, such as: farm yards, fuel storage areas, fence rows, rights-of-way, fallow land; soil bank land, barrier strips.

GENERAL SOIL TREATMENTS, such as: Manure, Composts, Cull piles, Mulches, soil application with no mention of crops to be grown (potting soil, tops soil)

PESTS

NeemAzal® T/S 1.2 EC may be used against the following pests:

Aphids (such as pea aphid, Rosy Apple Aphid), Beetles (such as Japanese beetle), Borers, (such as peachtree borers, peach twig borers), True Bugs, (such as Lygus bugs, stink bugs), Caterpillars, (such as leafrollers, cutworms, loopers, armyworms), Flies (such as walnut husk fly, leafminers and fungus gnats), Leafhoppers, Leafminers, Whiteflies, Mealy Bugs, Mites, Psyllids (such as pear psylla), Weevils, Scales (such as San Jose scale), Thrips, (such as western flower thrips).

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal

Pesticide Storage:	Store in original container in a dry, cool, well ventilated area.
Pesticide Disposal:	Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.
Container Disposal:	Nonrefillable container. Do not reuse or refill this container. Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill container $\frac{1}{4}$ full with water and recap. Shake 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Drain for 10 seconds after flow begins to drip. Repeat this procedure two more times.

NOTICE ON CONDITIONS OF SALE

The directions for use of this product are based upon tests believed to be reliable, and the directions for use must be followed. To the extent applicable by law the use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. To the extent applicable by law, the buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

Date Revised: 11/17/08

Sublabel 2

NeemAzal[®] T/S 1.2 EC

BOTANICAL

BOTANICAL INSECTICIDE, REPELLANT, ANTI FEEDANT AND INSECT GROWTH REGULATOR (IGR)

- BOTANICAL PRODUCT FOR CONTROL OF INSECTS ON INDOOR AND OUTDOOR TREES, SHRUBS, FLOWERS, FRUIT AND NUT TREES, GARDEN VEGETABLES AND PLANTS.
- INDOOR AND OUTDOOR VEGETABLES, ORNAMENTAL FLOWERS, TREES, TURFGRASS, SHRUBS AND PLANTS, INCLUDING PLANTS GROWN IN CONTAINERS, INTERIORSCAPES, HOME AND GARDEN USE.

ACTIVE INGREDIENT:

	% By Wt.
Azadirachtin	1.2%
OTHER INGREDIENTS	98.8%
	TOTAL 100.00%

Contains 0.0987 lb. (44.8 g) azadirachtin per gallon.

FOR ORGANIC PRODUCTION

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call poison control center or doctor for further treatment advice.
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.
If swallowed	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. CALL THE POISON CONTROL HOTLINE 24 HOURS A DAY AT 1-888-478-0798	

NET CONTENTS _1 qt, 1 or 2.5_ GALLONS
Batch Number: xxxxx

EPA Reg. No. 71908-1

EPA Est. No. 71908-IND-001
USA Phone Number: (916) 973-1351

EID PARRY (INDIA) LIMITED
Bioproducts Division
234 NSC Bose Road
Chennai (Madras) 600 001 India

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin or if inhaled. Avoid breathing vapor. Causes moderate eye irritation. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Wear chemical resistant gloves.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long sleeved shirt and long pants
- Chemical resistant gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates.

For Terrestrial Uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwater or rinsate.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170.

This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS).

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is

- Coveralls
- Chemical resistant gloves
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in the box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

This is an end use product. E.I.D. Parry (India) Ltd. does not intend that this product be reformulated or repackaged except under a toll repackaging agreement.

PRODUCT DESCRIPTION:

NeemAzal® T/S 1.2 EC Insecticide is a botanical product for control of insects on indoor and outdoor plants including ornamental trees, shrubs, flowers, garden vegetables, turfgrass, fruit trees and nut trees.

When used as a component of an Integrated Pest Management (IPM) program, **NeemAzal® T/S 1.2 EC** Insecticide provides an effective resistance management tool.

MODE OF ACTION:

NeemAzal® T/S 1.2 EC Insecticide controls target pests on contact or by ingestion. The product acts on pests by way of repellence, anti-feedance, and interference with the molting process.

Azadirachtin, an insect growth regulator (IGR), mimics the pests' hormones and disrupts distinct stages of growth and development of insects and mites. The primary mode of action of azadirachtin is an interference with synthesis and metabolism of ecdysone and the juvenile hormone. Ecdysone is the molting hormone of insects, and azadirachtin can regulate growth leading to death before or during molting.

INDOOR AND OUTDOOR ORNAMENTAL TREES, SHRUBS, FLOWERS, AND PLANTS ESTABLISHED IN RESIDENTIAL, LANDSCAPE PLANTINGS AROUND INSTITUTIONAL, PUBLIC, COMMERCIAL AND INDUSTRIAL BUILDINGS, PARKS, RECREATIONAL AREAS, GREENHOUSES, SHADECLOTHS, NURSERIES, AND ATHLETIC FIELDS.

NeemAzal® T/S 1.2 EC Insecticide has been evaluated for phytotoxicity on a wide range of ornamentals and crops. However, since testing on all plant varieties is not feasible, test a small portion of the area to be treated for phytotoxicity before treating the entire area. All possible combinations or sequences of pesticide sprays, including other fertilizers, surfactants, adjuvants and other pesticides have not been tested. Thus, testing for phytotoxicity of spray mixtures is recommended.

The professional user assumes the responsibility for determining the level of tolerance of treated plants to **NeemAzal® T/S 1.2 EC** Insecticide when applied alone or in tank-mix combinations under commercial growing conditions.

Waxy bloom on certain ornamental plants is reduced after a **NeemAzal® T/S 1.2 EC** Insecticide application.

Applications of **NeemAzal® T/S 1.2 EC** will remove the glaucous 'blue' coloring from evergreens such as Colorado blue spruce and Koster spruce.

Use **NeemAzal® T/S 1.2 EC** Insecticide on the following plants:

Ornamental Plants and Flowers including but not limited to:	Actinopteris, African violets*, ageratum, aglaonema, Algerian ivy, allamanda, alocasia, amaranthus, anthurium, aphelandra, arborvitae, Artemisia, aster, aucuba ilex, azalea, baby's breath, begonia, Boston fern, bougainvillea, boxwood, brachycome, cacti, calabrese, caladium, calathea, calendula, calla, camellia, carnation, ceanothus, chrysanthemum, cineraria, coleus, columbine, cotoneaster, cyclamen, daffodil, dahlia, daisy, daylily, delphinium, dianthus, dieffenbachia, dogwood, dusty miller, Easter lily, English ivy, euphorbia, fern, ficus, foliage plants, foxglove, freesia, fuchsia, gaillardia, gardenia, geranium, gerbera, gladiola, gloxinia, gypsophilla, hedera, hibiscus, hyacinth, hydrangea, ilex, impatiens, iris, ivy, jasmine, lilac, lily, maidenhair fern, mandevilla, marigold, narcissus, nasturtium, orchid*, pansy, pelargonium, peony, peperomia, petunia, philodendron, phlox, photinia, pinks, pittosporum, poinsettia*, pothos, portulaca, primrose, pyracantha, rhododendron, rose*, rosemary, rubber plant, salvia, schefflera, sedum, sempervivum, snapdragon, spathiphyllum, stock, syngonium, tulip, verbena, vinca, wandering jew, yucca, zinnia *Please note that when making applications to these species, spotting of plant foliage is possible.
Ornamental Trees and Shrubs including but not limited to:	Andromeda, arborvitae, ash, Austrian pine, azalea, beech, birch, birdsnest spruce, blue spruce, bougainvillea, boxwood, butternut, cedar, charmaecyparis, cherry, cotoneaster, crabapple, cypress, dogwood, Douglas fir, elm, euonymus, firethorn, forsythia, hackberry, hawthorn, hemlock, hickory, holly, honey locust, horse chestnut, juniper, larch, laurel, lilac, linden, London planetree, magnolia, mandevilla, maple, mimosa, mountain ash, myrtle, oak, pachysandra, peach, photinia, pine, planetree, poplar, privet, purpleleaf wintercreeper, quince, sage, spruce, sycamore, white cedar, white pine, yew

PESTS CONTROLLED OR SUPPRESSED

Use NeemAzal® T/S 1.2 EC Insecticide against the following pests presented in Table 1.

TABLE 1. TARGET PEST SPECIES OF NeemAzal® T/S 1.2 EC INSECTICIDE.

HEMIPTERA AND HOMOPTERA ok including but not limited to: true bugs including boxelder bugs, chinch bugs, lygus bugs and stink bug; lacebugs; leafhoppers including grape leafhopper, spittlebug, potato leafhopper and variegated leafhopper; mealy bugs including apple mealy bugs, citrus mealy bugs, grape mealy bugs; whiteflies including greenhouse whitefly, silverleaf whitefly and sweet potato whitefly and woolly whitefly; aphids including apple aphid, green peach aphid, melon aphid, pea aphid, potato aphid and rose aphid; psyllids including pear psyllids and scales including black scale, brown soft scale, California red scale, coffee scale, olive scale, San Jose scale, and cottony cushion scale.	LEPIDOPTERA Including but not limited to: Moths including European pine shoot moth, pine tip moth and Tussock moth; leafrollers including blueberry leafroller, filbert leafroller, fruitree leafroller, citrus leafminers, grape leafroller, oblique banded leafroller, omnivorous leafroller; Cutworms including black cutworm and citrus cutworm; Caterpillars and loopers including bagworms, budworms, cabbage looper, canker worms, case bearers, caseworms, corn earworm, diamondback moth, fruit worms, grapeleaf skeletonizer, gypsy moth, hornworms, imported cabbageworm, navel orangeworm, soybean looper, spruce budworm, tent caterpillar, tip moths, tent caterpillars, tobacco budworm, tobacco hornworm, tomato pinworm and tussock moth; Armyworms including beet armyworm, fall armyworm, lawn armyworm, southern armyworm and yellow striped armyworm; webworms and leaf perforators .
COLEOPTERA including but not limited to: beetles, grubs and weevils including Asian long-horned beetle, bark beetles, black vine weevil, Colorado potato beetle, elm bark beetle, European chafer, flea beetles, Japanese beetle, June beetle, leaf beetles, Mexican bean beetle, Northern masked chafer, rose chafer and Southern masked chafer and twig girders.	DIPTERA Including but not limited to: Flies including Caribbean fruit fly, cherry maggots, crane fly, fungus gnat , Hessian fly, oriental fruit fly, Mediterranean fruit fly, marsh crane flies, melon fly, shore fly and walnut husk fly; leafminers including citrus leafminers and serpentine leafminers.
THYSANOPTERA including but not limited to: thrips including citrus thrips, flower thrips, gladiolus thrips, onion thrips, thrips palmi and Western flower thrips.	*Not intended for use on public health pests
ORTHOPTERA including but not limited to: crickets; grasshoppers; locusts	ACARINA Including but not limited to: mites including, red spider mites, brown mite, clover mite, conifer spider mite, European red mite, spruce spider mite, and two-spotted spider mite.
NEMATODA nematodes (suppression)	*Not intended for use on public health pests
	HYMENOPTERA Including but not limited to: sawflies including European sawflies, pear sawflies, red-headed pine sawflies, yellow-headed pin sawflies.
	*Not intended for use on public health pests

SPRAY PREPARATION

NeemAzal® T/S 1.2 EC Insecticide is an emulsifiable concentrate to be diluted with water.

Water as diluent:

Add one-half the required amount of water to the spray tank, then add **NeemAzal® T/S 1.2 EC** Insecticide slowly with agitation, and complete filling the tank with water. To prevent separation of the emulsion, mix thoroughly and continue agitation while spraying.

This product forms an emulsion and can separate upon extended or prolonged standing. Re-agitate to assure uniformity of the spray mixture.

Adjusting the mixture pH to between 5 and 7 will provide optimal performance. Do not use tank additives that alter the pH of the spray solution above pH 7. Buffer the spray solution to alter the pH range as appropriate.

Prepare only the volume needed for the intended application, and use the spray mixture within 24 hours of preparation.

TANK MIXTURES

NeemAzal® T/S 1.2 EC Insecticide is an emulsifiable concentrate and is compatible with commonly used pesticides and fertilizers. Always check the physical compatibility using a jar test in the correct proportions if needed.

If a broader spectrum of control is required tank-mix **NeemAzal® T/S 1.2 EC** Insecticide with insecticides or miticides. If a rapid knockdown of heavy populations is necessary, then include an effective contact insecticide/miticide in combination with **NeemAzal® T/S 1.2 EC** Insecticide.

Tank mixtures are for use only in states where the companion product(s) and the application site are registered.

Always read and follow the directions for use, precautions and limitations for use on all product labels used in combination. *Applications must follow the precautions and limitations of the most restrictive product label in the mixture. Do not exceed the dosage rates of any product.*

Select the right companion products:

IPM uses a variety of control options including biological, chemical, and cultural practices. Azadirachtin is botanical with growth regulator effect on insects and mites. Companion products include pyrethroids, spinosyns, microbial toxins, and chloronicotinyls that can complement azadirachtin. Formulations of bifenthrin, spinosad, abamectins, and imidacloprid are effective for different pests. Select the product that has been proven to provide adequate performance for the pests you are trying to control.

Physical Incompatibility

Do not use **NeemAzal® T/S 1.2 EC** Insecticide with Captan, Bordeaux mixture, triphenyltin hydroxide, lime sulfur, Rayplex iron or other highly alkaline materials as they can cause phytotoxicity and/or reduced efficacy on some target pests. Tank-mix combinations with compounds known to be incompatible with oil-based formulations are not to be used or phytotoxicity will occur.

ADJUVANTS

The addition of adjuvants may enhance control under certain conditions; the use of adjuvants or oils will cause phytotoxicity and should be thoroughly tested prior to use. Do not add crop oils to spray mixtures intended for use on ornamental plants, flowers, trees, and shrubs.

APPLICATION EQUIPMENT

Ground Equipment

Apply **NeemAzal® T/S 1.2 EC** Insecticide with hand-operated (manual) or power spray equipment suitable for low volume and/or high volume applications. Follow the recommendations of the equipment manufacturer when using backpack sprayers, hose-end sprayers, compression (pump-up) sprayers, and other sprayers suitable for foliar applications of insecticides.

Chemigation and Subsurface Equipment

NeemAzal® T/S 1.2 EC Insecticide may also be applied through chemigation systems and sub-soil treatment equipment; always follow equipment manufacturer's directions.

APPLICATION SCHEDULE

For the most effective control, apply the product when pests are expected to appear or as soon as possible after pests appear and are in immature stages. Spray at an interval of seven (7) to ten (10) days or as the situation warrants.

During high pest infestation levels or when canopy is dense use higher dosage (use) rates and increase the spray frequency. Spraying in the morning or evening hours is recommended. Repeat spraying if rain occurs within two to three hours of spraying. For additional guidance, consult with the state agricultural experiment station or local extension horticulturalist/arborist for information on tactics and windows of application.

14/79

APPLICATION RATES

Use **NeemAzal® T/S 1.2 EC** Insecticide on ornamental pests as a spray concentration of 0.25 - 1.70% vol/vol per treatment with high volume applications in Table 2.

The application rates are specified as rate ranges depending upon the pest infestations:

Lower rate ranges with a spray concentration of 0.25 - 0.75% vol/vol: Use lower rate ranges for light infestations of lepidopterous insects, at the first sign or at the first observation of the early and uniform growth stages of the pest(s), and/or tank mixtures with contact insecticides.

Medium rate ranges with a spray concentration of 0.75 - 1.25% vol/vol: Use medium rate ranges for moderate infestations, when multiple growth stages of the pests are present, and/or heterogeneous pesticide populations are present.

Upper rate ranges with a spray concentration of 1.25 - 1.70% vol/vol: Use upper rate ranges for moderate to heavy pest populations of difficult-to-control pest species, for the late stages of larva/worms, for dense foliage, and/or when re-infestations occur.

High Volume Applications:

Apply **NeemAzal® T/S 1.2 EC** Insecticide at spray concentration of 0.25 - 1.70% v/v in sufficient amounts of water to achieve complete coverage. Use an adequate spray volume to wet the leaves (foliage) and stems. Spray volumes will vary with the plant size. Attempt to penetrate dense foliage. Thorough coverage of the upper and lower leaf surfaces is critical for effective levels of control.

Refer to Table 3 for the amounts of **NeemAzal® T/S 1.2 EC** Insecticide required to prepare spray concentrations of 0.25% to 1.70% for spray volumes of 1 gallon to 200 gallons.

Specialized Low Volume Applications:

Select a spray volume to achieve sufficient coverage. Uniform coverage of both upper and lower leaf surfaces is critical for effective insect control.

Apply **NeemAzal® T/S 1.2 EC** Insecticide in a *minimum* spray volume of 5 gallons per acre. Larger plants will require the higher spray volumes (20 - 25 gallons per acre) to obtain sufficient coverage. Do not exceed 20 grams active ingredient per acre per application or 57 fl. oz. of product per acre per application. Refer to Table 4 for the amounts of **NeemAzal® T/S 1.2 EC** Insecticide required to prepare spray concentrations of 0.25% to 1.70% for spray volumes of 5 - 25 gallons per acre.

15/79

TABLE 2. APPLICATION RATES FOR ORNAMENTALS ESTABLISHED IN RESIDENTIAL, LANDSCAPE PLANTINGS AROUND INSTITUTIONAL, PUBLIC, COMMERCIAL AND INDUSTRIAL BUILDINGS, PARKS, RECREATIONAL AREAS, GREENHOUSES, SHADECLOTHS, NURSERIES, AND ATHLETIC FIELDS

USE	PESTS	SPRAY CONCENTRATION%	Amounts of NeemAzal® T/S 1.2 EC Insecticide		
			Fluid ounces per gallon	Fluid ounces per 100 gallons	Quarts per 100 gallons
Including trees, shrubs, flowers, conifers, evergreens, herbaceous ornamentals, foliage plants, container-grown ornamentals, plants and groundcovers	Armyworms Azalea caterpillars Aphids Bagworms Black vine weevils Boxelder bugs Budworms Cankerworms Cutworms Eastern tent caterpillars Elm leaf beetles European sawflies Fall webworms Flea beetles Forest tent caterpillars Gypsy moth larvae Japanese beetles June beetles Lace bugs Leaf-feeding caterpillars Leafhoppers Leafminers Leaf rollers Leaf skeletonizers Oleander moth larvae Pine sawflies Pine shoot beetles Pinetip moths Plant bugs Sawflies (larva) Scale insects (crawlers) Spruce budworm Striped beetles Striped oakworms Thrips Tussock moth larvae Brown softscale California redscale (crawler) Clover mites Mealybugs Pineneedlescale (crawler) Spider mites Whiteflies and other species identified in Table 1.	Lower rate ranges of 0.25 - 0.75% vol/vol: Medium rate ranges of 0.75 - 1.25% vol/vol: Upper rate ranges of 1.25 - 1.70% vol/vol:	0.32-1.0 fl. oz. 1.00-1.60 fl. oz. 1.60-2.18 fl. oz.	32-96 fl. oz. 96-160 fl. oz. 160-218 fl. oz.	1.0-3.0 qts. 3.0-5.0 qts. 5.0-6.8 qts.

16/19
TABLE 3. SPRAY PREPARATION FOR HIGH VOLUME APPLICATIONS FOR SPRAY CONCENTRATIONS OF 0.25% to 1.70%.

Gallons Of Water	Amounts of NeemAzal® T/S 1.2 EC Insecticide For:						
	0.25%	0.50%	0.75%	1.00%	1.25%	1.50%	1.70%
1 gallon	0.32 fl. oz.	0.64 fl. oz.	0.96 fl. oz.	1.28 fl. oz.	1.60 fl. oz.	1.94 fl. oz.	2.18 fl. oz.
5 gallons	1.60 fl. oz.	3.2 fl. oz.	4.8 fl. oz.	6.4 fl. oz.	8.0 fl. oz.	9.7 fl. oz.	10.9 fl. oz.
10 gallons	3.2 fl. oz.	6.4 fl. oz.	9.6 fl. oz.	12.8 fl. oz.	16.0 fl. oz.	19.4 fl. oz.	21.8 fl. oz.
25 gallons	8.0 fl. oz.	16.0 fl. oz.	24.0 fl. oz.	32 fl. oz.	1.25 qts.	1.50 qts.	1.70 qts.
50 gallons	16.0 fl. oz.	32.0 fl. oz.	1.50 qts.	2.0 qts.	2.5 qts.	3.0 qts.	3.4 qts.
100 gallons	1.0 qt.	2.0 qts.	3.0 qts.	4.0 qts.	5.0 qts.	6.0 qts.	6.8 qts.
150 gallons	1.5 qts.	3.0 qts.	4.5 qts.	6.0 qts.	7.5 qts.	9.0 qts.	10.2 qts.
200 gallons	2.0 qts.	4.0 qts.	6.0 qts.	8.0 qts.	10.0 qts.	12.0 qts.	13.6 qts.

TABLE 4. SPECIALIZED SPRAY PREPARATION FOR LOW VOLUME APPLICATIONS OF 5 - 25 GALLONS PER ACRE WITH SPRAY CONCENTRATIONS OF 0.25% to 1.70%.

Spray Concentration Desired, % vol/vol	Spray Volume, Gallons Per Acre				
	5 gpa	10 gpa	15 gpa	20 gpa	25 gpa
0.25% v/v	1.6 fl. oz/acre	3.2 fl. oz/acre	4.9 fl. oz/acre	6.5 fl. oz/acre	8.0 fl. oz/acre
0.50% v/v	3.2 fl. oz/acre	6.4 fl. oz/acre	9.6 fl. oz/acre	12.8 fl. oz/acre	16.0 fl. oz/acre
0.75% v/v	4.8 fl. oz/acre	9.6 fl. oz/acre	14.4 fl. oz/acre	19.2 fl. oz/acre	24.0 fl. oz/acre
1.00% v/v	6.4 fl. oz/acre	12.8 fl. oz/acre	19.2 fl. oz/acre	25.5 fl. oz/acre	32.0 fl. oz/acre
1.25% v/v	8.0 fl. oz/acre	16.0 fl. oz/acre	24.0 fl. oz/acre	32.0 fl. oz/acre	40.0 fl. oz/acre
1.50% v/v	9.6 fl. oz/acre	19.2 fl. oz/acre	28.9 fl. oz/acre	38.5 fl. oz/acre	48.0 fl. oz/acre
1.70% v/v	10.8 fl. oz/acre	21.6 fl. oz/acre	32.5 fl. oz/acre	43.3 fl. oz/acre	54.0 fl. oz/acre

SPECIFIC USE INSTRUCTIONS:

Decision-making for IPM:

Scouting, monitoring, sampling, record-keeping, and predictive models are techniques to determine *if* and *when* insecticide/miticide applications are needed. The application schedule should coincide with the most vulnerable stage of the pest. For azadirachtin, target the most vulnerable stages of young larvae and young nymphs. The early larval stages and the early instar stages are more susceptible to this IGR than the later stages of the same pests.

For Lepidoptera:

- Armyworms: Apply when larvae are small.
- Bagworms: Apply when bags are small and larvae are actively feeding.
- Gypsy moth larvae: Apply when larvae are small and all eggs have hatched.
- Spruce budworms: Apply when larvae are exposed and actively feeding.

For Acarina:

- Spider mites: Apply when nymphs are first observed and before mite populations have become severe. Use multiple applications with 7-10 day intervals until infestation is controlled. Thorough coverage of both upper and lower leaf surfaces is needed.

For Thysanoptera:

- Thrips: Apply early at first signs of infestation and repeat until infestation is controlled.

For Hymenoptera:

- Sawfly: Apply when larvae are small. Refer to tree injection method of this label.

For Hemiptera and Homoptera:

- Leafhoppers: Apply when first observed and repeat applications at 5 - 7 day intervals.
- Mealybugs: Obtain thorough coverage of leaves and twigs.
- Scale: Obtain thorough coverage of leaves and twigs.

For Coleoptera:

- Beetles: Apply early at first signs of infestation and repeat applications at 7 - 10 day intervals.
- Japanese beetle (adults): Use foliar applications to repel adult feeding and treat at 5 - 7 day intervals.

For Diptera:

- Leafminers: Apply early to larvae when stippling or mining of leaves is first observed. Repeat applications at 7 - 10 day intervals until infestation is controlled.

17/79

TURFGRASS ESTABLISHED IN RESIDENTIAL (LAWNS), INSTITUTIONAL, PUBLIC, COMMERCIAL AND INDUSTRIAL SITES, PARKS, RECREATIONAL AREAS, GOLF COURSES, SOD FARMS, AND ATHLETIC FIELDS

Use **NeemAzal® T/S 1.2 EC** Insecticide to control the pests presented in Table 5. Dilute **NeemAzal® T/S 1.2 EC** Insecticide in water.

The most vulnerable stage to this product is young larvae and nymphs. Schedule treatments for the early larval stages and early instars when populations are established, but before turf damage becomes noticeable.

The maximum rate on turfgrass of **NeemAzal® T/S 1.2 EC** Insecticide is 57.0 fl. oz of product per acre per application or 1.3 fl. oz product per 1,000 sq. ft. per application. Apply at a rate up to 57 fl. oz of product per acre. Use the higher rate specified on this label for moderate to heavy infestations.

Irrigation:

Avoid (delay or postpone) irrigation for 12 - 24 hours after application of this product.

Mowing:

Avoid (delay or postpone) mowing of the treated area for 12 - 24 hours after treatment.

Degree day and plant phenology models can assist in developing the appropriate application schedule for the target pests. Consult your state university or local Cooperative Extension Service office for specific pest control application timing in your area.

NeemAzal® T/S 1.2 EC Insecticide can be tank mixed with other insecticide/miticides if a broader spectrum of pest control is required. Observe all precautionary statements and follow all label directions of companion product(s).

Specific Use Instructions:

1. Armyworms: Apply during the early morning or late afternoon to maximize control.
2. Sod webworm larvae: Applications in the late afternoon or early evening can maximize control.

TABLE 5. APPLICATION RATES FOR TURFGRASS ESTABLISHED IN RESIDENTIAL (LAWNS), INSTITUTIONAL, PUBLIC, COMMERCIAL AND INDUSTRIAL SITES, PARKS, RECREATIONAL AREAS, GOLF COURSES, SOD FARMS, AND ATHLETIC FIELDS.

USE	PESTS	Amount of NeemAzal® T/S 1.2 EC Insecticide		Spray Volumes		Number of Applications & Interval Days
		fl. oz./acre	fl. oz./1,000 sq. ft.	gals./acre	gals./1000 sq. ft.	
Cool-Season and Warm-Season Turfgrass	<i>Larvae and nymphs of these pests including but not limited to:</i> Armyworms Bermudagrass mite Cutworms Grasshopper Sod webworm	Up to 57.0 fl. oz	Up to 1.3 fl. oz	40 - 100 gpa	1-2 gal/1,000 sq. ft.	As needed, 7 days

DRENCH APPLICATION FOR GREENHOUSES, NURSERIES, INTERIORSCAPES AND FOR PLANTS GROWN IN CONTAINERS:

Use **NeemAzal® T/S 1.2 EC** Insecticide as a soil drench for effective control of soil-borne insect larvae, including soil-borne larvae of foliar pests, such as fungus gnats, nematodes, or soil borne thrips. When applying as a drench, avoid excessive leaching.

Preventive applications as a soil drench may be warranted for certain pests. Soil drench applications of azadirachtin will have a slower rate of activity because of soil absorption when compared to foliar applications of azadirachtin. Target the initial application of a soil drench treatment to coincide with the early stages of young larvae and young nymphs.

Dilute **NeemAzal® T/S 1.2 EC** Insecticide with water for concentrations of 0.4 to 0.8% vol/vol. Drench the soil in the pot with one (1) pint of finished spray per 1.0 gallon of soil. For fungus gnats, use the 0.4% spray concentration. For mushroom fly maggot control, use the 0.6% vol/vol spray concentration. For leafminers and other difficult to control pests, use the 0.8% vol/vol spray concentration. Two to three (2-3) applications should be scheduled at 10-14 day intervals until the pest pressure has ended.

DILUTION TABLE FOR DRENCH APPLICATIONS

Gallons of Water	Amount of NeemAzal® T/S 1.2 EC Insecticide			Application Interval	Number of Applications
	0.4%	0.6%	0.8%		
1 gallon	1 Tbs.	1.5 Tbs.	2.0 Tbs.	10 - 14 days	2 - 3
5 gallons	2.7 fl. oz	4.0 fl. oz	5.5 fl. oz	10 - 14 days	2 - 3
10 gallons	5.4 fl. oz	8.0 fl. oz	11.0 fl. oz	10 - 14 days	2 - 3
100 gallons	1.7 qts.	2.5 qts.	3.4 qts.	10 - 14 days	2 - 3

NeemAzal® T/S 1.2 EC Insecticide can also be applied through sub-surface treatment equipment. Always follow manufacturer's use directions.

TREE INJECTION

Inject **NeemAzal® T/S 1.2 EC** Insecticide into mature trees established in landscapes, residential settings, nurseries, and forestry sites.

Use appropriate tree injection equipment and follow the instructions provided by the equipment manufacturer.

Application Schedule For Tree Injections

Consult with your state agricultural experiment station, extension specialist, or your local U.S. Forest Service authority for information on the application schedule for specific pests in your area.

Pests Controlled and Hosts:

PESTS	HOSTS
Spruce budworm larva	White Spruce Black Spruce Balsam Fir
Pine false webworm	Eastern White Pine Red Pine
Pine sawfly larvae	White Pine
Cedar leafminer	White Cedar

Dosage Rate For Tree Injections

Use appropriate injection equipment. Inject at the rate of 0.37 - 0.74 fl. oz (11 - 22 ml) of product per inch tree trunk diameter at breast height. Or, inject at the rate of 0.127 - 0.25 grams azadirachtin per inch tree trunk diameter at breast height.

INSTRUCTIONS FOR GARDEN CROPS, VEGETABLES, HERBS AND SPICES, FRUITS, AND BERRIES

- For the most effective control, spray the product as soon as possible after pests appear and are in immature stages.
- Spray at an interval of seven to ten days or as the situation warrants. During high pest infestation levels use higher label rates and increase the spray frequency.
- For best results spray in the morning or evening hours.
- Repeat spraying if rain occurs within two to three hours of spraying.

SPRAY EQUIPMENT

Use any suitable application equipment to ensure uniform coverage.

USE RATES

Apply **NeemAzal® T/S 1.2 EC** Insecticide as directed to any food or non-food crop up to and including the day of harvest, at a maximum rate of 57 fl. ozs. (20 grams active ingredient) per acre (1.33 fl. ozs. per 1,000 sq. ft.) per application. Rates in Table 6 pertain to typical pest infestations.

Apply **NeemAzal® T/S 1.2 EC** Insecticide alone to food/garden crops on the day of harvest.

Dilute **NeemAzal® T/S 1.2 EC** Insecticide with water at 0.5 - 4.0 tablespoons (Tbs) per gallon of water. For hose end sprayers, set the RATE PER GALLON at the dial setting of 1 to 4 Tbs. depending on the crop and pests. Use the lower RATE PER GALLON for low to moderate infestations and use the higher specified RATE PER GALLON for severe infestations.

TABLE 6. USE RATES FOR GARDEN CROPS, VEGETABLES, HERBS AND SPICES, BERRIES AND FRUIT.

CROP	PESTS such as:	Dilution Rate for Sprayers	
		Fl. Ozs. of product per 1,000 Sq. Ft.	Tbs. of product per 1.0 gallon of water
<i>Leafy Vegetables including but not limited to: Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Endive, Kale, Lettuce, Spinach</i>	Leafrollers, Cutworms, Loopers, Armyworms	0.19 - 0.96 fl. ozs.	3/4 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Beetles, Weevils, Flies, Thrips, Mites	0.24 - 0.96 fl. ozs.	1 Tbs. - 4 Tbs./gal
<i>Root Vegetables, including but not limited to: Beet, Carrot, Horseradish, Parsnip, Potato, Radish, Sweet potato, Turnip, Yams</i>	Beetles, Weevils	0.11 - 0.25 fl. ozs.	1/2 Tbs. - 1 1/2 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Thrips, Mites	0.24 - 0.96 fl. ozs.	1 Tbs. - 4 Tbs./gal
	Beetles, Weevils	0.29 - 0.96 fl. ozs.	2 Tbs. - 4 Tbs./gal
<i>Fruiting Vegetables including but not limited to: Eggplant, Pepper, Tomatillo, Tomato</i>	Thrips	0.29 - 0.96 fl. ozs.	2 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl. ozs.	1 Tbs. - 4 Tbs./gal
	Beetles, Weevils	0.29 - 0.96 fl. ozs.	2 Tbs. - 4 Tbs./gal
<i>Cucurbit Vegetables including but not limited to: Cucumber, Gourd (edible), Muskmelon, Pumpkin, Squash, Watermelon, including Cantaloupe, Casaba, Gherkins, Melons (including hybrids), Zucchini</i>	Thrips	0.29 - 0.96 fl. ozs.	2 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl. ozs.	1 Tbs. - 4 Tbs./gal
	Beetles, Weevils	0.29 - 0.96 fl. ozs.	2 Tbs. - 4 Tbs./gal
<i>Legume Vegetables including but not limited to: Bean, Chickpea, Lentil, Pea</i>	Thrips	0.29 - 0.96 fl. ozs.	2 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl. ozs.	1 Tbs. - 4 Tbs./gal
	Beetles, Weevils	0.29 - 0.96 fl. ozs.	2 Tbs. - 4 Tbs./gal
<i>Bulb Vegetables including but not limited to: Garlic, Onion, Shallot</i>	Thrips	0.29 - 0.96 fl. ozs.	2 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl. ozs.	1 Tbs. - 4 Tbs./gal

CROP	PESTS such as:	Dilution Rate for Sprayers	
		Fl. Ozs. of product per 1,000 Sq. Ft.	Tbs. of product per 1.0 gallon of water
<i>Berries including but not limited to: Blackberry, Blueberry, Raspberry, Strawberry, others include: Boysenberry, Currants, Dewberry, Elderberry, Gooseberry, Loganberry.</i>	Beetles, Weevils	0.29 - 0.96 fl. ozs.	2 Tbs. - 4 Tbs./gal
	Thrips	0.29 - 0.96 fl. ozs.	2 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl. ozs.	1 Tbs. - 4 Tbs./gal
<i>Herbs and Spices including but not limited to: Chive, Dill, Fennel, Mustard, Sage, Sweet bay, others include: Anise, Balm, Basil, Black pepper, Borage, Caraway, Catnip, Chamomile, Coriander, Cumin, Curry leaf, Dandelion, Fenugreek, Horehound, Hyssop, Marjoram, Marigold, Mint, Nasturtium, Pennyroyal, Peppermint, Rosemary, Savory, Spearmint, Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood</i>	Beetles, Weevils	0.29 - 0.96 fl. ozs.	2 Tbs. - 4 Tbs./gal
	Thrips	0.29 - 0.96 fl. ozs.	2 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl. ozs.	1 Tbs. - 4 Tbs./gal
<i>Nut Trees including but not limited to: Almond, Brazil nut, Filbert, Hickory nut, Pecan, Pistachios, Walnut</i>	Beetles, Weevils	0.29 - 0.96 fl. ozs.	2 Tbs. - 4 Tbs./gal
	Thrips	0.29 - 0.96 fl. ozs.	2 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl. ozs.	1 Tbs. - 4 Tbs./gal
<i>Pome Fruits including but not limited to: Apple, Quince, or Pear (Comice varieties: DO NOT apply more than 24 fl oz/A. DO NOT apply after pink stage of flowering; test small areas of other varieties of pears for plant safety prior to full scale usage.)</i>	Beetles, Weevils	0.29 - 0.96 fl. ozs.	2 Tbs. - 4 Tbs./gal
	Thrips	0.29 - 0.96 fl. ozs.	2 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl. ozs.	1 Tbs. - 4 Tbs./gal
<i>Stone Fruits including but not limited to: Apricot, Cherry, Nectarine, Peach, Plum</i>	Beetles, Weevils	0.29 - 0.96 fl. ozs.	2 Tbs. - 4 Tbs./gal
	Thrips	0.29 - 0.96 fl. ozs.	2 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl. ozs.	1 Tbs. - 4 Tbs./gal
<i>Citrus Fruits including but not limited to: Grapefruit, Lemon, Lime, Orange others include: Citrus Citron, Mandarin (tangerine), Nectarine, Satsuma (orange mandarin), Tangerine</i>	Beetles, Weevils	0.29 - 0.96 fl. ozs.	2 Tbs. - 4 Tbs./gal
	Thrips	0.29 - 0.96 fl. ozs.	2 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl. ozs.	1 Tbs. - 4 Tbs./gal

CHEMIGATION GENERAL INFORMATION

Apply this product only through drip (trickle) or sprinkler irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues can result from non-uniform distribution of treated water. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. Direct your questions concerning calibration to your State Extension Service Specialist, the equipment manufacturer, or other expert. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of a responsible person, shall shut the system down and make necessary adjustments should the need arise.

Dilute NeemAzal® T/S 1.2 EC Insecticide with water before introduction into the system. Use the diluted solution within 8 hours. Do not apply in irrigation water if the pH exceeds 7.0. The optimum pH range for application is 5.5 to 6.5. The pH of the irrigation water can be adjusted by use of a

suitable buffering agent. Agitation is necessary. Apply at the specified rate using sufficient water to achieve an even distribution within an 8-hour period. Do not apply **NeemAzal® T/S 1.2 EC** Insecticide at a rate that exceeds 3.5 pints active ingredient per acre (57 fl. oz).

If the irrigation cycle will last longer than 8 hours and the **NeemAzal® T/S 1.2 EC** Insecticide is premixed in the supply tank, the tank mix must be buffered to a pH of 8 or lower.

For Chemigation Systems Connected to A Public Water System: *Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of a year.*

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone (RPZ), backflow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction.

There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top of overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speeds favor drift beyond the area intended for treatment.

Operation Of Sprinkler Chemigation Or Drip (Trickle) Utilizing A Pressurized Water And Pesticide Injection System: The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

STORAGE AND DISPOSAL Do not contaminate water, food, or feed by storage and disposal	
Pesticide Storage:	Store in original container in a dry, cool, well ventilated area.
Pesticide Disposal:	Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.
Container Disposal:	Nonrefillable container. Do not reuse or refill this container. Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill container $\frac{1}{4}$ full with water and recap. Shake 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Drain for 10 seconds after flow begins to drip. Repeat this procedure two more times.

NOTICE ON CONDITIONS OF SALE

The directions for use of this product are based upon tests believed to be reliable, and the directions for use must be followed. To the extent applicable by law the use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. To the extent applicable by law, the buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

Advertising claims that may be presented on the retail container label or on the labeling accompanying the product.

- Controls mites, caterpillars, whiteflies, thrips, aphids, and other insects as listed on this label.
- Organic certified
- Controls chewing and sucking insects
- Low-odor formulation
- Broad-spectrum control
- Can be applied the day of harvest
- For use on a wide variety of trees, shrubs, flowers, fruit and nut trees, garden vegetables and plants
- Low (Mild) odor
- Organic Materials Review Institute (OMRI) Certified.

EPA Text: 'NeemAzal® T/S 1.2 EC (06-13-06)
Revised 11-17-08

Sublabel 3

AzaMax™

Botanical Insecticide, Miticide, and Nematicide

REPELLENT, ANTI FEEDANT AND INSECT GROWTH REGULATOR (IGR)

- BOTANICAL PRODUCT FOR CONTROL OF INSECTS ON FRUIT AND NUT TREES
- INDOOR AND OUTDOOR VEGETABLES, ORNAMENTAL FLOWERS, TREES, SHRUBS AND PLANTS, INCLUDING PLANTS GROWN IN CONTAINERS, INTERIORSCAPES, HOME AND GARDEN USE.

See the Directions for Use for a Complete List of Insects Controlled.

ACTIVE INGREDIENT:

	% By Wt.
Azadirachtin	1.2%
OTHER INGREDIENTS	98.8%
	TOTAL 100.00%

Contains 0.35 grams azadirachtin per fluid ounce.



FOR ORGANIC PRODUCTION

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call poison control center or doctor for further treatment advice.
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.
If swallowed	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

CALL THE POISON CONTROL HOTLINE 24 HOURS A DAY AT 1-888-478-0798.

NET CONTENTS: 2 fl. oz, 4 fl. oz., 8 fl. oz., 16 fl. oz., 32 fl. oz. Or 128 fl. oz

EPA Reg. No. 71908-1
EPA Est. No. 71908-IND-001

E.I.D. PARRY (INDIA) LIMITED
Bioproducts Division
234 NSC Bose Road
Chennai (Madras) 600 001
India

USA Phone Number: (916) 973-1351
Batch Number:xxxxx

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS -

CAUTION: Harmful if absorbed through skin or if inhaled. Avoid breathing vapor. Causes moderate eye irritation. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Wear chemical resistant gloves.

Personal Protective Equipment (PPE)

Applicators and handlers must wear:

- Long-sleeved shirt
- Long pants
- Socks and shoes
- Chemical resistant gloves

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates.

For Terrestrial Uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwater or rinsate.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Do not apply this product through any type of irrigation system:

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses and handlers of agricultural products. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment (PPE), notification to workers and restricted entry interval. The requirements in this box apply to the uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow any worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, water, wear coveralls, chemical resistant gloves, shoes plus socks and protective eyewear.

MODE OF ACTION:

AzaMax™ controls target pests on contact or by ingestion. The product acts on pests by way of repellence, anti-feedance, and interference with the molting process.

Azadirachtin, an insect growth regulator (IGR), mimics the pests' hormones and disrupts distinct stages of growth and development of insects and mites. The primary mode of action of azadirachtin is an interference with synthesis and metabolism of ecdysone and the juvenile hormone. Ecdysone is the molting hormone of insects, and azadirachtin can regulate growth leading to death before or during molting.

I. USES FOR ORNAMENTALS AND LANDSCAPE PLANTINGS

Ornamental Plants and Flowers including but not limited to:	Actinopteris, African violets*, ageratum, aglaonema, Algerian ivy, allamanda, alocasia, amaranthus, anthurium, aphelandra, arborvitae, Artemisia, aster, aucuba ilex, azalea, baby's breath, begonia, Boston fern, bougainvillea, boxwood, brachycome, cacti, calabrese, caladium, calathea, calendula, calla, camellia, carnation, ceanothus, chrysanthemum, cineraria, coleus, columbine, cotoneaster, cyclamen, daffodil, dahlia, daisy, daylily, delphinium, dianthus, dieffenbachia, dogwood, dusty miller, Easter lily, English ivy, euphorbia, fern, ficus, foliage plants, foxglove, freesia, fuschia, gaillardia, gardenia, geranium, gerbera, gladiola, gloxinia, gypsophilla, hedera, hibiscus, hyacinth, hydrangea, ilex, impatiens, iris, ivy, jasmine, lilac, lily, maidenhair fern, mandevilla, marigold, narcissus, nasturtium, orchid*, pansy, pelargonium, peony, peperomia, petunia, philodendron, phlox, photinia, pinks, pittosporum, poinsettia*, pothos, portulaca, primrose, pyracantha, rhododendron, rose*, rosemary, rubber plant, salvia, schefflera, sedum, sempervivum, snapdragon, spathiphyllum, stock, syngonium, tulip, verbena, vinca, wandering jew, yucca, zinnia <i>* Please note that when making applications to these species, spotting of plant foliage and blossoms is possible.</i>
Ornamental Trees and Shrubs including but not limited to:	Andromeda, arborvitae, ash, Austrian pine, azalea, beech, birch, birdsnest spruce, blue spruce, bougainvillea, boxwood, butternut, cedar, charmaecyparis, cherry, cotoneaster, crabapple, cypress, dogwood, Douglas fir, elm, euonymus, firethorn, forsythia, hackberry, hawthorn, hemlock, hickory, holly, honey locust, horse chestnut, juniper, larch, laurel, lilac, linden, London planetree, magnolia, mandevilla, maple, mimosa, mountain ash, myrtle, oak, pachysandra, peach, photinia, pine, planetree, poplar, privet, purpleleaf wintercreeper, quince, sage, spruce, sycamore, white cedar, white pine, yew

Waxy bloom on certain ornamental plants will be reduced after an **AzaMax™** application.

Applications of **AzaMax™** will remove the glaucus 'blue' coloring from evergreens such as Colorado blue spruce and Koster spruce.

PESTS CONTROLLED OR SUPPRESSED

Use AzaMax™ against the following pests.

TARGET PEST SPECIES OF AzaMax™

HEMIPTERA AND HOMOPTERA including but not limited to: true bugs including boxelder bugs, chinch bugs, lygus bugs and stink bug; lacebugs; leafhoppers including grape leafhopper, spittlebug, potato leafhopper and variegated leafhopper; mealy bugs including apple mealy bugs, citrus mealy bugs, grape mealy bugs; whiteflies including greenhouse whitefly, silverleaf whitefly and sweet potato whitefly and woolly whitefly; aphids including apple aphid, green peach aphid, melon aphid, pea aphid, potato aphid and rose aphid; psyllids including pear psyllids and scales including black scale, brown soft scale, California red scale, coffee scale, olive scale, San Jose scale, and cottony cushion scale.	LEPIDOPTERA including but not limited to: moths including European pine shoot moth, pine tip moth and Tussock moth; leafrollers including blueberry leafroller, filbert leafroller, fruitree leafroller, citrus leafminers, grape leafroller, oblique banded leafroller, omnivorous leafroller; cutworms including black cutworm and citrus cutworm; caterpillars and loopers including bagworms, budworms, cabbage looper, canker worms, case bearers, caseworms, corn earworm, diamondback moth, fruit worms, grapeleaf skeletonizer, gypsy moth, hornworms, imported cabbageworm, navel orangeworm, soybean looper, spruce budworm, tent caterpillar, tip moths, tent caterpillars, tobacco budworm, tobacco hornworm, tomato pinworm and tussock moth; armyworms including beet armyworm, fall armyworm, lawn armyworm, southern armyworm and yellow striped armyworm; webworms and leaf perforators .
COLEOPTERA including but not limited to: beetles, grubs and weevils including Asian long-horned beetle, bark beetles, black vine weevil, Colorado potato beetle, elm bark beetle, European chafer, flea beetles, Japanese beetle, June beetle, leaf beetles, Mexican bean beetle, Northern masked chafer, rose chafer and Southern masked chafer and twig girders.	DIPTERA including but not limited to: flies including Caribbean fruit fly, cherry maggots, crane fly, fungus gnat , Hessian fly, oriental fruit fly, Mediterranean fruit fly, marsh crane flies, melon fly, shore fly and walnut husk fly; leafminers including citrus leafminers and serpentine leafminers.
THYSANOPTERA including but not limited to: thrips including citrus thrips, flower thrips, gladiolus thrips, onion thrips, thrips palmi and Western flower thrips.	*Not intended for use on public health pests
ORTHOPTERA including but not limited to: crickets; grasshoppers; locusts	ACARINA including but not limited to: mites including, red spider mites, brown mite, clover mite, conifer spider mite, European red mite, spruce spider mite, and two-spotted spider mite.
	*Not intended for use on public health pests
NEMATODA Nematodes (suppression)	HYMENOPTERA including but not limited to: sawflies including European sawflies, pear sawflies, red-headed pine sawflies, yellow-headed pin sawflies.
	*Not intended for use on public health pests

SPRAY PREPARATION

AzaMax™ is an emulsifiable concentrate to be diluted with water.

This product forms an emulsion and which separates upon extended or prolonged standing. Re-agitate to assure uniformity of the spray mixture.

Prepare only the volume needed for the intended application, and use the spray mixture within 24 hours of preparation.

TANK MIXTURES

AzaMax™ is an emulsifiable concentrate and is compatible with commonly used pesticides and fertilizers. Always check the physical compatibility using a jar test in the correct proportions if needed.

If a broader spectrum of control is required tank-mix **AzaMax™** with insecticides or miticides. If a rapid knockdown of heavy populations is necessary, then include an effective contact insecticide/miticide in combination with **AzaMax™**.

Always read and follow the directions for use, precautions and limitations for use on all product labels used in combination. *Applications must follow the precautions and limitations of the most restrictive product label in the mixture. Do not exceed the dosage rates of any product.*

Select the right companion products:

IPM uses a variety of control options including biological, chemical, and cultural practices. **AzaMax™** is botanical with growth regulator effect on insects and mites. Companion products include pyrethroids, spinosyns, microbial toxins, and chloronicotinyls that complement azadirachtin. Formulations of bifenthrin, spinosad, abamectins, and imidacloprid are effective for different pests. Select the product that has been proven to provide adequate performance for the pests you are trying to control.

Physical Incompatibility

Do not use **AzaMax™** with Captan, Bordeaux mixture, triphenyltin hydroxide, lime sulfur, Rayplex iron or other highly alkaline materials as they can cause phytotoxicity and/or reduced efficacy on some target pests. Phytotoxicity will occur if tank-mix combinations with compounds known to be incompatible with oil-based formulations are used.

APPLICATION EQUIPMENT

Ground Equipment

Apply **AzaMax™** with hand-operated (manual) or power spray equipment suitable for low volume and/or high volume applications. Follow the recommendations of the equipment manufacturer when using backpack sprayers, hose-end sprayers, compression (pump-up) sprayers, and other sprayers suitable for foliar applications of insecticides.

APPLICATION SCHEDULE

For the most effective control, apply the product when pests are expected to appear or as soon as possible after pests appear and are in immature stages. Spray at an interval of seven (7) to ten (10) days or as the situation warrants.

During high pest infestation levels or when canopy is dense use higher dosage (use) rates and increase the spray frequency. For best results, spray in the morning or evening hours. Repeat spraying if rain occurs within two to three hours of spraying. For additional guidance, consult with your state agricultural experiment station or local extension horticulturist/arborist for information on tactics and windows of application.

FOLIAR APPLICATION RATES FOR ORNAMENTALS and LANDSCAPE PLANTINGS

USE	SPRAY CONCENTRATION%	Amounts of
		AzaMax™
		Fluid Ounces Per Gallon
Including trees, shrubs, flowers, conifers, evergreens, herbaceous ornamentals, foliage plants, container-grown ornamentals, plants and groundcovers	Lower rate ranges of 0.25 - 0.75% vol/vol:	0.32-1.0 fl.oz.
	Medium rate ranges of 0.75 - 1.25% vol/vol:	1.00-1.60 fl.oz.
	Upper rate ranges of 1.25 - 1.70% vol/vol:	1.60-2.18 fl.oz.

DRENCH APPLICATION FOR INTERIORSCAPES AND FOR PLANTS GROWN IN CONTAINERS:

Use **AzaMax™** as a soil drench for effective control of soil-borne insect larvae, including soil-borne larvae of foliar pests, such as fungus gnats, nematodes, or soil borne thrips. When applying as a drench, avoid excessive leaching.

Preventive applications as a soil drench may be warranted for certain pests. Soil drench applications of azadirachtin will have a slower rate of activity because of soil absorption when compared to foliar applications of **AzaMax™**. Target the initial application of a soil drench treatment to coincide with the early stages of young larvae and young nymphs.

Dilute **AzaMax™** with water for concentrations of 0.4 to 0.8% vol/vol. Drench the soil in the pot with one (1) pint of finished spray per 1.0 gallon of soil. For fungus gnats, use the 0.4% spray concentration. For mushroom fly maggot control, use the 0.6% vol/vol spray concentration. For leafminers and other difficult to control pests, use the 0.8% vol/vol spray concentration. Make two to three (2-3) applications at 10-14 day intervals until the pest pressure has ended.

DILUTION TABLE FOR DRENCH APPLICATIONS

Gallons of Water	Amount of AzaMax™			Application Interval	Number of Applications
	0.4%	0.6%	0.8%		
1 gallon	1 Tbs.	1.5 Tbs.	2.0 Tbs.	10 - 14 days	2 - 3
5 gallons	2.7 fl.oz.	4.0 fl.oz.	5.5 fl.oz.	10 - 14 days	2 - 3
10 gallons	5.4 fl.oz.	8.0 fl.oz.	11.0 fl.oz.	10 - 14 days	2 - 3

AzaMax™ can also be applied through sub-surface treatment equipment. Always follow manufacturer's use directions.

II. USES FOR GARDEN CROPS, VEGETABLES, HERBS AND SPICES, FRUITS AND BERRIES

INSTRUCTIONS FOR GARDEN CROPS, VEGETABLES, HERBS AND SPICES, FRUITS, AND BERRIES

- For the most effective control, spray the product as soon as possible after pests appear and are in immature stages.
- Spray at an interval of seven to ten days or as the situation warrants. During high pest infestation levels use higher label rates and increase the spray frequency.
- For best results, spray in the morning or evening hours.
- Repeat spraying if rain occurs within two to three hours of spraying.

USE RATES

Apply **AzaMax™** as directed to any food or non-food crop up to and including the day of harvest, at a maximum rate of 1.33 fl.ozs. per 1,000 sq.ft. per application.

Apply **AzaMax™** alone to food/garden crops on the day of harvest.

Dilute **AzaMax™** with water at 0.5 - 4.0 tablespoons (Tbs) per gallon of water. For hose end sprayers, set the RATE PER GALLON at the dial setting of 1 to 4 Tbs. depending on the crop and pests. Use the lower RATE PER GALLON for low to moderate infestations and use the higher specified RATE PER GALLON for severe infestations.

29
79

USE RATES FOR GARDEN CROPS, VEGETABLES, HERBS AND SPICES, BERRIES AND FRUIT.

CROP	PESTS such as:	Dilution Rate for Sprayers	
		Fl.Ozs. of product per 1,000 Sq.Ft.	Tbs. of product per 1.0 gallon of water
<i>Leafy Vegetables including but not limited to: Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Endive, Kale, Lettuce, Spinach</i>	Leafrollers, Cutworms, Loopers, Armyworms	0.19 - 0.96 fl.ozs.	¾ Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Beetles, Weevils, Flies, Thrips, Mites	0.24 - 0.96 fl.ozs.	1 Tbs. - 4 Tbs./gal
<i>Root Vegetables, including but not limited to: Beet, Carrot, Horseradish, Parsnip, Potato, Radish, Sweet potato, Turnip, Yams</i>	Beetles, Weevils	0.11 - 0.25 fl.ozs.	½ Tbs. - 1 ½ Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Thrips, Mites	0.24 - 0.96 fl.ozs.	1 Tbs. - 4 Tbs./gal
<i>Fruiting Vegetables including but not limited to: Eggplant, Pepper, Tomatillo, Tomato</i>	Beetles, Weevils	0.29 - 0.96 fl.ozs.	2 Tbs. - 4 Tbs./gal
	Thrips	0.29 - 0.96 fl.ozs.	2 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl.ozs.	1 Tbs. - 4 Tbs./gal
<i>Cucurbit Vegetables including but not limited to: Cucumber, Gourd (edible), Muskmelon, Pumpkin, Squash, Watermelon, including Cantaloupe, Casaba, Gherkins, Melons (including hybrids), Zucchini</i>	Beetles, Weevils	0.29 - 0.96 fl.ozs.	2 Tbs. - 4 Tbs./gal
	Thrips	0.29 - 0.96 fl.ozs.	2 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl.ozs.	1 Tbs. - 4 Tbs./gal
<i>Legume Vegetables including but not limited to: Bean, Chickpea, Lentil, Pea</i>	Beetles, Weevils	0.29 - 0.96 fl.ozs.	2 Tbs. - 4 Tbs./gal
	Thrips	0.29 - 0.96 fl.ozs.	2 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl.ozs.	1 Tbs. - 4 Tbs./gal
<i>Bulb Vegetables including but not limited to: Garlic, Onion, Shallot</i>	Beetles, Weevils	0.29 - 0.96 fl.ozs.	2 Tbs. - 4 Tbs./gal
	Thrips	0.29 - 0.96 fl.ozs.	2 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl.ozs.	1 Tbs. - 4 Tbs./gal
<i>Berries including but not limited to: Blackberry, Blueberry, Raspberry, Strawberry, others include: Boysenberry, Currants, Dewberry, Elderberry, Gooseberry, Loganberry</i>	Beetles, Weevils	0.29 - 0.96 fl.ozs.	2 Tbs. - 4 Tbs./gal
	Thrips	0.29 - 0.96 fl.ozs.	2 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl.ozs.	1 Tbs. - 4 Tbs./gal
<i>Herbs and Spices including but not limited to: Chive, Dill, Fennel, Mustard, Sage, Sweet bay, others include: Anise, Balm, Basil, Black pepper, Borage, Caraway, Catnip, Chamomile, Coriander, Cumin, Curry leaf, Dandelion, Fenugreek, Horehound, Hyssop, Marjoram, Marigold, Mint,</i>	Beetles, Weevils	0.29 - 0.96 fl.ozs.	2 Tbs. - 4 Tbs./gal
	Thrips	0.29 - 0.96 fl.ozs.	2 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl.ozs.	1 Tbs. - 4 Tbs./gal

CROP	PESTS such as:	Dilution Rate for Sprayers	
		Fl.Ozs. of product per 1,000 Sq.Ft.	Tbs. of product per 1.0 gallon of water
Nasturtium, Pennyroyal, Peppermint, Rosemary, Savory, Spearmint, Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood			
Nut Trees including but not limited to: Almond, Brazil nut, Filbert, Hickory nut, Pecan, Pistachios, Walnut	Beetles, Weevils	0.29 - 0.96 fl.ozs.	2 Tbs. - 4 Tbs./gal
	Thrips	0.29 - 0.96 fl.ozs.	2 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl.ozs.	1 Tbs. - 4 Tbs./gal
Pome Fruits including but not limited to: Apple, Quince, or Pear (Comice varieties: DO NOT apply more than 24 fl oz/A. DO NOT apply after pink stage of flowering; test small areas of other varieties of pears for plant safety prior to full scale usage.)	Beetles, Weevils	0.29 - 0.96 fl.ozs.	2 Tbs. - 4 Tbs./gal
	Thrips	0.29 - 0.96 fl.ozs.	2 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl.ozs.	1 Tbs. - 4 Tbs./gal
Stone Fruits including but not limited to: Apricot, Cherry, Nectarine, Peach, Plum	Beetles, Weevils	0.29 - 0.96 fl.ozs.	2 Tbs. - 4 Tbs./gal
	Thrips	0.29 - 0.96 fl.ozs.	2 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl.ozs.	1 Tbs. - 4 Tbs./gal
Citrus Fruits including but not limited to: Grapefruit, Lemon, Lime, Orange others include: Citrus Citron, Mandarin (tangerine), Nectarine, Satsuma (orange mandarin), Tangerine	Beetles, Weevils	0.29 - 0.96 fl.ozs.	2 Tbs. - 4 Tbs./gal
	Thrips	0.29 - 0.96 fl.ozs.	2 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl.ozs.	1 Tbs. - 4 Tbs./gal

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original containers in a dry, cool, well-ventilated area.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill container 1/4 full with water and recap. Shake 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Drain for 10 seconds after flow begins to drip. Repeat this procedure two more times.

NOTICE ON CONDITIONS OF SALE

The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Plant injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of E.I.D. PARRY (INDIA) LIMITED. All such risks shall be assumed by the user or buyer.

Advertising claims that may be presented on the retail container label or on the labeling accompanying the product.

- Controls mites, caterpillars, whiteflies, thrips, aphids, and other insects as listed on this label.
- Organic certified
- Controls chewing and sucking insects
- Low-odor formulation
- Broad-spectrum control
- Can be applied the day of harvest
- For use on a wide variety of trees, shrubs, flowers, fruit and nut trees, garden vegetables and plants
- Low (Mild) odor
- Organic Materials Review Institute (OMRI) Certified.
- Spray and Drench
- Can be used in Organic Gardening  FOR ORGANIC PRODUCTION
- Controls chewing and sucking insects
- Broad spectrum insect and mite control
- Rose Spray
- For Use on Roses and Flowers
- Vegetable Garden Spray
- For Use on Tomatoes and Garden Vegetables
- Controls Japanese Beetle on Roses
- Insect and Mite Control for the Home Garden Vegetables
- Complete Insect and Mite Control for Flowers
- Three applications in 30 days provides complete plant protection
- Quadruple action insect control – Antifeedant, Insect Growth Regulator, Anti-ovipository, Repellant
- The Pure Power of PURER azadirachtin
- Single product; multiple action
- A proven botanical pesticide born out of a decade of intense global research
- Works effectively against a broad spectrum of pests on a stand alone basis on a variety of vegetable crops
- Controls insect pest during various stages of growth
- The benefits to your insect control program are multiple
- Controls Insects Systemically
- Taken up through the roots, controls insects on the leaves
- Insects stop feeding when they feed on treated plants
- For foliar and systemic insect control
- For Use on Flowers, Ornamentals, and Landscape Plantings
- For Use on Garden Crops, Vegetables, Herbs, Spices, Fruits, and Berries
- Three in one product – Insecticide, Miticide, and Nematicide
- Recommended by General Hydroponics Bringing Nature and Technology Together
- Brought to you Exclusively by General Hydroponics – Bringing Nature and Technology Together

EPA Text: NeemAzal® T/S 1.2 EC (06-13-06)
Revised 11-17-08

Sublabel 4

EcoGarden™

Botanical Insecticide, Miticide, and Nematicide

REPELLANT, ANTIFEEDANT AND INSECT GROWTH REGULATOR (IGR)

BOTANICAL PRODUCT FOR CONTROL OF INSECTS ON INDOOR AND OUTDOOR ORNAMENTAL FLOWERS, TREES, SHRUBS, VEGETABLES, FRUIT AND NUT TREES, PLANTS, INCLUDING PLANTS GROWN IN CONTAINERS, RECIRCULATORY, AEROPONIC, AND HYDROPONIC SYSTEMS, INTERIORSCAPES, HOME, AND GARDEN.

See the Directions for Use for a Complete List of Insects Controlled.

ACTIVE INGREDIENT:		% By Wt.
Azadirachtin	1.2%
OTHER INGREDIENTS.....	98.8%
		TOTAL 100.00%

Contains 0.35 grams azadirachtin per fluid ounce.



FOR ORGANIC PRODUCTION

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call poison control center or doctor for further treatment advice.
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.
If swallowed	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

CALL THE POISON CONTROL HOTLINE 24 HOURS A DAY AT 1-888-478-0798.

NET CONTENTS: 2 fl. oz., 4 fl. oz., 8 fl. oz., 16 fl. oz., 32 fl. oz., or 128 fl. oz.

EPA Reg. No. 71908-1
EPA Est. No. 71908-IND-001

E.I.D. PARRY (INDIA) LIMITED
Bioproducts Division
234 NSC Bose Road
Chennai (Madras) 600 001
India

USA Phone Number: (916) 973-1351

Batch Number: xxxx

33/19

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION - Harmful if absorbed through skin or if inhaled. Avoid breathing vapor. Causes moderate eye irritation. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Wear chemical resistant gloves.

Personal Protective Equipment (PPE)

Applicators and handlers must wear:

- Long-sleeved shirt
- Long pants
- Socks and shoes
- Chemical resistant gloves

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates.

For Terrestrial Uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwater or rinsate.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Do not apply this product through any type of irrigation system

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses and handlers of agricultural products. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment (PPE), notification to workers and restricted entry interval. The requirements in this box apply to the uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow any worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, water, wear coveralls, chemical resistant gloves, shoes plus socks and protective eyewear.

MODE OF ACTION:

EcoGarden™ Botanical Insecticide, Miticide, and Nematicide controls target pests on contact or by ingestion. The product acts on pests by way of repellence, anti-feedance, and interference with the molting process.

Azadirachtin, an insect growth regulator (IGR), mimics the pests' hormones and disrupts distinct stages of growth and development of insects and mites. The primary mode of action of azadirachtin is an interference with synthesis and metabolism of ecdysone and the juvenile hormone. Ecdysone is the molting hormone of insects, and azadirachtin can regulate growth leading to death before or during molting.

EcoGarden™ will provide control results comparable to the synthetic insecticide standards. **EcoGarden™** provides broad spectrum control with very low environmental impact. **EcoGarden™** provides all the benefits of azadirachtin, a proven anti-feedant, insect growth regulator (IGR), anti-ovipository, and repellent, as well as a toxin to soft bodied insect larvae.

The active ingredient in **EcoGarden™** - Azadirachtin – is a unique botanical insecticide, miticide and nematicide.

Mode of Action: Control of different orders of insects or insects in different phases of their life cycle is due to the complexity of the azadirachtin molecule and the many modes of action inherent in azadirachtin.	Anti-Feedant	Insects feed less or not at all on treated plants. Foliage is not damaged and insects ultimately starve to death.
	Insect Growth Regulator (IGR)	Insects fail to mature and reproduce, eliminating populations over time.
	Anti-ovipository	Insects do not lay eggs on treated plants. The likelihood of insect infestation is greatly decreased in treated plants. This adds a preventive aspect to your insect control.
	Repellant	Insects do not prefer treated plants.

PESTS CONTROLLED OR SUPPRESSED

Use EcoGarden™ against the following pests.

TARGET PEST SPECIES OF EcoGarden™

HEMIPTERA AND HOMOPTERA including but not limited to: true bugs including boxelder bugs, chinch bugs, lygus bugs and stink bug; lacebugs; leafhoppers including grape leafhopper, spittlebug, potato leafhopper and variegated leafhopper; mealy bugs including apple mealy bugs, citrus mealy bugs, grape mealy bugs; whiteflies including greenhouse whitefly, silverleaf whitefly and sweet potato whitefly and woolly whitefly; aphids including apple aphid, green peach aphid, melon aphid, pea aphid, potato aphid and rose aphid; psyllids including pear psyllids and scales including black scale, brown soft scale, California red scale, coffee scale, olive scale; San Jose scale, and cottony cushion scale.	LEPIDOPTERA including but not limited to: moths including European pine shoot moth, pine tip moth and Tussock moth; leafrollers including blueberry leafroller, filbert leafroller, fruittree leafroller, citrus leafminers, grape leafroller, oblique banded leafroller, omnivorous leafroller; cutworms including black cutworm and citrus cutworm; caterpillars and loopers including bagworms, budworms, cabbage looper, canker worms, case bearers, caseworms, corn earworm, diamondback moth, fruit worms, grapeleaf skeletonizer, gypsy moth, hornworms, imported cabbageworm, navel orangeworm, soybean looper, spruce budworm, tent caterpillar, tip moths, tent caterpillars, tobacco budworm, tobacco hornworm, tomato pinworm and tussock moth; armyworms including beet armyworm, fall armyworm, lawn armyworm, southern armyworm and yellow striped armyworm; webworms and leaf perforators .
COLEOPTERA including but not limited to: beetles, grubs and weevils including Asian long-horned beetle, bark beetles, black vine weevil, Colorado potato beetle, elm bark beetle, European chafer, flea beetles, Japanese beetle, June beetle, leaf beetles, Mexican bean beetle, Northern masked chafer, rose chafer and Southern masked chafer and twig girders.	DIPTERA including but not limited to: flies including Caribbean fruit fly, cherry maggots, crane fly, fungus gnat , Hessian fly, oriental fruit fly, Mediterranean fruit fly, marsh crane flies, melon fly, shore fly and walnut husk fly; leafminers including citrus leafminers and serpentine leafminers.
THYSANOPTERA including but not limited to: thrips including citrus thrips, flower thrips, gladiolus thrips, onion thrips, thrips palmi and Western flower thrips.	*Not intended for use on public health pests
ORTHOPTERA including but not limited to: crickets; grasshoppers; locusts	ACARINA Including but not limited to: mites including , red spider mites, brown mite, clover mite, conifer spider mite, European red mite, spruce spider mite, and two-spotted spider mite.
	*Not intended for use on public health pests
	HYMENOPTERA including but not limited to: sawflies including European sawflies, pear sawflies, red-headed pine sawflies, yellow-headed pin sawflies.
	*Not intended for use on public health pests
NEMATODA Nematodes (suppression)	

FOR USE ON ORNAMENTALS AND LANDSCAPE PLANTINGS

Ornamental Plants and Flowers including but not limited to:	Actinopteris, African violets*, ageratum, aglaonema, Algerian ivy, allamanda, alocasia, amaranthus, anthurium, aphelandra, arborvitae, Artemisia, aster, aucuba ilex, azalea, baby's breath, begonia, Boston fern, bougainvillea, boxwood, brachycome, cacti, calabrese, caladium, calathea, calendula, calla, camellia, carnation, ceanothus, chrysanthemum, cineraria, coleus, columbine, cotoneaster, cyclamen, daffodil, dahlia, daisy, daylily, delphinium, dianthus, dieffenbachia, dogwood, dusty miller, Easter lily, English ivy, euphorbia, fern, ficus, foliage plants, foxglove, freesia, fuchsia, gaillardia, gardenia, geranium, gerbera, gladiola, gloxinia, gypsophila, hedera, hibiscus, hyacinth, hydrangea, ilex, impatiens, iris, ivy, jasmine, lilac, lily, maidenhair fern, mandevilla, marigold, narcissus, nasturtium, orchid*, pansy, pelargonium, peony, peperomia, petunia, philodendron, phlox, photinia, pinks, pittosporum, poinsettia*, pothos, portulaca, primrose, pyracantha, rhododendron, rose*, rosemary, rubber plant, salvia, schefflera, sedum, sempervivum, snapdragon, spathiphyllum, stock, syngonium, tulip, verbena, vinca, wandering jew, yucca, zinnia *Please note that when making applications to these species, spotting of plant foliage and blossoms is possible.
Ornamental Trees and Shrubs including but not limited to:	Andromeda, arborvitae, ash, Austrian pine, azalea, beech, birch, birdsnest spruce, blue spruce, bougainvillea, boxwood, butternut, cedar, charmaecyparis, cherry, cotoneaster, crabapple, cyprus, dogwood, Douglas fir, elm, euonymus, firethorn, forsythia, hackberry, hawthorn, hemlock, hickory, holly, honey locust, horse chestnut, juniper, larch, laurel, lilac, linden, London planetree, magnolia, mandevilla, maple, mimosa, mountain ash, myrtle, oak, pachysandra, peach, photinia, pine, planetree, poplar, privet, purpleleaf wintercreeper, quince, sage, spruce, sycamore, white cedar, white pine, yew

Waxy bloom on certain ornamental plants will be reduced after an application.

Applications will remove the glaucus 'blue' coloring from evergreens such as Colorado blue spruce and Koster spruce.

FOR USE ON GARDEN CROPS, VEGETABLES, HERBS, SPICES, FRUITS AND BERRIES

Leafy Vegetables including but not limited to:	Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Endive, Kale, Lettuce, Spinach
Root Vegetables, including but not limited to:	Beet, Carrot, Horseradish, Parsnip, Potato, Radish, Sweet potato, Turnip, Yams
Fruiting Vegetables including but not limited to:	Eggplant, Pepper, Tomatillo, Tomato
Cucurbit Vegetables including but not limited to:	Cucumber, Gourd (edible), Muskmelon, Pumpkin, Squash, Watermelon, including Cantaloupe, Casaba, Gherkins, Melons (including hybrids), Zucchini
Legume Vegetables including but not limited to:	Bean, Chickpea, Lentil, Pea
Bulb Vegetables including but not limited to:	Garlic, Onion, Shallot
Berries including but not limited to:	Blackberry, Blueberry, Raspberry, Strawberry, others include: Boysenberry, Currants, Dewberry, Elderberry, Gooseberry, Loganberry
Herbs and Spices including but not limited to:	Chive, Dill, Fennel, Mustard, Sage, Sweet bay, others include: Anise, Balm, Basil, Black pepper, Borage, Caraway, Catnip, Chamomile, Coriander, Cumin, Curry leaf, Dandelion, Fenugreek, Horehound, Hyssop, Marjoram, Marigold, Mint, Nasturtium, Pennyroyal, Peppermint, Rosemary, Savory, Spearmint, Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood
Nut Trees including but not limited to:	Almond, Brazil nut, Filbert, Hickory nut, Pecan, Pistachios, Walnut
Pome Fruits including but not limited to:	Apple, Quince, or Pear (Comice varieties: DO NOT apply more than 24 fl oz/A. DO NOT apply after pink stage of flowering; test small areas of other varieties of pears for plant safety prior to full scale usage.)
Stone Fruits including but not limited to:	Apricot, Cherry, Nectarine, Peach, Plum
Citrus Fruits including but not limited to:	Grapefruit, Lemon, Lime, Orange others include: Citrus Citron, Mandarin (tangerine), Nectarine, Satsuma (orange mandarin), Tangerine

EcoGarden™ has been evaluated for phytotoxicity on a wide range of ornamentals and garden plants. However, since testing on all plant varieties is not feasible, test a small portion of the area to be treated for phytotoxicity before treating the entire area.

There are no restrictions on applying **EcoGarden™** up to the time of harvest.

SPRAY PREPARATION

EcoGarden™ is an emulsifiable concentrate to be diluted with water.

This product forms an emulsion and which separates upon extended or prolonged standing. Re-agitate to assure uniformity of the spray mixture.

Prepare only the volume needed for the intended application, and use the spray mixture within 24 hours of preparation.

TANK MIXTURES

EcoGarden™ is an emulsifiable concentrate and is compatible with commonly used pesticides and fertilizers. Always check the physical compatibility using a jar test in the correct proportions if needed.

If a broader spectrum of control is required tank-mix **EcoGarden™** with insecticides or miticides. If a rapid knockdown of heavy populations is necessary, then include an effective contact insecticide/miticide in combination with **EcoGarden™**.

Always read and follow the directions for use, precautions and limitations for use on all product labels used in combination. *Applications must follow the precautions and limitations of the most restrictive product label in the mixture. Do not exceed the dosage rates of any product.*

Select the right companion products:

IPM uses a variety of control options including biological, chemical, and cultural practices. **EcoGarden™** is botanical with growth regulator effect on insects and mites. Companion products include pyrethroids, spinosyns, microbial toxins, and chloronicotinyls that complement azadirachtin. Formulations of bifenthrin, spinosad, abamectins, and imidacloprid are effective for different pests. Select the product that has been proven to provide adequate performance for the pests you are trying to control.

Physical Incompatibility

Do not use **EcoGarden™** with Captan, Bordeaux mixture, triphenyltin hydroxide, lime sulfur, Rayplex iron or other highly alkaline materials as they can cause phytotoxicity and/or reduced efficacy on some target pests. Phytotoxicity will occur if tank-mix combinations with compounds known to be incompatible with oil-based formulations are used.

APPLICATION EQUIPMENT

Apply **EcoGarden™** with hand-operated (manual) or power spray equipment suitable for low volume and/or high volume applications. Follow the recommendations of the equipment manufacturer when using backpack sprayers, hose-end sprayers, compression (pump-up) sprayers, and other sprayers suitable for foliar applications of insecticides.

APPLICATION SCHEDULE

For the most effective control, apply **EcoGarden™** when pests are expected to appear or as soon as possible after pests appear and are in immature stages. Spray at an interval of seven (7) to ten (10) days or as the situation warrants.

During high pest infestation levels or when canopy is dense use higher dosage (use) rates and increase the spray frequency. For best results, spray in the morning or evening hours. Repeat spraying if rain occurs within two to three hours of spraying. For additional guidance, consult with your state agricultural experiment station or local extension horticulturalist/arborist for information on tactics and windows of application.

APPLICATION METHODS

Apply **EcoGarden™** as directed to any food or non-food crop up to and including the day of harvest, at a maximum rate of 1.33 fl. oz. per 1,000 sq. ft. per application.

Dilute **EcoGarden™** with water at a rate of 0.5 - 4.0 tablespoons (Tbs) per gallon of water. For hose end sprayers, set the RATE PER GALLON at the dial setting of 1 to 4 Tbs. depending on the crop and pests. Use the lower RATE PER GALLON for low to moderate infestations and use the higher specified RATE PER GALLON for severe infestations.

FOLIAR APPLICATION

USE	SPRAY CONCENTRATION%	Amounts of	Amounts of
		EcoGarden™	EcoGarden™
		Fluid Ounces (Tbs.) Per Quart	Fluid Ounces (Tbs.) Per Gallon
Including trees, shrubs, flowers, conifers, evergreens, herbaceous ornamentals, foliage plants, container-grown ornamentals & garden plants and groundcovers	Lower rate ranges of 0.25 - 0.75% vol/vol: Medium rate ranges of 0.75 - 1.25% vol/vol: Upper rate ranges of 1.25 - 1.70% vol/vol:	0.08 - 0.25 fl. oz. (1/6 - 1/2 Tbs.) 0.25 - 0.40 fl. oz. (1/2 - 5/6 Tbs.) 0.40 - 0.50 fl. oz. (5/6 - 1.0 Tbs.)	0.32 - 1.0 fl. oz. (2/3 - 2.0 Tbs.) 1.0 - 1.6 fl. oz. (2.0 Tbs. - 1 2/3 Tbs.) 1.6 - 2.0 fl. oz. (1 2/3 - 4 Tbs.)

DRENCH APPLICATION

Use **EcoGarden™** as a soil drench for effective control of soil-borne insect larvae, including soil-borne larvae of foliar pests, such as fungus gnats, nematodes, or soil borne thrips. When applying as a drench, avoid excessive leaching.

Preventive applications as a soil drench may be warranted for certain pests. Soil drench applications of azadirachtin will have a slower rate of activity because of soil absorption when compared to foliar applications of **EcoGarden™**. Target the initial application of a soil drench treatment to coincide with the early stages of young larvae and young nymphs.

Dilute **EcoGarden™** with water for concentrations of 0.4 to 0.8% volume/volume. See use rate table below. Add the required amount of **EcoGarden™** to a clean bucket with at least one-half of the water to be drenched. Agitate the mixture thoroughly and then fill with the remaining water and continue agitation until the product is thoroughly dispersed.

Drench the soil in the pot with one (1) pint of finished product dilution per 1.0 gallon of soil. For fungus gnats, use the 0.4% spray concentration. For mushroom fly maggot control, use the 0.6% volume/volume spray concentration. For leafminers and other difficult to control pests, use the 0.8% volume/volume spray concentration. Make two to three (2 - 3) applications at 10 - 14 day intervals until pest pressure has ended. With high insect pressure make applications every 5 to 6 days. Additional applications of **EcoGarden™** may be required with increased and prolonged pest infestation.

DILUTION TABLE FOR DRENCH APPLICATIONS

Gallons of Water	Amount of EcoGarden™			Application Interval
	0.4%	0.6%	0.8%	
1 gallon	1 Tbs.	1.5 Tbs.	2.0 Tbs.	10 - 14 days
1 gallon	0.5 fl. oz.	0.8 fl. oz.	1.0 fl. oz.	10 - 14 days
5 gallons	2.5 fl. oz.	4.0 fl. oz.	5.0 fl. oz.	10 - 14 days
10 gallons	5.0 fl. oz.	8.0 fl. oz.	10.0 fl. oz.	10 - 14 days

RECIRCULATORY, AEROPONIC, AND HYDROPONIC APPLICATION

Use **EcoGarden™** in recirculatory, aeroponic, or hydroponic systems for the control of foliar pests, soil borne insect larvae, including soil borne larvae of foliar pests such as fungus gnats, nematodes or soil borne thrips for interiorscapes, hydroponic, aeroponic and container plants.

Dilute **EcoGarden™** with water for concentrations of 0.1% to 0.8% volume/volume in a recirculatory or in a hydroponic liquid system. See use rate table below. Agitate the mixture thoroughly until the product is thoroughly dispersed.

For fungus gnats, use the 0.6% volume/volume concentration. For mushroom fly, maggot control, use the 0.6% volume/volume concentration. For leafminers and other difficult to control pests, use the 0.8% volume/volume concentration. Make two to three (2-3) applications at 10-14 day intervals until the pest pressure has ended. With high insect pressure applications make applications every 5 to 7 days. Additional applications of **EcoGarden™** may be required with increased and prolonged pest infestation.

DILUTION TABLE FOR RECIRCULATORY, AEROPONIC, AND HYDROPONIC APPLICATIONS

Gallons of Water	Amount of EcoGarden™					Application Interval
	0.1%	0.2%	0.4%	0.6%	0.8%	
1 gallon	1/4 Tbs.	1/2 Tbs.	1 Tbs.	1.5 Tbs.	2.0 Tbs.	7 - 14 days
1 gallon	0.14 fl. oz.	0.25 fl. oz.	0.5 fl. oz.	0.8 fl. oz.	1.0 fl. oz.	7 - 14 days
5 gallons	0.7 fl. oz.	1.3 fl. oz.	2.5 fl. oz.	4.0 fl. oz.	5.0 fl. oz.	7 - 14 days
10 gallons	1.4 fl. oz.	2.6 fl. oz.	5.0 fl. oz.	8.0 fl. oz.	10.0 fl. oz.	7 - 14 days

Preventive applications as a recirculatory system application may be warranted for certain pests.

EcoGarden™ can also be applied through sub-surface treatment equipment. Always follow manufacturer's use directions.

FOLIAR APPLICATION – SPECIFIC PESTS OF GARDEN CROPS, VEGETABLES, HERBS AND SPICES, BERRIES AND FRUIT

CROP	PESTS such as:	Dilution Rate for Sprayers	
		Fl. Oz. of product per 1,000 Sq. Ft.	Tbs. of product per 1.0 gallon of water
<i>Leafy Vegetables including but not limited to: Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Endive, Kale, Lettuce, Spinach</i>	Leafrollers, Cutworms, Loopers, Armyworms	0.19 - 0.96 fl. oz.	3/4 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Beetles, Weevils, Flies, Thrips, Mites	0.24 - 0.96 fl. oz.	1 Tbs. - 4 Tbs./gal
<i>Root Vegetables, including but not limited to: Beet, Carrot, Horseradish, Parsnip, Potato, Radish, Sweet potato, Turnip, Yams</i>	Beetles, Weevils	0.11 - 0.25 fl. oz.	1/2 Tbs. - 1 1/2 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Thrips, Mites	0.24 - 0.96 fl. oz.	1 Tbs. - 4 Tbs./gal
<i>Fruiting Vegetables including but not limited to: Eggplant, Pepper, Tomatillo, Tomato</i>	Beetles, Weevils	0.29 - 0.96 fl. oz.	2 Tbs. - 4 Tbs./gal
	Thrips	0.29 - 0.96 fl. oz.	2 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl. oz.	1 Tbs. - 4 Tbs./gal
<i>Cucurbit Vegetables including but not limited to: Cucumber, Gourd (edible), Muskmelon, Pumpkin, Squash, Watermelon, including Cantaloupe, Casaba, Gherkins, Melons (including hybrids); Zucchini</i>	Beetles, Weevils	0.29 - 0.96 fl. oz.	2 Tbs. - 4 Tbs./gal
	Thrips	0.29 - 0.96 fl. oz.	2 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl. oz.	1 Tbs. - 4 Tbs./gal
<i>Legume Vegetables including but not limited to: Bean, Chickpea, Lentil, Pea</i>	Beetles, Weevils	0.29 - 0.96 fl. oz.	2 Tbs. - 4 Tbs./gal
	Thrips	0.29 - 0.96 fl. oz.	2 Tbs. - 4 Tbs./gal

CROP	PESTS such as:	Dilution Rate for Sprayers	
		Fl. Oz. of product per 1,000 Sq. Ft.	Tbs. of product per 1.0 gallon of water
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl. oz.	1 Tbs. - 4 Tbs./gal
<i>Bulb Vegetables including but not limited to: Garlic, Onion, Shallot</i>	Beetles, Weevils	0.29 - 0.96 fl. oz.	2 Tbs. - 4 Tbs./gal
	Thrips	0.29 - 0.96 fl. oz.	2 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl. oz.	1 Tbs. - 4 Tbs./gal
<i>Berries including but not limited to: Blackberry, Blueberry, Raspberry, Strawberry, others include: Boysenberry, Currants, Dewberry, Elderberry, Gooseberry, Loganberry</i>	Beetles, Weevils	0.29 - 0.96 fl. oz.	2 Tbs. - 4 Tbs./gal
	Thrips	0.29 - 0.96 fl. oz.	2 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl. oz.	1 Tbs. - 4 Tbs./gal
<i>Herbs and Spices including but not limited to: Chive, Dill, Fennel, Mustard, Sage, Sweet bay, others include: Anise, Balm, Basil, Black pepper, Borage, Caraway, Catnip, Chamomile, Coriander, Cumin, Curry leaf, Dandelion, Fenugreek, Horehound, Hyssop, Marjoram, Marigold, Mint, Nasturtium, Pennyroyal, Peppermint, Rosemary, Savory, Spearmint, Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood</i>	Beetles, Weevils	0.29 - 0.96 fl. oz.	2 Tbs. - 4 Tbs./gal
	Thrips	0.29 - 0.96 fl. oz.	2 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl. oz.	1 Tbs. - 4 Tbs./gal
<i>Nut Trees including but not limited to: Almond, Brazil nut, Filbert, Hickory nut, Pecan, Pistachios, Walnut</i>	Beetles, Weevils	0.29 - 0.96 fl. oz.	2 Tbs. - 4 Tbs./gal
	Thrips	0.29 - 0.96 fl. oz.	2 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl. oz.	1 Tbs. - 4 Tbs./gal
<i>Pome Fruits including but not limited to: Apple, Quince, or Pear (Comice varieties: DO NOT apply more than 24 fl oz/A. DO NOT apply after pink stage of flowering; test small areas of other varieties of pears for plant safety prior to full scale usage.)</i>	Beetles, Weevils	0.29 - 0.96 fl. oz.	2 Tbs. - 4 Tbs./gal
	Thrips	0.29 - 0.96 fl. oz.	2 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl. oz.	1 Tbs. - 4 Tbs./gal
<i>Stone Fruits including but not limited to: Apricot, Cherry, Nectarine, Peach, Plum</i>	Beetles, Weevils	0.29 - 0.96 fl. oz.	2 Tbs. - 4 Tbs./gal
	Thrips	0.29 - 0.96 fl. oz.	2 Tbs. - 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl. oz.	1 Tbs. - 4 Tbs./gal
<i>Citrus Fruits including but not limited to: Grapefruit, Lemon, Lime, Orange others include: Citrus Citron, Mandarin</i>	Beetles, Weevils	0.29 - 0.96 fl. oz.	2 Tbs. - 4 Tbs./gal
	Thrips	0.29 - 0.96 fl. oz.	2 Tbs. - 4 Tbs./gal

CROP	PESTS such as:	Dilution Rate for Sprayers	
		Fl. Oz. of product per 1,000 Sq. Ft.	Tbs. of product per 1.0 gallon of water
(tangerine), Nectarine, Satsuma (orange mandarin), Tangerine	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 - 0.96 fl. oz.	.1 Tbs. - 4 Tbs./gal

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal..

PESTICIDE STORAGE: Store in original containers in a dry, cool, well-ventilated area.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill container 1/4 full with water and recap. Shake 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Drain for 10 seconds after flow begins to drip. Repeat this procedure two more times.

NOTICE ON CONDITIONS OF SALE

The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Plant injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of E.I.D. PARRY (INDIA) LIMITED. All such risks shall be assumed by the user or buyer.

Revised 11-17-08

- Controls mites, caterpillars, whiteflies, thrips, aphids, and other insects as listed on this label.
- Organic Certified
- Controls chewing and sucking insects
- Low-odor formulation
- Broad-spectrum control
- Can be applied the day of harvest
- For use on a wide variety of trees, shrubs, flowers, fruit and nut trees, garden vegetables and plants
- Low (Mild) odor
- Organic Materials Review Institute (OMRI) Certified.
- Spray and Drench
- For use with recirculatory, aeroponic, and hydroponic systems
- Contains azadirachtin from E.I.D. PARRY (INDIA) LIMITED
- 100% Vegetable Based
- Can be used in Organic Gardening  FOR ORGANIC PRODUCTION
- Tough Pests. Easy Solution.
- Controls chewing and sucking insects
- Broad spectrum insect and mite control
- Can be applied up to and on the day of harvest for all food & non food use plants
- Rose Spray
- For use on Roses and Flowers
- Vegetable Garden Spray
- For use on Tomatoes and Garden Vegetables
- Controls Japanese Beetle on Roses
- Insect and Mite Control for the Home Garden Vegetables
- Complete Insect and Mite Control for Flowers
- Three applications in 30 days provides complete plant protection
- Quadruple action insect control – Antifeedant, Insect Growth Regulator, Anti-ovipository, Repellant
- The Pure Power of PURER azadirachtin
- Single product; multiple action
- A proven botanical pesticide born out of a decade of intense global research
- Works effectively against a broad spectrum of pests on a stand alone basis on a variety of vegetable crops
- Controls insect pest during various stages of growth
- The Benefits to your insect control program are multiple
- Controls Insects Systemically
- Taken up through the roots, controls insects on the leaves
- Insects stop feeding when they feed on treated plants
- For foliar and systemic insect control
- For Use on Flowers, Ornamentals, and Landscape Plantings
- For Use on Garden Crops, Vegetables, Herbs, Spices, Fruits, and Berries
- Three in one product – Insecticide, Miticide, and Nematicide
- Recommended by General Hydroponics – Bringing Nature and Technology Together
- Brought to you Exclusively by General Hydroponics – Bringing Nature and Technology Together

EcoGarden™

Botanical Insecticide, Miticide, and Nematicide

Single Dose Delivery Cartridge (patent pending)

Read This Entire Label Before Use

ACTIVE INGREDIENT:

Azadirachtin..... 1.2%

OTHER INGREDIENTS..... 98.8%

TOTAL 100.00%

Contains 0.35 grams azadirachtin per fluid ounce

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

If on skin or clothing Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

CALL THE POISON CONTROL HOTLINE 24 HOURS A DAY AT 1-888-478-0798

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION - Harmful if absorbed through the skin. Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after handling and before eating drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse. Wear chemical resistant gloves.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water when disposing of washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

NET CONTENTS: 4 Single Dose Delivery Cartridges

Single Dose Delivery Cartridge – 2.5 milliliters (0.084 fluid ounce or 1/6 Tbs.) or 5 milliliters (0.169 fluid ounce or 1/3 Tbs.)

EPA Reg. No. 71908-1
EPA Est. No. 71908-IND-001
USA Phone Number: (913) 973-1351
Batch Number: xxxx

E.I.D. PARRY (INDIA) LIMITED
Bioproducts Division
234 NSC Bose Road
Chennai (Madras) 600 001
India

GENERAL INFORMATION

EcoGarden™ Botanical Insecticide, Miticide, & Nematicide will provide control results comparable to the synthetic insecticide standards. **EcoGarden™** provides broad spectrum control with very low environmental impact. **EcoGarden™** provides all the benefits of azadirachtin, a proven anti-feedant, insect growth regulator (IGR), anti-ovipository, and repellent, as well as a toxin to soft bodied insect larvae.

The active ingredient in **EcoGarden™** - Azadirachtin – is a unique botanical insecticide, miticide and nematicide.

Mode of Action: Control of different orders of insects or insects in different phases of their life cycle is due to the complexity of the azadirachtin molecule and the many modes of action inherent in azadirachtin.	Anti-Feedant	Insects feed less or not at all on treated plants. Foliage is not damaged and insects ultimately starve to death.
	Insect Growth Regulator (IGR)	Insects fail to mature and reproduce, eliminating populations over time.
	Anti-ovipository	Insects do not lay eggs on treated plants. The likelihood of insect infestation is greatly decreased in treated plants. This adds a preventive aspect to your insect control.
	Repellant	Insects do not prefer treated plants.

45
7a

PESTS CONTROLLED OR SUPPRESSED

HEMIPTERA AND HOMOPTERA including but not limited to:

true bugs including boxelder bugs, chinch bugs, lygus bugs and stink bug; **lacebugs**; **leafhoppers** including grape leafhopper, spittlebug, potato leafhopper and variegated leafhopper; **mealy bugs** including apple mealy bugs, citrus mealy bugs, grape mealy bugs; **whiteflies** including greenhouse whitefly, silverleaf whitefly and sweet potato whitefly and woolly whitefly; **aphids** including apple aphid, green peach aphid, melon aphid, pea aphid, potato aphid and rose aphid; **psyllids** including pear psyllids and **scales** including black scale, brown soft scale, California red scale, coffee scale, olive scale, San Jose scale, and cottony cushion scale

COLEOPTERA including but not limited to:

beetles, grubs and weevils including Asian long-horned beetle, bark beetles, black vine weevil, Colorado potato beetle, elm bark beetle, European chafer, flea beetles; Japanese beetle, June beetle, leaf beetles, Mexican bean beetle, Northern masked chafer, rose chafer and Southern masked chafer and twig girders.

THYSANOPTERA including but not limited to:

thrips including citrus thrips, flower thrips, gladiolus thrips, onion thrips, thrips palmi and Western flower thrips.

LEPIDOPTERA including but not limited to:

moths including European pine shoot moth, pine tip moth and Tussock moth; **leafrollers** including blueberry leafroller, filbert leafroller, fruitree leafroller, citrus leafminers, grape leafroller, oblique banded leafroller, omnivorous leafroller; **cutworms** including black cutworm and citrus cutworm; **caterpillars and loopers** including bagworms, budworms, cabbage looper, canker worms, case bearers, caseworms, corn earworm, diamondback moth, fruit worms, grapeleaf skeletonizer, gypsy moth, hornworms, imported cabbageworm, navel orangeworm, soybean looper, spruce budworm, tent caterpillar, tip moths, tent caterpillars, tobacco budworm, tobacco hornworm, tomato pinworm and tussock moth; **armyworms** including beet armyworm, fall armyworm, lawn armyworm, southern armyworm and yellow striped armyworm; **webworms and leaf perforators**.

DIPTERA including but not limited to*:

flies including Caribbean fruit fly, cherry maggots, crane fly, **fungus gnat**, Hessian fly, oriental fruit fly, Mediterranean fruit fly, marsh crane flies, melon fly, shore fly and walnut husk fly; **leafminers** including citrus leafminers and serpentine leafminers

*Not intended for use on public health pests

HYMENOPTERA including but not limited to*:

sawflies including European sawflies, pear sawflies, red-headed pine sawflies, yellow-headed pine sawflies.

*Not intended for use on public health pests

ORTHOPTERA including but not limited to:

crickets; grasshoppers; locusts

ACARINA including but not limited to*:

Mites including, **red spider mites**, brown mite, clover mite, conifer spider mite, European red mite, spruce spider mite, and two-spotted spider mite.

*Not intended for use on public health pests

NEMATODA:

Nematodes (suppression)

FOR USE ON FLOWERS, ORNAMENTALS AND LANDSCAPE PLANTINGS

46/79

Ornamental Plants and Flowers including but not limited to:	Actinopteris, African violets*, ageratum, aglaonema, Algerian ivy, allamanda, alocasia, amaranthus , anthurium, aphelandra, arborvitae, Artemisia, aster, aucuba ilex, azalea , baby's breath, begonia, Boston fern, bougainvillea, boxwood, brachycome, cacti, calabrese, caladium, calathea, calendula, calla, camellia, carnation , ceanothus, chrysanthemum, cineraria, coleus, columbine, cotoneaster, cyclamen, daffodil, dahlia, daisy, daylily, delphinium, dianthus, dieffenbachia, dogwood, dusty miller, Easter lily, English ivy, euphorbia, fern , ficus, foliage plants, foxglove, freesia, fuchsia, gaillardia, gardenia, geranium, gerbera, gladiola, gloxinia, gypsophilla, hedera, hibiscus, hyacinth, hydrangea, ilex, impatiens, iris, ivy, jasmine, lilac, lily, maidenhair fern, mandevilla, marigold, narcissus, nasturtium, orchid *, pansy, pelargonium, peony , peperomia, petunia, philodendron, phlox, photinia, pinks, pittosporum, poinsettia*, pothos, portulaca, primrose, pyracantha, rhododendron, rose *, rosemary, rubber plant, salvia, schefflera, sedum, sempervivum, snapdragon , spathiphyllum, stock, syngonium, tulip, verbena, vinca, wandering jew, yucca, zinnia * Please note that when making applications to these species, spotting of plant foliage and blossoms is possible.
Ornamental Trees and Shrubs including but not limited to:	Andromeda, arborvitae, ash, Austrian pine, azalea , beech, birch, birdsnest spruce, blue spruce, bougainvillea, boxwood, butternut, cedar, charmaecyparis, cherry, cotoneaster, crabapple , cyprus, dogwood, Douglas fir , elm, euonymus, firethorn, forsythia, hackberry, hawthorn, hemlock, hickory, holly, honey locust, horse chestnut, juniper, larch, laurel, lilac, linden, London planetree, magnolia , mandevilla, maple , mimosa, mountain ash, myrtle, oak , pachysandra, peach, photinia, pine, planetree, poplar, privet, purpleleaf wintercreeper, quince, sage, spruce, sycamore , white cedar, white pine, yew

Waxy bloom on certain ornamental plants will be reduced after an application.

Applications will remove the glaucus 'blue' coloring from evergreens such as Colorado blue spruce and Koster spruce.

FOR USE ON GARDEN CROPS, VEGETABLES, HERBS, SPICES, FRUITS AND BERRIES

Leafy Vegetables including but not limited to:	Broccoli , Brussels Sprouts, Cabbage, Cauliflower, Collards, Endive, Kale, Lettuce , Spinach
Root Vegetables, including but not limited to:	Beet, Carrot , Horseradish, Parsnip, Potato, Radish, Sweet potato, Turnip, Yams
Fruiting Vegetables including but not limited to:	Eggplant, Pepper, Tomatillo, Tomato
Cucurbit Vegetables including but not limited to:	Cucumber , Gourd (edible), Muskmelon, Pumpkin, Squash, Watermelon, including Cantaloupe, Casaba, Gherkins, Melons (including hybrids), Zucchini
Legume Vegetables including but not limited to:	Bean, Chickpea, Lentil, Pea
Bulb Vegetables including but not limited to:	Garlic, Onion , Shallot
Berries including but not limited to:	Blackberry, Blueberry, Raspberry, Strawberry , others include: Boysenberry, Currants, Dewberry, Elderberry, Gooseberry, Loganberry
Herbs and Spices including but not limited to:	Chive, Dill, Fennel, Mustard , Sage, Sweet bay, others include: Anise, Balm, Basil , Black pepper, Borage, Caraway, Catnip, Chamomile, Coriander, Cumin, Curry leaf, Dandelion, Fenugreek, Horehound, Hyssop , Marjoram, Marigold , Mint, Nasturtium, Pennyroyal, Peppermint, Rosemary, Savory, Spearmint, Tansy, Tarragon, Thyme , Wintergreen, Woodruff, Wormwood
Nut Trees including but not limited to:	Almond , Brazil nut, Filbert, Hickory nut, Pecan, Pistachios, Walnut
Pome Fruits including but not limited to:	Apple , Quince, or Pear (Comice varieties: DO NOT apply more than 24 fl oz/A. DO NOT apply after pink stage of flowering; test small areas of other varieties of pears for plant safety prior to full scale usage.)
Stone Fruits including but not limited to:	Apricot, Cherry , Nectarine, Peach, Plum
Citrus Fruits including but not limited to:	Grapefruit , Lemon, Lime, Orange others include: Citrus Citron, Mandarin (tangerine), Nectarine , Satsuma (orange mandarin), Tangerine

EcoGarden™ has been evaluated for phytotoxicity on a wide range of ornamentals and garden plants. However, since testing on all plant varieties is not feasible, test a small portion of the area to be treated for phytotoxicity before treating the entire area.

There are no restrictions on applying **EcoGarden™** up to the time of harvest.

APPLICATION METHODS

EcoGarden™ is an emulsifiable concentrate to be diluted with water. The product forms an emulsion and requires agitation to assure uniformity of the spray mixture.

For Use on Flower, Ornamental, Garden, and Landscape Plantings For Use on Garden Crops, Vegetables, Herbs, Spices, Fruits, and Berries

Instructions for Using Single Dose Sprayer System™

Single Dose Delivery Cartridge – 2.5 milliliters (0.084 fluid ounce or 1/6 Tbs.)

or

Single Dose Delivery Cartridge – 5 milliliters (0.169 fluid ounce or 1/3 Tbs.)

EcoGarden™ Single Dose Sprayer System*

Remove access cap from Single Dose Sprayer bottle.

Fill sprayer bottle with 32 ounces of tap water (see fill line on bottle).

Open box that contains the Single Dose Delivery Cartridges.

Remove a Single Dose Delivery Cartridge.

Insert Cartridge into access opening & tighten down the Cartridge until the inner-module detaches from Cartridge and then falls into the sprayer bottle.

Shake sprayer bottle vigorously for 20 seconds or until product is thoroughly dispersed.

Apply Product Spray Solution.

*Patent Pending -- Single Dose Delivery Cartridges can be purchased separately.

To remove inner module from sprayer bottle, take off attached trigger sprayer, shake the module through the neck of the sprayer bottle, and then out.

For the most effective control, spray **EcoGarden™** as soon as possible after pests appear and are in the immature stages. Spray at an interval of seven to ten days or as the situation warrants. During high pest infestation levels increase the spray frequency. Apply **EcoGarden™** so as to thoroughly cover both sides of the foliage. Foliar applications offer locally systemic activity against insect pests. Repeat spraying if rain occurs within two to three hours of spraying. **EcoGarden™** can be applied as directed to any food or non-food plant up to and including the day of harvest.

An application of **EcoGarden™** does not provide the quick "knock-down" of a contact poison. Usage experience has shown that with three treatments over a period of 21 to 30 days (spray every 7 to 10 days), pest control is comparable to the synthetic insecticide "standards". If **EcoGarden™** is used as a preventative treatment throughout the season, prior to insect infestation, the goal of protecting your plants will be accomplished. With high insect pressure applications should be made every 5 to 7 days. Additional applications of **EcoGarden™** may be required with increased and prolonged pest infestation.

How to Use:

- ◆ Adjust nozzle to desired spray pattern
- ◆ Hold Sprayer about 12 inches from foliage
- ◆ Spray leaves, stems, and tender new shoots
- ◆ To obtain good insect & mite control, spray both upper & lower surface of leaves. Apply spray until leaves are evenly coated but not to run off.

The spray mixture should be used within 24 hours of preparation. Any un-used solution can be drenched over the base of the plants – please refer to the drench application section of this label.

DRENCH APPLICATION FOR INDOOR AND OUTDOOR PLANTS GROWN IN CONTAINERS

Use **EcoGarden™** as a soil drench for effective control of soil-borne insect larvae, including soil-borne larvae of foliar pests, such as fungus gnats, nematodes, or soil borne thrips. When applying as a drench, avoid excessive leaching.

Preventive applications as a soil drench may be warranted for certain pests. Soil drench applications of **EcoGarden™** will have a slower rate of activity because of soil absorption when compared to foliar applications methods. Target the initial application of a soil drench treatment to coincide with the early stages of young larvae and young nymphs.

Drench the soil in the pot with one (1) pint of finished spray per 1.0 gallon of soil. Make two to three (2-3) applications at 10 to 14 day intervals until the pest pressure has ended.

48
79

STORAGE & DISPOSAL

PESTICIDE STORAGE:

Do not store this product above 105°F or below -15°F for extended periods of time. Keep containers tightly closed and in original containers when not in use. Store in original container in a cool, dry place, out of reach of children.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved water disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container.

Offer for recycling, if available.

NOTICE ON CONDITIONS OF SALE

The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Plant injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of E.I.D. PARRY (INDIA) LIMITED. All such risks shall be assumed by the user or buyer.

Revised 11-17-08

49/79
Advertising claims that may be presented on the retail container label or on the labeling accompanying the product.

- ◆ Single Dose Sprayer System (Patent Pending)
- ◆ Just add water, attach Single Dose Delivery Cartridge, and spray
- ◆ Spray and Drench
- ◆ Contains azadirachtin from E.I.D. PARRY (INDIA) LIMITED
- ◆ Packaging designed for energy and environmental conservation
- ◆ Can be used in Organic Gardening  FOR ORGANIC PRODUCTION
- ◆ Tough Pests. Easy Solution
- ◆ Organic Certified
- ◆ Organic Materials Review Institute (OMRI) Certified
- ◆ Controls chewing and sucking insects
- ◆ Broad spectrum insect and mite control
- ◆ Can be applied up to and on the day of harvest for all food & non food use plants
- ◆ Can be used on a wide variety of trees, shrubs flowers, fruit and nut trees, garden vegetables and plants.
- ◆ Rose Spray
- ◆ For use on Rose and Flowers
- ◆ Vegetable Garden Spray
- ◆ For use on Tomatoes and Garden Vegetables
- ◆ Controls Japanese Beetle on Roses
- ◆ Insect and Mite Control for the Home Garden Vegetables
- ◆ Complete Insect and Mite Control for Flowers
- ◆ Three applications in 30 days provides complete plant protection
- ◆ Quadruple action insect control – Antifeedant, Insect Growth Regulator, Anti-ovipository, Repellant
- ◆ The Pure Power of PURER azadirachtin
- ◆ Single product; multiple action
- ◆ Single product; multiple pest control
- ◆ A proven botanical pesticide born out of a decade of intense global research
- ◆ Works effectively against a broad spectrum of pests on a stand alone basis on a variety of vegetable crops
- ◆ Controls insect pest during various stages of growth
- ◆ The Benefits to your insect control program are multiple
- ◆ Low Odor Formulation
- ◆ Controls Insects Systemically
- ◆ Taken up through the roots, controls insects on the leaves
- ◆ Insects stop feeding when they feed on treated plants
- ◆ Easy to Use
- ◆ For foliar and systemic insect control
- ◆ No Measuring
- ◆ Ready to Prepare and Use
- ◆ Make fresh biopesticide spray in a matter of minutes
- ◆ Single Dose Delivery Cartridges can be purchased separately
- ◆ Ready to Make and Spray
- ◆ Eco Friendly Reusable Sprayer Bottle with 360 Degree Sprayer
- ◆ Three in one product – Insecticide, Miticide, and Nematicide
- ◆ Recommended by General Hydroponics – Bringing Nature and Technology Together

EcoGarden™

Botanical Insecticide, Miticide, and Nematicide

Single Dose Delivery Sachet
Read This Entire Label Before Use

ACTIVE INGREDIENT:	% By Wt.
Azadirachtin.....	1.2%
OTHER INGREDIENTS.....	98.8%
	TOTAL 100.00%

Contains 0.35 grams azadirachtin per fluid ounce

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
CALL THE POISON CONTROL HOTLINE 24 HOURS A DAY AT 1-888-478-0798	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution: - Harmful if absorbed through the skin. Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after handling and before eating drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse. Wear chemical resistant gloves.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water when disposing of washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

NET CONTENTS: Four Single Dose Delivery Sachets
Single DoseDelivery Sachet – 2.5 milliliters (0.084 fluid ounce or 1/6 Tbs.) and 5 milliliters (0.169 fluid ounce or 1/3 Tbs.)

EPA Reg. No. 71908-1
EPA Est. No. 71908-IND-001
USA Phone Number: (916) 973-1351
Batch Number:xxx

E.I.D. PARRY (INDIA) LIMITED
Bioproducts Division
234 NSC Bose Road
Chennai (Madras) 600 001
India

GENERAL INFORMATION

EcoGarden™ Botanical Insecticide, Miticide, & Nematicide will provide control results comparable to the synthetic insecticide standards. **EcoGarden™** provides broad spectrum control with very low environmental impact. **EcoGarden™** provides all the benefits of azadirachtin, a proven anti-feedant, insect growth regulator (IGR), anti-ovipository, and repellent, as well as a toxin to soft bodied insect larvae.

The active ingredient in **EcoGarden™** - Azadirachtin – is a unique botanical insecticide, miticide and nematicide.

Mode of Action: Control of different orders of insects or insects in different phases of their life cycle is due to the complexity of the azadirachtin molecule and the many modes of action inherent in azadirachtin.	Anti-Feedant	Insects feed less or not at all on treated plants. Foliage is not damaged and insects ultimately starve to death.
	Insect Growth Regulator (IGR)	Insects fail to mature and reproduce, eliminating populations over time.
	Anti-ovipository	Insects do not lay eggs on treated plants. The likelihood of insect infestation is greatly decreased in treated plants. This adds a preventive aspect to your insect control.
	Repellant	Insects do not prefer treated plants.

52/9

PESTS CONTROLLED OR SUPPRESSED

HEMIPTERA AND HOMOPTERA including but not limited to:

true bugs including boxelder bugs, chinch bugs, lygus bugs and stink bug; **lacebugs**; **leafhoppers** including grape leafhopper, spittlebug, potato leafhopper and variegated leafhopper; **mealy bugs** including apple mealy bugs, citrus mealy bugs, grape mealy bugs; **whiteflies** including greenhouse whitefly, silverleaf whitefly and sweet potato whitefly and woolly whitefly; **aphids** including apple aphid, green peach aphid, melon aphid, pea aphid, potato aphid and rose aphid; **psyllids** including pear psyllids and **scales** including black scale, brown soft scale, California red scale, coffee scale, olive scale, San Jose scale, and cottony cushion scale

COLEOPTERA including but not limited to:

beetles, grubs and weevils including Asian long-horned beetle, bark beetles, black vine weevil, Colorado potato beetle, elm bark beetle, European chafer, flea beetles, Japanese beetle, June beetle, leaf beetles, Mexican bean beetle, Northern masked chafer, rose chafer and Southern masked chafer and twig girders.

THYSANOPTERA including but not limited to:

thrips including citrus thrips, flower thrips, gladiolus thrips, onion thrips, thrips palmi and Western flower thrips.

LEPIDOPTERA including but not limited to:

moths including European pine shoot moth, pine tip moth and Tussock moth; **leafrollers** including blueberry leafroller, filbert leafroller, fruitree leafroller, citrus leafminers, grape leafroller, oblique banded leafroller, omnivorous leafroller; **cutworms** including black cutworm and citrus cutworm; **caterpillars and loopers** including bagworms, budworms, cabbage looper, canker worms, case bearers, caseworms, corn earworm, diamondback moth, fruit worms, grapeleaf skeletonizer, gypsy moth, hornworms, imported cabbageworm, navel orangeworm, soybean looper, spruce budworm, tent caterpillar, tip moths, tent caterpillars, tobacco budworm, tobacco hornworm, tomato pinworm and tussock moth; **armyworms** including beet armyworm, fall armyworm, lawn armyworm, southern armyworm and yellow striped armyworm; **webworms and leaf perforators**.

DIPTERA including but not limited to*:

flies including Caribbean fruit fly, cherry maggots, crane fly, **fungus gnat**, Hessian fly, oriental fruit fly, Mediterranean fruit fly, marsh crane flies, melon fly, shore fly and walnut husk fly; **leafminers** including citrus leafminers and serpentine leafminers.

*Not intended for use on public health pests

HYMENOPTERA including but not limited to*:

sawflies including European sawflies, pear sawflies, red-headed pine sawflies, yellow-headed pin sawflies.

*Not intended for use on public health pests

ORTHOPTERA including but not limited to: **crickets; grasshoppers; locusts**

ACARINA including but not limited to*:

Mites including, **red spider mites**, brown mite, clover mite, conifer spider mite, European red mite, spruce spider mite, and two-spotted spider mite.

*Not intended for use on public health pests

NEMATODA:

Nematodes (suppression)

FOR USE ON FLOWERS, ORNAMENTALS AND LANDSCAPE PLANTINGS

53/79

Ornamental Plants and Flowers including but not limited to:	Actinopteris, African violets*, ageratum, aglaonema, Algerian ivy, allamanda, alocasia, amaranthus , anthurium, aphelandra, arborvitae, Artemisia, aster, aucubá ilex, azalea , baby's breath, begonia, Boston fern, bougainvillea, boxwood, brachycome, cacti, calabrese, caladium, calathea, calendula, calla, camellia, carnation , ceanothus, chrysanthemum, cineraria, coleus, columbine, cotoneaster, cyclamen, daffodil, dahlia, daisy, daylily, delphinium, dianthus, dieffenbachia, dogwood, dusty miller, Easter lily, English ivy, euphorbia, fern , ficus, foliage plants, foxglove, freesia, fuchsia, gaillardia, gardenia, geranium, gerbera, gladiola, gloxinia, gypsophila, hedera, hibiscus, hyacinth, hydrangea, ilex, impatiens, iris, ivy, jasmine, lilac, lily, maidenhair fern, mandevilla, marigold, narcissus, nasturtium, orchid*, pansy, pelargonium, peony , peperomia, petunia, philodendron, phlox, photinia, pinks, pittosporum, poinsettia*, pothos, portulaca, primrose, pyracantha, rhododendron, rose *, rosemary, rubber plant, salvia, schefflera, sedum, sempervivum, snapdragon , spathiphyllum, stock, syngonium, tulip, verbena, vinca, wandering jew, yucca, zinnia * Please note that when making applications to these species, spotting of plant foliage and blossoms is possible.
Ornamental Trees and Shrubs including but not limited to:	Andromeda, arborvitae, ash, Austrian pine, azalea , beech, birch, birdsnest spruce, blue spruce, bougainvillea, boxwood, butternut, cedar, charmaecyparis, cherry, cotoneaster, crabapple , cyprus, dogwood, Douglas fir , elm, euonymus, firethorn, forsythia, hackberry, hawthorn, hemlock, hickory, holly, honey locust, horse chestnut, juniper, larch, laurel, lilac, linden, London planetree, magnolia , mandevilla, maple , mimosa, mountain ash, myrtle, oak , pachysandra, peach, photinia, pine, planetree, poplar, privet, purpleleaf wintercreeper, quince, sage, spruce, sycamore , white cedar, white pine, yew

Waxy bloom on certain ornamental plants will be reduced after an application.

Applications will remove the glaucus 'blue' coloring from evergreens such as Colorado blue spruce and Koster spruce.

FOR USE ON GARDEN CROPS, VEGETABLES, HERBS, SPICES, FRUITS AND BERRIES

Leafy Vegetables including but not limited to:	Broccoli , Brussels Sprouts, Cabbage, Cauliflower, Collards, Endive, Kale, Lettuce , Spinach
Root Vegetables, including but not limited to:	Beet, Carrot , Horseradish, Parsnip, Potato, Radish, Sweet potato, Turnip, Yams
Fruiting Vegetables including but not limited to:	Eggplant, Pepper, Tomatillo, Tomato
Cucurbit Vegetables including but not limited to:	Cucumber , Gourd (edible), Muskmelon, Pumpkin, Squash, Watermelon, including Cantaloupe, Casaba, Gherkins, Melons (including hybrids), Zucchini
Legume Vegetables including but not limited to:	Bean, Chickpea, Lentil, Pea
Bulb Vegetables including but not limited to:	Garlic, Onion , Shallot
Berries including but not limited to:	Blackberry, Blueberry, Raspberry, Strawberry , others include: Boysenberry, Currants, Dewberry, Elderberry, Gooseberry, Loganberry
Herbs and Spices including but not limited to:	Chive, Dill, Fennel, Mustard , Sage, Sweet bay, others include: Anise, Balm, Basil , Black pepper, Borage, Caraway, Catnip, Chamomile, Coriander, Cumin, Curry leaf, Dandelion, Fenugreek, Horehound, Hyssop , Marjoram, Marigold , Mint, Nasturtium, Pennyroyal, Peppermint, Rosemary, Savory, Spearmint, Tansy, Tarragon, Thyme , Wintergreen, Woodruff, Wormwood
Nut Trees including but not limited to:	Almond , Brazil nut, Filbert, Hickory nut, Pecan, Pistachios, Walnut
Pome Fruits including but not limited to:	Apple , Quince, or Pear (Comice varieties: DO NOT apply more than 24 fl oz/A. DO NOT apply after pink stage of flowering; test small areas of other varieties of pears for plant safety prior to full scale usage.)
Stone Fruits including but not limited to:	Apricot, Cherry , Nectarine, Peach, Plum
Citrus Fruits including but not limited to:	Grapefruit , Lemon, Lime, Orange others include: Citrus Citron, Mandarin (tangerine), Nectarine , Satsuma (orange mandarin), Tangerine

EcoGarden™ has been evaluated for phytotoxicity on a wide range of ornamentals and garden plants. However, since testing on all plant varieties is not feasible, test a small portion of the area to be treated for phytotoxicity before treating the entire area.

There are no restrictions on applying **EcoGarden™** up to the time of harvest.

APPLICATION METHODS

EcoGarden™ is an emulsifiable concentrate to be diluted with water. The product forms an emulsion and requires agitation to assure uniformity of the spray mixture.

For Use on Flower, Ornamental, Garden, and Landscape Plantings For Use on Garden Crops, Vegetables, Herbs, Spices, Fruits, and Berries

Instructions for Using Single Dose Sprayer System™

Single Dose Delivery Sachet – 2.5 milliliters (0.084 fluid ounce or 1/6 Tbs.)

or

Single Dose Delivery Sachet – 5 milliliters (0.169 fluid ounce or 1/3 Tbs.)

EcoGarden™ Single Dose Sprayer System*

Remove access cap from Single Dose Sprayer bottle.

Fill sprayer bottle with 32 ounces of tap water (see fill line on bottle).

Open box that contains the Single Dose Delivery Sachet.

Remove a Single Dose Delivery Sachet and then tear open outer sac to remove Water Soluble Sachet. ----- Note →

Insert Water Soluble Sachet into access opening & replace access cap.

Let the sprayer bottle set for 1 minute while the Water Soluble Sachet begins to dissolve.

Shake sprayer bottle for 30 seconds or until the Water Soluble Sachet is completely dissolved and product is thoroughly dispersed.

Apply Product Spray Solution.

The clear inner sachet containing the product is water soluble. Do not allow the sachet to become wet prior to adding to water. Do not handle with wet hands or wet gloves. Rough handling may cause breakage.

*Patent Pending -- Single Dose Delivery Sachets can be purchased separately.

For the most effective control, spray **EcoGarden™** as soon as possible after pests appear and are in the immature stages. Spray at an interval of seven to ten days or as the situation warrants. During high pest infestation levels increase the spray frequency. Apply **EcoGarden™** so as to thoroughly cover both sides of the foliage. Foliar applications offer locally systemic activity against insect pests. Repeat spraying if rain occurs within two to three hours of spraying. **EcoGarden™** can be applied as directed to any food or non-food plant up to and including the day of harvest.

An application of **EcoGarden™** does not provide the quick "knock-down" of a contact poison. Usage experience has shown that with three treatments over a period of 21 to 30 days (spray every 7 to 10 days), pest control is comparable to the synthetic insecticide "standards". If **EcoGarden™** is used as a preventative treatment throughout the season, prior to insect infestation, the goal of protecting your plants will be accomplished. With high insect pressure applications should be made every 5 to 7 days. Additional applications of **EcoGarden™** may be required with increased and prolonged pest infestation.

How to Use:

- ◆ Adjust nozzle to desired spray pattern
- ◆ Hold Sprayer about 12 inches from foliage
- ◆ Spray leaves, stems, and tender new shoots
- ◆ To obtain good insect & mite control, spray both upper & lower surface of leaves. Apply spray until leaves are evenly coated but not to run off.

The spray mixture should be used within 24 hours of preparation. Any un-used solution can be drenched over the base of the plants – please refer to the drench application section of this label.

DRENCH APPLICATION FOR INDOOR AND OUTDOOR PLANTS GROWN IN CONTAINERS

Use **EcoGarden™** as a soil drench for effective control of soil-borne insect larvae, including soil-borne larvae of foliar pests, such as fungus gnats, nematodes, or soil borne thrips. When applying as a drench, avoid excessive leaching.

Preventive applications as a soil drench may be warranted for certain pests. Soil drench applications of **EcoGarden™** will have a slower rate of activity because of soil absorption when compared to foliar applications methods. Target the initial application of a soil drench treatment to coincide with the early stages of young larvae and young nymphs.

Drench the soil in the pot with one (1) pint of finished spray per 1.0 gallon of soil. Make two to three (2-3) applications at 10 to 14 day intervals until the pest pressure has ended.

54
79

55/79

STORAGE & DISPOSAL

PESTICIDE STORAGE:

Do not store this product above 105°F or below -15°F for extended periods of time. Keep containers tightly closed and in original containers when not in use. Store in original container in a cool, dry place, out of reach of children.

PESTICIDE DISPOSAL:

Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Nonrefillable container. Do not reuse or refill this container.

Dispose of the empty outer packet or outer capsule in the trash, as long as the water soluble sachet is unbroken.

NOTICE ON CONDITIONS OF SALE

The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Plant injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of E.I.D. PARRY (INDIA) LIMITED. All such risks shall be assumed by the user or buyer.

Revised 11-17-08

56/79
Advertising claims that may be presented on the retail container label or on the labeling accompanying the product.

- ◆ Single Dose Sprayer System (Patent Pending)
- ◆ Just add water, insert Single Dose Delivery Sachet, and spray
- ◆ Spray and Drench
- ◆ Contains azadirachtin from E.I.D. PARRY (INDIA) LIMITED
- ◆ Packaging designed for energy and environmental conservation
- ◆ Can be used in Organic Gardening  FOR ORGANIC PRODUCTION
- ◆ Tough Pests. Easy Solution
- ◆ Organic Certified
- ◆ Organic Materials Review Institute (OMRI) Certified
- ◆ Controls chewing and sucking insects
- ◆ Broad spectrum insect and mite control
- ◆ Can be applied up to and on the day of harvest for all food & non food use plants
- ◆ Can be used on a wide variety of trees, shrubs flowers, fruit and nut trees, garden vegetables and plants.
- ◆ Rose Spray
- ◆ For use on Rose and Flowers
- ◆ Vegetable Garden Spray
- ◆ For use on Tomatoes and Garden Vegetables
- ◆ Controls Japanese Beetle on Roses
- ◆ Insect and Mite Control for the Home Garden Vegetables
- ◆ Complete Insect and Mite Control for Flowers
- ◆ Three applications in 30 days provides complete plant protection
- ◆ Quadruple action insect control – Antifeedant, Insect Growth Regulator, Anti-ovipository, Repellant
- ◆ The Pure Power of PURER azadirachtin
- ◆ Single product; multiple action
- ◆ Single product; multiple pest control
- ◆ A proven botanical pesticide born out of a decade of intense global research
- ◆ Works effectively against a broad spectrum of pests on a stand alone basis on a variety of vegetable crops
- ◆ Controls insect pest during various stages of growth
- ◆ The Benefits to your insect control program are multiple
- ◆ Low Odor Formulation
- ◆ Controls Insects Systemically
- ◆ Taken up through the roots, controls insects on the leaves
- ◆ Insects stop feeding when they feed on treated plants
- ◆ Easy to Use
- ◆ For foliar and systemic insect control
- ◆ No Measuring
- ◆ Ready to Prepare and Use
- ◆ Make fresh biopesticide spray in a matter of minutes
- ◆ Single Dose Delivery Sachets can be purchased separately
- ◆ Ready to Make and Spray
- ◆ Eco Friendly Reusable Sprayer Bottle with 360 Degree Sprayer
- ◆ Three in one product – Insecticide, Miticide, and Nematicide
- ◆ Recommended by General Hydroponics – Bringing Nature and Technology Together

EcoGarden™

Botanical Insecticide, Miticide, and Nematicide

Single Dose Delivery Vial

Read This Entire Label Before Use

ACTIVE INGREDIENT:	% By Wt.
Azadirachtin.....	1.2%
OTHER INGREDIENTS.....	98.8%
	TOTAL 100.00%

Contains 0.35 grams azadirachtin per fluid ounce

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
-------------------------------	---

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

CALL THE POISON CONTROL HOTLINE 24 HOURS A DAY AT 1-888-478-0798

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION - Harmful if absorbed through the skin. Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after handling and before eating drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse. Wear chemical resistant gloves.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water when disposing of washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

NET CONTENTS: Four Single Dose Delivery Vials

Single Dose Delivery Vial – 2.5 milliliters (0.084 fluid ounce or 1/6 Tbs.) and 5 milliliters (0.169 fluid ounce or 1/3 Tbs.)

EPA Reg. No. 71908-1
 EPA Est. No. 71908-IND-001
 USA Phone Number: (913) 973-1351
 Batch Number:xxxx

E.I.D. PARRY (INDIA) LIMITED
 Bioproducts Division
 234 NSC Bose Road
 Chennai (Madras) 600.001
 India

GENERAL INFORMATION

EcoGarden™ Botanical Insecticide, Miticide, & Nematicide will provide control results comparable to the synthetic insecticide standards. **EcoGarden™** provides broad spectrum control with very low environmental impact. **EcoGarden™** provides all the benefits of azadirachtin, a proven anti-feedant, insect growth regulator (IGR), anti-ovipository, and repellent, as well as a toxin to soft bodied insect larvae.

The active ingredient in **EcoGarden™** - Azadirachtin – is a unique botanical insecticide, miticide and nematicide.

Mode of Action: Control of different orders of insects or insects in different phases of their life cycle is due to the complexity of the azadirachtin molecule and the many modes of action inherent in azadirachtin.	Anti-Feedant	Insects feed less or not at all on treated plants. Foliage is not damaged and insects ultimately starve to death.
	Insect Growth Regulator (IGR)	Insects fail to mature and reproduce, eliminating populations over time.
	Anti-ovipository	Insects do not lay eggs on treated plants. The likelihood of insect infestation is greatly decreased in treated plants. This adds a preventive aspect to your insect control.
	Repellant	Insects do not prefer treated plants.

59
79

PESTS CONTROLLED OR SUPPRESSED

HEMIPTERA AND HOMOPTERA including but not limited to:

true bugs including boxelder bugs, chinch bugs, lygus bugs and stink bug; **lacebugs**; **leafhoppers** including grape leafhopper, spittlebug, potato leafhopper and variegated leafhopper; **mealy bugs** including apple mealy bugs, citrus mealy bugs, grape mealy bugs; **whiteflies** including greenhouse whitefly, silverleaf whitefly and sweet potato whitefly and woolly whitefly; **aphids** including apple aphid, green peach aphid, melon aphid, pea aphid, potato aphid and rose aphid; **psyllids** including pear psyllids and **scales** including black scale, brown soft scale, California red scale, coffee scale, olive scale, San Jose scale, and cottony cushion scale

COLEOPTERA including but not limited to:

beetles, grubs and weevils including Asian long-horned beetle, bark beetles, black vine weevil, Colorado potato beetle, elm bark beetle, European chafer, flea beetles, Japanese beetle, June beetle, leaf beetles, Mexican bean beetle, Northern masked chafer, rose chafer and Southern masked chafer and twig girders.

THYSANOPTERA including but not limited to:

thrips including citrus thrips, flower thrips, gladiolus thrips, onion thrips, thrips palmi and Western flower thrips.

LEPIDOPTERA including but not limited to:

moths including European pine shoot moth, pine tip moth and Tussock moth; **leafrollers** including blueberry leafroller, filbert leafroller, fruitree leafroller, citrus leafminers, grape leafroller, oblique banded leafroller, omnivorous leafroller; **cutworms** including black cutworm and citrus cutworm; **caterpillars and loopers** including bagworms, budworms, cabbage looper, canker worms, case bearers, caseworms, corn earworm, diamondback moth, fruit worms, grapeleaf skeletonizer, gypsy moth, hornworms, imported cabbageworm, navel orangeworm, soybean looper, spruce budworm, tent caterpillar, tip moths, tent caterpillars, tobacco budworm, tobacco hornworm, tomato pinworm and tussock moth; **armyworms** including beet armyworm, fall armyworm, lawn armyworm, southern armyworm and yellow striped armyworm; **webworms and leaf perforators**.

DIPTERA including but not limited to:*

flies including Caribbean fruit fly, cherry maggots, crane fly, **fungus gnat**, Hessian fly, oriental fruit fly, Mediterranean fruit fly, marsh crane flies, melon fly, shore fly and walnut husk fly; **leafminers** including citrus leafminers and serpentine leafminers

*Not intended for use on public health pests

HYMENOPTERA including but not limited to"**

sawflies including European sawflies, pear sawflies, red-headed pine sawflies, yellow-headed pin sawflies.

*Not intended for use on public health pests

ORTHOPTERA including but not limited to:

crickets; grasshoppers; locusts

ACARINA including but not limited to:*

Mites including, **red spider mites**, brown mite, clover mite, conifer spider mite, European red mite, spruce spider mite, and two-spotted spider mite.

*Not intended for use on public health pests

NEMATODA:

Nematodes (suppression)

FOR USE ON FLOWERS, ORNAMENTALS AND LANDSCAPE PLANTINGS

60/79

Ornamental Plants and Flowers including but not limited to:	Actinopteris, African violets*, ageratum, aglaonema, Algerian ivy, allamanda, alocasia, amaranthus , anthurium, aphelandra, arborvitae, Artemisia, aster, aucuba ilex, azalea , baby's breath, begonia, Boston fern, bougainvillea, boxwood, brachycome, cacti, calabrese, caladium, calathea, calendula, calla, camellia, carnation , ceanothus, chrysanthemum, cineraria, coleus, columbine, cotoneaster, cyclamen, daffodil, dahlia, daisy, daylily, delphinium, dianthus, dieffenbachia, dogwood, dusty miller, Easter lily, English ivy, euphorbia, fern , ficus, foliage plants, foxglove, freesia, fuschia, gaillardia, gardenia, geranium, gerbera, gladiola, gloxinia, gypsophilla, heder, hibiscus, hyacinth, hydrangea, ilex, impatiens, iris, ivy, jasmine, lilac, lily, maidenhair fern, mandevilla, marigold, narcissus, nasturtium, orchid*, pansy, pelargonium, peony , peperomia, petunia, philodendron, phlox, photinia, pinks, pittosporum, poinsettia*, pothos, portulaca, primrose, pyracantha, rhododendron, rose *, rosemary, rubber plant, salvia, schefflera, sedum, sempervivum, snapdragon , spathiphyllum, stock, syngonium, tulip, verbena, vinca, wandering jew, yucca, zinnia
	* Please note that when making applications to these species, spotting of plant foliage and blossoms is possible.

Waxy bloom on certain ornamental plants will be reduced after an application.

Applications will remove the glaucous 'blue' coloring from evergreens such as Colorado blue spruce and Koster spruce.

FOR USE ON GARDEN CROPS, VEGETABLES, HERBS, SPICES, FRUITS AND BERRIES

Leafy Vegetables including but not limited to:	Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Endive, Kale, Lettuce, Spinach
Root Vegetables, including but not limited to:	Beet, Carrot , Horseradish, Parsnip, Potato, Radish, Sweet potato, Turnip, Yams
Fruiting Vegetables including but not limited to:	Eggplant, Pepper, Tomatillo, Tomato
Cucurbit Vegetables including but not limited to:	Cucumber , Gourd (edible), Muskmelon, Pumpkin, Squash, Watermelon, including Cantaloupe, Casaba, Gherkins, Melons (including hybrids), Zucchini
Legume Vegetables including but not limited to:	Bean, Chickpea, Lentil, Pea
Bulb Vegetables including but not limited to:	Garlic, Onion , Shallot
Berries including but not limited to:	Blackberry, Blueberry, Raspberry, Strawberry , others include: Boysenberry, Currants, Dewberry, Elderberry, Gooseberry, Loganberry
Herbs and Spices including but not limited to:	Chive, Dill, Fennel, Mustard , Sage, Sweet bay, others include: Anise, Balm, Basil , Black pepper, Borage, Caraway, Catnip, Chamomile, Coriander, Cumin, Curry leaf, Dandelion, Fenugreek, Horehound, Hyssop , Marjoram, Marigold , Mint, Nasturtium, Pennyroyal, Peppermint, Rosemary, Savory, Spearmint, Tansy, Tarragon, Thyme , Wintergreen, Woodruff, Wormwood
Nut Trees including but not limited to:	Almond , Brazil nut, Filbert, Hickory nut, Pecan, Pistachios, Walnut
Pome Fruits including but not limited to:	Apple , Quince, or Pear (Comice varieties: <i>DO NOT apply more than 24 fl oz/A. DO NOT apply after pink stage of flowering; test small areas of other varieties of pears for plant safety prior to full scale usage.</i>)
Stone Fruits including but not limited to:	Apricot, Cherry , Nectarine, Peach, Plum
Citrus Fruits including but not limited to:	Grapefruit , Lemon, Lime, Orange others include: Citrus Citron, Mandarin (tangerine), Nectarine , Satsuma (orange mandarin), Tangerine

EcoGarden™ has been evaluated for phytotoxicity on a wide range of ornamentals and garden plants. However, since testing on all plant varieties is not feasible, test a small portion of the area to be treated for phytotoxicity before treating the entire area.

There are no restrictions on applying **EcoGarden™** up to the time of harvest.

APPLICATION METHODS

EcoGarden™ is an emulsifiable concentrate to be diluted with water. The product forms an emulsion and requires agitation to assure uniformity of the spray mixture.

For Use on Flower, Ornamental, Garden, and Landscape Plantings **For Use on Garden Crops, Vegetables, Herbs, Spices, Fruits, and Berries**

Instructions for Using Single Dose Sprayer System™

Single Dose Delivery Vial – 2.5 milliliters (0.084 fluid ounce or 1/6 Tbs.)

or

Single Dose Delivery Vial – 5 milliliters (0.169 fluid ounce or 1/3 Tbs.)

EcoGarden™ Single Dose Sprayer System*

Remove access cap from Single Dose Sprayer bottle.

Fill sprayer bottle with 32 ounces of tap water (see fill line on bottle).

Open box that contains the Single Dose Delivery Vials.

Remove a Single Dose Delivery Vial.

Pour contents of the Vial into the sprayer bottle through the access opening.

Replace access cap on the Single Dose Sprayer bottle.

Shake sprayer bottle vigorously for 20 seconds or until product is thoroughly dispersed.

Apply Product Spray Solution.

*Patent Pending – Single Dose Delivery Vials can be purchased separately.

For the most effective control, spray **EcoGarden™** as soon as possible after pests appear and are in the immature stages. Spray at an interval of seven to ten days or as the situation warrants. During high pest infestation levels increase the spray frequency. Apply **EcoGarden™** so as to thoroughly cover both sides of the foliage. Foliar applications offer locally systemic activity against insect pests. Repeat spraying if rain occurs within two to three hours of spraying. **EcoGarden™** can be applied as directed to any food or non-food plant up to and including the day of harvest.

An application of **EcoGarden™** does not provide the quick "knock-down" of a contact poison. Usage experience has shown that with three treatments over a period of 21 to 30 days (spray every 7 to 10 days), pest control is comparable to the synthetic insecticide "standards". If **EcoGarden™** is used as a preventative treatment throughout the season, prior to insect infestation, the goal of protecting your plants will be accomplished. With high insect pressure applications should be made every 5 to 7 days. Additional applications of **EcoGarden™** may be required with increased and prolonged pest infestation.

How to Use:

- ◆ Adjust nozzle to desired spray pattern
- ◆ Hold Sprayer about 12 inches from foliage
- ◆ Spray leaves, stems, and tender new shoots
- ◆ To obtain good insect & mite control, spray both upper & lower surface of leaves. Apply spray until leaves are evenly coated but not to run off.

The spray mixture should be used within 24 hours of preparation. Any un-used solution can be drenched over the base of the plants – please refer to the drench application section of this label.

DRENCH APPLICATION FOR INDOOR AND OUTDOOR PLANTS GROWN IN CONTAINERS

Use **EcoGarden™** as a soil drench for effective control of soil-borne insect larvae, including soil-borne larvae of foliar pests, such as fungus gnats, nematodes, or soil borne thrips. When applying as a drench, avoid excessive leaching.

Preventive applications as a soil drench may be warranted for certain pests. Soil drench applications of **EcoGarden™** will have a slower rate of activity because of soil absorption when compared to foliar applications methods. Target the initial application of a soil drench treatment to coincide with the early stages of young larvae and young nymphs.

Drench the soil in the pot with one (1) pint of finished spray per 1.0 gallon of soil. Make two to three (2-3) applications at 10 to 14 day intervals until the pest pressure has ended.

62
79

STORAGE & DISPOSAL

PESTICIDE STORAGE:

Do not store this product above 105°F or below -15°F for extended periods of time. Keep containers tightly closed and in original containers when not in use. Store in original container in a cool, dry place, out of reach of children.

PESTICIDE DISPOSAL:

Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Nonrefillable container. Do not reuse or refill this container.

Offer for recycling, if available.

NOTICE ON CONDITIONS OF SALE

The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Plant injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of E.I.D. PARRY (INDIA) LIMITED. All such risks shall be assumed by the user or buyer.

Submitted to EPA: 10-6-08

63/79

Advertising claims that may be presented on the retail container label or on the labeling accompanying the product.

- ◆ Spray and Drench
- ◆ Contains azadirachtin from E.I.D. PARRY (INDIA) LIMITED
- ◆ Packaging designed for energy and environmental conservation
- ◆ Can be used in Organic Gardening  FOR ORGANIC PRODUCTION
- ◆ Tough Pests. Easy Solution
- ◆ Organic Certified
- ◆ Organic Materials Review Institute (OMRI) Certified
- ◆ Controls chewing and sucking insects
- ◆ Broad spectrum insect and mite control
- ◆ Can be applied up to and on the day of harvest for all food & non food use plants
- ◆ Can be used on a wide variety of trees, shrubs flowers, fruit and nut trees, garden vegetables and plants.
- ◆ Rose Spray
- ◆ For use on Rose and Flowers
- ◆ Vegetable Garden Spray
- ◆ For use on Tomatoes and Garden Vegetables
- ◆ Controls Japanese Beetle on Roses
- ◆ Insect and Mite Control for the Home Garden Vegetables
- ◆ Complete Insect and Mite Control for Flowers
- ◆ Three applications in 30 days provides complete plant protection
- ◆ Quadruple action insect control – Antifeedant, Insect Growth Regulator, Anti-ovipository, Repellent
- ◆ The Pure Power of PURER azadirachtin
- ◆ Single product; multiple action
- ◆ Single product; multiple pest control
- ◆ A proven botanical pesticide born out of a decade of intense global research
- ◆ Works effectively against a broad spectrum of pests on a stand alone basis on a variety of vegetable crops
- ◆ Controls insect pest during various stages of growth
- ◆ The Benefits to your insect control program are multiple
- ◆ Low Odor Formulation
- ◆ Controls Insects Systemically
- ◆ Taken up through the roots, controls insects on the leaves
- ◆ Insects stop feeding when they feed on treated plants
- ◆ Easy to Use
- ◆ For foliar and systemic insect control
- ◆ No Measuring
- ◆ Ready to Prepare and Use
- ◆ Make fresh biopesticide spray in a matter of minutes
- ◆ Single Dose Delivery Vials can be purchased separately
- ◆ Ready to Make and Spray
- ◆ Eco Friendly Reusable Sprayer Bottle with 360 Degree Sprayer
- ◆ Three in one product – Insecticide, Miticide, and Nematicide
- ◆ Recommended by General Hydroponics – Bringing Nature and Technology Together

Revised 11-17-08

EcoGarden™
Botanical Insecticide, Miticide, and Nematicide

Water Soluble Sachet
Read This Entire Label Before Use

ACTIVE INGREDIENT:	% By Wt.
Azadirachtin.....	1.2%
OTHER INGREDIENTS.....	98.8%
	TOTAL 100.00%

Contains 0.35 grams azadirachtin per fluid ounce

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

If on skin or clothing Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

CALL THE POISON CONTROL HOTLINE 24 HOURS A DAY AT 1-888-478-0798

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through the skin. Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after handling and before eating drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse. Wear chemical resistant gloves.

Personal Protective Equipment (PPE)

Applicators and handlers must wear:

- Long-sleeved shirt
- Long pants
- Socks and shoes
- Chemical resistant gloves

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

NET CONTENTS:

Water Soluble Sachet – 10 milliliters (0.338 fluid ounce or 2/3 Tbs.)
6, 12, 24, 48 or 96 numbers of sachets per package.

EPA Reg. No. 71908-1
EPA Est. No. 71908-IND-001
USA Phone Number: (916) 973-1351
Batch Number: xxxx

E.I.D. PARRY (INDIA) LIMITED
Bioproducts Division
234 NSC Bose Road
Chennai (Madras) 600 001
India

USER SAFETY RECOMMENDATIONS

65/19
Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. For terrestrial uses: Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

GENERAL INFORMATION

EcoGarden™ Botanical Insecticide, Miticide, & Nematicide will provide control results comparable to the synthetic insecticide standards. **EcoGarden™** provides broad spectrum control with very low environmental impact. **EcoGarden™** provides all the benefits of azadirachtin, a proven anti-feedant, insect growth regulator (IGR), anti-ovipository, and repellent, as well as a toxin to soft bodied insect larvae. Do not exceed 120 grams of active ingredient (azadirachtin) per acre per application. Each sachet contains 0.1183 grams of active ingredient, azadirachtin. Do not apply this product through any type of irrigation system.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment (PPE), notification to workers and restricted entry interval. The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow any worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

For early entry into the treated areas during the restricted entry interval (REI) of 4 hours.

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, water, wear: coveralls, chemical resistant gloves and shoes plus socks.

The active ingredient in **EcoGarden™** - Azadirachtin – is a unique botanical insecticide, miticide and nematicide.

Mode of Action: Control of different orders of insects or insects in different phases of their life cycle is due to the complexity of the azadirachtin molecule and the many modes of action inherent in azadirachtin.	Anti-Feedant	Insects feed less or not at all on treated plants. Foliage is not damaged and insects ultimately starve to death.
	Insect Growth Regulator (IGR)	Insects fail to mature and reproduce, eliminating populations over time.
	Anti-ovipository	Insects do not lay eggs on treated plants. The likelihood of insect infestation is greatly decreased in treated plants. This adds a preventive aspect to your insect control.
	Repellant	Insects do not prefer treated plants.

66
79

PESTS CONTROLLED OR SUPPRESSED

HEMIPTERA AND HOMOPTERA including but not limited to:

true bugs including boxelder bugs, chinch bugs, lygus bugs and stink bug; **lacebugs**; **leafhoppers** including grape leafhopper, spittlebug, potato leafhopper and variegated leafhopper; **mealy bugs** including apple mealy bugs, citrus mealy bugs, grape mealy bugs; **whiteflies** including greenhouse whitefly, silverleaf whitefly and sweet potato whitefly and woolly whitefly; **aphids** including apple aphid, green peach aphid, melon aphid, pea aphid, potato aphid and rose aphid; **psyllids** including pear psyllids and **scales** including black scale, brown soft scale, California red scale, coffee scale, olive scale, San Jose scale, and cottony cushion scale

COLEOPTERA including but not limited to:

beetles, grubs and weevils including Asian long-horned beetle, bark beetles, black vine weevil, Colorado potato beetle, elm bark beetle, European chafer, flea beetles, Japanese beetle, June beetle, leaf beetles, Mexican bean beetle, Northern masked chafer, rose chafer and Southern masked chafer and twig girders.

THYSANOPTERA including but not limited to:

thrips including citrus thrips, flower thrips, gladiolus thrips, onion thrips, thrips palmi and Western flower thrips.

LEPIDOPTERA including but not limited to:

moths including European pine shoot moth, pine tip moth and Tussock moth; **leafrollers** including blueberry leafroller, filbert leafroller, fruitree leafroller, citrus leafminers, grape leafroller, oblique banded leafroller, omnivorous leafroller; **cutworms** including black cutworm and citrus cutworm; **caterpillars and loopers** including bagworms, budworms, cabbage looper, canker worms, case bearers, caseworms, corn earworm, diamondback moth, fruit worms, grapeleaf skeletonizer, gypsy moth, hornworms, imported cabbageworm, navel orangeworm, soybean looper, spruce budworm, tent caterpillar, tip moths, tent caterpillars, tobacco budworm, tobacco hornworm, tomato pinworm and tussock moth; **armyworms** including beet armyworm, fall armyworm, lawn armyworm, southern armyworm and yellow striped armyworm; **webworms** and **leaf perforators**.

DIPTERA including but not limited to: *

flies including Caribbean fruit fly, cherry maggots, crane fly, **fungus gnat**, Hessian fly, oriental fruit fly, Mediterranean fruit fly, marsh crane flies, melon fly, shore fly and walnut husk fly; **leafminers** including citrus leafminers and serpentine leafminers

*Not intended for use on public pests

HYMENOPTERA including but not limited to: *

sawflies including European sawflies, pear sawflies, red-headed pine sawflies, yellow-headed pin sawflies.

*Not intended for use on public pests

ORTHOPTERA including but not limited to:

crickets; grasshoppers; locusts

ACARINA including but not limited to: *

Mites including, **red spider mites**, brown mite, clover mite, conifer spider mite, European red mite, spruce spider mite, and two-spotted spider mite.

*Not intended for use on public pests

NEMATODA:**Nematodes (suppression)**67
79**FOR USE ON FLOWERS, ORNAMENTALS AND LANDSCAPE PLANTINGS**

Ornamental Plants and Flowers including but not limited to:	Actinopteris, African violets*, ageratum, aglaonema, Algerian ivy, allamanda, alocasia, amaranthus , anthurium, aphelandra, arborvitae, Artemisia, aster, aucuba ilex , azalea , baby's breath, begonia, Boston fern, bougainvillea, boxwood, brachycome, cacti, calabrese, caladium, calathea, calendula, calla, camellia, carnation , ceanothus, chrysanthemum, cineraria, coleus, columbine, cotoneaster, cyclamen, daffodil, dahlia, daisy, daylily, delphinium, dianthus, dieffenbachia, dogwood, dusty miller, Easter lily, English ivy, euphorbia, fern , ficus , foliage plants, foxglove, freesia, fuchsia, gaillardia, gardenia, geranium, gerbera, gladiola, gloxinia, gypsophila, hedera, hibiscus, hyacinth, hydrangea, ilex , impatiens, iris, ivy, jasmine, lilac, lily, maidenhair fern, mandevilla, marigold, narcissus, nasturtium, orchid*, pansy, pelargonium, peony , peperomia, petunia, philodendron, phlox, photinia, pinks, pittosporum, poinsettia*, pothos, portulaca, primrose, pyracantha, rhododendron, rose *, rosemary, rubber plant, salvia, schefflera, sedum, sempervivum, snapdragon , spathiphyllum, stock, syngonium, tulip, verbena, vinca, wandering jew, yucca, zinnia * Please note that when making applications to these plants spotting of foliage and blossoms is possible.
Ornamental Trees and Shrubs including but not limited to:	Andromeda, arborvitae, ash, Austrian pine, azalea , beech, birch, birdsnest spruce, blue spruce, bougainvillea, boxwood, butternut, cedar, charmaecyparis, cherry, cotoneaster, crabapple , cyprus, dogwood, Douglas fir , elm, euonymus, firethorn, forsythia, hackberry, hawthorn, hemlock, hickory, holly, honey locust, horse chestnut, juniper, larch, laurel, lilac, linden, London planetree, magnolia , mandevilla, maple , mimosa, mountain ash, myrtle, oak , pachysandra, peach, photinia, pine, planetree, poplar, privet, purpleleaf wintercreeper, quince, sage, spruce, sycamore , white cedar, white pine, yew

Waxy bloom on certain ornamental plants will be reduced after an application.

Applications will remove the glaucous 'blue' coloring from evergreens such as Colorado blue spruce and Koster spruce.

FOR USE ON GARDEN CROPS, VEGETABLES, HERBS, SPICES, FRUITS AND BERRIES

Leafy Vegetables including but not limited to:	Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Endive, Kale, Lettuce , Spinach
Root Vegetables, including but not limited to:	Beet, Carrot , Horseradish, Parsnip, Potato, Radish, Sweet potato, Turnip, Yams
Fruiting Vegetables including but not limited to:	Eggplant, Pepper, Tomatillo, Tomato
Cucurbit Vegetables including but not limited to:	Cucumber , Gourd (edible), Muskmelon, Pumpkin, Squash, Watermelon, including Cantaloupe, Casaba, Gherkins, Melons (including hybrids), Zucchini
Legume Vegetables including but not limited to:	Bean, Chickpea, Lentil, Pea
Bulb Vegetables including but not limited to:	Garlic, Onion , Shallot
Berries including but not limited to:	Blackberry, Blueberry, Raspberry, Strawberry , others include: Boysenberry, Currants, Dewberry, Elderberry, Gooseberry, Loganberry
Herbs and Spices including but not limited to:	Chive, Dill, Fennel, Mustard , Sage, Sweet bay, others include: Anise, Balm, Basil , Black pepper, Borage, Caraway, Catnip, Chamomile, Coriander, Cumin, Curry leaf, Dandelion, Fenugreek, Horehound, Hyssop , Marjoram, Marigold , Mint, Nasturtium, Pennyroyal, Peppermint, Rosemary, Savory, Spearmint, Tansy, Tarragon, Thyme , Wintergreen, Woodruff, Wormwood
Nut Trees including but not limited to:	Almond, Brazil nut, Filbert, Hickory nut, Pecan, Pistachios, Walnut
Pome Fruits including but not limited to:	Apple , Quince, or Pear (Comice varieties: DO NOT apply more than 24 fl oz/A. DO NOT apply after pink stage of flowering; test small areas of other varieties of pears for plant safety prior to full scale usage.)
Stone Fruits including but not limited to:	Apricot, Cherry , Nectarine, Peach, Plum
Citrus Fruits including but not limited to:	Grapefruit , Lemon, Lime, Orange others include: Citrus Citron, Mandarin (tangerine), Nectarine , Satsuma (orange mandarin), Tangerine

68/79
EcoGarden™ has been evaluated for phytotoxicity on a wide range of ornamentals and garden plants. However, since testing on all plant varieties is not feasible, test a small portion of the area to be treated for phytotoxicity before treating the entire area.

There are no restrictions on applying **EcoGarden™** up to the time of harvest.

APPLICATION METHODS

EcoGarden™ is an emulsifiable concentrate to be diluted with water. The product forms an emulsion and requires agitation to assure uniformity of the spray mixture.

For Use on Flower, Ornamental, Garden, and Landscape Plantings

For Use on Garden Crops, Vegetables, Herbs, Spices, Fruits, and Berries:

Instructions for Using Water Soluble Sachet

For Foliar Applications

For Foliar Applications apply **EcoGarden™** with hand-operated (manual) or power spray equipment suitable for low volume. Follow the recommendations of the equipment manufacturer when using backpack sprayers, hose-end sprayers, compression (pump-up) sprayers, and other sprayers suitable for foliar applications of insecticides.

Open the outer packet that contains the **EcoGarden™** Water Soluble Sachet. Within each outer packet is an inner sachet containing the **EcoGarden™** Water Soluble Sachet. The clear inner sachet containing the product is water soluble. Do not allow the sachet to become wet prior to adding to water. Do not handle with wet hands or wet gloves. Rough handling may cause breakage.

Drop the required number (see use rates below) of **EcoGarden™** Water Soluble Sachet into a clean bucket or sprayer tank with at least one-half of the water to be sprayed. Agitate the mixture thoroughly for 60 seconds and then fill with the remaining water and continue agitation for another 60 seconds or until the Water Soluble Sachet is dissolved and product is thoroughly dispersed. Prepare only the volume needed for the intended application and use spray mixture with 24 hours of preparation.

For the most effective control, spray **EcoGarden™** as soon as possible after pests appear and are in the immature stages. Spray at an interval of seven to ten days or as the situation warrants. During high pest infestation levels increase the spray frequency. Apply **EcoGarden™** so as to thoroughly cover both sides of the foliage. Foliar applications offer locally systemic activity against insect pests. Repeat spraying if rain occurs within two to three hours of spraying. **EcoGarden™** can be applied as directed to any food or non-food plant up to and including the day of harvest.

An application of **EcoGarden™** does not provide the quick "knock-down" of a contact poison. Usage experience has shown that with three treatments over a period of 21 to 30 days, pest control is comparable to the synthetic insecticide "standards". If **EcoGarden™** is used as a preventative treatment throughout the season, prior to insect infestation, the goal of protecting your plants will be accomplished. Additional applications of **EcoGarden™** may be required with increased and prolonged pest infestation.

Any un-used solution can be drenched over the base of the plants – please refer to the drench application section of this label.

Use Rates - Add one (1) to two (2) Water Soluble Sachets for each one (1) gallon of water solution used

For Drench Applications

Use **EcoGarden™** as a soil drench for effective control of foliar pests, and soil-borne insect larvae, including soil-borne larvae of foliar pests such as fungus gnats, nematodes, or soil borne thrips in interiorscapes, pot, container and garden plants.

Drop the required number (see use rate table below) of **EcoGarden™** Water Soluble Sachet into a clean bucket with at least one-half of the water to be drenched. Agitate the mixture thoroughly for 60 seconds and then fill with the remaining water and continue agitation for another 60 seconds or until the Water Soluble Sachet is dissolved and product is thoroughly dispersed.

Drench the soil in the pot with one (1) pint of finished product dilution per one (1) gallon of soil. For fungus gnats, use the 0.6% volume/volume concentration. For mushroom fly maggot control, use the 0.6% volume/volume concentration. For leafminers and other difficult to control pests, use the 0.8% volume/volume concentration. Make two to three (2-3) applications at 10-14 day intervals until the pest pressure has ended. With high insect pressure make applications every 5 to 7 days. Additional applications of **EcoGarden™** may be required with increased and prolonged pest infestation.

69/79

Use Rate Table - Dilutions for Water Soluble Sachets

Gallons of Water	Amount of EcoGarden™			Application Interval
	0.3%	0.6%	0.8%	
1 gallon	1 sachet	2 sachets	3 sachets	10 - 14 days
5 gallons	6 sachets	12 sachets	15 sachets	10 - 14 days
10 gallons	12 sachets	24 sachets	30 sachets	10 - 14 days

Preventive applications as a soil drench application may be warranted for certain pests. Soil drench applications of EcoGarden™ will have a slower rate of activity because of soil absorption when compared to foliar applications of EcoGarden™. Target the initial application of a soil drench treatment to coincide with the early stages of young larvae and young nymphs. When applying as a soil drench, avoid excessive leaching.

For Recirculatory, Aeroponic, and Hydroponic Applications

Use EcoGarden™ in recirculatory, aeroponic, or hydroponic systems for the control of foliar pests, soil borne insect larvae, including soil borne larvae of foliar pests such as fungus gnats, nematodes or soil borne thrips for interiorscapes, hydroponic, aeroponic, and container plants.

Drop the required number (see use rate table below) of EcoGarden™ Water Soluble Sachet into the recirculatory or in the hydroponic liquid system. Agitate the mixture thoroughly for 60 seconds or until the Water Soluble Sachet is dissolved and product is thoroughly dispersed.

For fungus gnats, use the 0.6% volume/volume concentration. For mushroom fly maggot control, use the 0.6% volume/volume concentration. For leafminers and other difficult to control pests, use the 0.8% volume/volume concentration. Make two to three (2-3) applications at 10-14 day intervals until the pest pressure has ended. With high insect pressure make applications every 5 to 7 days. Additional applications of EcoGarden™ may be required with increased and prolonged pest infestation.

Use Rate Table - Dilutions for Water Soluble Sachets

Gallons of Water	Amount of EcoGarden™					Application Interval
	0.1%	0.2%	0.4%	0.6%	0.8%	
1 1/2 gallons		1 sachet	2 sachets	3 sachets	4 sachets	7 - 14 days
3 gallons	1 sachet	2 sachets	5 sachets	7 sachets	9 sachets	7 - 14 days
6 gallons	2 sachets	4 sachets	10 sachets	14 sachets	18 sachets	7 - 14 days
9 gallons	3 sachets	6 sachets	15 sachets	21 sachets	27 sachets	7 - 14 days
12 gallons	4 sachets	8 sachets	20 sachets	28 sachets	36 sachets	7 - 14 days

Preventive applications as a recirculatory or hydroponic system application may be warranted for certain pests. Target the initial application of a recirculatory or hydroponic system treatment to coincide with the early stages of young larvae and young nymphs.

EcoGarden™ can also be applied through sub-surface treatment equipment. Always follow manufacturer's use directions.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal

PESTICIDE STORAGE:

Do not store this product above 105°F or below -15°F for extended periods of time. Keep containers tightly closed and in original containers when not in use. Store in original container in a cool, dry place, out of reach of children.

PESTICIDE DISPOSAL:

Wastes resulting from the use of this product must be disposed of on site or at an approved waste facility.

CONTAINER DISPOSAL:

Nonrefillable container. Do not reuse or refill this container.

Dispose of the empty outer packet or outer capsule in the trash, as long as the water soluble sachet is unbroken.

NOTICE ON CONDITIONS OF SALE

The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Plant injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of E.I.D. PARRY (INDIA) LIMITED. All such risks shall be assumed by the user or buyer.

71/79
Advertising claims that may be presented on the retail container label or on the labeling accompanying the product.

- ◆ Spray and Drench
- ◆ Contains azadirachtin from E.I.D. PARRY (INDIA) LIMITED
- ◆ Packaging designed for energy and environmental conservation
- ◆ Can be used in Organic Gardening  FOR ORGANIC PRODUCTION
- ◆ Tough Pests. Easy Solution.
- ◆ Organic Certified
- ◆ Organic Materials Review Institute (OMRI) Certified
- ◆ Controls chewing and sucking insects
- ◆ Broad spectrum insect and mite control
- ◆ Can be applied up to and on the day of harvest for all food & non food use plants
- ◆ Can be used on a wide variety of trees, shrubs flowers, fruit and nut trees, garden vegetables and plants.
- ◆ Rose Spray
- ◆ For use on Rose and Flowers
- ◆ Vegetable Garden Spray
- ◆ For use on Tomatoes and Garden Vegetables
- ◆ Controls Japanese Beetle on Roses
- ◆ Insect and Mite Control for the Home Garden Vegetables
- ◆ Complete Insect and Mite Control for Flowers
- ◆ Three applications in 30 days provides complete plant protection
- ◆ Quadruple action insect control – Antifeedant, Insect Growth Regulator, Anti-ovipository, Repellent
- ◆ The Pure Power of PURER azadirachtin
- ◆ Single product; multiple action
- ◆ Single product; multiple pest control
- ◆ A proven botanical pesticide born out of a decade of intense global research
- ◆ Works effectively against a broad spectrum of pests on a stand alone basis on a variety of vegetable crops
- ◆ Controls insect pest during various stages of growth
- ◆ The Benefits to your insect control program are multiple
- ◆ Low Odor Formulation
- ◆ Controls Insects Systemically
- ◆ Taken up through the roots, controls insects on the leaves
- ◆ Insects stop feeding when they feed on treated plants
- ◆ Easy to Use
- ◆ For foliar and systemic insect control
- ◆ No Measuring
- ◆ Ready to Prepare and Use
- ◆ Make fresh biopesticide spray in a matter of minutes
- ◆ Ready to Make and Spray
- ◆ Three in one product – Insecticide, Miticide, and Nematicide
- ◆ Recommended by General Hydroponics – Bringing Nature and Technology Together

Revised 11-17-08

EcoGarden™

Botanical Insecticide, Miticide, and Nematicide

Water Soluble Capsule
Read This Entire Label Before Use

ACTIVE INGREDIENT:	% By Wt.
Azadirachtin.....	1.2%
OTHER INGREDIENTS.....	98.8%
	TOTAL 100.00%

KEEP OUT OF REACH OF CHILDREN

Contains 0.35 grams azadirachtin per fluid ounce

CAUTION

FIRST AID

If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
------------------------	---

Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
---	--

CALL THE POISON CONTROL HOTLINE 24 HOURS A DAY AT 1-888-478-0798	
--	--

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION - Harmful if absorbed through the skin. Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after handling and before eating drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse. Wear chemical resistant gloves.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water when cleaning equipment or disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

NET CONTENTS:

Water Soluble Capsule – 1 milliliter (0.034 fluid ounce or 1/15 Tbs.)
20, 40, 60, 80 or 100 number of capsules per package

EPA Reg. No. 71908-1
EPA Est. No. 71908-IND-001
USA Phone Number: (916) 973-1351
Batch Number:xxxx

E.I.D. PARRY (INDIA) LIMITED
Bioproducts Division
234 NSC Bose Road
Chennai (Madras) 600 001
India

GENERAL INFORMATION

EcoGarden™ Botanical Insecticide, Miticide, & Nematicide will provide control results comparable to the synthetic insecticide standards. **EcoGarden™** provides broad spectrum control with very low environmental impact. **EcoGarden™** provides all the benefits of azadirachtin, a proven anti-feedant, insect growth regulator (IGR), anti-ovipository, and repellent, as well as a toxin to soft bodied insect larvae.

The active ingredient in **EcoGarden™** - Azadirachtin – is a unique botanical insecticide, miticide and nematicide.

Mode of Action: Control of different orders of insects or insects in different phases of their life cycle is due to the complexity of the azadirachtin molecule and the many modes of action inherent in azadirachtin.	Anti-Feedant	Insects feed less or not at all on treated plants. Foliage is not damaged and insects ultimately starve to death.
	Insect Growth Regulator (IGR)	Insects fail to mature and reproduce, eliminating populations over time.
	Anti-ovipository	Insects do not lay eggs on treated plants. The likelihood of insect infestation is greatly decreased in treated plants. This adds a preventive aspect to your insect control.
	Repellant	Insects do not prefer treated plants.

74
79

PESTS CONTROLLED OR SUPPRESSED

HEMIPTERA AND HOMOPTERA including but not limited to:

true bugs including boxelder bugs, chinch bugs, lygus bugs and stink bug; **lacebugs**; **leafhoppers** including grape leafhopper, spittlebug, potato leafhopper and variegated leafhopper; **mealy bugs** including apple mealy bugs, citrus mealy bugs, grape mealy bugs; **whiteflies** including greenhouse whitefly, silverleaf whitefly and sweet potato whitefly and woolly whitefly; **aphids** including apple aphid, green peach aphid, melon aphid, pea aphid, potato aphid and rose aphid; **psyllids** including pear psyllids and **scales** including black scale, brown soft scale, California red scale, coffee scale, olive scale, San Jose scale, and cottony cushion scale

COLEOPTERA including but not limited to:

beetles, grubs and weevils including Asian long-horned beetle, bark beetles, black vine weevil, Colorado potato beetle, elm bark beetle, European chafer, flea beetles, Japanese beetle, June beetle, leaf beetles, Mexican bean beetle, Northern masked chafer, rose chafer and Southern masked chafer and twig girders.

THYSANOPTERA including but not limited to:

thrips including citrus thrips, flower thrips, gladiolus thrips, onion thrips, thrips palmi and Western flower thrips.

LEPIDOPTERA including but not limited to:

moths including European pine shoot moth, pine tip moth and Tussock moth; **leafrollers** including blueberry leafroller, filbert leafroller, fruitree leafroller, citrus leafminers, grape leafroller, oblique banded leafroller, omnivorous leafroller; **cutworms** including black cutworm and citrus cutworm; **caterpillars and loopers** including bagworms, budworms, cabbage looper, canker worms, case bearers, caseworms, corn earworm, diamondback moth, fruit worms, grapeleaf skeletonizer, gypsy moth, hornworms, imported cabbageworm, navel orangeworm, soybean looper, spruce budworm, tent caterpillar, tip moths, tent caterpillars, tobacco budworm, tobacco hornworm, tomato pinworm and tussock moth; **armyworms** including beet armyworm, fall armyworm, lawn armyworm, southern armyworm and yellow striped armyworm; **webworms and leaf perforators**.

DIPTERA including but not limited to:*

flies including Caribbean fruit fly, cherry maggots, crane fly, **fungus gnat**, Hessian fly, oriental fruit fly, Mediterranean fruit fly, marsh crane flies, melon fly, shore fly and walnut husk fly; **leafminers** including citrus leafminers and serpentine leafminers

*Not intended for use on public health pests

HYMENOPTERA including but not limited to:*

sawflies including European sawflies, pear sawflies, red-headed pine sawflies, yellow-headed pin sawflies.

*Not intended for use on public health pests

ORTHOPTERA including but not limited to:

crickets; grasshoppers; locusts

ACARINA including but not limited to:*

Mites including, **red spider mites**, brown mite, clover mite, conifer spider mite, European red mite, spruce spider mite, and two-spotted spider mite.

*Not intended for use on public health pests

NEMATODA:

Nematodes (suppression)

FOR USE ON FLOWERS, ORNAMENTALS AND LANDSCAPE PLANTINGS

Ornamental Plants and Flowers including but not limited to:	Actinopteris, African violets, ageratum, aglaonema, Algerian ivy, allamanda, alocasia, amaranthus , anthurium, aphelandra, arborvitae, Artemisia, aster, aucuba ilex , azalea , baby's breath, begonia, Boston fern, bougainvillea, boxwood, brachycome, cacti, calabrese, caladium, calathea, calendula, calla, camellia, carnation , ceanothus, chrysanthemum, cineraria, coleus, columbine, cotoneaster, cyclamen, daffodil, dahlia, daisy, daylily, delphinium, dianthus, dieffenbachia, dogwood, dusty miller, Easter lily, English ivy, euphorbia, fern , ficus , foliage plants, foxglove, freesia, fuchsia, gaillardia, gardenia, geranium, gerbera, gladiola, gloxinia, gypsophila, hedera, hibiscus, hyacinth, hydrangea, ilex , impatiens , iris, ivy, jasmine, lilac, lily, maidenhair fern, mandevilla, marigold, narcissus, nasturtium, orchid, pansy, pelargonium, peony , peperomia, petunia, philodendron, phlox, photinia, pinks, pittosporum, poinsettia, pothos, portulaca, primrose, pyracantha, rhododendron, rose , rosemary, rubber plant, salvia, schefflera, sedum, sempervivum, snapdragon , spathiphyllum, stock, syngonium, tulip, verbena, vinca, wandering jew, yucca, zinnia
Ornamental Trees and Shrubs including but not limited to:	Andromeda, arborvitae, ash, Austrian pine, azalea , beech, birch, birdsnest spruce, blue spruce, bougainvillea, boxwood, butternut, cedar, charmaecyparis, cherry, cotoneaster, crabapple , cyprus, dogwood, Douglas fir , elm, euonymus, firethorn, forsythia, hackberry, hawthorn, hemlock, hickory, holly, honey locust, horse chestnut, juniper, larch, laurel, lilac, linden, London planetree, magnolia , mandevilla, maple , mimosa, mountain ash, myrtle, oak , pachysandra, peach, photinia, pine, planetree, poplar, privet, purpleleaf wintercreeper, quince, sage, spruce, sycamore , white cedar, white pine, yew

FOR USE ON GARDEN CROPS, VEGETABLES, HERBS, SPICES, FRUITS AND BERRIES

Leafy Vegetables including but not limited to:	Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Endive, Kale, Lettuce , Spinach
Root Vegetables, including but not limited to:	Beet, Carrot , Horseradish, Parsnip, Potato, Radish, Sweet potato, Turnip, Yams
Fruiting Vegetables including but not limited to:	Eggplant, Pepper, Tomatillo, Tomato
Cucurbit Vegetables including but not limited to:	Cucumber , Gourd (edible), Muskmelon, Pumpkin, Squash, Watermelon, including Cantaloupe, Casaba, Gherkins, Melons (including hybrids), Zucchini
Legume Vegetables including but not limited to:	Bean, Chickpea, Lentil, Pea
Bulb Vegetables including but not limited to:	Garlic, Onion , Shallot
Berries including but not limited to:	Blackberry, Blueberry, Raspberry, Strawberry , others include: Boysenberry, Currants, Dewberry, Elderberry, Gooseberry, Loganberry
Herbs and Spices including but not limited to:	Chive, Dill, Fennel, Mustard , Sage, Sweet bay, others include: Anise, Balm, Basil , Black pepper, Borage, Caraway, Catnip, Chamomile, Coriander, Cumin, Curry leaf, Dandelion, Fenugreek, Horehound, Hyssop , Marjoram, Marigold , Mint, Nasturtium, Pennyroyal, Peppermint, Rosemary, Savory, Spearmint, Tansy, Tarragon, Thyme , Wintergreen, Woodruff, Wormwood
Nut Trees including but not limited to:	Almond , Brazil nut, Filbert, Hickory nut, Pecan, Pistachios, Walnut
Pome Fruits including but not limited to:	Apple , Quince, or Pear (Comice varieties: DO NOT apply more than 24 fl oz/A. DO NOT apply after pink stage of flowering; test small areas of other varieties of pears for plant safety prior to full scale usage.)
Stone Fruits including but not limited to:	Apricot, Cherry , Nectarine, Peach, Plum
Citrus Fruits including but not limited to:	Grapefruit, Lemon, Lime, Orange others include: Citrus Citron, Mandarin (tangerine), Nectarine , Satsuma (orange mandarin), Tangerine

There are no restrictions on applying **EcoGarden™** up to the time of harvest.

APPLICATION METHODS

Instructions for using Water Soluble Capsule

Water Soluble Capsule – 1 milliliter (0.034 fluid ounce or 1/15 Tbs.)

Use three (3) to five (5) EcoGarden™ Water Soluble Capsules per plant up to five gallon container size

Use six (6) to ten (10) EcoGarden™ Water Soluble Capsules per plant for container larger than five gallon size

Use **EcoGarden™** Water Soluble Capsules for effective control of all foliar pests, soil-borne insect larvae, including soil-borne larvae of foliar pests, such as fungus gnats, nematodes, or soil-borne thrips on indoor and outdoor plants and garden vegetables.

EcoGarden™ is a systemic product that is translocated upward into the plant system from root uptake. For best results, capsules must be placed around the plant in container or garden plants deep into the soil near growing portion of the target plant root structure. Water Soluble Capsule applications of **EcoGarden™** will have a slower rate of activity since it takes some time for the product to translocate throughout the plant. Target **EcoGarden™** application to coincide with the early stages of insect development. Repeat **EcoGarden™** application every 1 to 6 weeks depending on pest infestation levels. When pest infestation is low, **EcoGarden™** applications can be made at intervals of every 8 to 12 weeks.

Early application as soon as a pest population is identified brings good level of pest control. When applying as a soil drench, avoid excessive leaching

EcoGarden™ Water Soluble Capsule

Open the cover pack and remove Mid-Size Capsule.

Open the Mid-Size Capsule and remove the inner liquid filled Water Soluble Capsules.

Do not handle Water Soluble Capsules with wet hands or wet gloves. Rough handling may cause breakage.

Insert a liquid filled Water Soluble Capsule into surface of soil.

Distribute & push remaining liquid filled Water Soluble Capsules into surface of soil.

Drench the soil with water.

76
79

77
79

STORAGE & DISPOSAL

PESTICIDE STORAGE:

Do not store this product above 105°F or below -15°F for extended periods of time. Keep containers tightly closed and in original containers when not in use. Store in original container in a cool, dry place, out of reach of children.

PESTICIDE DISPOSAL:

Nonrefillable container. Do not reuse or refill this container.

Dispose of the empty outer capsule in the trash, as long as the water soluble capsule is unbroken.

NOTICE ON CONDITIONS OF SALE

The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Plant injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of E.I.D. PARRY (INDIA) LIMITED. All such risks shall be assumed by the user or buyer.

Revised 11-17-08

78/19

Advertising claims that may be presented on the retail container label or on the labeling accompanying the product.

- ◆ Contains azadirachtin from E.I.D. PARRY (INDIA) LIMITED
- ◆
- ◆ Packaging designed for energy and environmental conservation
- ◆ Can be used in Organic Gardening  FOR ORGANIC PRODUCTION
- ◆ Tough Pests. Easy Solution
- ◆ Organic Certified
- ◆ Organic Materials Review Institute (OMRI) Certified
- ◆ Controls chewing and sucking insects
- ◆ Broad spectrum insect and mite control
- ◆ Can be applied up to and on the day of harvest for all food & non food use plants
- ◆ Can be used on a wide variety of flowers, garden vegetables and plants.
- ◆ For use on Rose and Flowers
- ◆ For use on Tomatoes and Garden Vegetables
- ◆ Controls Japanese Beetle on Roses
- ◆ Insect and Mite Control for the Home Garden Vegetables
- ◆ Complete Insect and Mite Control for Flowers
- ◆ Quadruple action insect control – Antifeedant, Insect Growth Regulator, Anti-ovipository, Repellent
- ◆ Single product; multiple action
- ◆ Single product; multiple pest control
- ◆ A proven botanical pesticide born out of a decade of intense global research
- ◆ Works effectively against a broad spectrum of pests on a stand alone basis on a variety of vegetable crops
- ◆ Controls insect pest during various stages of growth
- ◆ The Benefits to your insect control program are multiple
- ◆ Controls Insects Systemically
- ◆ Taken up through the roots, controls insects on the leaves
- ◆ Insects stop feeding when they feed on treated plants
- ◆ Easy to Use
- ◆ No Measuring
- ◆
- ◆ Three in one product – Insecticide, Miticide, and Nematicide
- ◆
- ◆ Recommended by General Hydroponics – Bringing Nature and Technology Together

7a
/1a

Sublabels 5, 6, 7, 8, and 9
Label for Individual Unit

EcoGarden™ Botanical Insecticide, Miticide and Nemacide

Active Ingredient:	% by Weight
Azadirachtin.....	1.2%
Other Ingredients.....	98.8%
Total.....	100.0%

KEEP OUT OF REACH OF CHILDREN
CAUTION

EPA Reg. No. 71908-1

See accompanying label for complete directions for use.