

71962-1

9/30/2014

1/72



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

September 30, 2014

SciReg, Inc.  
12733 Director's Loop  
Woodbridge, VA 22192  
Attn: Frederick T. Smith

Subject: Notification of Product name change in accordance with PR Notice 98-10.

Product Name: Agri-Fos Systemic Fungicide.

**New Product Name: Agri-Fos 400®.**

EPA Reg. No: **71962-1.**

Your Submission dated: July 17, 2014.

Dear Mr. Smith:

The Biopesticides and Pollution Prevention Division is in receipt of your application for Notification under Pesticide Registration Notice (PRN) 98-10 dated above. A screen of this request has been conducted for its applicability under PRN 98-10, and it has been determined that the action requested falls within the scope of this Notice. Our records have been duly noted, and the application, EPA Form 8750-1 submitted, has been stamped "Notification Accepted" and will be placed accordingly in our records.

Three (3) copies of final printed labeling must be submitted to the Agency before your product as modified, may be sold or distributed [PR Notice 82-2 and 40 CFR 156.10(a) (6)].

If you have any questions regarding this action, you may contact Mr. Sylvester George at (703) 603-0688 or via e-mail at [george.sylvester@epa.gov](mailto:george.sylvester@epa.gov).

Sincerely,

*Linda A. Hollis*

Linda A. Hollis, Chief  
Biochemical Pesticide Branch  
Biopesticides and Pollution  
Prevention Division (7511P)

Master label encompassing Sub-label A – Commercial and Agricultural  
Uses, Sub-label B – Turf and Ornamental Uses and Sub-label C –  
Residential Uses

2/72

**AGRI-FOS 400®**

SYSTEMIC FUNGICIDE FOR THE EFFECTIVE CONTROL OF VARIOUS PLANT DISEASES INCLUDING BLACK SPOT OR SCAB IN APPLE, ROOT ROT IN AVOCADO, BUD ROT AND NUT FALL IN COCONUT, ROOT ROT IN CITRUS AND CUCURBITS, DOWNY MILDEW IN CUCURBITS, GRAPE, LETTUCE, AND ONION, ANTHRACNOSE IN MANGO, ROOT AND HEART ROT IN PINEAPPLE, LATE BLIGHT IN POTATO, ROOT AND COLLAR ROT IN STONEFRUIT, LEATHER ROT AND PHYTOPHTHORA DISEASES IN STRAWBERRY, LATE BLIGHT IN TOMATO, DOWNY MILDEW, PHYTOPHTHORA & PYTHIUM IN ORNAMENTALS, INTERIORSAPES, BEDDING PLANTS, VEGETABLE TRANSPLANTS, AND HERBS AND SPICES, PHYTOPHTHORA AND FUSARIUM IN CONIFERS, PYTHIUM IN TURF, AND PHYTOPHTHORA AND PYTHIUM DISEASES ASSOCIATED WITH STEM AND CANKER BLIGHT (SUDDEN OAK DEATH) AND GENERAL BEECH DECLINE AND VARIOUS TREE STEM CANKERS AND PHLOEM CANKERS.

ACTIVE INGREDIENTS: Mono- and di-potassium salts of Phosphorous Acid*	45.8%
OTHER INGREDIENTS:	54.2%
TOTAL	100.0%

\*Contains 5.17 lbs/gallon of the active ingredients, mono- and di-potassium salts of Phosphorous Acid.  
Equivalent to 3.35 lbs Phosphorous Acid/gallon

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

EPA Reg. No. 71962-1

EPA Est. No. 71962-AUS-001  
72499-GA-001  
71975-WA-001  
68253-CO-001

Manufactured for:  
**LIQUID FERTILISER PTY. LTD. (Trading as Agrichem)**  
2 Hovey Road  
Yatala, Queensland 4207  
Australia  
320-222-3026

Notification Accepted

Date: 9/20/2014

Reviewer: *Subrata Gargal*

Net Contents: 1 pt., 1 qt., 1 gal., 5 gal., 25 gal., 55 gal., 250 gal.

Sub-label A – Commercial and Agricultural Uses

3/72

**AGRI-FOS 400®**

SYSTEMIC FUNGICIDE FOR THE EFFECTIVE CONTROL OF VARIOUS PLANT DISEASES INCLUDING BLACK SPOT OR SCAB IN APPLE, ROOT ROT IN AVOCADO, BUD ROT AND NUT FALL IN COCONUT, ROOT ROT IN CITRUS AND CUCURBITS, DOWNY MILDEW IN CUCURBITS, GRAPE, LETTUCE, AND ONION, ANTHRACNOSE IN MANGO, ROOT AND HEART ROT IN PINEAPPLE, LATE BLIGHT IN POTATO, ROOT AND COLLAR ROT IN STONEFRUIT, LEATHER ROT AND PHYTOPHTHORA DISEASES IN STRAWBERRY, LATE BLIGHT IN TOMATO, AND DOWNY MILDEW, PHYTOPHTHORA & PYTHIUM IN HERBS AND SPICES.

ACTIVE INGREDIENTS: Mono- and di-potassium salts of Phosphorous Acid*	45.8%
OTHER INGREDIENTS:	54.2%
TOTAL	100.0%

\*Contains 5.17 lbs/gallon of the active ingredients, mono- and di-potassium salts of Phosphorous Acid.  
Equivalent to 3.35 lbs Phosphorous Acid/gallon

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

Consult page 2 of this booklet for First Aid and Precautionary Statements

EPA Reg. No. 71962-1


EPA Est. No. 71962-AUS-001  
72499-GA-001  
71975-WA-001  
68253-CO-001

Manufactured for:  
**LIQUID FERTILISER PTY. LTD. (Trading as Agrichem)**  
2 Hovey Road  
Yatala, Queensland 4207  
Australia  
320-222-3026

Batch No.:

Date of Manufacture:

Net Contents: 1 qt., 1 gal., 5 gal., 25 gal., 55 gal., or 250 gal.



4/  
72

<b>FIRST AID</b>	
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice</li><li>• Have person sip a glass of water if able to swallow</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor</li><li>• Do not give anything to an unconscious person</li></ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye</li><li>• Call poison control center or doctor for treatment advice</li></ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes</li><li>• Call a poison control center or doctor for treatment advice</li></ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"><li>• Move person to fresh air</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible</li><li>• Call a poison control center or doctor for further treatment advice</li></ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment. Hotline Number: National Poison Control, 1-800-222-1222	

### **PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION.** Harmful if swallowed, absorbed through skin or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist or vapors. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Wear the appropriate Personal Protective Equipment (PPE).

#### **PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Applicators, mixers, loaders, and other handlers must wear:

- protective eyewear
- long pants and long-sleeved shirt
- waterproof gloves
- shoes plus socks.

Follow manufacturer's instructions for maintaining/cleaning personal protective equipment (PPE). If no such instructions for washables, use hot water and detergent. Keep and wash PPE separately from other laundry.

When handlers use closed systems, aircraft or enclosed cabs in a manner that meets the requirements listed in the worker protection standard (WPS) for agricultural pesticides (40 CFR 170.240 (d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

#### **USER SAFETY RECOMMENDATIONS**

Users should:

- Wash hands before drinking, eating, chewing gum, using tobacco or using the toilet.
- Remove PPE clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

5/  
72

For Terrestrial Use: Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the state or tribal agency responsible for pesticide registration.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, CFR 40 part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI) of 4 hours**, unless wearing the appropriate PPE.

PPE required for early entry to treated areas that are permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soils or water, is: coveralls worn over short-sleeved shirt and short pants, waterproof gloves, shoes plus socks and protective eyewear.

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Do not enter treated areas without protective clothing until sprays have dried.

## CHEMIGATION

Use of AGRI-FOS 400® through chemigation is not permitted in California.

Apply this product only through center pivot, solid set or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

6/72

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

#### SPRINKLER AND DRIP (TRICKLE) IRRIGATION SYSTEMS:

The irrigation system must contain a functional check valve, vacuum relief valve and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

(Sprinkler Chemigation Only) Do not apply when wind speed favors drift beyond the area intended for treatment.

Apply AGRI-FOS 400® continuously for the duration of the water application. After treatment with AGRI-FOS 400® has been completed, avoid further irrigation of the treated area until foliage is dry or for 24 to 48 hours.

#### GENERAL APPLICATION INSTRUCTIONS

Apply AGRI-FOS 400® by various application methods, including foliar spray (aerial and ground), soil drench, soil incorporation, basal bark application and bare root dip. For foliar sprays, apply AGRI-FOS 400® with sufficient water volumes for adequate coverage of foliage, according to crop and growth stage. To ensure good coverage, spray to wetness, but avoid run-off. **This product does not have a pre-harvest interval (PHI). Harvest when dry.**

When using AGRI-FOS 400® with Supa Stik PN, adhere to all applicable label directions. Only use Supa Stik PN with basal bark applications.

#### MIXING INSTRUCTIONS

1. Fill the spray tank with ¼ - ½ of the volume of water required before adding AGRI-FOS 400®
2. Add AGRI-FOS 400® slowly to the tank and agitate by mechanical or hydraulic means
3. Continue agitating as tank fills with water to the desired volume
4. Maintain agitation during application

#### COMPATIBILITY

AGRI-FOS 400® is compatible with most products used in agriculture. However, individual crop sensitivity to these mixtures may vary. Mixtures of AGRI-FOS 400® with some foliar fertilizers and copper products are not always compatible or cause phytotoxicity to some plants. If these combinations or others have not been used previously, do not tank mix without first testing the compatibility of the tank mix. Do not apply tank mixture without first assessing its safety to the

7/72

crop (phytotoxicity). Tank mix AGRI-FOS 400® with fertilizers only if crop safety has been established and the AGRI-FOS 400® use rates are followed.

Due to AGRI-FOS 400®'s acidic nature, do not use acidifying-type compatibility agents. If spray adjuvants are used test them before use to confirm compatibility with AGRI-FOS 400®.

Use a jar test to confirm compatibility with AGRI-FOS 400®. In a clean jar using the same water source that is normally used to fill spray tank, add the same proportions of each product and the appropriate quantity of water and mix thoroughly. Let stand for 3 minutes. If mixture remains in solution or is remixed readily the tank mix is compatible.

Spray the solution that results from the above compatibility test onto a few plants and inspect for visual effects of phytotoxicity (leaf burn) 3 days later.

### GENERAL HYDROPONIC APPLICATIONS\*

For use with plants grown in recirculating hydroponic systems.

Disease	Rate	Application Program
General root rots (Pythium and Phytophthora spp) and root diseases	1 to 2 quarts in 5300 gallons of nutrient solution	Apply every 4 to 6 weeks in the summer and every 8 weeks in the fall. The application time interval may be modified depending on crop load, water quality, and disease pressure.

\*Not for use in California.

### AGRICULTURAL APPLICATIONS

#### APPLES, CRAB APPLES, LOQUATS, PEARS & QUINCE\*

Use AGRI-FOS 400® for effective control of black spot, root and collar rot and fire blight in apples, crab apples, loquats, pears, and quinces.

Disease	Application Method	Rate	Application Program
Apple black spot and scab (Venturia inaequalis)	Foliar spray Ground	Apply 3 to 4 pints in 100 gallons water per acre.	Apply in combination with a mancozeb-containing product at 3 lbs/acre. Apply at ¼ - ½ inch green tip through first cover at 7 to 10 day intervals or according to forecasted infection events. Continue with AGRI-FOS 400® and Captan in the remaining applications.
Apple black spot and scab (Venturia inaequalis)	Basal bark spray at bud swell or silver tip stage of growth in early spring	62.4 fl. oz. + 62.4 fl. oz. of water + 1.5 to 3.0 fl. oz. Supa Stik PN Bark Penetrating Surfactant	Spray a combination of AGRI-FOS 400® and Supa Stik PN on the complete circumference of the tree trunk until saturation/runoff. Spray from ground level up to 5 feet above the soil line, including the base of the first scaffolding limbs, if present (treatment generally lasts 8-12 weeks depending on

8/72

			pathogen levels. Higher disease pressure will shorten the length of control).
Root and collar rot (Phytophthora cactorum) Fire Blight (Erwinia amylovora)	Foliar spray Ground	2½ to 5 pints in 100 gallons of water per acre	Thorough spray coverage of plant is required. Start applications when conditions favor disease development. Apply at one to two month intervals. Use the low rate on the shorter interval and the high rate on the longer interval.
	Basal bark spray treat in spring and fall for best results	62.4 fl. oz. + 62.4 fl. oz. of water + 1.5 to 3.0 fl. oz. Supa Stik PN Bark Penetrating Surfactant	Spray a combination of AGRI-FOS 400® and Supa Stik PN around the complete circumference of the tree trunk until saturation/runoff. Spray from ground level up to 5 feet above the soil line, including the base of the first scaffolding limbs, if present. (Treatment generally lasts 8-12 weeks depending on pathogen levels. Higher disease pressure will shorten the length of control).

\*Not for use in California except basal bark spray on apples and crab apples.

**ASPARAGUS\***

Use AGRI-FOS 400® for effective control of crown rot & asparagus spear slime disease in asparagus.

Disease	Application Method	Rate	Application Program
Crown rot & Asparagus spear slime (Phytophthora spp)	Foliar spray Ground	1¼ quarts per acre in 25 gallons water to 2½ quarts per acre in 250 gallons water	Apply to ferns that have 2 to 3 inches of new growth. Do not apply to ferns that are starting to die down (senesce). Established plantings, start applications when conditions are favorable to disease (cool wet conditions). Ensure thorough coverage.

\*Not for use in California.

9/72

**AVOCADOS**

Use AGRI-FOS 400® for effective control of root rot, trunk cankers and downy mildew disease in avocado.

Disease	Application Method	Rate	Application Program
Root rot (Phytophthora cinnamomi)	Tree injection	Skeletal trees 1st year: ¼ fl. oz. undiluted product per yard of canopy diameter. Other situations: ¾ teaspoon diluted with ½ fl. oz. of water per yard of canopy diameter.	Inject trees at spring flush maturity. Repeat post vegetative at summer flush maturity. Drill holes 3/16 inch (5 mm) in diameter and 1 inch (25 mm) to 2 inches (50 mm) deep with slight downward angle in trunk. Place syringes in the main trunk of the tree and space evenly around the circumference of the trunk. Suitable for use with Chemjet tree injectors, Ag-murf gun, Smart Shot injector, or hydraulic tree injection. Do not prune back trees before injection process as burning of new growth may occur. Do not inject trees in winter months. Do not cut back the canopy of injected trees. Do not add any material, other than water, to AGRI-FOS 400® by trunk injection. Do not inject more liquid in a lesser number of syringes than directed.
	Foliar spray Ground	2 quarts in 100 gallons of water per acre	Spray to run off at 2 – 2½ gallons of spray solution per adult tree. Start applications in spring, up to 4 applications a year at two-month intervals. Ensure thorough coverage.
Canker (Phytophthora citricola)	Trunk spray	1¼ to 2½ quarts mixed with 5 gallons of water with 3.0 fl. oz. of Supa Stik PN Bark Penetrating Surfactant	Apply to trunk lesions using sufficient spray volume to completely wet the trunk and lesions. If lesions absent, apply to trunk from soil level to two feet up trunk. If lesions present use higher rate.
Downy mildew	Foliar spray Ground	¾ pints in 100 to 500 gallons water	Spray to run-off, as required for disease control.

10/72

**BERRIES\***

Use AGRI-FOS 400® for effective control of root rot, general leaf and berry diseases such as those caused by *Septoria* spp., and downy mildew, and for suppression of Anthracnose spp. in bush and cane berries such as, but not limited to, bingleberries, blackberries, black satin berries, blueberries, boysenberries, Cherokee blackberries, Chesterberries, Cheyenne blackberries, coryberries, cranberries, darrowberries, dewberries, Dirksen thornless berries, elderberries, Himalayaberries, huckleberries, hullberries, Lavacaberries, loganberries, lowberries, lucretiaberries, mammoth blackberries, marionberries, mulberries, nectarberries, ollalieberries, Oregon evergreen berries, raspberries (red, black, hybrids/cultivars), and youngberries.

Disease	Application Method	Rate	Application Program
Root rot ( <i>Phytophthora</i> spp.)	Foliar spray	1-3 quarts per 100 gallons of water per acre. Ensure foliage is completely wet.	New plantings: start application when new growth is 2 to 3 inches long. Established plantings start applications when cool wet conditions occur which favor disease. West of Rocky Mountains: Autumn applications, apply when conditions favor disease, repeat in 4 weeks. Spring applications, first application after bud break and repeat in 4 weeks. East of Rocky Mountains: First application spring post bud break (2 to 3 inches new growth) and repeat at 50 to 60 day intervals. Do not exceed 4 applications per season.  For blueberries - First application in spring at pink bud and then on a regular schedule of application at two to three intervals.
	Ground		
General leaf and berry diseases such as those caused by <i>Septoria</i> spp and Anthracnose spp	Root dip	Mix a 2.5% v/v solution (1 quart per 10 gallons of water)	Apply as pre-plant dip to the roots for 2-3 minutes. Plant within 48 hours after dipping. Mix a fresh solution daily.
	Chemigation Overhead	.1 to 2 quarts in 1000 gallons of water per acre	Apply with normal irrigation schedule. Do not apply more than 4 times per crop cycle.
	Low volume	2 to 3 quarts in 100 gallons of water per acre	
Downy mildew	Foliar spray Aerial	1½ to 2 quarts in 20 gallons of water per acre	Low Disease Pressure: Apply lower rate at first onset of disease. Repeat applications at 1 to 3 week intervals. Do not apply more than 6 times per crop cycle.
	Ground	1½ to 2 quarts in up to 100 gallons of water per acre	

11/72

			<p>High Disease Pressure: Apply higher rate at first onset of disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.</p> <p>AGRI-FOS 400® is best when used in combination with conventional registered fungicides to increase the performance of the disease control program.</p>
--	--	--	---

\*Not for use in California.

**BRASSICAS\***

Use AGRI-FOS 400® for effective control of damping-off, root rot, bacterial, and fungal diseases, and downy mildew in brassicas such as, but not limited to, broccoli, Brussels sprouts, cabbage, cauliflower, cavalo broccolo, collards, Chinese cabbage, Chinese mustard cabbage, kale, kohlrabi, mizuna, mustard greens, mustard spinach and rape greens.

Disease	Application Method	Rate	Application Program
Root rot and damping off (Phytophthora spp and Pythium spp)	Foliar spray	1 to 1½ quarts in 10 gallons of water per acre	<p>Begin application after plants are established and conditions favor disease development.</p> <p>Low Disease Pressure: Apply lower rate at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle.</p> <p>High Disease Pressure: Apply higher rate at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.</p>
	Aerial	1 to 2 quarts in 30 gallons of water per acre	
	Ground	1 quart in 100 gallons of water to nursery plants in seedling trays 1 to 7 days prior to out planting	
	Pre-plant seedling nursery treatment	3 pints at planting or to newly planted seedlings by side dressing or shank application	
	Transplant and furrow application		
Bacterial diseases	Chemigation	1 to 2 quarts in 1000 gallons of water per acre	<p>Apply with normal irrigation schedule.</p> <p>Low Disease Pressure: Apply lower rate at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle.</p> <p>High Disease Pressure: Apply higher rate at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.</p>
	Overhead:	2 to 3 quarts in 100 gallons of water per acre	
	Low volume:		

12/72

Downy mildew (Peronospora parasitica)	Foliar spray Ground	1¼ quarts per acre in 25 gallons of water per acre to 2½ quarts per acre in 250 gallons of water per acre  California: 1¼ quarts per acre in 100 gallons of water to 2½ quarts per acre in 185 gallons of water	1 to 3 week intervals between applications when conditions favor disease development (cool, moist weather). Use higher rates and shorter intervals when disease pressure increases.
Diseases caused by Septoria, Collectotrichum, Alternaria, and Powdery mildew	Foliar spray Ground	1 to 2 quarts in 30 gallons of water per acre	<u>Low Disease Pressure:</u> Apply lower rate at first onset of disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle.  <u>High Disease Pressure:</u> Apply higher rate at first onset of disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.

\*Use in California limited to control of downy mildew.

**CEREAL GRAINS\***

Apply AGRI-FOS 400® to crops such as, but not limited to, field corn, ornamental corn, sweet corn, Indian corn, wheat, rye, barley, oats, triticale, and sorghum (milo). Use AGRI-FOS 400® for effective control of damping-off and root rot diseases.

Disease	Application Method	Rate	Application Program
Phytophthora and Pythium spp	Foliar spray Ground	1¼ quarts per acre in 25 gallons of water to 2 quarts per acre in 250 gallons of water	Corn: Apply at 14-day intervals from 4-leaf stage, as needed. Assure good coverage.  Other grains: Apply at 14-21 day intervals, as needed. Assure good coverage.

\*Not for use in California.

**CITRUS - Mature trees**

Use AGRI-FOS 400® for effective control of root rot and collar rot diseases in citrus.

Disease	Application Method	Rate	Application Program
Brown rot and root rot Phytophthora spp	Foliar spray Ground	2½ quarts per acre in a maximum of 250 gallons of water	<b>When conditions favor disease.</b> Spray trees to run off ensure even coverage. Do not apply at high temperatures (above 95°F) particularly if humidity is low or to moisture-stressed trees.

13/72

Root rot and collar rot (Phytophthora spp. Phytophthora nicotianae and Phytophthora citrophthora)	Trunk spray	Mix 1¼ to 2½ quarts in a minimum of 5 gallons of water with 1 to 3 fl. oz. of Supa Stik PN Bark Penetrating Surfactant	Spray trunk lesions with enough spray volume to ensure lesions are completely wet. When disease levels are high use higher rate.
* Pre-harvest blue and green mold and brown rot (Phytophthora citricola)	Foliar spray Ground	2 quarts in up to 100 gallons of water per acre	Apply 2 to 4 weeks prior to harvest. Ensure fruit is thoroughly covered by the spray application.

\* Not for use in California.

**COCONUTS**

Use AGRI-FOS 400® for effective control of bud rot and nut fall in coconuts.

Disease	Application Method	Rate	Application Program
Bud rot – Nut fall (Phytophthora palmivora)	Injection	2 tsp. to 1 fl. oz. per tree	Dilute AGRI-FOS 400® with water to give final injection volume of 1 fl. oz. to 2 fl. oz. of water and AGRI-FOS 400®. Inject into the trunk or root system.

**COTTON\***

Use AGRI-FOS 400® for effective control of damping-off and root rot diseases in cotton.

Disease	Application Method	Rate	Application Program
Phytophthora and Pythium spp	Foliar spray Ground	1 to 2 quarts in 30 to 50 gallons of water per acre	Apply at crop emergence every 21 days or during wet conditions that favor pathogen development.

\*Not for use in California.

**CUCURBITS**

Use AGRI-FOS 400® for effective control of sudden wilt, gummy stem blight, downy mildew, powdery mildew, Alternaria leaf blight, and bottom soft root complex, and for suppression of Anthracnose in cucurbits grown in field situations such as, but not limited to, cucumber, Chinese waxgourd, citron melon, gherkin rock melon, honeydew melon, pumpkin, zucchini, watermelon and squash (summer and winter), momordica spp balsam apple, balsam pear, bitter melon, and Chinese cucumber.

Disease	Application Method	Rate	Application Program
Sudden wilt – Root and fruit rot (Phytophthora spp) Phytophthora blight (Phytophthora capici)	Foliar spray Ground	1-3 quarts per 100 gallons.	Entire spray coverage of plant is required. Do not exceed a total of 6 applications per season.
Gummy stem blight (Mycosphaerella melonis)		Apply 1¼ quarts per acre in 40 gallons of water. (CA only)	Apply when disease is evident. Continue applications at 21 day intervals until cure is apparent. Do not exceed a total of 6 applications per season.

14/72

Downy mildew (Pseudoperonospora cubensis)			Apply within 7 to 10 days of infection. Repeat as necessary.  Do not exceed a total of 6 applications per season.
* Powdery mildew and other leaf diseases such as Anthracnose and Alternaria leaf blight	Foliar spray Ground	2 to 2½ quarts in 30 gallons of water per acre	<u>Low Disease Pressure:</u> Apply lower rate at first onset of disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle.  <u>High Disease Pressure:</u> Apply higher rate at first onset of disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.
* Bottom soft rot complex	Foliar spray Ground	2 to 2½ quarts in up to 100 gallons of water per acre	Apply after fruit set and during bulking 3 times during the growing season.

\* Not for use in California.

**CUCURBITS, TANK MIXTURES\***

For the effective control of downy mildew diseases, tank-mix AGRI-FOS 400® with a mancozeb-containing fungicide and apply to cucurbits.

Product	Disease	Rate Per Acre	Application Program
AGRI-FOS 400® + mancozeb-containing fungicide	Downy mildew	Light to medium foliage cover: Apply 1¼-2 quarts of product per acre.  Heavy foliage cover: Apply 3 quarts of product per acre + Label rate of mancozeb	Apply as a ground foliar spray the indicated quantity and dilution rates with water of both products.  For best results apply AGRI-FOS 400® as a tank mix with protectant fungicides such as mancozeb, copper oxychloride, etc., to ensure both pre- and post-infection activity.

\*Not for use in California.

15/72

**FRUITING VEGETABLES\***

Use AGRI-FOS 400® for effective control of damping-off, root rot, gummy stem blight diseases, bacterial diseases, downy mildew, powdery mildew, and Alternaria leaf blight, and for suppression of Anthracnose in eggplant, tomatoes, tomatillos, and peppers such as, but not limited to, bell, chili, cooking, pimento, and sweet.

Disease	Application Method	Rate	Application Program
Eggplant: Pythium and Phytophthora spp, and Gummy stem blight (Mycosphaerella melonis)	Foliar spray Ground	Apply 1½ quarts per acre in 40 gallons of water.	Entire spray coverage of plant is required. Do not exceed a total of 6 applications per season.  Apply when disease is evident. Continue applications at 21-day intervals until cure is apparent.  Do not exceed a total of 6 applications per season.
Peppers: Late blight and root rot (Phytophthora infestans and Phytophthora spp)		1¼ quarts per acre in 25 gallons of water to 2 quarts per acre in 250 gallons of water.	First application at transplant or when direct seeded crops are at 2-4 true leaf, then at one to two week intervals as required to control disease. In high disease situations use higher rates and shorter spray intervals.
Tomatoes/Tomatillos: Late blight and root rot (Phytophthora infestans and Phytophthora spp)		1½ quarts per acre in 25 gallons of water to 2 quarts per acre in 250 gallons of water  California: 1¼ quarts per acre in 90 gallons of water	First application at transplant or when direct seeded crops are at 2-4 true leaf, then at one to two week intervals as required to control disease. In high disease situations use higher rates and shorter spray intervals.
Root rot and damping off (Phytophthora spp and Pythium spp)  Bacterial diseases	Foliar spray Aerial  Ground  Pre-plant seedling nursery treatment  Transplant and furrow application	1 to 1½ quarts in 10 gallons of water per acre  1 to 2 quarts in 30 gallons of water per acre  1 quart in 100 gallons of water to nursery plants in seedling trays 1 to 7 days prior to out planting  3 pints at planting or to newly planted seedlings by side dressing or shank application	Begin application after plants are established and conditions favor disease development.  <u>Low Disease Pressure:</u> Apply lower rate at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle.  <u>High Disease Pressure:</u> Apply higher rate at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.  For control of bacterial leaf spot of tomatoes, apply the high rate of AGRI-FOS 400® with registered bacteriacides.

16/72

	Chemigation Overhead  Low volume	1 to 2 quarts in 1000 gallons of water per acre  2 to 3 quarts in 100 gallons of water per acre	Apply with normal irrigation schedule.  <u>Low Disease Pressure:</u> Apply lower rate at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle.  <u>High Disease Pressure:</u> Apply higher rate at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.
Downy mildew	Foliar spray Aerial  Ground	1 to 1½ quarts in 10 gallons of water per acre  1 to 2 quarts in 30 gallons of water per acre	<u>Low Disease Pressure:</u> Apply lower rate at first onset of disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle.  <u>High Disease Pressure:</u> Apply higher rate at first onset of disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.
Powdery mildew and other leaf diseases such as Anthracnose and Alternaria leaf blight	Foliar spray Aerial  Ground	1 to 1½ quarts in 10 gallons of water per acre  1 to 2 quarts in 30 gallons of water per acre	<u>Low Disease Pressure:</u> Apply lower rate at first onset of disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle.  <u>High Disease Pressure:</u> Apply higher rate at first onset of disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.

\* Use in California limited to control of late blight and root rot in tomatoes and tomatillos.

**GRAPES\***

Use AGRI-FOS 400® for effective control of root rot, downy mildew, and powdery mildew diseases in grapes.

Disease	Application Method	Rate	Application Program
Downy mildew (Plasmopara viticola)	Foliar spray Ground	Early season small canopy, 1¼ quarts per acre in 100 gallons water  Late season/large canopy, 2 quarts per acre in 150 gallons water to 2½ quarts per acre in 200 gallons water	It is essential that the rate of AGRI-FOS 400® be adjusted to the vine- row volume, i.e., the volume of vine foliage per acre. Spray timing is critical. Apply AGRI- FOS 400® at times of high disease risk, especially between the time that conditions are conducive to downy mildew infection and the appearance of oil spots.

17/72

			Ensure spray coverage is adequate and that the appropriate rate of AGRI-FOS 400® is applied to match vine growth, particularly from mid-season onwards, and especially where grapes are grown on root stock.
Phytophthora and Pythium root rots	Foliar spray Ground	1½ to 2 quarts in 50 gallons of water per acre	Apply to vines that have stressed root system that can lead to root rots. Mitigating factors such as nematode pressure, water logging, and compaction contribute to vine declines.  Table Grapes: Begin application in the Spring at the 4 to 6 inch shoot stage. Continue applications at 1 to 2 week intervals until flowering. Resume applications in the Fall after harvest. Do not apply more than 4 times per crop cycle.  Wine and Raisin Grapes: Begin applications in the Spring at the 4 to 6 inch shoot stage. Continue applications at 1 to 2 week intervals through flowering. Do not apply more than 4 times per crop cycle.
Downy mildew (Plasmopara viticola)	Foliar spray Ground	1½ to 2 quarts in 50 gallons of water per acre  AGRI-FOS 400® is most effectively applied to control downy mildew when tank mixed with other registered fungicides.	Apply a bud break with additional applications at 7 to 10 day intervals in a rotational program with other registered fungicides. Use higher rate and volume based on disease severity and density of canopy. Do not apply more than 6 times per crop cycle.
Powdery mildew (Uncinular necator)	Foliar spray Ground	2 to 2½ quarts in 30 gallons of water per acre	Low Disease Pressure: Apply lower rate at first onset of disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle.  High Disease Pressure: Apply higher rate at first onset of disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6

18/72

			times per crop cycle.
--	--	--	-----------------------

\* Use in California limited to control of downy mildew.

**GRAPES, TANK MIXTURES\***

For the effective control of downy mildew diseases, tank-mix AGRI-FOS 400® with a mancozeb-containing fungicide and apply to grapes.

Product	Disease	Rate Per Acre	Application Program
AGRI-FOS 400® + mancozeb-containing fungicide	Downy mildew	Early season small canopy, 1¼ quarts per acre in 100 gallons water  Late season/large canopy, 2 quarts per acre in 150 gallons water to 2½ quarts per acre in 200 gallons water  + Label rate of mancozeb	Apply as a foliar spray the indicated quantity and dilution rates with water of both products.  For best results apply AGRI-FOS 400® as a tank mix with protectant fungicides such as mancozeb, copper oxychloride, etc., to ensure both pre- and post-infection activity.

\*Not for use in California.

**HERBS AND SPICES\***

Use AGRI-FOS 400® for effective control of Downy mildew, Phytophthora spp and Pythium spp diseases of Herbs and Spices grown in fields, nurseries and greenhouses. Apply AGRI-FOS 400® to plants such as, but not limited to, Anise, Balm, Basil, Chamomile, Caraway, Catnip, Celery, Chives, Coriander, Cumin, Curry Leaf, Dill, Fennel, Marjoram, Mint, Nasturtium, Rosemary, Sage, Savory, Sweet Bay, Tarragon, Thyme, and Wintergreen.

Make applications before disease development and in conjunction with good cultural management practices. Use higher rate of application when disease pressure is severe. Do not exceed indicated application rates or apply more frequently than stated on label in order to avoid plant injury. Do not apply to plants that are heat or moisture stressed.

Disease	Application Method	Rate	Application Program
Downy Mildew	Foliar spray	1¼ to 2½ quarts per 100 gallons of water	Apply spray to thoroughly wet all foliage. Application intervals: 14 to 21 days. Repeat as required.
	Ground	Or ½ to 1 1/8 fl. oz. per gallon of water	
Phytophthora spp and Pythium spp	Foliar spray	1 to 2 quarts per 100 gallons of water	Apply spray to thoroughly wet all foliage. Application intervals: 14 to 21 days. Repeat as required. Note do not apply more than 500 gallons of spray solution per acre.
	Ground	Or 2 to 4 tsp. per gallon of water	
	Soil drench	6¼ to 12¾ fl. oz. per 100 gallons of water	Apply each 25 gallons of solution to an area of 100 square feet. Follow application with irrigation. Repeat as required. Limit of one application per month.

\*Not for use in California.

**HOPS\***

Use AGRI-FOS 400® for effective control of downy mildew in hops.

19/72

Disease	Application Method	Rate	Application Program
Downy mildew	Foliar spray by ground equipment only.	1-3 quarts per 100 gallons of water per acre	When conditions favor disease, apply when A. Shoots are ½ to 1 foot long. B. Post-training when vines are 6 feet high. C. 21 days post-application (B). D. During bloom.

\*Not for use in California.

**LEAFY VEGETABLES\***

Use AGRI-FOS 400® for effective control of root rot, damping-off, downy mildew, powdery mildew, and other fungal leaf diseases, and for suppression of Anthracnose in leafy vegetables such as, but not limited to, amaranth, arugula, cardoon, celery, chervil, chrysanthemum, corn salad, cress, dandelion, dock, endive, fennel, lettuce, orach, parsley, purslane, radicchio, rhubarb, spinach, and Swiss chard. Excludes Brassica vegetables.

Disease	Application Method	Rate	Application Program
Downy mildew (Bremia lactucae)	Foliar spray	½ gallon per acre in 40 gallons water	Ensure spray coverage is adequate to wet the whole plant.  During warm, wet conditions repeat application at 7 to 10 day intervals, if needed.
	Ground		
Root rots and damping off (Phytophthora and Pythium spp)	Foliar spray	Apply 1 to 2 quarts in 40 gallons of water per acre	Begin application after plants are established and conditions favor disease development.  Low Disease Pressure: Apply lower rate at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle. High Disease Pressure: Apply higher rate at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.
	Ground		
	Pre-plant nursery application	1 quart in 100 gallons of water to nursery plants in seedling trays 1 to 7 days prior to out planting	
	Chemigation Overhead	Apply 1 to 2 quarts per acre in a minimum of 1000 gallons per acre of water	Apply with normal irrigation schedule.  Low Disease Pressure: Apply lower rate at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle. High Disease Pressure: Apply higher rate at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.
	Low Volume	Apply 1 to 2 quarts per acre in a minimum of 100 gallons per acre of water	
	Transplant and furrow application	3 pints at planting or to newly planted seedlings by side dressing or shank application	

20/  
72

Powdery mildew and leaf diseases such as Anthracnose (Collectotrichum spp), leaf blights (Septoria and Cercospora spp), and bacterial rots (Erwinia spp)	Foliar spray Ground	1½ to 2 quarts in a 30 gallons of water per acre	<p><b>Low Disease Pressure:</b> Apply lower rate at first onset of disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle.</p> <p><b>High Disease Pressure:</b> Apply higher rate at first onset of disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.</p>
--	------------------------	--	---

\*Use in California limited to control of downy mildew in lettuce.

**LEGUMES\***

Use AGRI-FOS 400® for effective control of damping-off, root rot, Fusarium, Rhizoctonia, downy mildew, powdery mildew, and other fungal leaf diseases, and for suppression of Anthracnose in legumes (succulent and dried) such as, but not limited to, green beans, soybeans, wax beans, field beans, navy beans, lima beans, fava beans, kidney beans, pinto bean, mung beans, broad beans, lentils, chickpeas, English peas, snow peas, sugar snap peas, black-eyed peas, cow peas, and pigeon peas.

Disease	Application Method	Rate	Application Program
Phytophthora and Pythium spp  Fusarium and Rhizoctonia	Foliar spray Ground  Pre-plant nursery application  Transplant and furrow application	1½ to 2½ quarts in 30 gallons of water per acre  1 quart in 100 gallons of water to nursery plants in seedling trays 1 to 7 days prior to out planting  3 pints at planting or to newly planted seedlings by side dressing or shank application	Begin application after plants are established and conditions favor disease development.  Low Disease Pressure: Apply lower rate at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle.  High Disease Pressure: Apply higher rate at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.
Downy mildew	Foliar spray Ground	1 to 3 quarts in 30 gallons of water per acre	Low Disease Pressure: Apply lower rate at the first onset of the disease. Repeat applications at 1-2 week intervals. Do not apply more than 6 times per crop cycle.  High Disease Pressure: Apply higher rate at the first onset of the disease. Repeat applications at 7-10 day intervals. Do not apply more than 6 times per crop cycle.
Powdery mildew and leaf diseases such as Anthracnose (Collectotrichum spp), leaf blights (Septoria and	Foliar spray Ground	2 to 2½ quarts in 30 gallons of water per acre	Low Disease Pressure: Apply lower rate at the first onset of the disease. Repeat applications at 1 to 2

21/72

Cercospora spp), and bacterial rots (Erwinia spp)			week intervals. Do not apply more than 6 times per crop cycle.  High Disease Pressure: Apply higher rate at the first onset of the disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.
---	--	--	--

\*Not for use in California.

**MANGOS**

Use AGRI-FOS 400® for suppression of anthracnose in mangos.

Disease	Application Method	Rate	Application Program
Anthracnose (Colletotrichum gloeosporoides)	Foliar spray Ground	2 tsp. per gallon of water	Spray tree every 14 days during blossom period, then monthly until harvest. Spray to the point of run-off.

**MISCELLANEOUS CROPS\***

Use AGRI-FOS 400® for effective control of damping-off, root rot, downy mildew, powdery mildew, and other bacterial and leaf diseases, and for suppression of Anthracnose in coffee, okra, papaya, and persimmon.

Disease	Application Method	Rate	Application Program
Bacterial and leaf diseases such as coffee berry disease (Collectotrichum spp), Anthracnose (Collectotrichum spp), and various leaf spots (Septoria and Cercospora spp)	Foliar spray Aerial  Foliar spray Ground	1 to 2 quarts in 10 gallons of water per acre   1 to 2 quarts in 100 gallons of water per acre	Begin application after plants are established and conditions favor disease development.  Low Disease Pressure: Apply lower rate at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle.  High Disease Pressure: Apply higher rate at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.
	Root dip	Mix a 0.25% v/v solution (1/3 fl. oz. with 1 gallon of water)	Apply as pre-plant dip to transplants immediately prior to planting. Dip plants momentarily and plant within 48. Mix a fresh solution daily.

22/72

	Chemigation Overhead	2 to 3 quarts in 1000 gallons of water per acre	Apply with normal irrigation schedule.
	Low volume	2 to 4 quarts in 100 gallons of water per acre	Low Disease Pressure: Apply lower rate at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle.
	Transplant and furrow application	3 pints at planting or to newly planted seedlings by side dressing or shank application	Apply higher rate at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.
Downy mildew	Foliar spray Aerial	1 to 3 quarts in 10 gallons of water per acre	Low Disease Pressure: Apply lower rate at the first onset of the disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle.  High Disease Pressure: Apply higher rate at the first onset of the disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.
	Foliar spray Ground	1 to 3 quarts in 100 gallons of water per acre	
Powdery mildew	Foliar spray Aerial	1 to 3 quarts in 10 gallons of water per acre	Low Disease Pressure: Apply lower rate at the first onset of the disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle.  High Disease Pressure: Apply higher rate at the first onset of the disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.
	Foliar spray Ground	1 to 3 quarts in 100 gallons of water per acre	
Pseudomonas garcae	Foliar spray Ground	Mix a 0.50% v/v solution (2 quarts per 100 gallons of water)	Apply prior to onset of disease. Apply spray to saturation or runoff.

\*Not for use in California.

**NONGRASS ANIMAL FEED\***

Use AGRI-FOS 400® for effective control of damping-off and root rot diseases in forage crops such as, but not limited to, alfalfa, clover, and vetch.

Disease	Application Method	Rate	Application Program
Phytophthora and Pythium spp	Foliar spray	1 1/2 quarts per acre in 25 gallons of water to 2 quarts per acre in 250 gallons of water.	Apply at 14-day intervals after plant emergence, as needed. Assure good coverage.

\*Not for use in California.

23/72

**ONIONS**

Use AGRI-FOS 400® for effective control of downy mildew disease in onions, garlic, shallots, and leeks. Use as a preventative control program for best results.

Disease	Application Method	Rate	Application Program
Downy mildew (Peronospora destructor)	Foliar spray Ground	2 quarts per acre in 100 gallons water per acre	As a regular preventative control program or when disease first appears.

**ONIONS, TANK MIXTURES\***

For the effective control of downy mildew diseases, tank-mix AGRI-FOS 400® with a mancozeb-containing fungicide and apply to onions, garlic and shallots.

Product	Disease	Rate Per Acre	Application Program
AGRI-FOS 400® + mancozeb-containing fungicide	Downy mildew	2 quarts per acre in 100 gallons water per acre + Label rate of mancozeb	Apply as a foliar spray the indicated quantity and dilution rates with water of both products.  For best results, apply AGRI-FOS 400® as a tank mix with protectant fungicides such as mancozeb to ensure both pre- and post-infection activity.

\*Not for use in California.

**PEANUTS\***

Use AGRI-FOS 400® for effective control of damping-off and root rot disease and for suppression of Anthracnose in peanuts.

Disease	Application Method	Rate	Application Program
Phytophthora and Pythium spp	Foliar spray	1½ quarts per acre in 25 gallons of water to 2 quarts per acre in 250 gallons of water	Apply at 14-day intervals, as necessary. Ensure thorough coverage.
Leaf and crown diseases, Anthracnose (Collectothricum)	Transplant and furrow application	3 pints at planting or to newly planted seedlings by side dressing or shank application	Low Disease Pressure: Apply lower rate at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle.  High Disease Pressure: Apply higher rate at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.

\*Not for use in California.

**PINEAPPLES**

Use AGRI-FOS 400® for effective control of Phytophthora root and heart rot diseases in pineapples.

Disease	Application Method	Rate	Application Program
Phytophthora root and heart rot (Phytophthora cinnamomi and parasitica spp)	Foliar spray	2½ quarts in 25-50 gallons water per acre.	Apply to tops, 14 days prior to harvest of planting material.
	Pre-plant dip	1¼ quarts in 100 gallons	Will treat enough slips to plant one acre.

24/  
72

	Foliar spray	water 2 quarts in 100 gallons water	Established plantings when conditions favor disease. Apply at 90 day intervals.  Ensure thorough coverage of plants.
--	--------------	---	--

**POTATOES, POST-HARVEST\***

Use AGRI-FOS 400® only on russet-skinned potatoes intended for processing for suppression of late blight (*Phytophthora infestans*), pink rot (*Phytophthora erythroseptica*), and powdery mildew. Apply 16.5 fl. oz. of AGRI-FOS 400® per ton of tubers in one half gallon of water as a mist spray. For the best control, be sure the tubers are completely and evenly covered.

Disease	Application Method	Rate	Application Program
Potatoes, Post-harvest: Late blight ( <i>Phytophthora infestans</i> ) and pink rot ( <i>Phytophthora erythroseptica</i> )	Spray on tubers	16.5 fl. oz. in ½ gallon of water/ton of tubers	Apply only to russet- skinned potatoes intended for processing. For best results, be sure tubers are thoroughly and evenly covered.
Late blight ( <i>Phytophthora infestans</i> ) and pink rot ( <i>Phytophthora erythroseptica</i> )	Foliar spray Ground	2 to 2½ quarts in 50 gallons of water per acre	Begin application after plants are established and conditions favor disease development.  Low Disease Pressure: Apply lower rate at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle.  High Disease Pressure: Apply higher rate at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.  Late blight management using AGRI-FOS 400® requires the high application rate and is most effectively applied when tank mixed with other registered fungicides.
	Seed piece spray  Tuber spray	Mix a 15% v/v solution (0.3 quart with 2 quarts of water). Treat 2 tons of tubers per 1 gallon of solution.	Treat seed pieces with a full coverage spray.
Powdery mildew	Foliar spray Ground	1 to 3 quarts in 50 gallons of water per acre	Low Disease Pressure: Apply lower rate at the first onset of the disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 4 times per crop cycle.  High Disease Pressure: Apply higher rate at the first onset of the

25/72

			disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 4 times per crop cycle.
--	--	--	---

\*Not for use in California

**ROOT AND TUBER VEGETABLES**

Use AGRI-FOS 400® for effective control of foliar and root rot in ginseng, damping-off and root rot diseases in carrots, late blight disease and storage diseases such as pink rot and pythium leak, and downy mildew in potatoes, sweet potatoes, and yams.

Disease	Application Method	Rate	Application Program
* Ginseng: Foliar and root rot (Phytophthora cactorum)	Foliar spray Ground	2½ quarts in 100 gallons of water	In cool wet conditions that favor Phytophthora. Apply at 7 day intervals.  Do not exceed a total of 8 applications per season.
* Carrots: Phytophthora and Pythium spp	Foliar spray Ground	1½ quarts per acre in 25 gallons of water to 2 quarts per acre in 250 gallons of water	Apply at 14-day intervals after plant emergence, as needed. Assure good coverage.
Potatoes, Sweet Potatoes, Yams: Pink rot and Pythium leak (Phytophthora erythroseptica and Pythium spp)	In-furrow spray	2-5 quarts per acre in 10 gallons water	Apply in a band spray directly over top of potato seed just before row is closed.
Potatoes, Sweet Potatoes, Yams: Late blight, Pink rot and Pythium leak (Phytophthora infestans, Phytophthora erythroseptica and Pythium spp)	Foliar spray Ground	1¼ quarts per acre in 90-375 gallons water per acre	Apply at 5 to 14 day intervals subject to disease incidence.
* Downy mildew	Foliar spray Aerial  Foliar spray Ground	1 to 3 quarts in 10 gallons of water per acre  1 to 3 quarts in 30 gallons of water per acre	Low Disease Pressure: Apply lower rate at the first onset of the disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle.  High Disease Pressure: Apply higher rate at the first onset of the disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.

\* Not for use in California.

26/72

**STONE FRUIT\***

Use AGRI-FOS 400® for effective control of root, collar rot, almond pruning wound, and various canker disease in stone fruit such as, but not limited to, apricots, sweet and tart cherries, nectarines, peaches, plums, and fresh prunes.

Disease	Application Method	Rate	Application Program
Root and collar rot (Phytophthora spp)	Foliar spray  Basal bark spray, apply in spring and fall	2½ quarts per acre in 250 gallons water  62.4 fl. oz. + 62.4 fl. oz. of water + 3.2 fl. oz. Supa Stik PN Bark Penetrating Surfactant	Three treatments are required 1. Spring 2. Mid summer 3. Fall, post harvest  Spray a combination of AGRI-FOS 400® and Supa Stik PN around the complete circumference of the tree trunk until saturation/runoff. Spray from ground level up to 5 feet above the soil line, including the base of the first scaffolding limbs, if present. Treat in spring and fall for best results.
Almond pruning – wound cankers (Phytophthora syringae)	Paint or spray	1¼-2½ quarts in 100 gallons water	Apply to pruning wound and surrounding area, ensure area is thoroughly wet. In high disease situations use higher rate.
Suppression of armillaria root rot (Armillaria luteobublina)	Basal bark spray	1½ to 2 quarts + 2 quarts water + 1% Supa Stik PN Bark Penetrating Surfactant	Spray a combination of AGRI-FOS 400® and Supa Stik PN around the complete circumference of the tree trunk until saturation/runoff. Spray from ground level up to 5 feet above the soil line, including the base of the first scaffolding limbs, if present. For large trees larger than 18 inches DBH (Diameter at Breast Height, 4.5 feet above ground) that have been previously attacked by Armillaria root rot, make a fall application prior to leaf senesce and a spring application for best results. For trees less than 18 inches DBH, make an early spring application.

\*Not for use in California.

27/72

**STRAWBERRIES\***

Use AGRI-FOS 400® for effective control of leather rot, red stele, Phytophthora, foliar and bacterial disease in strawberries and suppression of powdery mildew.

Disease	Application Method	Rate	Application Program
Phytophthora and Pythium spp (red stele, leather rot, and root rot)  Foliar fungal and bacterial diseases (Rhizopus and Xanthomonas)	Pre-planting dip	1¼ quarts in 100 gallons water	Dip planting material in this solution for 30 minutes, then plant within one day. Use program for annual and perennial varieties.
	Foliar spray Ground	1 to 3 quarts in 50 to 100 gallons of water per acre	Annual crops, first treatment 14 to 21 days post planting, repeat at 1-2 month intervals when disease is evident.  Perennial crops, first treatment during spring growth flush, repeat at 1-2 month intervals when disease is evident. For susceptible varieties us higher rates and shorter spray intervals.
	Transplant and furrow application	3 pints at planting or to newly planted seedlings by side dressing or shank application	For leather rot, apply at 10% bloom and early fruit set, then 1 to 2 week intervals as required for disease control. In high disease situations, use higher rate and shorter spray intervals.  Grey mold and Anthracnose suppression using AGRI-FOS 400® requires use of high application rates and is most effective when tank mixed with other registered fungicides.
Phytophthora spp	Dip	Mix a 0.25% v/v solution (1 quart in 100 gallons of water)	Dip runners in the solution for 1 to 2 minutes. Plant within 48 hours. Mix a fresh solution daily.

28/72

Suppression of powdery mildew	Foliar spray Ground	2 to 2½ quarts in 50 gallons of water per acre	<p>Low Disease Pressure: Apply lower rate at the first onset of the disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle.</p> <p>High Disease Pressure: Apply higher rate at the first onset of the disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.</p>
-------------------------------	------------------------	--	---

\* Use in California limited to control of red stele, leather rot, and root rot.

**TREE NUTS\***

Use AGRI-FOS 400® for effective control of root and collar rot, and almond pruning-wound canker disease in tree nuts such as, but not limited to, almonds, beech nuts, Brazil nuts, butternuts, cashews, chestnuts, chinquapin, hazelnuts, hickory nuts, macadamia nuts, pecans, pistachios, and walnuts.

Disease	Application Method	Rate	Application Program
Other than macadamia nuts: Root and collar rot (Phytophthora spp)	Foliar spray Ground	1¼ quarts per acre in 125 gallons water	<p>Three treatments are required</p> <ol style="list-style-type: none"> <li>1. Spring</li> <li>2. Mid summer</li> <li>3. Fall, post harvest</li> </ol>
Other than macadamia nuts: Almond pruning – wound canker (Phytophthora syringae)	Paint or spray	2½ quarts in 100 gallons water	Apply to pruning wound and surrounding area, ensure area is thoroughly wet.
Macadamia nuts: Raceme blight (Phytophthora spp)	Foliar spray Ground	3¼ quarts per acre in 250 gallons of water	<p>Apply when disease is first seen and reapply at 3 week intervals.</p> <p>Spray to the point of run-off.</p>
Root rot, crown rot, trunk cankers, foliar blights (Phytophthora and Pythium spp)	Foliar spray Ground	2 to 2½ quarts in 100 gallons of water per acre	<p>Begin application after plants are established and conditions favor disease development.</p> <p>Low Disease Pressure: Apply lower rate at 3 month intervals. Do not apply more than 4 times per crop cycle.</p> <p>High Disease Pressure: Apply higher rate at monthly intervals. Do not apply more than 4 times per crop cycle.</p>
	Root dip	Mix a 0.5% v/v solution (2 quarts in 100 gallons of water)	Dip roots in the solution for 30 seconds. Plant within 48 hours. Mix a fresh solution daily.
Pecan scab	Foliar spray Ground	2 quarts in 50 to 100 gallons of water per acre	Apply preventatively with other products

29/72

			shown to be effective against pecan scab.
Pruning wound, crown and trunk cankers (Phytophthora spp)	Trunk spray	2 quarts + 2 quarts of water + 1% Supa Stik PN Bark Penetrating Surfactant	Apply higher rate when lesions are present. Clean wound sites and apply on and around the lesions using enough spray volume to thoroughly wet the lesions. Apply to the trunk from the soil line to 5 feet up the trunk. Apply one time in the Spring, Summer, and Fall.
Downy mildew (Phytophthora and Peronospora spp)  Powdery mildew	Foliar spray Aerial  Ground	1 to 2 quarts in 20 gallons of water per acre  1 to 2 quarts in 100 gallons of water per acre	Low Disease Pressure: Apply lower rate at first onset of disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle.  High Disease Pressure: Apply higher rate at first onset of disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.
Tree cankers	Basal bark spray	1½ to 2 quarts + 2 quarts water + 1% Supa Stik PN Bark Penetrating Surfactant	Spray a combination of AGRI-FOS 400® and Supa Stik PN around the complete circumference of the tree trunk until saturation/runoff. Spray from ground level up to 5 feet above the soil line, including the base of the first scaffolding limbs, if present. For large trees larger than 18 inches DBH (Diameter at Breast Height, 4.5 feet above ground) that have been previously attacked by Armillaria root rot, make a fall application prior to leaf senesce and a spring application for best results. For trees less than 18 inches DBH, make an early spring application.

\*Not for use in California.

30/  
72

### SEED TREATMENT APPLICATIONS\*

Use AGRI-FOS 400® for effective control of Phytophthora and Pythium diseases on agricultural crop seeds from crops listed elsewhere on this label. Do not use treated seed for food, feed, or oil. Dye used to color treated seed must be an EPA approved dye (refer to 40 CFR 153-155 (c)). Seed treatment on agricultural establishment in hopper-box, planter box, or other seed treatment application at or immediately before planting is within the scope of WPS, while commercial treatment of seeds is not within the scope of WPS.

Disease	Application Method	Rate
Phytophthora, Pythium and Fusarium spp	Can be applied at-planting or in commercial seed treatment operations.	8-24 fl. oz. AGRI-FOS 400® per 100 lbs of seed or 4-10 quarts AGRI-FOS 400® per ton of seed, depending on the size of the seeds to be treated.

\*Not for use in California.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

#### PESTICIDE STORAGE:

Keep this product in containers stored upright and secured with the original closure. Do not store this product near any heat source. Do not store near strong oxidants. If transfer to another container becomes necessary, ensure that the container is clearly labeled, the container is a type suitable for the product, and is clean and free of other materials.

#### PESTICIDE DISPOSAL:

Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

#### CONTAINER HANDLING:

Nonrefillable container. Do not reuse or refill this container.

Containers with a capacity of equal to or less than 5 gallons: Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Containers with a capacity greater than 5 gallons: Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

31/72

### WARRANTY AND DISCLAIMER

To the extent consistent with applicable law, all conditions and warranties and statutory or other rights of action which buyer or any other user may have against Liquid Fertiliser Pty. Ltd. are hereby excluded. To the extent consistent with applicable law, Liquid Fertiliser Pty. Ltd. hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information provided by Liquid Fertiliser Pty. Ltd. or on its behalf unless it is established that such information or advice was provided negligently and that the product has been used strictly as directed. To the extent consistent with applicable law, Liquid Fertiliser Pty. Ltd.'s liability shall in all circumstances be limited to replacement of product or a refund of the purchase price thereof.

Sub-label B – Turf and Ornamental Uses

32/72

**AGRI-FOS 400®**

SYSTEMIC FUNGICIDE FOR THE EFFECTIVE CONTROL OF VARIOUS PLANT DISEASES INCLUDING DOWNY MILDEW, PHYTOPHTHORA & PYTHIUM IN ORNAMENTALS, INTERIORSCAPES, BEDDING PLANTS, AND VEGETABLE TRANSPLANTS, PHYTOPHTHORA AND FUSARIUM IN CONIFERS, PYTHIUM IN TURF, AND PHYTOPHTHORA AND PYTHIUM DISEASES ASSOCIATED WITH STEM AND CANCKER BLIGHT (SUDDEN OAK DEATH) AND GENERAL BEECH DECLINE AND VARIOUS TREE STEM CANCKERS AND PHLOEM CANCKERS.

ACTIVE INGREDIENTS: Mono- and di-potassium salts of Phosphorous Acid*	45.8%
OTHER INGREDIENTS:	54.2%
TOTAL	100.0%

\*Contains 5.17 lbs/gallon of the active ingredients, mono- and di-potassium salts of Phosphorous Acid.  
Equivalent to 3.35 lbs Phosphorous Acid/gallon

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

Consult page 2 of this booklet for First Aid and Precautionary Statements

EPA Reg. No. 71962-1

EPA Est. No. 71962-AUS-001  
72499-GA-001  
71975-WA-001  
68253-CO-001

Manufactured for:  
**LIQUID FERTILISER PTY. LTD. (Trading as Agrichem)**  
2 Hovey Road  
Yatala, Queensland 4207  
Australia  
320-222-3026

Batch No.:

Date of Manufacture:

Net Contents: 1 qt., 1 gal., 5 gal., 25 gal., 55 gal., or 250 gal.

<b>FIRST AID</b>	
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice</li> <li>• Have person sip a glass of water if able to swallow</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor</li> <li>• Do not give anything to an unconscious person</li> </ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye</li> <li>• Call a poison control center or doctor for treatment advice</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes</li> <li>• Call a poison control center or doctor for treatment advice</li> </ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible</li> <li>• Call a poison control center or doctor for further treatment advice</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. Hotline Number: National Poison Control, 1-800-222-1222</p>	

### **PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION.** Harmful if swallowed, absorbed through skin or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist or vapors. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Wear the appropriate Personal Protective Equipment (PPE).

#### **PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Applicators, mixers, loaders, and other handlers must wear:

- protective eyewear
- long pants and long-sleeved shirt
- waterproof gloves
- shoes plus socks.

Follow manufacturer's instructions for maintaining/cleaning personal protective equipment (PPE). If no such instructions for washables, use hot water and detergent. Keep and wash PPE separately from other laundry.

When handlers use closed systems, aircraft or enclosed cabs in a manner that meets the requirements listed in the worker protection standard (WPS) for agricultural pesticides (40 CFR 170.240 (d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

<p><b>USER SAFETY RECOMMENDATIONS</b></p> <p>Users should:</p> <ul style="list-style-type: none"> <li>• Wash hands before drinking, eating, chewing gum, using tobacco or using the toilet.</li> <li>• Remove PPE clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.</li> <li>• Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.</li> </ul>
---

34/  
72

## ENVIRONMENTAL HAZARDS

For Terrestrial Use: Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the state or tribal agency responsible for pesticide registration.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, CFR 40 part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI) of 4 hours**, unless wearing the appropriate PPE.

PPE required for early entry to treated areas that are permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soils or water, is: coveralls worn over short-sleeved shirt and short pants, waterproof gloves, shoes plus socks and protective eyewear.

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170): The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Do not enter treated areas without protective clothing until sprays have dried.

## CHEMIGATION

Use of AGRI-FOS 400® through chemigation is not permitted in California.

Apply this product only through center pivot, solid set or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

35/72

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

#### SPRINKLER AND DRIP (TRICKLE) IRRIGATION SYSTEMS:

The irrigation system must contain a functional check valve, vacuum relief valve and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

(Sprinkler Chemigation Only) Do not apply when wind speed favors drift beyond the area intended for treatment.

Apply AGRI-FOS 400® continuously for the duration of the water application. After treatment with AGRI-FOS 400® has been completed, avoid further irrigation of the treated area until foliage is dry or for 24 to 48 hours.

#### GENERAL APPLICATION INSTRUCTIONS

Apply AGRI-FOS 400® by various application methods, including foliar spray (aerial and ground), soil drench, soil incorporation, basal bark application and bare root dip. For foliar sprays, apply AGRI-FOS 400® with sufficient water volumes for adequate coverage of foliage, according to crop and growth stage. To ensure good coverage, spray to wetness, but avoid run-off. **This product does not have a pre-harvest interval (PHI). Harvest when dry.**

When using AGRI-FOS 400® with Supa Stik PN adhere to all applicable label directions. Only use Supa Stik PN with basal bark applications.

#### MIXING INSTRUCTIONS

5. Fill the spray tank with ¼ - ½ of the volume of water required before adding AGRI-FOS 400®
6. Add AGRI-FOS 400® slowly to the tank and agitate by mechanical or hydraulic means
7. Continue agitating as tank fills with water to the desired volume
8. Maintain agitation during application

#### COMPATIBILITY

AGRI-FOS 400® is compatible with most products used in agriculture. However, individual crop sensitivity to these mixtures may vary. Mixtures of AGRI-FOS 400® with some foliar fertilizers and copper products are not always compatible or cause phytotoxicity to some plants. If these combinations or others have not been used previously, do not tank mix without first testing the compatibility of the tank mix. Do not apply tank mixture without first assessing its safety to the

36/72

crop (phytotoxicity). Tank mix AGRI-FOS 400® with fertilizers only if crop safety has been established and the AGRI-FOS 400® use rates are followed.

Due to AGRI-FOS 400®'s acidic nature, do not use acidifying-type compatibility agents. If spray adjuvants are used test them before use to confirm compatibility with AGRI-FOS 400®.

Use a jar test to confirm compatibility with AGRI-FOS 400®. In a clean jar using the same water source that is normally used to fill spray tank, add the same proportions of each product and the appropriate quantity of water and mix thoroughly. Let stand for 3 minutes. If mixture remains in solution or is remixed readily the tank mix is compatible.

Spray the solution that results from the above compatibility test onto a few plants and inspect for visual effects of phytotoxicity (leaf burn) 3 days later.

**GRASS GROWN FOR SEED PRODUCTION\***

Use AGRI-FOS 400® for effective control of damping-off and root rot diseases in turf grasses such as, but not limited to, Bermuda, fescue, bent, blue, rye, zoysia, buffalo, and poa annua.

Disease	Application Method	Rate	Application Program
Phytophthora and Pythium spp	Foliar spray Ground	1½ quarts per acre in 25 gallons of water to 2 quarts per acre in 250 gallons of water	Apply at 14-21 day intervals, as necessary. Ensure thorough coverage.

\*Not for use in California.

**LANDSCAPE, GOLF COURSE, NURSERY, FORESTRY, AND PARK APPLICATIONS\***

Use AGRI-FOS 400® for effective control of Phytophthora and Pythium and other diseases associated with Stem and Canker Blight (Sudden Oak Death, Phytophthora ramorum), Beech Decline, and general tree decline syndromes, and for suppression of Anthracnose and Verticillium wilt in landscapes, nurseries, golf courses, forests, and parks. Apply AGRI-FOS 400® to trees such as, but not limited to, Ash, Azalea, Beech, Black Locust, Buckeye, Catalpa, Cedar, Cherry – Stonefruits, Chestnut, Coffee Tree, Crab Apple, Cork Tree, Dogwood - All, Elder, Elm, Fir, Hawthorne, Honey Locust, Juniper, Lilac, Linden, Magnolia, Maples – All, Pine, Oaks – All, Olives, Ornamental Pear, Plum, Pyracantha, RedBud, Smoke Tree, Sumac, Sweet Birch, Sweet Gum, Sycamore, Viburnum, White Pine, White Cedar, and Willow.

Make applications before disease development and in conjunction with good cultural management practices. Use higher rate of application when disease pressure is severe. Do not exceed indicated application rates or apply more frequently than stated on label or tree injury may occur. Do not apply to trees that are heat or moisture stressed. Do not exceed indicated spray intervals or label rates in order to avoid tree injury. When applying to indoor plants do not overspray and use care to apply only to target plants. If meeting these conditions is not possible, remove plants to an outdoor location for treatment and drying before bringing back indoors.

Disease	Application Method	Rate	Application Program
Phytophthora and Pythium spp, and Phytophthora ramorum	Injection	11 fl. oz. per 21 fl. oz. of water Or 1/2 tsp. per tsp. of water	Drill holes 3/16 inch (5 mm) in diameter <u>into live sapwood</u> (depth dependent upon age of tree) with downward angle into trunk uniformly around the tree circumference, using a slow drill. Do not inject into areas of obvious decay, canker or mechanical injury that appear on the tree trunk. Calculate the amount of product required by measuring the trees by one of the following 3 methods, and use the highest calculated number of injections. 1) 1 injection per square yard of canopy; 2) 1 injection per yard of diameter of canopy measured at the drip-line; 3) 1 injection per 6 inches of trunk circumference measured 4 feet above soil level. Make injections with applicators that maintain positive pressure differential such as ChemJet®, Sidewinder®, Ag-murf Gun®, Marley® Injector, Smart Shot injector, or hydraulic applicator type equipment that forces solution into the sapwood of the tree.
Phytophthora and Pythium spp, and Phytophthora ramorum	Basal bark spray (all other tree species) apply in spring and fall Best for thin bark trees such as maples, lindens, sycamores, and dogwoods	62.4 fl. oz. + 62.4 fl. oz. of water + 1.5 to 3.0 fl. oz. Supa Stik PN Bark Penetrating Surfactant	Spray a combination of AGRI-FOS 400® and Supa Stik PN around the complete circumference of the tree trunk until saturation/runoff. Spray from ground level up to 5 feet above the soil line, including the base of the first scaffolding limbs, if present. Treat in spring and fall for best results.

<p>Fusarium subglutinans (Pine Pitch Canker)</p>	<p>Basal bark spray (pines) apply anytime active growth is observed</p>	<p>1 gallon of AGRI-FOS 400® + 2 gallons of water + 4 fl. oz. of Supa Stik PN Bark Penetrating Surfactant</p>	<p>Apply uniformly to 5 to 6 feet of trunk circumference. Spray from top down to ground level from either first branch or from as high as possible without exposing applicator to drift. Spray to saturation/ runoff. Various types of application equipment can be used such as hydraulic sprayers, handheld pump-type sprayers, and backpack sprayers.</p>
<p>Fusarium subglutinans (Pine Pitch Canker) and Gnomonia platani (Sycamore Anthracnose)</p>	<p>Injection</p>	<p>4 tsp per tree of a 1 gallon AGRI-FOS-400® + 2 gallons of water + 4 fl. oz. of Supa Stik PN solution.</p>	<p>Drill holes 3/16 inch (5 mm) in diameter and 1 inch (25 mm) to 2 inches (50 mm) deep with slight downward angle in trunk. Place syringes in the main trunk of the tree and space evenly around the circumference of the trunk. Suitable for use with Chemjet tree injectors, Ag-murf gun, Smart Shot injector, or positive pressure hydraulic tree injection. Trees should be at least 10" diameter at breast height.</p>
<p>Fire blight</p>	<p>Foliar spray</p>	<p>1½ to 2 quarts per 100 gallons of water</p>	<p>First application pre bloom (bud swell or sliver tip stage). Application intervals: 7 days until end of bloom period.  Apply spray to thoroughly wet all foliage.</p>
<p>** Suppression of Anthracnose</p>	<p>Foliar spray Ground</p>	<p>2 quarts per 100 gallons of water per acre</p>	<p>Apply at pre bloom (bud swell or green tip stage) with a supplemental application 14 days later with AGRI-FOS 400® or other fungicide compound with efficacy on Anthracnose.</p>

39/72

<p><b>** Suppression of Anthracnose</b></p>	<p>Basal bark spray apply early spring at bud swell until green tip stage of growth</p>	<p>62.4 fl. oz. + 62.4 fl. oz. of water + 1.5 to 3.0 fl. oz. Supa Stik PN Bark Penetrating Surfactant</p>	<p>Spray a combination of AGRI-FOS 400® and Supa Stik PN around the complete circumference of the tree trunk until saturation/runoff. Spray from ground level up to 5 feet above the soil line, including the base of the first scaffolding limbs, if present. For large trees larger than 18 inches DBH (Diameter at Breast Height, 4.5 feet above ground) that have been previously attacked by Anthracnose, make a fall application prior to leaf senesce and a spring application for best results. For trees less than 18 inches DBH, make an early spring application.</p>
<p><b>** Suppression of Verticillium wilt</b></p>	<p>Foliar spray Ground</p>	<p>2 quarts in 100 gallons of water per acre</p>	<p>For trees previously identified with infections, apply first application pre-bloom. Repeat applications between 21 to 30 days.</p>
<p><b>** Suppression of Verticillium wilt</b></p>	<p>Basal bark spray</p>	<p>1½ to 2 quarts + 2 quarts of water + 2 fl. oz. Supa Stik PN Bark Penetrating Surfactant</p>	<p>Spray a combination of AGRI-FOS 400® and Supa Stik PN around the complete circumference of the tree trunk until saturation/runoff. Spray from ground level up to 5 feet above the soil line, including the base of the first scaffolding limbs, if present. For large trees larger than 24 inches DBH (Diameter at Breast Height, 4.5 feet above ground) that have been previously attacked by Verticillium wilt, make a fall application prior to leaf senesce and a spring application for best results. For trees less than 24 inches DBH, make an early spring application.</p>
<p><b>**Black spot (Diplocarpon spp)</b></p>	<p>Foliar spray Ground</p>	<p>1 to 2 quarts per 100 gallons of water</p>	<p>Apply uniformly to foliage preventatively or at first sign of disease.</p>

\*Use in California limited to Oaks (Coastal, Live, Shreve, Black, Canyon), Tan Oaks, and other tree species that are host to *P. ramorum*. Do not apply to any other tree species without consulting the list of *P. ramorum* host species listed at the following website: <http://nature.berkeley.edu/comtf/index.html>. Make applications by injection and basal bark spray of pines, basal bark spray of apples and crabapples, and injection of Sycamores for control of Sycamore anthracnose. For questions contact your local extension office.

\*\* Not for use in California.

40/72

## ORNAMENTAL APPLICATIONS

Use AGRI-FOS 400® for effective control of Bacterial blight, Downy mildew, powdery mildew, Phytophthora spp, and Pythium spp diseases, and for suppression of Anthracnose of Ornamentals in landscapes, nurseries, golf courses, parks, interiorscapes, and greenhouses. Apply AGRI-FOS 400® to plants such as, but not limited to, Aglaonema, Anthurium, Aphelandra, Arborvitae, Azaleas, Bougainvillea, Boxwood, Cattelya skinneri, Ceanothus, Cotoneaster, Cissus, Diefenbachia, English ivy, Eucalyptus, Ficus, Hibiscus, Japanese andromeda, Japanese Holly, Leather leaf Fern, Peperomia, Photinia, Pittosporum, Philodendron, Pieris, Pothos, Rhododendron, Roses (container, field, landscape, and mini varieties), Schefflera, Sedum, Sempervivum, Syngonium, Spathiphyllum, Taxus media, and Zygocactus.

Make applications before disease development and in conjunction with good cultural management practices. Use higher rate of application when disease pressure is severe. Do not exceed indicated application rates or apply more frequently than stated on label or plant injury may occur. Do not apply to plants that are heat or moisture stressed. Do not apply to plants that are in a state of dormancy. Do not exceed indicated spray intervals or label rates in order to avoid plant injury. When applying to indoor plants do not overspray and use care to apply only to target plants. If meeting these conditions is not possible, remove plants to an outdoor location for treatment and drying before bringing back indoors.

Disease	Application Method	Rate	Application Program
Bacterial blight (Xanthomonas campestris) pathogens: dieffenbachiae, fici hederiae, and syngonli  Downy mildew  Phytophthora spp, Phytophthora ramorum, and Pythium spp	Foliar spray Ground	2-4 pints per 100 gallons of water Or 2-4 tsp. per gallon of water	Apply spray to thoroughly wet all foliage. Application intervals: 7 to 14 days Repeat as required.
	Foliar spray Ground	1-2 quarts per 100 gallons of water Or 1-2 tsp. per gallon of water	Apply spray to thoroughly wet all foliage. Application intervals: 14 to 21 days. Repeat as required.
	Soil drench	6¼-12¼ fl. oz. per 100 gallons of water	Apply each 25 gallons of solution to an area of 100 square feet. Follow application with irrigation. Repeat as required. Limit of one application per month.
	Soil incorporation	1-2 pints per cubic yard of soil	Just prior to potting, mix 1 to 2 pts. of AGRI-FOS 400® into each cubic yard of growing media. If disease pressure is high, apply foliar spray or soil drench.
	Bare rooted dipping of transplants	2 pints per 100 gallons of water Or 2 tsp. per gallon	Immediately before transplanting, dip transplants for two minutes, keep roots submerged, ensure root mass is thoroughly wet.
* Powdery mildew	Foliar spray Ground	1½ to 2½ quarts in 50 gallons of water per acre	Low Disease Pressure: Apply lower rate at the first onset of the disease. Repeat

41/72

			<p>applications at 1-2 week intervals.</p> <p>High Disease Pressure: Apply higher rate at the first onset of the disease. Repeat applications at 7-10 day intervals. Do not apply more than 6 times per crop cycle.</p> <p>Application amount depends upon plant type, maturity, and application technique/method.</p>
* Suppression of Anthracnose	Foliar spray Ground	2 to 2½ quarts in 50 gallons of water per acre	Apply prior to onset of disease. Apply spray to saturation or runoff.
* Black spot (Diplocarpon spp)	Foliar spray Ground	1 to 2 quarts per 100 gallons of water	Apply uniformly to foliage preventatively or at first sign of disease.

\* Not for use in California.

### NURSERY AND BEDDING PLANTS

Use AGRI-FOS 400® for effective control of Downy mildew, Phytophthora spp and Pythium spp diseases of BEDDING PLANTS grown in landscapes, nurseries and golf courses, parks, interiorscapes, and greenhouses. Apply AGRI-FOS 400® to plants such as, but not limited to; Ageratum, Aglaonema, Algerian Ivy, Anthurium, Aphelandra, Arborvitae, Artemesia, Aster, Azaleas, Begonia, Baby's Breath, Bougainvillea, Boxwood, Caladium, Carnation, Cattelya Skinneri, Ceanothos, Cissus, Chrysanthemum, Coleus, Columbine, Cotoneaster, Daisy, Delphinium, Dieffenbachia, Dogwood, Easter Lily, English Ivy, Ficus, Foxglove, Gaillardia, Geranium, Gloxinia, Hibiscus, Impatiens, Japanese Holly, Juniper, Lavender, Leather Leaf Fern, Marigold, Monterey Pink, Pansy, Pieris, Peperomia, Petunia, Philodendron, Phlox, Photinia, Pinks, Pittosporum, Poinsettia, Pothos, Primrose, Prostrate Rosemary, Rosemary, Rhododendron, Salvia, Schefflera, Sedium, Sempervivum, Snapdragon, Spathiphyllum, Taxus Media, Verbena, Vinca, White Cedar, White Pine, Zinnia, and Zygocactus. AGRI-FOS 400® may also be used for Vegetable Transplants grown in Greenhouse, Lathhouse, or Shade house sites.

Make applications before disease development and in conjunction with good cultural management practices. Use higher rate of application when disease pressure is severe. Do not exceed indicated application rates or apply more frequently than stated on label in order to avoid plant injury. Do not apply to plants that are heat or moisture stressed. When applying to indoor plants do not overspray and use care to apply only to target plants. If meeting these conditions is not possible, remove plants to an outdoor location for treatment and drying before bringing back indoors.

Disease	Application Method	Rate	Application Program
Downy Mildew	Foliar Spray Ground	1¼ to 2½ quarts per 100 gallons of water Or ½ to 1 1/8 fl. oz. per gallon of water	Apply spray to thoroughly wet all foliage. Application intervals: 14 to 21 days. Repeat as required.
Phytophthora spp and Pythium spp	Foliar spray Ground	1 to 2 quarts per 100 gallons of water Or 2 to 4 tsp. per gallon of water	Apply spray to thoroughly wet all foliage. Application intervals: 14 to 21 days. Repeat as required. Note do not apply more than 500 gallons of

42/72

	Soil drench	6¼ to 12¾ fl. oz. per 100 gallons of water	spray solution per acre. Apply each 25 gallons of solution to an area of 100 square feet. Follow application with irrigation. Repeat as required. Limit of one application per month.
Phytophthora spp*	Foliar spray Ground	2 quarts per acre	Apply spray in 20-60 gallons per acre.
Phytophthora spp*	Hand gun	2 quarts per 100 gallons of water	Apply spray to thoroughly wet all foliage.

\*Lavender applications.

**CONIFERS IN COMMERCIAL NURSERIES, PLANTATIONS AND FORESTS\*  
(INCLUDING CHRISTMAS TREES)**

Apply AGRI-FOS 400® in conjunction with good cultural management practices for effective control of root rot (Phytophthora spp) in CONIFERS including, but not limited to, Pines, Spruce and Douglas Fir. Use higher rate of application when disease pressure is severe. Do not exceed indicated application rates or apply more frequently than stated on label in order to avoid plant injury. Do not apply to CONIFERS that are moisture or heat stressed.

Disease	Application Method	Rate	Application Program
Phytophthora	Foliar spray Ground	1 to 2 quarts per 100 gallons of water Or 2 to 4 tsp. per gallon of water	Apply spray to thoroughly wet all foliage. Application intervals: 14 to 21 days. Repeat as required.
	Soil drench	1 to 2 quarts per 100 gallons of water Or 2 to 4 tsp. per gallon of water	Apply one gallon of solution per sq yd. Follow application with irrigation. Application intervals: 14 to 21 days. Repeat as required.
	Bare root dipping at transplanting	1 quart per 100 gallons of water Or 2 tsp. per gallon of water	Immediately before transplanting, dip transplants for two minutes; keep roots submerged and ensure root mass is thoroughly wet.
Fusarium subglutinans (Pine Pitch Canker)	Basal bark spray apply any time active growth is observed	1 gallon AGRI-FOS 400® + 2 gallons of water + 4 fl. oz. of Supa Stik PN Bark Penetrating Surfactant	Spray a combination of AGRI-FOS 400® and Supa Stik PN around the complete circumference of the tree trunk until saturation/runoff. Spray from ground level up to 5 feet above the soil line.
Fusarium subglutinans (Pine Pitch Canker)	Injection	4 tsp. per tree of a 1 gallon AGRI-FOS 400® + 2 gallons of water	Drill holes 3/16 inch (5 mm) in diameter and 1 inch (25 mm) to 2 inches (50 mm) deep with slight downward angle in trunk. Place syringes in the main trunk of the tree and space evenly around the circumference of the trunk. Suitable for use with Chemjet tree injectors, Ag-murf gun, or hydraulic tree injection. Trees should

43/  
72

		be at least 10" diameter at breast height.
--	--	--

\*Not for use in California except as an injection and/or basal bark spray on pines.  
Do not graze livestock in treated areas of conifer nurseries or plantations. Do not feed forage from treated areas of plantations and or nurseries.

**TURF\***

Use AGRI-FOS 400® for the effective control of Rhizoctonia; Pythium and damping-off diseases, and for suppression of Anthracnose of turf grasses on golf courses, parks, commercial landscapes, commercial turf production, and sod farms. When conditions favor disease, begin preventive applications and repeat at indicated intervals. Use higher rate of application when disease pressure is severe.

Disease	Application Method	Rate	Application Program
Pythium	Foliar spray Ground	5 to 10 fl. oz. per 1000 sq. ft.	Apply indicated quantity of product in 1 to 2 gallons of water per 1000 sq. ft. Ensure foliage is thoroughly wet.  Application intervals: 14-21 days. Repeat as required. Do not irrigate or mow treated areas until spray has completely dried.
Suppression of Anthracnose	Foliar spray Ground	5 to 10 fl. oz. in 2 gallons of water per 1000 sq. ft.	Apply prior to onset of disease. Apply every 7 to 14 days.
Rhizoctonia	Foliar spray Ground	5 to 10 fl. oz. in 2 gallons of water per 1000 sq. ft.	Apply prior to onset of disease. Apply every 7 to 14 days.

\* Use in California limited to control of pythium.

**TURF TANK MIXTURES**

For the effective control of summer stress complex caused by a complex of Rhizoctonia and Pythium diseases, tank-mix AGRI-FOS 400® with Fore® WP [or Protect T/O or mancozeb-containing] fungicide and apply to turf grasses on golf courses, parks, commercial landscapes, commercial turf production, and sod farms.

Product	Disease	Rate per 1000 sq. ft.	Application Program
AGRI-FOS 400® + FORE® WP** [or Protect T/O or mancozeb-containing fungicide]	Summer Stress Complex (Rhizoctonia and Pythium spp.)	5 to 10 fl. oz. AGRI-FOS 400® + 4 to 8 fl. oz. FORE® WP** [or Protect T/O or mancozeb-containing fungicide]	Apply indicated quantity of product in 1 to 5 gal. of water per 1000 sq. ft. as a foliar spray. Start as a preventive spray at two-week intervals and repeat as required. Do not irrigate or mow treated areas until spray has completely dried.

\*\*Registered trade mark of Rohm & Haas.

Do not graze animals on treated areas of turf. Do not feed treated turf clippings to poultry or livestock.

44/72

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

#### **PESTICIDE STORAGE:**

Keep this product in containers stored upright and secured with the original closure. Do not store this product near any heat source. Do not store near strong oxidants. If transfer to another container becomes necessary, ensure that the container is clearly labeled, the container is a type suitable for the product, and is clean and free of other materials.

#### **PESTICIDE DISPOSAL:**

Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

#### **CONTAINER HANDLING:**

Nonrefillable container. Do not reuse or refill this container.

Containers with a capacity of equal to or less than 5 gallons: Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Containers with a capacity greater than 5 gallons: Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### WARRANTY AND DISCLAIMER

To the extent consistent with applicable law, all conditions and warranties and statutory or other rights of action which buyer or any other user may have against Liquid Fertiliser Pty. Ltd. are hereby excluded. To the extent consistent with applicable law, Liquid Fertiliser Pty. Ltd. hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information provided by Liquid Fertiliser Pty. Ltd. or on its behalf unless it is established that such information or advice was provided negligently and that the product has been used strictly as directed. To the extent consistent with applicable law, Liquid Fertiliser Pty. Ltd.'s liability shall in all circumstances be limited to replacement of product or a refund of the purchase price thereof.

45/72

Sub-label C – Residential Uses

**AGRI-FOS 400®**

FOR RESIDENTIAL USE ONLY

SYSTEMIC FUNGICIDE FOR THE EFFECTIVE CONTROL OF VARIOUS PLANT DISEASES INCLUDING BLACK SPOT OR SCAB IN APPLE, ROOT ROT IN AVOCADO, BUD ROT AND NUT FALL IN COCONUT, ROOT ROT IN CITRUS AND CUCURBITS, DOWNY MILDEW IN CUCURBITS, GRAPE, LETTUCE, AND ONION, ANTHRACNOSE IN MANGO, ROOT AND HEART ROT IN PINEAPPLE, LATE BLIGHT IN POTATO, ROOT AND COLLAR ROT IN STONEFRUIT, LEATHER ROT AND PHYTOPHTHORA DISEASES IN STRAWBERRY, LATE BLIGHT IN TOMATO, DOWNY MILDEW, PHYTOPHTHORA & PYTHIUM IN ORNAMENTALS, INTERIORSAPES, BEDDING PLANTS, VEGETABLE TRANSPLANTS, AND HERBS AND SPICES, PHYTOPHTHORA AND FUSARIUM IN CONIFERS; PYTHIUM IN TURF AND LAWNS IN ALL AREAS, AND PHYTOPHTHORA AND PYTHIUM DISEASES ASSOCIATED WITH STEM AND CANER BLIGHT (SUDDEN OAK DEATH) AND GENERAL BEECH DECLINE AND VARIOUS TREE STEM CANKERS AND PHLOEM CANKERS.

[FOR USE ON: ORNAMENTALS, TURF, FRUIT & MORE, AS LISTED]

[CONTROLS DAMPING-OFF AND ROOT ROT IN FRUITING VEGETABLES, ORNAMENTALS, AND BEDDING PLANTS, AS LISTED.]

[CONTROLS: ROOT ROT, DOWNY MILDEW, FIRE BLIGHT, BLACK SPOT & MORE, AS LISTED]

[Systemic Action: fungicide moves throughout the entire plant to treat diseases, as listed.]

ACTIVE INGREDIENTS: Mono- and di-potassium salts of Phosphorous Acid*	45.8%
OTHER INGREDIENTS:	54.2%
TOTAL	100.0%

\*Contains 5.17 lbs/gallon of the active ingredients, mono- and di-potassium salts of Phosphorous Acid.

Equivalent to 3.35 lbs Phosphorous Acid/gallon

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

Consult page 2 of this booklet for First Aid and Precautionary Statements

EPA Reg. No. 71962-1

EPA Est. No. 71962-AUS-001  
72499-GA-001  
71975-WA-001  
68253-CO-001

Manufactured for:  
**LIQUID FERTILISER PTY. LTD. (Trading as Agrichem)**  
2 Hovey Road  
Yatala, Queensland 4207  
Australia  
320-222-3026

Batch No.:

Date of Manufacture:

Net Contents: 1 pt., 1 qt., 1 gal., or 5 gal.

46/72

<b>FIRST AID</b>	
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice</li><li>• Have person sip a glass of water if able to swallow</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor</li><li>• Do not give anything to an unconscious person</li></ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye</li><li>• Call poison control center or doctor for treatment advice</li></ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes</li><li>• Call a poison control center or doctor for treatment advice</li></ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"><li>• Move person to fresh air</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible</li><li>• Call a poison control center or doctor for further treatment advice</li></ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment. Hotline Number: National Poison Control, 1-800-222-1222	

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS.**

**CAUTION.** Harmful if swallowed, absorbed through skin or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist or vapors. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

**ENVIRONMENTAL HAZARDS**

For Terrestrial Use: To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read entire label before using this product. For use only in home gardens, on home lawns, and on home ornamentals and related home plants.

When using AGRI-FOS 400® with Supa Stik PN adhere to all applicable label directions. Only use Supa Stik PN with basal bark applications.

**GENERAL APPLICATION INSTRUCTIONS**

Apply AGRI-FOS 400® by various application methods, including foliar spray, soil drench, soil incorporation, basal bark application and bare root dip. For foliar sprays, apply AGRI-FOS 400® with sufficient water volumes for adequate coverage of foliage, according to plant type and growth stage. To ensure good coverage, spray to wetness, but avoid run-off. **Harvest when dry.**

47/  
72

When applying AGRI-FOS 400® to plant species for the first time, spray a limited number of plants first and wait for 3 days. Then check for leaf burn (phytotoxicity).

**MIXING INSTRUCTIONS**

1. Fill the spray tank with ¼ - ½ of the volume of water required before adding AGRI-FOS 400®
2. Add AGRI-FOS 400® slowly to the tank and agitate
3. Fill tank with balance of water to the desired volume
4. Agitate during application

Conversion Table			
1/8 fl. oz.	=	¾ teaspoon (tsp.)	
¼ fl. oz.	=	1½ tsp.	
1/3 fl. oz.	=	2 tsp.	
½ fl. oz.	=	3 tsp.	
2/3 fl. oz.	=	4 tsp.	
¾ fl. oz.	=	4½ tsp.	
1 fl. oz.	=	2 tablespoons (Tbs.)	= 6 tsp.

**CITRUS, FRUIT, NUT AND VEGETABLE APPLICATIONS**

**APPLES, CRAB APPLES, LOQUATS, PEARS & QUINCE\***

Use AGRI-FOS 400® for effective control of black spot, root and collar rot and fire blight in apples, crab apples, loquats, pears, and quinces.

Disease	Application Method	Rate	Application Program
Apple black spot and scab (Venturia inaequalis)	Foliar spray	3-4 tsp. per gallon of water	Apply at ¼ - ½ inch green tip through first cover at 7 to 10 day intervals or according to forecasted infection events. Continue with AGRI-FOS 400® and Captan in the remaining applications.
Apple black spot and scab (Venturia inaequalis)	Basal bark spray apply early spring at bud swell or at silver tip stage of growth	16 fl. oz. + 16 fl. oz. of water + 1 fl. oz. Supa Stik PN Bark Penetrating Surfactant	Spray a combination of AGRI-FOS 400® and Supa Stik PN around the complete circumference of the tree trunk until saturation/runoff. Spray from ground level up to 5 feet above the soil line, including the base of the first scaffolding limbs, if present (treatment generally lasts 8-12 weeks depending on pathogen levels. Higher disease pressure will shorten the length of control). Various types of application equipment can be used such as hydraulic sprayers, handheld pump-type sprayers, backpack



49/  
72

**AVOCADOS**

Use AGRI-FOS 400® for effective control of root rot, trunk cankers and downy mildew disease in Avocado.

Disease	Application Method	Rate	Application Program
Root rot (Phytophthora cinnamomi)	Tree injection	Skeletal trees 1st year: ¼ fl. oz. undiluted product per yard of canopy diameter. Other situations: 1/8 fl. oz. diluted with ½ fl. oz. of water per yard of canopy diameter	Inject trees at spring flush maturity. Repeat treatment in February or March. Drill holes 3/16 inch (5 mm) in diameter and 1 inch (25 mm) to 2 inches (50 mm) deep with slight downward angle in trunk. Place syringes in the main trunk of the tree and space evenly around the circumference of the trunk. Suitable for use with Chemjet tree injectors, Ag-murf gun, Smart Shot injector, or hydraulic tree injection. Do not prune back trees before injection process as burning of new growth may occur. Do not inject trees in winter months. Do not cut back the canopy of injected trees. Do not add any material, other than water, to AGRI-FOS 400® by trunk injection. Do not inject more liquid in a lesser number of syringes than directed.
	Foliar spray	1/3 fl. oz. per gallon of water	Spray to run off at 2-2½ gallons of spray solution per adult tree. Start applications in spring, up to 4 applications a year at two-month intervals. Ensure thorough coverage
Canker (Phytophthora citricola)	Trunk spray	8 to 16 fl. oz. with one gallon of water with 1/2 fl. oz. of Supa Stik PN Bark Penetrating Surfactant	Apply to trunk lesions using sufficient spray volume to completely wet the trunk and lesions. If lesions absent, apply to trunk from soil level to two feet up trunk. If lesions present use higher rate
Downy mildew	Foliar spray	¾ teaspoon per gallon of water	Spray to run-off, as required for disease control

**BERRIES\***

Use AGRI-FOS 400® for effective control of root rot, general leaf and berry diseases such as those caused by Septoria spp., and downy mildew, and for suppression of Anthracnose spp. in bush and cane berries such as, but not limited to, bingleberries, blackberries, black satin berries, blueberries, boysenberries, Cherokee blackberries, Chesterberries, Cheyenne blackberries, coryberries, cranberries, darrowberries, dewberries, Dirksen thornless berries, elderberries, Himalayaberries, huckleberries, hullberries, Lavacaberries, loganberries, lowberries, lucretiaberries, mammoth blackberries, marionberries, mulberries, nectarberries, olallieberries, Oregon evergreen berries, raspberries (red, black, hybrids/cultivars), and youngberries.

Disease	Application Method	Rate	Application Program
<p>Root rot (Phytophthora spp.)</p>	<p>Foliar spray</p>	<p>2-6 tsp. per gallon of water. Ensure foliage is completely wet.</p>	<p>New plantings: start application when new growth is 2 to 3 inches long. Established plantings start applications when cool wet conditions occur which favor disease. West of Rocky Mountains: Autumn applications, apply when conditions favor disease, repeat in 4 weeks. Spring applications, first application after bud break and repeat in 4 weeks. East of Rocky Mountains: First application spring post bud break (2 to 3 inches new growth) and repeat at 50 to 60 day intervals. Do not exceed 4 applications per growing season.</p> <p>For blueberries - First application in spring at pink bud and then on a regular schedule of application at two to three intervals.</p>
<p>General leaf and berry diseases such as those caused by Septoria spp and Anthracnose spp</p>	<p>Root dip</p>	<p>Mix a 2.5% v/v solution (3 fl. oz./1 gallon of water)</p>	<p>Apply as pre-plant dip to the roots for 2-3 minutes. Plant within 48 hours after dipping. Mix a fresh solution daily.</p>
<p>Downy mildew</p>	<p>Foliar spray</p>	<p>2 to 4 tsp. per gallon of water</p>	<p>Low Disease Pressure: Apply lower rate at first onset of disease. Repeat applications at 1 to 3 week intervals. Do not apply more than 6 times per growing season.</p> <p>High Disease Pressure: Apply higher rate at first onset of disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per growing</p>

51/72

			<p>season.</p> <p>AGRI-FOS 400® is best when used in combination with conventional registered fungicides to increase the performance of the disease control program.</p>
--	--	--	--

\*Not for use in California.

**BRASSICAS\***

Use AGRI-FOS 400® for effective control of downy mildew, damping-off, root rot, bacterial, and fungal diseases in brassicas such as, but not limited to, broccoli, Brussels sprouts, cabbage, cauliflower, cavalo broccolo, collards, Chinese cabbage, Chinese mustard cabbage, kale, kohlrabi, mizuna, mustard greens, mustard spinach and rape greens.

Disease	Application Method	Rate	Application Program
Root rot and damping off (Phytophthora spp and Pythium spp)	Foliar spray	1 to 2 fl. oz. per gallon of water	<p>Begin application after plants are established and conditions favor disease development.</p> <p>Low Disease Pressure: Apply lower rate at 1 to 2 week intervals. Do not apply more than 6 times per growing season.</p> <p>High Disease Pressure: Apply higher rate at 7 to 10 day intervals. Do not apply more than 6 times per growing season.</p>
	Pre-plant seedling nursery treatment	2 tsp. in 1 gallon of water to plants in seedling trays 1 to 7 days prior to out planting	
	Transplant and furrow application	3 pints at planting or to newly planted seedlings by side dressing	
Downy mildew (Peronospora parasitica)	Foliar spray	<p>2 tsp. to 2 fl. oz. per gallon of water</p> <p>California: 1/3 to 1/2 fl. oz. per gallon of water</p>	<p>1 to 3 week intervals between applications when conditions favor disease development (cool, moist weather). Use higher rates and shorter intervals when disease pressure increases.</p>
Diseases caused by Septoria, Collectotrichum, Alternaria, and powdery mildew	Foliar spray, ground	1 to 2 fl. oz. per gallon of water	<p>Low Disease Pressure: Apply lower rate at first onset of disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per growing season.</p> <p>High Disease Pressure: Apply higher rate at first onset of disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per growing season.</p>

\* Use in California limited to control of downy mildew.

52/72

**CITRUS - Mature trees**

Use AGRI-FOS 400® for effective control of root rot and collar rot diseases in citrus.

Disease	Application Method	Rate	Application Program
Brown rot and foot rot Phytophthora spp	Foliar spray	1 teaspoon per gallon of water	<b>When conditions favor disease.</b> Spray trees to run off ensure even coverage. Do not apply at high temperatures (above 95°F) particularly if humidity is low or to moisture-stressed trees
Root rot and collar rot (Phytophthora spp nicotianae and Phytophthora citrophthora)	Trunk spray	8 to 16 fl. oz. per gallon of water with ¼ to ½ fl. oz. of Supa Stik PN Bark Penetrating Surfactant	Spray trunk lesions with enough spray volume to ensure lesions are completely wet. When disease levels are high use higher rate
* Pre-harvest blue and green mold and brown rot (Phytophthora citricola)	Foliar spray, ground	4 tsp. per gallon of water	Apply 2 to 4 weeks prior to harvest. Ensure fruit is thoroughly covered by the spray application.

\*Not for use in California.

**COCONUTS**

Use AGRI-FOS 400® for effective control of bud rot and nut fall in coconuts.

Disease	Application Method	Rate	Application Program
Bud rot – Nut fall (Phytophthora palmivora)	Injection	1/3 to 1 fl. oz. per tree	Dilute AGRI-FOS 400® with water to give final injection volume of 1 fl. oz. to 2 fl. oz. of water and AGRI-FOS 400®.  Inject into the trunk or root system.

**CUCURBITS**

Use AGRI-FOS 400® for effective control of sudden wilt, gummy stem blight, downy mildew, powdery mildew, Alternaria leaf blight, and bottom soft root complex, and for suppression of Anthracnose in cucurbits such as, but not limited to, cucumber, Chinese waxgourd, citron melon, gherkin rock melon, honeydew melon, pumpkin, zucchini, watermelon and squash (summer and winter), momordica spp balsam apple, balsam pear, bitter melon, and Chinese cucumber.

Disease	Application Method	Rate	Application Program
Sudden wilt – Root and fruit rot (Phytophthora spp)	Foliar spray	2-6 tsp. per gallon of water	Entire spray coverage of plant is required. Do not exceed a total of 6 applications per season.
Gummy stem blight (Mycosphaerella melonis)		Apply 1 fl. oz. per gallon of water <b>(CA only)</b>	Apply when disease is evident. Continue applications at 21 day intervals until cure is apparent.  Do not exceed a total of 6 applications per growing season.

Downy mildew (Pseudoperonospora cubensis)			Apply within 7 to 10 days of infection. Repeat as necessary.  Do not exceed a total of 6 applications per growing season.
* Powdery mildew and other leaf diseases such as Anthracnose and Alternaria leaf blight	Foliar spray	2 to 3 fl. oz. per gallon of water	Low Disease Pressure: Apply lower rate at first onset of disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per growing season.  High Disease Pressure: Apply higher rate at first onset of disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per growing season.
* Bottom soft rot complex	Foliar spray	4-5 tsp. per gallon of water	Apply after fruit set and during bulking 3 times during the growing season.

\* Not for use in California.

**FRUITING VEGETABLES\***

Use AGRI-FOS 400® for effective control of damping-off, root rot, gummy stem blight diseases, bacterial diseases, downy mildew, powdery mildew, and Alternaria leaf blight, and for suppression of Anthracnose in eggplant, tomatoes, tomatillos, and peppers such as, but not limited to, bell, chili, cooking, pimento, and sweet.

Disease	Application Method	Rate	Application Program
Eggplant: Pythium and Phytophthora spp, and Gummy stem blight (Mycosphaerella melonis)		Apply 1 fl. oz. per gallon of water.	Entire spray coverage of plant is required. Do not exceed a total of 6 applications per growing season.  Apply when disease is evident. Continue applications at 21-day intervals until cure is apparent.  Do not exceed a total of 6 applications per growing season.
Peppers: Late blight and root rot (Phytophthora infestans and Phytophthora spp)	Foliar spray	2 tsp. to 2 fl. oz. per gallon of water	First application at transplant or when direct seeded crops are at 2-4 true leaf, then at one to two week intervals as required to control disease. In high disease situations use higher rates and shorter spray intervals.

<p>Tomatoes/Tomatillos: Late blight and root rot (Phytophthora infestans and Phytophthora spp)</p>		<p>2 tsp. to 2 fl. oz. per gallon of water</p> <p>California: 3 tsp. per gallon of water</p>	<p>First application at transplant or when direct seeded crops are at 2-4 true leaf, then at one to two week intervals as required to control disease. In high disease situations use higher rates and shorter spray intervals.</p>
<p>Root rot and damping off (Phytophthora spp and Pythium spp)</p> <p>Bacterial diseases</p>	<p>Foliar spray</p> <p>Pre-plant seedling nursery treatment</p> <p>Transplant and furrow application</p>	<p>1 to 2 fl. oz. per gallon of water</p> <p>1 tsp in 1 gallon of water to plants in seedling trays 1 to 7 days prior to out planting</p> <p>3 pints at planting or to newly planted seedlings by side dressing</p>	<p>Begin application after plants are established and conditions favor disease development.</p> <p>Low Disease Pressure: Apply lower rate at 1 to 2 week intervals. Do not apply more than 6 times per growing season.</p> <p>High Disease Pressure: Apply higher rate at 7 to 10 day intervals. Do not apply more than 6 times per growing season.</p> <p>For control of bacterial leaf spot of tomatoes, apply the high rate of AGRI-FOS 400® with registered bactericides.</p>
<p>Downy mildew</p>	<p>Foliar spray</p>	<p>1 to 2 fl. oz. per gallon of water</p>	<p>Low Disease Pressure: Apply lower rate at first onset of disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per growing season.</p> <p>High Disease Pressure: Apply higher rate at first onset of disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per growing season.</p>

55/72

Powdery mildew and other leaf diseases such as Anthracnose and Alternaria leaf blight	Foliar spray	1 to 2 fl. oz. per gallon of water	<p>Low Disease Pressure: Apply lower rate at first onset of disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per growing season.</p> <p>High Disease Pressure: Apply higher rate at first onset of disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per growing season.</p>
---	--------------	------------------------------------	---

\*Use in California limited to control of late blight and root rot in tomatoes and tomatillos.

**GRAPES\***

Use AGRI-FOS 400® for effective control of root rot, downy mildew, and powdery mildew diseases in grapes.

Disease	Application Method	Rate	Application Program
Downy mildew (Plasmopara viticola)	Foliar spray	½ fl. oz per gallon of water	It is essential that the rate of AGRI-FOS 400® be adjusted to the vine-row volume, i.e., the volume of vine foliage per acre. Spray timing is critical. Apply AGRI-FOS 400® at times of high disease risk, especially between the time that conditions are conducive to downy mildew infection and the appearance of oil spots. Ensure spray coverage is adequate and that the appropriate rate of AGRI-FOS 400® is applied to match vine growth, particularly from mid-season onwards, and especially where grapes are grown on root stock.
Phytophthora and Pythium root rots	Foliar spray	1 to 1½ fl. oz. per gallon of water	<p>Apply to vines that have stressed root system that can lead to root rots. Mitigating factors such as nematode pressure, water logging, and compaction contribute to vine declines.</p> <p>Table Grapes: Begin application in the Spring at the 4 to 6 inch shoot stage. Continue applications at 1 to 2 week intervals until</p>

56/72

			<p>flowering. Resume applications in the Fall after harvest. Do not apply more than 4 times per growing season.</p> <p>Wine and Raisin Grapes: Begin applications in the Spring at the 4 to 6 inch shoot stage. Continue applications at 1 to 2 week intervals through flowering. Do not apply more than 4 times per growing season.</p>
Downy mildew	Foliar spray	<p>1 to 1½ fl. oz. per gallon of water</p> <p>AGRI-FOS 400® is most effectively applied to control downy mildew when mixed with other registered fungicides.</p>	<p>Apply a bud break with additional applications at 7 to 10 day intervals in a rotational program with other registered fungicides. Use higher rate and volume based on disease severity and density of canopy. Do not apply more than 6 times per growing season.</p>
Powdery mildew	Foliar spray	<p>2 to 3 fl. oz. per gallon of water</p>	<p>Low Disease Pressure: Apply lower rate at first onset of disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per growing season.</p> <p>High Disease Pressure: Apply higher rate at first onset of disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per growing season.</p>

\* Use in California limited to control of downy mildew.

**HERBS AND SPICES\***

Use AGRI-FOS 400® for effective control of Downy mildew, Phytophthora spp and Pythium spp diseases of Herbs and Spices grown in fields, nurseries and greenhouses. Apply AGRI-FOS 400® to plants such as, but not limited to, Anise, Balm, Basil, Chamomile, Caraway, Catnip, Celery, Chives, Coriander, Cumin, Curry Leaf, Dill, Fennel, Marjoram, Mint, Nasturtium, Rosemary, Sage, Savory, Sweet Bay, Tarragon, Thyme, and Wintergreen.

Make applications before disease development and in conjunction with good cultural management practices. Use higher rate of application when disease pressure is severe. Do not exceed indicated application rates or apply more frequently than stated on label in order to avoid plant injury. Do not apply to plants that are heat or moisture stressed.

Disease	Application Method	Rate	Application Program
Downy Mildew	Foliar Spray	½ to 1 1/8 fl. oz. per gallon of water	Apply spray to thoroughly wet all foliage. Application intervals: 14 to 21 days. Repeat as required.

57/72

Phytophthora spp and Pythium spp	Foliar spray	2-4 tsp. per gallon of water	Apply spray to thoroughly wet all foliage. Application intervals: 14 to 21 days. Repeat as required.
	Soil drench	1/8 tsp. per gallon of water	Apply each gallons of solution to an area of 4 square feet. Follow application with irrigation. Repeat as required. Limit of one application per month.

\*Not for use in California.

**HOPS\***

Use AGRI-FOS 400® for effective control of downy mildew in hops.

Disease	Application Method	Rate	Application Program
Downy mildew	Foliar spray by ground equipment only.	2-6 tsp. per gallon of water	When conditions favor disease, apply when A. Shoots are ½ to 1 foot long. B. Post-training when vines are 6 feet high. C. 21 days post-application (B) D. During bloom

\*Not for use in California.

**LEAFY VEGETABLES\***

Use AGRI-FOS 400® for effective control of root rot, damping-off, downy mildew, powdery mildew, and other fungal leaf diseases, and for suppression of Anthracnose in leafy vegetables such as, but not limited to, amaranth, arugula, cardoon, celery, chervil, chrysanthemum, corn salad, cress, dandelion, dock, endive, fennel, lettuce, orach, parsley, purslane, radicchio, rhubarb, spinach, and Swiss chard. Excludes Brassica vegetables.

Disease	Application Method	Rate	Application Program
Downy mildew (Bremia lactucae)	Foliar spray	1 2/3 fl. oz. per gallon of water	Ensure spray coverage is adequate to wet the whole plant.  During warm, wet conditions repeat application at 7 to 10 day intervals, if needed.
Root rots and damping off (Phytophthora and Pythium spp)	Foliar spray  Pre-plant nursery application	1 to 1½ fl. oz. per gallon of water  2 tsp. in 1 gallon of water to plants in seedling trays 1 to 7 days prior to out planting	Begin application after plants are established and conditions favor disease development.  Low Disease Pressure: Apply lower rate at 1 to 2 week intervals. Do not apply more than 6 times per growing season.
Powdery mildew and leaf diseases such as Anthracnose (Collectotrichum spp), leaf blights (Septoria and Cercospora spp), and bacterial rots (Erwinia	Foliar spray, ground	1½ to 2 fl. oz. per gallon of water	High Disease Pressure: Apply higher rate at 7 to 10 day intervals. Do not apply more than

58/72

spp)			6 times per growing season.
------	--	--	-----------------------------

\*Use in California limited to control of downy mildew in lettuce.

**LEGUMES\***

Use AGRI-FOS 400® for effective control of damping-off, root rot, Fusarium, Rhizoctonia, downy mildew, powdery mildew, and other fungal leaf diseases, and for suppression of Anthracnose in legumes (succulent and dried) such as, but not limited to, green beans, soybeans, wax beans, field beans, navy beans, lima beans, fava beans, kidney beans, pinto bean, mung beans, broad beans, lentils, chickpeas, English peas, snow peas, sugar snap peas, black-eyed peas, cow peas, and pigeon peas.

Disease	Application Method	Rate	Application Program
Phytophthora and Pythium spp  Fusarium and Rhizoctonia	Foliar spray  Pre-plant nursery application  Transplant and furrow application	1 to 2 quarts in 30 to 50 gallons of water per acre  2 tsp. in 1 gallon of water to plants in seedling trays 1 to 7 days prior to out planting  3 pints at planting or to newly planted seedlings by side dressing	Apply at crop emergence or during periods of crop stress caused Summer Stress Syndrome by or wet conditions that favor disease development.  Begin application after plants are established and conditions favor disease development.  Low Disease Pressure: Apply lower rate at 1 to 2 week intervals. Do not apply more than 6 times per growing season.  High Disease Pressure: Apply higher rate at 7 to 10 day intervals. Do not apply more than 6 times per growing season.
Downey mildew	Foliar spray, ground	1 to 2 fl. oz. per gallon of water	Low Disease Pressure: Apply lower rate at the first onset of the disease. Repeat applications at 1-2 week intervals. Do not apply more than 6 times per growing season.  High Disease Pressure: Apply higher rate at the first onset of the disease. Repeat applications at 7-10 day intervals. Do not apply more than 6 times per growing season.
Powdery mildew and leaf diseases such as Anthracnose (Collectotrichum spp),	Foliar spray, ground	2 to 3 fl. oz. per gallon of water	Low Disease Pressure: Apply lower rate at the first onset of the disease. Repeat

59/72

leaf blights (Septoria and Cercospora spp), and bacterial rots (Erwinia spp)			<p>applications at 1-2 week intervals. Do not apply more than 6 times per growing season.</p> <p>High Disease Pressure: Apply higher rate at the first onset of the disease. Repeat applications at 7-10 day intervals. Do not apply more than 6 times per growing season.</p>
--	--	--	--

\*Not for use in California.

**MANGOS**

Use AGRI-FOS 400® for suppression of anthracnose in mangos.

Disease	Application Method	Rate	Application Program
Anthracnose (Colletotrichum gloeosporoides)	Foliar spray	2 tsp. per gallon of water	<p>Spray tree every 14 days during blossom period, then monthly until harvest. Spray to the point of run-off.</p>

**MICELLANEOUS CROPS\***

Use AGRI-FOS 400® for effective control of damping-off, root rot, downy mildew, powdery mildew, and other bacterial and leaf diseases, and for suppression of Anthracnose in coffee, okra, papaya, and persimmon.

Disease	Application Method	Rate	Application Program
Phytophthora and Pythium spp	Foliar spray	2 to 4 tsp. per gallon of water	<p>Begin application after plants are established and conditions favor disease development.</p> <p>Low Disease Pressure: Apply lower rate at 1 to 2 week intervals. Do not apply more than 6 times per growing season.</p> <p>High Disease Pressure: Apply higher rate at 7 to 10 day intervals. Do not apply more than 6 times per growing season.</p>
	Root dip	Mix a 0.25% v/v solution (1/3 fl. oz. with 1 gallon of water)	Apply as pre-plant dip to transplants immediately prior to planting. Dip plants momentarily and plant within 48. Mix a fresh solution daily.
Downey mildew	Foliar spray	2 to 6 tsp. per gallon of water	Low Disease Pressure: Apply lower rate at the first onset of the disease. Repeat applications at 1 to 2 week intervals. Do not

60/72

			<p>apply more than 6 times per growing season.</p> <p>High Disease Pressure: Apply higher rate at the first onset of the disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per growing season.</p>
Powdery mildew	Foliar spray	2 to 6 tsp. per gallon of water	<p>Low Disease Pressure: Apply lower rate at the first onset of the disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per growing season.</p> <p>High Disease Pressure: Apply higher rate at the first onset of the disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per growing season.</p>
Pseudomonas garacae	Foliar spray	Mix a 0.50% v/v solution (4 tsp. per gallon of water)	Apply prior to onset of disease. Apply spray to saturation or runoff.

\*Not for use in California.

**ONIONS**

Use AGRI-FOS 400® for effective control of downy mildew disease in onions, garlic, shallots, and leeks. Use as a preventative control program for best results.

Disease	Application Method	Rate	Application Program
Downy mildew (Peronospora destructor)	Foliar spray	4 tsp. per gallon of water	As a regular preventative control program or when disease first appears.

**PEANUTS\***

Use AGRI-FOS 400® for effective control of damping-off and root rot disease and for suppression of Anthracnose in peanuts.

Disease	Application Method	Rate	Application Program
Root rots, pod rots, damping off, wilt (Phytophthora and Pythium spp)	Foliar spray	4 to 5 tsp. per gallon of water	Begin application after plants are established and conditions favor disease development.
Leaf and crown diseases, Anthracnose (Collectothricum)	Transplant and furrow application	3 pints at planting or to newly planted seedlings by side dressing or shank application	<p>Low Disease Pressure: Apply lower rate at 1 to 2 week intervals. Do not apply more than 6 times per growing season.</p> <p>High Disease Pressure: Apply higher rate at 7 to 10 day intervals. Do</p>

61/72

			not apply more than 6 times per growing season.
--	--	--	---

\*Not for use in California.

**PINEAPPLES**

Use AGRI-FOS 400® for effective control of Phytophthora root and heart rot diseases in pineapples.

Disease	Application Method	Rate	Application Program
Phytophthora root and heart rot (Phytophthora cinnamomi and parasitica spp)	Foliar spray	1 2/3 to 3 1/3 fl. oz. per gallon of water	Apply to tops, 14 days prior to harvest of planting material.
	Pre-plant dip	2 tsp. per gallon of water	Established plantings when conditions favor disease. Apply at 90 day intervals.
	Foliar spray	2/3 fl. oz. per gallon of water	Ensure thorough coverage of plants.

**ROOT AND TUBER VEGETABLES**

Use AGRI-FOS 400® for effective control of foliar and root rot in ginseng, damping-off and root rot diseases in carrots, late blight disease and storage diseases such as pink rot and pythium leak, and downy mildew in potatoes, sweet potatoes, and yams.

Disease	Application Method	Rate	Application Program
* Ginseng: Foliar and root rot (Phytophthora cactorum)	Foliar spray	4 1/2 tsp. per gallon of water	In cool wet conditions that favor Phytophthora. Apply at 7 day intervals.  Do not exceed a total of 8 applications per season.
* Carrots: Phytophthora and Pythium spp	Foliar spray	2 tsp. to 2 fl. oz. per gallon of water	Apply at 14-day intervals after plant emergence, as needed. Assure good coverage.
Potatoes, Sweet Potatoes, Yams: Pink rot and Pythium leak (Phytophthora erythroseptica and Pythium spp)	In-furrow spray	6 1/2 to 16 fl. oz. per gallon of water	Apply in a band spray directly over top of potato seed just before row is closed.
Potatoes, Sweet Potatoes, Yams: Late blight, Pink rot and Pythium leak (Phytophthora infestans, Phytophthora erythroseptica and Pythium spp)	Foliar spray	2/3 to 3 tsp. per gallon of water	Apply at 5 to 14 day intervals subject to disease incidence.
* Downy mildew	Foliar spray	1 to 2 fl. oz. per gallon of water	Low Disease Pressure: Apply lower rate at the first onset of the disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per growing season.  High Disease Pressure: Apply higher rate at the

62/72

			first onset of the disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per growing season.
--	--	--	--

\* Not for use in California.

**STONE FRUIT\***

Use AGRI-FOS 400® for effective control of root, collar rot, almond pruning wound, and various canker disease in stone fruit such as, but not limited to, apricots, sweet and tart cherries, nectarines, peaches, plums, and fresh prunes.

Disease	Application Method	Rate	Application Program
Root and collar rot (Phytophthora spp)	Foliar spray  Basal bark spray apply spring and fall for best results	1/3 fl. oz. per gallon of water  16 fl. oz. + 16 fl. oz. of water + 1 fl. oz. Supa Stik PN Bark Penetrating Surfactant	Three treatments are required 1. Spring 2. Mid summer 3. Fall, post harvest  Spray a combination of AGRI-FOS 400® and Supa Stik PN around the complete circumference of the tree trunk until saturation/runoff. Spray from ground level up to 5 feet above the soil line, including the base of the first scaffolding limbs, if present. (Treatment generally lasts 8-12 weeks depending on pathogen levels. Higher disease pressure will shorten the length of control.)
Almond pruning – wound cankers (Phytophthora syringae)	Paint or spray	1/3 to 3/4 fl. oz. per gallon of water	Apply to pruning wound and surrounding area, ensure area is thoroughly wet. In high disease situations use higher rate.
Suppression of armillaria root rot (Armillaria luteobublina)	Basal bark spray	1 1/2 to 2 quarts + 2 quarts water + 1% Supa Stik PN Bark Penetrating Surfactant	Spray a combination of AGRI-FOS 400® and Supa Stik PN around the complete circumference of the tree trunk until saturation/runoff. Spray from ground level up to 5 feet above the soil line, including the base of the first scaffolding limbs, if present. For large trees larger than 18 inches DBH (Diameter at Breast Height, 4.5 feet above ground) that have

63/72

			been previously attacked by Armillaria root rot, make a fall application prior to leaf senesce and a spring application for best results. For trees less than 18 inches DBH, make an early spring application.
--	--	--	--

\*Not for use in California.

**STRAWBERRIES\***

Use AGRI-FOS 400® for effective control of leather rot, red stelè, Phytophthora, foliar and bacterial disease in strawberries and suppression of powdery mildew.

Disease	Application Method	Rate	Application Program
Phytophthora and Pythium spp (red stele, leather rot, and root rot)  Suppression of rhizopus	Pre-planting dip	1/3 fl. oz. per gallon of water	Dip planting material in this solution for 30 minutes, then plant within one day. Use program for annual and perennial varieties
	Foliar spray	2 to 6 tsp. per gallon of water	Annual crops, first treatment 14 to 21 days post planting, repeat at 1-2 month intervals when disease is evident.  Perennial crops, first treatment during spring growth flush, repeat at 1-2 month intervals when disease is evident. For susceptible varieties us higher rates and shorter spray intervals.
	Transplant and furrow application	3 pints at planting or to newly planted seedlings by side dressing or shank application	For leather rot, apply at 10% bloom and early fruit set, then 1 to 2 week intervals as required for disease control, In high disease situations, use higher rate and shorter spray intervals.  Grey mold and Anthracnose suppression using AGRI-FOS 400® requires use of high application rates and is most effective when tank mixed with other registered fungicides.
Phytophthora spp	Dip	Mix a 0.25% v/v solution (1/3 fl. oz. with 1 gallon of water)	Dip runners in the solution for 1 to 2 minutes. Plant within 48 hours. Mix a fresh solution daily.

64/72

Suppression of powdery mildew	Foliar spray	1 to 1½ fl. oz. per gallon of water	<p>Low Disease Pressure: Apply lower rate at the first onset of the disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per growing season.</p> <p>High Disease Pressure: Apply higher rate at the first onset of the disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per growing season.</p>
-------------------------------	--------------	-------------------------------------	---

\* Use in California limited to control of red stele, leather rot, and root rot.

**TREE NUTS\***

Use AGRI-FOS 400® for effective control of root and collar rot, and almond pruning-wound canker disease in tree nuts such as, but not limited to, almonds, beech nuts, Brazil nuts, butternuts, cashews, chestnuts, chinquapin, hazelnuts, hickory nuts, macadamia nuts, pecans, pistachios, and walnuts.

Disease	Application Method	Rate	Application Program
Other than macadamia nuts: Root and collar rot (Phytophthora spp)	Foliar spray	2 tsp. per gallon of water	Three treatments are required 1. Spring 2. Mid summer 3. Fall, post harvest
Other than macadamia nuts: Almond pruning – wound canker (Phytophthora syringae)	Paint or spray	4½ tsp. per gallon of water	Apply to pruning wound and surrounding area, ensure area is thoroughly wet.
Macadamia nuts: Raceme blight (Phytophthora spp)	Foliar spray	3 tsp. per gallon of water	Apply when disease is first seen and reapply at 3 week intervals. Spray to the point of run-off
Root rot, crown rot, trunk cankers, foliar blights (Phytophthora and Pythium spp)	Foliar spray	4 to 5 tsp. per gallon of water	Begin application after plants are established and conditions favor disease development. Low Disease Pressure: Apply lower rate at 3 month intervals. Do not apply more than 4 times per growing season. High Disease Pressure: Apply higher rate at monthly intervals. Do not apply more than 4 times per growing season.
	Root dip	Mix a 0.5% v/v solution (4 tsp. in 1 gallon of water)	Dip roots in the solution for 30 seconds. Plant within 48 hours. Mix a fresh solution daily.

65/72

Pecan scab	Foliar spray	1½ fl. oz. per gallon of water	Apply preventatively at 21 to 30 day intervals.
Pruning wound, crown and trunk cankers (Phytophthora spp)	Trunk spray	1½ quarts + 2 quarts of water + 1 Supa Stik PN. Bark Penetrating Surfactant	Apply higher rate when lesions are present. Clean wound sites and apply on and around the lesions using enough spray volume to thoroughly wet the lesions. Apply to the trunk from the soil line to 5 feet up the trunk. Apply one time in the Spring, Summer, and Fall.
Downy mildew (Phytophthora and Peronospora spp)  Powdery mildew	Foliar spray	2 to 4 tsp. per gallon of water	Low Disease Pressure: Apply lower rate at first onset of disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per growing season.  High Disease Pressure: Apply higher rate at first onset of disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per growing season.
Tree cankers	Basal bark spray	1½ to 2 quarts + 2 quarts of water + 1% Supa Stik PN Bark Penetrating Surfactant	Spray a combination of AGRI-FOS 400® and Supa Stik PN around the complete circumference of the tree trunk until saturation/runoff. Spray from ground level up to 5 feet above the soil line, including the base of the first scaffolding limbs, if present. For large trees larger than 18 inches DBH (Diameter at Breast Height, 4.5 feet above ground) that have been previously attacked by Armillaria root rot, make a fall application prior to leaf senesce and a spring application for best results. For trees less than 18 inches DBH, make an early spring application.

\*Not for use in California.

66/72

## LANDSCAPE APPLICATIONS\*

Use AGRI-FOS 400® for effective control of Phytophthora and Pythium spp and other diseases associated with Sudden Oak Death, Beech Decline, and general tree decline syndromes, and for suppression of Anthracnose and Verticillium wilt. Apply AGRI-FOS 400® to trees such as, but not limited to, Ash, Azalea, Beech, Black Locust, Buckeye, Catalpa, Cedar, Cherry – Stonefruits, Chestnut, Coffee Tree, Crab Apple, Cork Tree, Dogwood – All, Elder, Elm, Fir, Hawthorne, Honey Locust, Juniper, Lilac, Linden, Magnolia, Maples – All, Pine, Oaks – All, Olives, Ornamental Pear, Plum, Pyracantha, RedBud, Smoke Tree, Sumac, Sweet Birch, Sweet Gum, Sycamore, Viburnum, White Pine, White Cedar, and Willow.

Make applications before disease development and in conjunction with good cultural management practices. Use higher rate of application when disease pressure is severe. Do not exceed indicated application rates or apply more frequently than stated on label or tree injury may occur. Do not apply to trees that are heat or moisture stressed. Do not apply to trees that are in a state of dormancy. Do not exceed indicated spray intervals or label rates in order to avoid tree injury.

Disease	Application Method	Rate	Application Program
Phytophthora and Pythium spp, and Phytophthora ramorum	Injection	11 fl. oz. per 21 fl. oz. of water Or ½ tsp. per tsp. of water	Drill holes 3/16 inch (5 mm) in diameter <u>into live sapwood</u> (depth dependent upon age of tree) with downward angle into trunk uniformly around the tree circumference, using a slow drill. Do not inject into areas of obvious decay, canker or mechanical injury that appear on the tree trunk. Calculate the amount of product required by measuring the trees by one of the following 3 methods, and use the highest calculated number of injections. 1) 1 injection per square yard of canopy; 2) 1 injection per yard of diameter of canopy measured at the drip-line; 3) 1 injection per 6 inches of trunk circumference measured 4 feet above soil level. Make injections with applicators that maintain positive pressure differential such as ChemJet®, Sidewinder®, Ag-murph Gun®, Marley® Injector, Smart Shot injector, or hydraulic applicator type equipment that forces solution into the sapwood of the tree.
Phytophthora and	Basal bark spray	16 fl. oz. + 16 fl. oz. of	Apply uniformly to 6-9

67/72

<p>Pythium spp, and Phytophthora ramorum</p>	<p>(all other tree species) apply spring and fall for best results or anytime active growth occurs Best for thin bark trees such as maples, lindens, sycamores, and dogwoods</p>	<p>water + 1 fl. oz. Supa Stik PN Bark Penetrating Surfactant</p>	<p>feet of trunk circumference. Spray from top down to ground level from either first branch or from as high as possible without exposing applicator to drift. Spray to saturation/runoff. Various types of application equipment can be used such as hydraulic sprayers, handheld pump-type sprayers, and backpack sprayers.</p>
<p>Fusarium subglutinans (Pine Pitch Canker)</p>	<p>Basal bark spray (Pines). Apply anytime active growth is observed.</p>	<p>16 fl. oz. AGRI-FOS 400® + 32 fl. oz. water + 0.5 fl. oz. of Supa Stik PN Bark Penetrating Surfactant</p>	<p>Spray a combination of AGRI-FOS 400® and Supa Stik PN around the complete circumference of the tree trunk until saturation/runoff. Spray from ground level up to 5 feet above the soil line, including the base of the first scaffolding limbs, if present. (Treatment generally lasts 8-12 weeks depending on pathogen levels. Higher disease pressure will shorten the length of control.)</p>
<p>Fusarium subglutinans (Pine Pitch Canker) and Gnomonia platani (Sycamore Anthracnose)</p>	<p>Injection</p>	<p>4 tsp per tree of a 16 fl. oz. AGRI-FOS 400 + 32 fl. oz. of water + 0.5 fl. oz. of Supa Stik PN Bark Penetrating Surfactant solution.</p>	<p>Drill holes 3/16 inch (5 mm) in diameter and 1 inch (25 mm) to 2 inches (50 mm) deep with slight downward angle in trunk. Place syringes in the main trunk of the tree and space evenly around the circumference of the trunk. Suitable for use with Chemjet tree injectors, Ag-murf gun, Smart Shot injector, or hydraulic tree injection. Trees should be at least 10" diameter at breast height.</p>
<p>** Suppression of Anthracnose</p>	<p>Foliar spray, ground</p>	<p>3-4 tsp. per gallon of water</p>	<p>Apply at pre bloom (bud swell or green tip stage) with a supplemental application 14 days later with AGRI-FOS 400® or other fungicide compound with efficacy on Anthracnose.</p>
<p>** Suppression of Anthracnose</p>	<p>Basal bark spray apply early spring at bud swell until green tip stage of growth</p>	<p>16 fl. oz. + 16 fl. oz. of water + 1 fl. oz. Supa Stik PN Bark Penetrating Surfactant</p>	<p>Spray a combination of AGRI-FOS 400® and Supa Stik PN around the complete circumference of the tree trunk until</p>

68/172

			saturation/runoff. Spray from ground level up to 5 feet above the soil line, including the base of the first scaffolding limbs, if present. For large trees larger than 18 inches DBH (Diameter at Breast Height, 4.5 feet above ground) that have been previously attacked by Anthracnose, make a fall application prior to leaf senesce and a spring application for best results. For trees less than 18 inches DBH, make an early spring application.
** Suppression of Verticillium wilt	Foliar spray	3-4 tsp. per gallon of water	For trees previously identified with infections, apply first application pre-bloom. Repeat applications between 21 to 30 days.
** Suppression of Verticillium wilt	Basal bark spray	1½ to 2 quarts + 2 quarts of water + 2 fl. oz. Supa Stik PN Bark Penetrating Surfactant	Spray a combination of AGRI-FOS 400® and Supa Stik PN around the complete circumference of the tree trunk until saturation/runoff. Spray from ground level up to 5 feet above the soil line, including the base of the first scaffolding limbs, if present. For large trees larger than 24 inches DBH (Diameter at Breast Height, 4.5 feet above ground) that have been previously attacked by Verticillium wilt, make a fall application prior to leaf senesce and a spring application for best results. For trees less than 24 inches DBH, make an early spring application.

\*Use in California limited to Oaks (Coastal, Live, Shreve, Black, Canyon), Tan Oaks, and other tree species that are host to P. ramorum. Do not apply to any other tree species without consulting the list of P. ramorum host species listed at the following website: <http://nature.berkeley.edu/comtf/index.html>. Make applications by injection and basal bark spray of pines, basal bark spray of apples and crabapples, and injection of Sycamores for control of Sycamore anthracnose. For questions contact your local extension office.

\*\* Not for use in California.

69/172

## ORNAMENTAL APPLICATIONS

Use AGRI-FOS 400® for effective control of Bacterial blight, Downy mildew, powdery mildew, Phytophthora spp [Root Rot], and Pythium spp [Damping-off] diseases, and Sudden Oak Death, and for suppression of Anthracnose of Ornamentals in landscapes and interiorscapes. Apply AGRI-FOS 400® to plants such as, but not limited to, Aglaonema, Aphelandra, Arborvitae, Azaleas, Bougainvillea, Boxwood, Cattelya skinneri, Ceanothus, Cotoneaster, Cissus, Diffenbachia, English ivy, Eucalyptus, Ficus, Hibiscus, Japanese andromeda, Japanese Holly, Leather leaf Fern, Peperomia, Photinia, Pittosporum, Philodendron, Pieris, Pothos, Rhododendron, Roses (container, landscape, mini varieties), Schefflera, Sedum, Sempervivum, Syngonium, Spathiphyllum, Taxus media, and Zygocactus.

Make applications before disease development and in conjunction with good cultural management practices. Use higher rate of application when disease pressure is severe. Do not exceed indicated application rates or apply more frequently than stated on label in order to avoid plant injury. Do not apply to plants that are heat or moisture stressed. Do not apply to plants that are in a state of dormancy. When applying to indoor plants do not overspray and use care to apply only to target plants. If meeting these conditions is not possible, remove plants to an outdoor location for treatment and drying before bringing back indoors.

Disease	Application Method	Rate	Application Program
Bacterial blight (Xanthomonas campestris) pothovars: dieffenbachiae, fici hederiae and syngonli	Foliar spray	2-4 tsp. per gallon of water	Apply spray to thoroughly wet all foliage. Application intervals: 7 to 14 days Repeat as required.
* Black spot (Diplocarpon spp)	Foliar spray	2-4 tsp. per gallon of water	Apply uniformly to foliage preventatively or at first sign of disease.
Downy mildew	Foliar spray	2-4 tsp. per gallon of water	Apply uniformly to foliage at disease onset and repeat applications every 14 days.
Phytophthora spp [Root Rot], Phytophthora ramorum, and Pythium spp [Damping-off]	Foliar spray	2-4 tsp. per gallon of water	Apply uniformly to foliage to point of runoff.
	Soil drench	Mix a 0.5% to 1.0% v/v solution (½ to 6½ fl. oz. in 1 to 5 gallons of water)	Apply uniformly to soil at base of plant and surrounding soil.
	Soil incorporation	1-2 pints per cubic yard of soil	Just prior to potting, mix 1 to 2 pts of AGRI-FOS 400® into each cubic yard of growing media. If disease pressure is high, make application by foliar spray or soil drench.
	Bare rooted dipping of transplants	2 tsp. per gallon of water	Immediately before transplanting, dip transplants for two minutes, keep roots submerged, ensure root mass is thoroughly wet.
* Powdery mildew	Foliar spray	2 to 4 tsp. per gallon of water	Apply uniformly to foliage at disease onset and repeat applications every 14 days.

70/  
72

* Suppression of Anthracnose	Foliar spray	1½ fl. oz. per gallon of water	Apply prior to onset of disease. Apply spray to saturation or runoff.
------------------------------	--------------	--------------------------------	---

\* Not for use in California.

### RESIDENTIAL NURSERY, GREENHOUSE, AND BEDDING PLANTS

Use AGRI-FOS 400® for effective control of Downy mildew, Phytophthora spp [Root Rot] and Pythium spp [Damping-off] diseases of BEDDING PLANTS and in interiorscapes. Apply AGRI-FOS 400® to plants such as, but not limited to, Ageratum, Aglaonema, Algerian Ivy, Anthurium, Aphelandra, Arborvitae, Artemesia, Aster, Azaleas, Begonia, Baby's Breath, Bougainvillea, Boxwood, Caladium, Carnation, Cattelya Skinneri, Ceanothos, Cissus, Chrysanthemum, Coleus, Columbine, Cotoneaster, Daisy, Delphinium, Dieffenbachia, Dogwood, Easter Lily, English Ivy, Ficus, Foxglove, Gaillardia, Geranium, Gloxinia, Hibiscus, Impatiens, Japanese Holly, Juniper, Lavender, Leather Leaf Fern, Marigold, Moterey Pink, Pieris, Peperomia, Petunia, Philodendron, Pansy, Phlox, Pinks, Pittosporum, Poinsettia, Pothos, Primrose, Prostrate Rosemary, Rosemary, Salvia, Schefflera, Sedium, Sempervivum, Snapdragon, Spathiphyllum, Taxus Media, Verbena, Vinca, White Cedar, White Pine, Zinnia, and Zygocactus. AGRI-FOS 400® may also be used for Vegetable Transplants grown in Greenhouse, Lathouse, or Shade house sites.

Make applications to outdoor or indoor plants before disease development and in conjunction with good cultural management practices. Use higher rate of application when disease pressure is severe. Do not exceed indicated application rates or apply more frequently than stated on label in order to avoid plant injury. Do not apply to plants that are heat or moisture stressed. When applying to indoor plants do not overspray and use care to apply only to target plants. If meeting these conditions is not possible, remove plants to an outdoor location for treatment and drying before bringing back indoors.

Disease	Application Method	Rate	Application Program
Downy Mildew	Foliar Spray	½ to 1 1/8 fl. oz. per gallon of water	Apply spray to thoroughly wet all foliage. Application intervals: 14 to 21 days. Repeat as required.
Phytophthora spp [Root Rot] and Pythium spp [Damping-off]	Foliar spray	2 to 4 tsp. per gallon of water	Apply spray to thoroughly wet all foliage. Application intervals: 14 to 21 days. Repeat as required.
	Soil drench	1/8 tsp. per gallon of water	Apply each gallon of solution to an area of 4 square feet. Follow application with irrigation. Repeat as required. Limit of one application per month.
Phytophthora spp* [Root Rot]	Foliar spray	¼ tsp. per 100 sq. ft.	Apply spray in 1 pint of water per 100 sq.ft.
Phytophthora spp* [Root Rot]	Hand gun	2 quarts per 100 gallons of water	Apply spray to thoroughly wet all foliage

\* Lavender applications.

## CONIFERS\*

71/  
72

Apply AGRI-FOS 400® in conjunction with good cultural management practices for effective control of root rot (Phytophthora spp) in CONIFERS including, but not limited to, Pines, Spruce and Douglas Fir. Use higher rate of application when disease pressure is severe. Do not exceed indicated application rates or apply more frequently than stated on label in order to avoid plant injury. Do not apply to CONIFERS that are moisture or heat stressed.

Disease	Application Methods	Rate	Application Program
Phytophthora spp	Foliar spray, ground	2 to 4 tsp. per gallon of water	Apply spray to thoroughly wet all foliage. Application intervals: 14 to 21 days. Repeat as required.
	Soil drench	2 to 4 tsp. per gallon of water	Apply one gallon of solution per sq yd. Follow application with irrigation. Application intervals: 14 to 21 day. Repeat as required.
	Bare root dipping of transplants	2 tsp. per gallon of water	Immediately before transplanting, dip transplants for two minutes; keep roots submerged and ensure root mass is thoroughly wet.
Fusarium subglutinans (Pine Pitch Canker)	Basal bark spray apply any time active growth occurs	16 fl. oz. AGRI-FOS 400® + 32 fl. oz. of water + 0.5 fl. oz. of Supa Stik PN Bark Penetrating Surfactant	Apply uniformly to 6-9 feet of trunk circumference. Spray from top down to ground level from either first branch or from as high as possible without exposing applicator to drift. Spray to saturation/runoff. Various types of application equipment can be used such as hydraulic sprayers, handheld pump-type sprayers, and backpack sprayers.
Fusarium subglutinans (Pine Pitch Canker)	Injection	4 tsp per tree of a 16 fl. oz. AGRI-FOS 400® + 32 fl. oz. of water + 0.5 fl. oz. of Supa Stik PN Bark Penetrating Surfactant solution.	Drill holes 3/16 inch (5 mm) in diameter and 1 inch (25 mm) to 2 inches (50 mm) deep with slight downward angle in trunk. Place syringes in the main trunk of the tree and space evenly around the circumference of the trunk. Suitable for use with Chemjet tree injectors, Ag-murf gun, or hydraulic tree injection. Trees should be at least 10" diameter at breast height.

\*Use in California limited to injection and/or basal bark spray of pines.

72/72

### TURF AND HOME LAWNS\*

Use AGRI-FOS 400® for the effective control of Rhizoctonia, Pythium and damping-off diseases, and for suppression of Anthracnose of turf and lawn grasses. When conditions favor disease, begin preventive applications and repeat at indicated intervals. Use higher rate of application when disease pressure is severe.

Disease	Application Method	Rate	Application Program
Pythium	Foliar spray	5 to 10 fl. oz. per 1000 sq. ft.	Apply indicated quantity of product in 1 to 5 gal. of water per 1000 sq. ft. Ensure foliage is thoroughly wet. Application intervals: 14-21 days. Repeat as required. Do not irrigate or mow treated areas until spray has completely dried.
Suppression of Anthracnose	Foliar spray	5 to 10 fl. oz. in 2 gallons of water per 1000 sq. ft.	Apply prior to onset of disease. Apply every 7 to 14 days.
Rhizoctonia	Foliar spray	5 to 10 fl. oz. in 2 gallons of water per 1000 sq. ft.	Apply prior to onset of disease. Apply every 7 to 14 days.

Do not graze animals on treated areas of lawn. Do not feed treated lawn clippings to poultry or livestock.

\* Use in California limited to control of pythium.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

##### PESTICIDE STORAGE:

Keep this product in containers stored upright and secured with the original closure. Do not store this product near any heat source. Do not store near strong oxidants. If transfer to another container becomes necessary, ensure that the container is clearly labeled, the container is a type suitable for the product, and is clean and free of other materials.

##### PESTICIDE DISPOSAL and CONTAINER HANDLING:

Nonrefillable container. Do not reuse or refill this container. If empty: Offer for recycling if available or discard in a sanitary landfill. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

#### WARRANTY AND DISCLAIMER

To the extent consistent with applicable law, all conditions and warranties and statutory or other rights of action which buyer or any other user may have against Liquid Fertiliser Pty. Ltd. are hereby excluded. To the extent consistent with applicable law, Liquid Fertiliser Pty. Ltd. hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information provided by Liquid Fertiliser Pty. Ltd. or on its behalf unless it is established that such information or advice was provided negligently and that the product has been used strictly as directed. To the extent consistent with applicable law, Liquid Fertiliser Pty. Ltd.'s liability shall in all circumstances be limited to replacement of product or a refund of the purchase price thereof.