Herbicide 3D RTU

- · Kills listed weeds
- · Kills hard-to-control weeds
- · All-in-one lawn weed killer
- · Gets the root
- · Kills over 200 types of weeds
- Systemic
- Kills weeds not lawns
- Won't harm lawns
- Kills tough weeds like ground ivy, chickweed, clover and nutgrass
- · Kills both broadleaf and grassy weeds
- · Kills dandelions, crabgrass, clover, plus other listed weeds
- · Kills Broadleaf and Grassy weeds in one step
- · Won't Harm Your Lawn
- Contains SmartTrack™, a non-staining marker that indicates where you've sprayed
- · Kills lawn weeds plus crabgrass in one step
- Rainproof in hours

ACTIVE INGREDIENTS:

Monosodium acid methanearsonate	0.36	%
2,4-D, dimethylamine salt	0.12	%
Mecoprop-p, dimethylamine salt	0.06	%
Dicamba, dimethylamine salt	0.02	%
OTHER INGREDIENTS: 9	99.44	%

THIS PRODUCT CONTAINS:

Total Arsenic (as elemental) all in water soluble form 0.17%.

0.10 % 2,4-D acid equivalent

0.05 % mecoprop-p acid equivalent

0.02 % dicamba acid equivalent

Isomer Specific by AOAC Method.

Net Contents: 16 fl. oz. (473 mL) or 24 fl. oz. (709 mL) or 1 Gal. (3.78 L)

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel Booklet for Additional Ingredients and Precautionary Statements

ACCEPTED

OCT 1 7 2001

Under the Federal Insectacide, Pungloide, and Rodenticide Act, as amended, for the perticide registered under, RPA Reg. No. 72/55-/

(BACK PANEL)

Herbicide 3D RTU kills both broadleaf and grassy weeds in one step. This convenient formula kills Dandelion, Chickweeds, Clovers, Crabgrass and Nutgrass/Nutsedge without harming your lawn. Controlling weeds in your lawn has never been easier.

Herbicide 3D RTU kills hard to control weeds in one step. This convenient formula kills grassy weeds such as Nutgrass/Nutsedge and broadleaf weeds such as Ground lvy, Chickweeds and Clovers without harming your lawn. Controlling weeds in your lawn has never been easier.

READ THE LABEL FIRST!

QUICK FACTS Kills Broadleaf AND Grassy Weeds Nor-Staining Marker Indicates Where You've Sprayed Kills weeds not lawns Kills the Weed, Root and All	
KILLS	Broadleaf Weeds: Dandelion, Chickweeds and Clovers
	Grassy Weeds: Crabgrass, Nutgrass/Nutsedge (and other listed broadleaf and grassyweeds)
WHERE TO USE	On Lawns (NOT for use on St. Augustine [including Floratam], Bahia , Bentgrass, Carpetgrass, Centipedegrass, or Dichondra).
WHEN TO USE	When weeds are actively growing
™	For questions or comments, call toll-free 1-877 BayerAL or visit us on-line at www.BayerAdvanced.com

FOR OUTDOOR RESIDENTIAL USE ONLY

ACTIVE INGREDIENTS:

Monosodium acid methanearsonate ().36	%
2,4-D, dimethylamine salt ().12	%
Mecoprop-p, dimethylamine sait		
Dicamba, dimethylamine salt		
OTHER INGREDIENTS*:	.44	%
100	0.00	%

*Other ingredients include: Water, suspending aids (to keep the active ingredients dispersed in water), surfactants (to keep the active ingredients dispersed in water), pigment (non-staining marker), antifoam, and preservative

THIS PRODUCT CONTAINS:

Total Arsenic (as elemental) all in water soluble form 0.17%. 0.10 % 2,4-D acid equivalent 0.05 % mecoprop-p acid equivalent 0.02 % dicamba acid equivalent lsomer Specific by AOAC Method.

EPA Reg. No. 72155-1

EPA Est. No. XXXX-AA-XX

DIRECTIONS FOR USE It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. For best results, read and follow all label directions. Do not apply near lakes, streams, rivers or **BEFORE** YOU USE Read and Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees follow directions and other desirable plants. when using. Apply when wind is calm. Spray when most of the day is expected to remain cooler than; 85° F. - Buffalograss, Kentucky Bluegrass, Perennial Ryegrass, and Fescue 90° F. - Bermuda, and Zoysia · Adjust spray nozzle to give a coarse spray. **HOW TO** Hold sprayer about 12 inches from weed. **APPLY** Aim at center of weed. Shake well Non-Staining marker indicates where before use. product was applied. To this empty container (24 fl. oz.), add 5 tsp. REFILL of Lawn Herbicide 3D Concentrate or 7 tsp. of Lawn Herbicide 3D-40 Concentrate, then fill with water very slowly to avoid foaming. Refill does not contain non-staining marker. Apply when weeds are young and actively **FOR BEST** RESULTS If the lawn is dry, water 24 hours before using this product. Rain or watering 6 to 8 hours after application will not wash away effectiveness. Apply 1 to 3 days after mowing Certain hard-to-kill weeds may need to be retreated in 14 days. Bare spots may be seeded 3 to 4 weeks after application. Do not apply to new lawns until the grass has reached a height of at least 2 inches. HERBICIDE 3D RTU is a selective herbicide -HOW it selectively kills weeds in the lawn but won't HERBICIDE harm the lawn itself. It is a blend of proven 3D RTU weed killers designed to get broadleaf weeds WORKS such as dandelion, clover and chickweed and grassy weeds such as crabgrass and

nutgrass/nutsedge.

KILLS	Annual valleur	Daisy English
ING	Annual yellow sweetclover	Daisy, English Daisy, fleabane
	Aster	Daisy, oxeye
	Austrian fieldcress	Dallisgrass
	Barnyardgrass	Dandelion Dishandra
	Bedstraw Beggarticks	Dichondra Dogbane
	Betony, Florida	Dogfennel
	Bindweed, field	Dollarweed
l	Bird vetch	Elderberry
	Bitter wintercress Bittercress, hairy	False dandelion False flax
	Bitterweed	Faise sunflower
	Black-eyed Susan	Fiddleneck
	Black medic	Florida pusley
	Black mustard Blackseed plantain	Foxtail Frenchweed
	Blessed thistle	Galinsoga
	Bloodflower	Goathead
	milkweed	Goldenrod
	Blue lettuce Blue vervain	Goosegrass Ground ivy
	Bracted plantain	Gumweed
	Brassbuttons	Hairy fleabane
	Bristly oxtongue	Hawkweed Healall
	Broadleaf dock Broadleaf plantain	Heartleaf drymary
	Broomweed	Heathaster
	Buckhorn	Hedge bindweed
	Buckhorn plantain	Hedge mustard Hemp
	Bulbous buttercup Bull thistle	Henbit
	Bullnettle	Hoary cress
	Burclover	Hoary plantain
	Burdock Burning nettle	Hoary vervain Horsenettle
	Burweed	Jimsonweed
	Buttercup	Johnsongrass
	Buttonweed Canada thistle	Knawel Knotweed
	Canada thistle Carolina geranium	Kochia
	Carpetweed	Lambsquarters
	Catchweed	Lespedeza
	bedstraw Catnip	Mallow Matchweed
	Catsear	Mexicanweed
	Chickweed,	Milk vetch
	Chickwood	Morningglory
	Chickweed, mouseear	Mouseear hawkweed
	Chicory	Mugwort
	Cinquefoil	Musk thistle
	Clover, crimson	Mustard Narrowleaf plantain
	Clover, hop Clover, red	Narrowleaf vetch
	Clover, strawberry	Nettle
	Clover, sweet	Nutgrass/Nutsedge
	Clover, white Cockle	Orange hawkweed Oriental cocklebur
	Cocklebur	Oxalis
	Common mullein	Parsley-piert
	Crabgrass	Parsnip
	Creeping charlie	Pearlwort Pennycress
	Creeping jenny Cudweed	Pennywort
	Curly dock	Peppergrass
		Pepperweed
Continued in nex	t page	

KILLS LIST con	tinued	
KILLS	Dimund	Charles - Latin
	Pigweed	Strawberry, India
	Pineywoods	mock
	bedstraw	Tall nettle
	Plains coreopsis	Tall vervain
	(tickseed)	Tansy mustard
	Plantain	Tansy ragwort
	Poison ivy	Tanweed
	Poison oak	Thistle
	Pokeweed	Trailing crownvetch
	Poorjoe	Tumble mustard
	Prairie sunflower	Tumble pigweed
	Prickly lettuce	Velvetleaf
	Prickly sida	Venice mallow
	Prostrate knotweed	Virginia buttonweed
	Prostrate pigweed	Virginia creeper
	Prostrate spurge	Virginia
	Prostrate vervain	pepperweed
	Puncturevine	Water pennywort
	Purstane, common	Wavyleaf bullthistle
	Ragweed	Western clematis
	Red sorrel	Western salsify
	Redroot pigweed	White mustard
	Redstern filaree	Wild aster
	Rough cinquefoil	Wild buckwheat
	Rough fleabane	Wild carrot
	Russian pigweed	Wild four-o'clock
	Russian thistle	Wild gartic
	Sandbur	Wild geranium
	Scarlet pimpernel	Wild lettuce
	Scotch thistle	Wild marigold
	Sheep sorrel	Wild mustard
	Shepherdspurse	Wild onion
	Slender plantain	Wild parsnip
į	Smailflower	Wild radish
	galinsoga	Wild rape
	Smooth dock	Wild strawberry
	Smooth pigweed	Wild sweet potato
	Sorrei	Wild vetch
	Sowthistle	Woodsorrei
	Spanishneedles	Wooly croton
	Speedwell	Woolly morningglory
	Spiny amaranth	Woolly plantain
	Spiny cocklebur	Wormseed
ļ	Spiny sowthistle	Yarrow
	Spotted catsear	Yellow rocket
ĺ	Spotted spurge	Yellowflower
	Spurweed	pepperweed
	Stinging nettle	and other broadleaf
		weeds



Children and pets may re-enter the treated area after it has dried.

IMPORTA	NT: Dire	ctions for Storage and Disposal	
Storage Conditions:		Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing.	
Disposal:		Do not reuse empty container (except to refill with other 3D Concentrate products).	
	-16	Wrap in paper or plastic bag and place in trash.	

Important: It is best to use all of the product in accordance with label directions. If it is necessary to dispose of unused product, please follow any applicable state and local guidelines.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

- Causes moderate eye irritation.
- Avoid contact with eyes, skin, or clothing.
- When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

After using this product:

- · Rinse gloves before removing and wash hands.
- If product gets on clothing, remove clothing as soon as possible and launder clothing separately before reuse.

FIRST AID		
IN CASE OF ACCIDENTAL	IMMEDIATELY	
Swallowing	Call a poison control center or doctor immediately for treatment advice.	
<u> </u>	Have person sip a glass of water if able to swallow.	
	Do not induce vomiting unless told to do so by a poison control center or doctor.	
	Do not give anything by mouth to an unconscious person.	
Contact with eyes	Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.	

NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 877-229-3763 for medical emergency information. Active Ingredients: 0.36 % Monosodium acid methanearsonate [CAS # 2163-80-6], 0.12% 2,4-D, dimethylamine salt [CAS # 2008-39-1], 0.06% Mecoprop-p, dimethylamine salt single isomer [CAS # 66423-09-4], and 0.02% Dicamba, dimethylamine salt [CAS # 2300-66-5].

ENVIRONMENTAL HAZARDS

- · This product is toxic to aquatic invertebrates.
- · Drift or runoff may harm fish and plants.
- Do not apply directly to water.

NOTICE: Research and testing have determined that the "Directions For Use" are appropriate for the proper use of HERBICIDE 3D RTU under expected conditions. The Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.

D - 9617 091301

Bayer Advanced LLC 1500 Urban Center Drive Birmingham, AL 35242