

U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7504P) Ariel Rios Building 1200 Pennsylvania Ave., NW Washington, D.C. 20460

EPA Registration Number: Date of Issuance:

72155-2

APR 29 2011

Term of Issuance:

Unconditional

Name of Pesticide Product:

Lawn Herbicide JD Ready-To-Use

NOTICE OF PESTICIDE:

X Registration

X Reregistration

(under FIFRA, as amended)

Bayer Advanced

P.O. Box 12014, 2 T.W. Alexander Drive

Research Triangle Park, North Carolina 27709

Name and Address of Registrant (include ZIP Code):

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

EPA received a label amendment request submitted on April 6, 2011. EPA grants this request under the authority of section 3(c)(5) of the Federal Insecticide, Fungicide and Rodenticide Act, as amended. With this accepted labeling, all requirements set forth in the Reregistation Eligibility Decision (RED) for **2,4-D, MCPP-p, and Dicamba** have been satisfied. Therefore, EPA reregisters the product listed above. This action is taken under the authority of section 4(g)(2)(c) of the Federal Insecticide, Fungicide, and Rodenticide Act, as amended. Reregistration under this section does not eliminate the need for continual reassessment of pesticides. EPA may require submission of data at any time to maintain the registration of your product.

Submit one (1) copy of final printed labeling. Amended labeling will supersede all previously accepted labels. A copy of your label stamped "Accepted" is enclosed for your records. Products shipped after twelve (12) months from the date of this Notice or the next printing of your label, whichever occurs first, must bear the new revised label.

If you have any questions regarding this Notice, please contact Beth Benbow at (703) 347-8072 or at benbow.bethany@epa.gov

lethyn VM (v

Signature of Approving Official:

Kathryn V. Montague/ Product Manager 23/

Herbicide Branch

Registration Division (7505P)

Date:

APR 29 2011

(FRONT PANEL)

Lawn Herbicide JD Ready-To-Use

[Bayer Advanced Southern Weed Killer for Lawns Ready-To-Use]

- Kills_listed weeds...
- Kills hard-to-control weeds
- Won't harm lawns†
- Gets the root
- · Kills over 200 types of weeds
- Systemic
- Kills the weeds, not the lawn†
- Kills Broadleaf weeds in St. Augustine & other Southern Lawns
- SmartTrack™ Marker Shows Treated Weeds
- · Kills Broadleaf weeds
- 100% Satisfaction Guaranteed* (or your money back)
- Kills 200+ Weeds
- For St. Augustine & Other Southern Lawns
- Kills Broadleaf Weeds in Southern Lawns
- . Ideal for Spot Treatment
- Rainproof in one hour

ACCEPTED

APR 29 2011

Under the Federal Insecticide, Fungioide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No 7-2(55-2

† When used as directed

ACTIVE INGREDIENTS:	
2,4-D, dimethylamine salt	0.311 %
Mecoprop-p, dimethylamine salt	0.075 %
Dicamba, dimethylamine salt	
OTHER INGREDIENTS:	99.580 %

Net Contents: 16 fl. oz. (473 mL) or 24 fl. oz. (709 mL) or 1 Gal. (3.78 L)

KEEP OUT OF REACH OF CHILDREN CAUTION

See Back Panel [Booklet] for Additional Ingredients and Precautionary Statements

[] Information is optional

1

(BACK PANEL)

This product kills broadleaf weeds in St. Augustine and other Southern lawns. This formula kills Dollarweed, Spurge, Oxalis and Dandelion without harming your lawn. Controlling weeds in your lawn has never been easier.

READ THE LABEL FIRST!

QUICK FACTS

- · Kills Broadleaf weeds in Southern Grasses
- [Contains SmartTrack™, a non-staining marker that indicates where you've sprayed]
- Kills the Weed, to the root
- Won't Harm Your Lawn†

KILLS	Broadleaf weeds such as Dollarweed, Spurge, Oxalis, Chickweeds, Clovers, Dandelion and other listed weeds
WHERE TO USE	On Lawns † (When used as directed; do not use on Carpetgrass, Dichondra or Flortam variety of St. Augustinegrass.)
WHEN TO USE	When weeds are actively growing
	For questions or comments, call toll-free 1-877 BayerAG or visit us on-line at www.Bayeradvanced.com

FOR OUTDOOR RESIDENTIAL USE ONLY

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	
Mecoprop-p, dimethylamine salt	
Dicamba, dimethylamine salt	
OTHER INGREDIENTS*:	<u>99.580 %</u>
	100.000 %

EPA Est. No. XXXX-AA-XX

EPA Reg. No. 72155-2

IMPORTANT! READ BEFORE USE

[For 1 gallon]



ATTACH SPRAYER

- Remove sprayer from bottle and extend hose.
- · Open flip-cap on bottle.
- Insert hose and plug into flip-cap opening until it snaps into place.

SPRAY

- Adjust nozzle to stream or spray position.
- Point sprayer downward and squeeze trigger repeatedly until liquid is released.

AFTER USE

- Turn sprayer nozzle to X (off) position.
- Do not unplug hose.
- Return sprayer to bottle. Sprayer will snap into place.

	DIRECTIONS FOR USE			
It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.				
For best results, read and follow all label directions.				
BEFORE YOU USE Read and follow directions when using.	 Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated areas to the point of runoff or apply when-raining or when rain is expected that day. Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other 			
	desirable plants. • Apply when wind is calm.			
	Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.			
HOW TO APPLY	Adjust spray nozzle to give a coarse spray.			
SHAKE WELL BEFORE USE	 Hold sprayer about 12 inches from weed. Aim at center of weed. NOTE: Over-applying could damage Bentgrass, Centipede and St. Augustine (including Floratam) grasses. Do not squirt the same weed more than twice (two pulls on the trigger sprayer) in a four-week period. This is enough to control weeds without harming sensitive lawn grasses. 			
	[Non-Staining marker indicates where product was applied.]			
FOR BEST RESULTS	 Apply when weeds are actively growing. Rainfall or watering one hour after application will not wash away effectiveness. 			
HOW THIS PRODUCT WORKS	This product is a <i>selective</i> [lawn] herbicide – it <i>selectively</i> kills weeds in the lawn but won't harm the lawn itself. It is a blend of proven weed killers designed to get broadleaf weeds such as dollarweed, spurge and oxalis. [This product works on broadleaf weeds by speeding up their growth, the new tissues block the food transport system (phloem), certain parts of the plants starve e.g., the roots, and the plants die.]			

				
KILLS	Annual yellow sweetclover	Cockle	Mugwort	Spiny amaranth
[CONTROLS]	Aster	Cocklebur	Musk thistle	Spiny cocklebur
	Austrian fieldcress	Common mullein	Mustard	Spiny sowthistle
	Bedstraw	Creeping charlie	Narrowleaf plantain	Spotted catsear
	Beggarticks	Creeping jenny	Narrowleaf vetch	Spotted spurge
	Betony, Florida	Cudweed	·Nettle	Spurweed
	Bindweed, field	Curly dock	Orange hawkweed	Stinging nettle
	Bird vetch	Daisy, English	Oriental cocklebur	Strawberry, India mock
	Bitter wintercress	Daisy, fleabane	Oxalis	Tall nettle
	Bittercress, hairy	Daisy, oxeye	Parsley-piert	Tall vervain
	Bitterweed	Dandelion	Parsnip	Tansy mustard
	Black-eyed Susan	Dichondra	Pearlwort	Tanweed
	Black medic	Dogbane	Pennycress	Thistle
	Black mustard	Dogfennel	Pennywort	Trailing crownvetch
	Blackseed plantain	Dollarweed	Pepperweed	Tumble mustard
	Blessed thistle	Elderberry	Pigweed	Tumble pigweed
	Bloodflower milkweed	False dandelion	Pineywoods bedstraw	Velvetleaf
	Blue lettuce	False flax	Plains coreopsis (tickseed)	Venice mallow
	Blue vervain	False sunflower	Plantain	Virginia buttonweed
	Bracted plantain	Fiddleneck	Poison ivy	Virginia creeper
	Brassbuttons	Florida pusley	Poison oak	Virginia pepperweed
	Bristly oxtongue	Frenchweed	Pokeweed	Water pennywort
	Broadleaf dock	Galinsoga	Poorjoe	Wavyleaf bullthistle
	Broadleaf plantain	Goathead	Prairie sunflower	Western clematis
	Broomweed	Goldenrod	Prickly lettuce	Western salsify
	Buckhorn	Ground ivy	Prickly sida	White mustard
	Buckhorn plantain	Gumweed	Prostrate knotweed	Wild aster
	Bulbous buttercup	Hairy fleabane	Prostrate pigweed	Wild buckwheat
	Bull thistle	Hawkweed	Prostrate spurge	Wild carrot
	Bulinettle	Healall	Prostrate vervain	Wild four-o'clock
	Burclover	Heartleaf drymary	Puncturevine	Wild garlic
	Burdock	Heath aster	Purslane, common	Wild geranium
	Burning nettle	Hedge bindweed	Ragweed	Wild lettuce
	Burweed	Hedge mustard	Red sorrel	Wild marigold
,	Buttercup	Hemp	Redroot pigweed	Wild mustard
	Buttonweed	Henbit	Redstem filaree	Wild onion
	Canada thistle	Hoary cress	Rough cinquefoil	Wild parsnip
	Carolina geranium	Hoary plantain	Rough fleabane	Wild radish
	Carpetweed	Hoary vervain	Russian pigweed	Wild rape
	Catchweed bedstraw	Horsenettle	Russian thistle	Wild strawberry
	Catnip	Jimsonweed	Scarlet pimpernel	Wild sweet potato
	Catsear	Knawel	Scotch thistle	Wild vetch
	Chickweed, common	Knotweed	Sheep sorrel	Woodsorrel
	Chickweed, mouseear	Kochia	Shepherdspurse	Wooly croton
	Chicory	Lambsquarters	Slender plantain	Woolly morningglory
	Cinquefoil	Lespedeza	Smallflower galinsoga	Woolly plantain
	Clover, crimson	Mallow	Smooth dock	Wormseed
	Clover, hop	Matchweed	Smooth pigweed	Yarrow
	Clover, red	Mexicanweed	Sorrel	Yellow rocket
	Clover, strawberry	Milk vetch	Sowthistle	Yellowflower pepperweed
	Clover, sweet	Morningglory	Spanishneedles	and other broadleaf
	Clover, white	Mouseear hawkweed	Speedwell	weeds



• Do not allow people or pets to enter the treated area until sprays have dried.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

- · Harmful if swallowed. Harmful if absorbed through skin.
- · Avoid contact with skin, eyes, or clothing. Causes moderate eye irritation.
- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.
- Wear long-sleeved shirt, long pants, shoes, socks, and gloves.
- · Remove and wash contaminated clothing before reuse.

	FIRST AID		
If in Eyes	 Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for further treatment advice. 		
If Swallowed	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. 		
If on Skin or Clothing	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 minutes. Call a poison control center or doctor for further treatment advice. 		

NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 877-229-3763 for medical emergency information. Active Ingredients: 0.311% 2,4-D, dimethylamine salt [CAS # 2008-39-1], 0.258% acid equivalent; 0.075% Mecoprop-p, dimethylamine salt [CAS # 66423-09-4], 0.062% acid equivalent; and 0.034% Dicamba, dimethylamine salt [CAS # 2300-66-5], 0.028% acid equivalent.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment wash waters or rinsate.

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal		
PESTICIDE STORAGE	Store in original container in a cool dry place, out of the reach of children, preferably a locked storage cabinet.	
PESTICIDE DISPOSAL AND CONTAINER HANDLING:	If empty: Non-refillable container. Do not refill or reuse this container. Place empty container in trash or offer for recycling if available. If partly filled:	
	Call your local solid waste agency toll free 1-877-229-3724 for disposal instructions.	
	Never place unused product down any indoor or outdoor drain.	

NOTICE: Research and testing have determined that the "Directions For Use" are appropriate for the proper use of this product under expected conditions. To the extent consistent with applicable law, the Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.



www.bayeraduanced.com
A Business Unit of Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709

Lawn Herbicide JD RTU (PENDING) 03/24/11, Resubmitted 04/05/11