

72155-4

08/30/2000

1/13



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

NOTICE OF PESTICIDE:

 x Registration
 Reregistration

(under FIFRA, as amended.)

EPA Reg.
Number:

72155-4

Date of Issuance:

AUG 30 2000

Term of Issuance:

Conditional

Name of Pesticide Product:

Lawn Herbicide JD Concentrate

Name and Address of Registrant (include ZIP Code):

Bayer-Pursell, LLC
1500 Urban Center Pkwy., Suite 520
Birmingham, AL 35242

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit/cite all data required for the registration/reregistration of your product when the Agency requires all registrants of similar products to submit data.
2. Make the following label changes:
 - a. Revise the EPA Registration Number to read, "EPA Reg. No. 72155-4".
 - b. Under the section entitled "ENVIRONMENTAL HAZARDS" on page (6) six of your product labeling, please incorporate the following groundwater contamination statement:

"Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D and MCPP pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination."

Signature of Approving Official:

Date:

AUG 30 2000

page 2

EPA Reg. No. 72155-4

- c. Under the headings entitled "Direction For Use" within the re-entry restriction sections, delete the following statement: "Children and pets may re-enter the treated area after it has dried." The aforementioned statement must be superceded by the following re-entry restriction statement: "Do not allow people (other than applicator) on treatment area during application. Do not enter treatment areas until spray has dried or dust has settled."

3. Submit (2) two copies of the revised final printed label for the record.

A stamped copy of the label is enclosed for your records.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Joanne I. Miller
Product Manger (23)
Herbicide Branch
Registration Division (7505C)

Enclosures

(FRONT PANEL)

Lawn Herbicide JD Concentrate

This label contains separate label statements for **Outdoor Residential Pump Sprayer/Hose-end Sprayer Concentrate** use product and **Outdoor Residential Ready-To-Spray** use product.

- Effective Weed Protection
- Effective Broadleaf Weed Protection
- Effectively Protects Lawns from Broadleaf Weeds
- Kills All Major Broadleaf Weeds
- Kills Weeds - Won't Harm Lawns
- Kills to the Root
- Protects Against All Major Broadleaf Weeds
- Tough Weed Protection
- Weed Protection System
- Systemic
- 1 Quart treats up to 16,000 square feet
- Convenient MEASURE-UP™ Cup

ACCEPTED
with COMMENTS
In EPA Letter Dated

AUG 30 2000

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
7255-4

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	7.59 %
Mecoprop, dimethylamine salt	3.66 %
Dicamba, dimethylamine salt	0.84 %
OTHER INGREDIENTS:	87.91 %

Net Contents: 16 fl. oz. (473 mL) or 24 fl. oz. (709 mL) or
32 fl. oz. (946 mL) or 1 Gal. (3.78 L)

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel [Booklet] for Additional Ingredients and
Precautionary Statements

[] Information is optional

(FRONT PANEL)

Lawn Herbicide JD Concentrate

- Effective Weed Protection
- Effective Broadleaf Weed Protection
- Effectively Protects Lawns from Broadleaf Weeds
- Kills All Major Broadleaf Weeds
- Kills Weeds - Won't Harm Lawns
- Kills to the Root
- Protects Against All Major Broadleaf Weeds
- Tough Weed Protection
- Weed Protection System
- Systemic
- 1 Quart treats up to 16,000 square feet
- Convenient MEASURE-UP™ Cup

(Pump Sprayer/Hose-end Sprayer label)

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	7.59 %
Mecoprop, dimethylamine salt	3.66 %
Dicamba, dimethylamine salt	0.84 %
OTHER INGREDIENTS:	87.91 %

Net Contents: 16 fl. oz. (473 mL) or 24 fl. oz. (709 mL) or
32 fl. oz. (946 mL) or 1 Gal. (3.78 L)



KEEP OUT OF REACH OF CHILDREN**CAUTION**

See Back Panel [Booklet] for Additional Ingredients and
Precautionary Statements

[] Information is optional

(BACK PANEL)

READ THE LABEL FIRST!

QUICK FACTS [Marketing claims positioned here]	
KILLS	Broadleaf weeds such as dandelion, clover, chickweed and other listed weeds
WHERE TO USE	Lawns (Not for use on Carpetgrass, Dichondra, Floratam variety of St. Augustinegrass or desirable clovers.)
WHEN TO USE	<ul style="list-style-type: none"> • When weeds are actively growing • Do not apply when temperature exceeds 85 °F.
HOW MUCH TO USE	Northern Lawns: Fescue, Kentucky Bluegrass, and Perennial Ryegrasses and Hardy Southern Lawns: Bermuda and Zoysia grasses 2 fl. oz. per 500 sq. ft. [1 qt treats up to 8,000 sq. ft.] Southern lawns: Bahia, Centipede and St. Augustinegrass grasses 1 fl. oz. per 500 sq. ft. [1 qt treats up to 16,000 sq. ft.]
 	For questions or comments, call toll-free 1-877 BayerAL or visit us on-line at www.AdvancedLawn.com

FOR OUTDOOR RESIDENTIAL USE ONLY

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt 7.59 %
Mecoprop, dimethylamine salt 3.66 %
Dicamba, dimethylamine salt 0.84 %
OTHER INGREDIENTS*: 87.91 %

100.00 %

*Other ingredients include: Water and surfactants (to keep the active ingredients dispersed in water).

THIS PRODUCT CONTAINS:

6.30% 2,4-D acid equivalent
3.02% Mecoprop acid equivalent
0.69% Dicamba acid equivalent
Isomer specific by AOAC Method

EPA Reg. No. 72155-__

EPA Est. No. __AA-__

DIRECTIONS FOR USE	
It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. For best results, read and follow all label directions.	
BEFORE YOU USE Read and follow directions when using.	<ul style="list-style-type: none"> Do not apply near lakes, streams, rivers or ponds. Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Apply when wind is calm.
HOW TO USE	<ul style="list-style-type: none"> Determine amount of product to use based on grass type, sprayer type and size of area to be treated. Prepare product for use according to How to Apply. Apply to lawn.
HOW MUCH TO APPLY	<p>Northern Lawns: Fescue, Kentucky Bluegrass, and Perennial Rye And Hardy Southern Lawns: Bermuda and Zoysia grasses 2 fl. oz. (4 Tbs.) per gallon of water. Each gallon of solution treats 500 sq. ft.</p> <p>Southern Lawns: Bahia, Centipede, and St. Augustinegrass 1 fl. oz. (2 Tbs.) per gallon of water. Each gallon of solution treats 500 sq. ft.</p> <p>Apply as a broadcast treatment no more than two times a year.</p> <p>TIP: Measure length and width of area to be treated in feet. Multiply these 2 numbers together. This is the area to be treated in square feet (sq. ft.)</p>
HOW TO APPLY	
Pump Style Sprayers	<ol style="list-style-type: none"> Add the appropriate amounts of water and concentrate to the sprayer tank. Close sprayer, shake well and pressurize. Adjust nozzle to a coarse spray pattern. Large Areas – Apply evenly over area. Individual Weeds – Hold tip of wand about 12 inches from weed, aim at center of weed and spray. Occasionally re-pressurize the sprayer if needed to maintain a good spray pattern.
Continued in next column	

Directions For Use continued	
Dial Style Hose-End Sprayers	<ol style="list-style-type: none"> Fill sprayer jar with enough concentrate to cover area to be treated. Close sprayer and set dial. TIP: The numbers on Dial-Style Hose-End sprayers are ounces of concentrate per gallon of spray. Attach sprayer to hose and apply. To avoid contact with the treated area, begin by spraying area farthest from faucet walking backwards as you spray.
Concentrate Plus Water Hose-End Sprayer	<ol style="list-style-type: none"> Pour concentrate into sprayer jar to the required fluid ounce level. Add the appropriate amount of water to the gallons level. Replace sprayer top and shake well before applying. To avoid contact with the treated area, begin by spraying area farthest from faucet walking backwards as you spray.
FOR BEST RESULTS	<ul style="list-style-type: none"> If the lawn is dry, water 24 hours before using this product. Do not water for 24 hours after application. Do not use this product if rain is likely within 24 hours. Apply after mowing [(mowing encourages weed growth which enhances uptake of the weed killer [herbicide])]. Certain hard-to-kill weeds may need to be spot-treated in 14 days. Do not apply to new lawns until the grass has reached a height of at least 2 inches.
HOW LAWN HERBICIDE JD CONCENTRATE WORKS	<p>LAWN HERBICIDE JD CONCENTRATE is a selective [lawn] herbicide - it selectively kills weeds in the lawn but won't harm the lawn itself. It is a blend of proven weed killers designed to get broadleaf weeds such as dandelion, clover and chickweed. [This product works on broadleaf weeds by speeding up their growth, the new tissues block the food transport system (phloem), certain parts of the plants starve e.g., the roots, and the plants die.]</p>



Children and pets may re-enter the treated area after it has dried.

CONTROLS [KILLS]		
Annual yellow sweetclover	Daisy, oxeye	
Aster	Dandelion	
Austrian fieldcress	Dichondra	
Bedstraw	Dogbane	
Beggaricks	Dogfennel	
Betony, Florida	Dollarweed	
Bindweed, field	Elderberry	
Bird vetch	False dandelion	
Bitter wintercress	False flax	
Bittercress, hairy	False sunflower	
Bitterweed	Fiddleneck	
Black-eyed Susan	Florida pusley	
Black medic	Frenchweed	
Black mustard	Galinsoga	
Blackseed plantain	Goathead	
Blessed thistle	Goldenrod	
Bloodflower milkweed	Ground ivy	
Blue lettuce	Gumweed	
Blue vervain	Hairy fleabane	
Bracted plantain	Hawkweed	
Brassbuttons	Healall	
Bristly oxtongue	Heartleaf drymary	
Broadleaf dock	Heath aster	
Broadleaf plantain	Hedge bindweed	
Broomweed	Hedge mustard	
Buckhorn	Hemp	
Buckhorn plantain	Henbit	
Bulbous buttercup	Hoary cress	
Bull thistle	Hoary plantain	
Bullnettle	Hoary vervain	
Burclover	Horsenettle	
Burdock	Jimsonweed	
Burning nettle	Knawel	
Burweed	Knotweed	
Buttercup	Kochia	
Buttonweed	Lambsquarters	
Canada thistle	Lespedeza	
Carolina geranium	Mallow	
Carpetweed	Matchweed	
Catchweed	Mexicanweed	
bedstraw	Milk vetch	
Catnip	Morningglory	
Catsear	Mouseear	
Chickweed, common	hawkweed	
Chickweed, mouseear	Mugwort	
Chicory	Musk thistle	
Cinquefoil	Mustard	
Clover, crimson	Narrowleaf plantain	
Clover, hop	Narrowleaf vetch	
Clover, red	Nettle	
Clover, strawberry	Orange hawkweed	
Clover, sweet	Oriental cocklebur	
Clover, white	Oxalis	
Cockle	Parsley-piert	
Cocklebur	Parsnip	
Common mullein	Pearlwort	
Creeping Charlie	Pennycress	
Creeping Jenny	Pennywort	
Cudweed	Pepperweed	
Curly dock	Pigweed	
Daisy, English	Pineywoods bedstraw	
Daisy, fleabane	Plains coreopsis (tickseed)	
	Plantain	
	Poison ivy	

Continued in next column

CONTROLS LIST continued		
CONTROLS [KILLS]		
Poison oak	Tansy mustard	
Pokeweed	Tanweed	
Poorjoe	Thistle	
Prairie sunflower	Trailing crownvetch	
Prickly lettuce	Tumble mustard	
Prickly sida	Tumble pigweed	
Prostrate knotweed	Velvetleaf	
Prostrate pigweed	Venice mallow	
Prostrate spurge	Virginia buttonweed	
Prostrate vervain	Virginia creeper	
Puncturevine	Virginia pepperweed	
Purslane, common	Water pennywort	
Ragweed	Wavyleaf bullthistle	
Red sorrel	Western clematis	
Redroot pigweed	Western salsify	
Redstem filaree	White mustard	
Rough cinquefoil	Wild aster	
Rough fleabane	Wild buckwheat	
Russian pigweed	Wild carrot	
Russian thistle	Wild four-o'clock	
Scarlet pimpernel	Wild garlic	
Scotch thistle	Wild geranium	
Sheep sorrel	Wild lettuce	
Shepherdspurse	Wild marigold	
Slender plantain	Wild mustard	
Smallflower galinsoga	Wild onion	
Smooth dock	Wild parsnip	
Smooth pigweed	Wild radish	
Sorrel	Wild rape	
Sowthistle	Wild strawberry	
Spanishneedles	Wild sweet potato	
Speedwell	Wild vetch	
Spiny amaranth	Woodsorrel	
Spiny cocklebur	Woolly croton	
Spiny sowthistle	Woolly morningglory	
Spotted catsear	Woolly plantain	
Spotted spurge	Wormseed	
Spurweed	Yarrow	
Stinging nettle	Yellow rocket	
Strawberry, India mock	Yellowflower pepperweed	
Tall nettle	and other broadleaf weeds	
Tall vervain		

IMPORTANT: Directions for Storage and Disposal

Storage Conditions: Store in original container out of the reach of children, preferably in a locked storage cabinet. Protect from freezing.

- Disposal:**
- Do not reuse empty container.
 - Rinse the empty product container thoroughly and disperse the rinse water on weeds as part of application.
 - Place empty container in trash.

Important: It is best to use all of the product in accordance with label directions. If it is necessary to dispose of unused product, please follow any applicable state and local guidelines.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS**

CAUTION

- Harmful if swallowed.
- Causes moderate eye irritation.
- Avoid contact with eyes, skin, or clothing.
- When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

After using this product:

- Rinse gloves before removing.
- Wash hands thoroughly with soap and water.
- If product gets on clothing, remove clothing as soon as possible and launder separately before reuse.

FIRST AID	
IN CASE OF ACCIDENTAL...	IMMEDIATELY...
Swallowing	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything to an unconscious person.
Contact with eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
Contact with skin	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 to 20 minutes.• Call a poison control center or doctor for treatment advice.
Inhalation	<ul style="list-style-type: none">• Move person to fresh air.• If not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 877-229-3763 for medical emergency information. Active Ingredients: 7.59% 2,4-D, dimethylamine salt [CAS # 2008-39-1], 3.66% Mecoprop, dimethylamine salt [CAS # 32351-70-5], and 0.84% Dicamba, dimethylamine salt [CAS # 2300-66-5].	

ENVIRONMENTAL HAZARDS

- This product is toxic to aquatic invertebrates. Do not apply directly to lakes, streams, rivers or ponds
- Drift or runoff may harm aquatic invertebrates and plants.
- Do not contaminate water when disposing of equipment washwaters.
- Do not apply in windy conditions.

NOTICE: Research and testing have determined that the "Directions For Use" are appropriate for the proper use of LAWN HERBICIDE JD CONCENTRATE under expected conditions. The Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.

D - 9443-1 050900

Bayer-Pursell, LLC
1500 Urban Center Drive
Birmingham, AL 35242

(FRONT PANEL)

Lawn Herbicide JD Concentrate

- Effective Weed Protection
- Effective Broadleaf Weed Protection
- Effectively Protects Lawns from Broadleaf Weeds
- Kills All Major Broadleaf Weeds
- Kills Weeds - Won't Harm Lawns
- Kills to the Root
- Protects Against All Major Broadleaf Weeds
- Tough Weed Protection
- Weed Protection System
- Systemic
- 1 Quart treats up to 16,000 square feet
- Convenient MEASURE-UP™ Cup

(Ready - To - Spray label)

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	7.59 %
Mecoprop, dimethylamine salt	3.66 %
Dicamba, dimethylamine salt	0.84 %
OTHER INGREDIENTS:	87.91 %

Net Contents: 16 fl. oz. (473 mL) or 24 fl. oz. (709 mL) or
32 fl. oz. (946 mL) or 1 Gal. (3.78 L)

KEEP OUT OF REACH OF CHILDREN



CAUTION

See Back Panel [Booklet] for Additional Ingredients and
Precautionary Statements

[] Information is optional

(BACK PANEL)

READ THE LABEL FIRST!

QUICK FACTS [Marketing claims positioned here]	
KILLS	Broadleaf weeds such as dandelion, clover, chickweed and other listed weeds
WHERE TO USE	Lawns (Not for use on Carpetgrass, Dichondra, Floratam variety of St. Augustinegrass or desirable clovers.)
WHEN TO USE	<ul style="list-style-type: none"> • When weeds are actively growing • Do not apply when temperature exceeds 85 °F.
HOW MUCH TO USE	Northern Lawns: Fescue, Kentucky Bluegrass, and Perennial Ryegrasses and Hardy Southern Lawns: Bermuda and Zoysia grasses One qt treats up to 8,000 sq. ft. Southern lawns: Bahia, Centipede and St. Augustinegrass grasses One qt treats up to 16,000 sq. ft. No measuring needed. Dilutes automatically.
  For questions or comments, call toll-free 1-877 BayerAL or visit us on-line at www.AdvancedLawn.com	

FOR OUTDOOR RESIDENTIAL USE ONLY

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	7.59 %
Mecoprop, dimethylamine salt	3.66 %
Dicamba, dimethylamine salt	0.84 %
OTHER INGREDIENTS*:	87.91 %

100.00 %

*Other ingredients include: Water and surfactants (to keep the active ingredients dispersed in water).

THIS PRODUCT CONTAINS:

6.30% 2,4-D acid equivalent
3.02% Mecoprop acid equivalent
0.69% Dicamba acid equivalent
Isomer specific by AOAC Method


EPA Reg. No. 72155-__

EPA Est. No. __-AA-__

DIRECTIONS FOR USE	
It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. For best results, read and follow all label directions.	
BEFORE YOU USE Read and follow directions when using.	<ul style="list-style-type: none"> Do not apply near lakes, streams, rivers or ponds. Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Apply when wind is calm.
HOW TO USE	<ul style="list-style-type: none"> Determine amount of product to use based on grass type and size of area to be treated. See How Much To Use section of Quick Facts Box. Prepare and apply product according to the following section, How To Attach and Spray.
HOW TO ATTACH & SPRAY	<p>This product is ready to spray. Just attach the bottle to your garden hose according to the following directions. Water automatically mixes with the concentrate as you spray.</p> <ol style="list-style-type: none"> 1. Connect your garden hose to the bottle. Be sure the blue water control valve on top of the spray nozzle is in the OFF position. 2. Turn on water at faucet. Twist the small, star-shaped, blue product-control button to align the flat portion with the lock tab. Push the button in to open. 3. To begin spraying, point nozzle toward the area to be treated and turn the blue water control valve to the ON position. Tab will be horizontal. 4. To stop spraying, turn the blue water control valve on top of the spray nozzle to OFF. To close, push in the small round button opposite the star-shaped product-control button. Turn off faucet. 5. The unused portion can be saved and used at a later date. <p>Apply as a broadcast treatment no more than two times a year.</p>

continued in next column

TIPS FOR APPLYING	
HOW TO APPLY	<ol style="list-style-type: none"> 1. Extend hose to the farthest area to be treated and work back toward the faucet to avoid walking over wet, treated grass. 2. Walk at a steady pace while spraying, using an even sweeping motion, slightly overlapping the treated areas.
FOR BEST RESULTS	<ul style="list-style-type: none"> If the lawn is dry, water 24 hours before using this product. Do not water for 24 hours after application. Do not use this product if rain is likely within 24 hours. Apply after mowing [(mowing encourages weed growth which enhances uptake of the weed killer [herbicide])]. Certain hard-to-kill weeds may need to be spot-treated in 14 days. Do not apply to new lawns until the grass has reached a height of at least 2 inches.
HOW LAWN HERBICIDE JD CONCENTRATE WORKS	<p>LAWN HERBICIDE JD CONCENTRATE is a selective [lawn] herbicide - it selectively kills weeds in the lawn but won't harm the lawn itself. It is a blend of proven weed killers designed to get broadleaf weeds such as dandelion, clover and chickweed. [This product works on broadleaf weeds by speeding up their growth, the new tissues block the food transport system (phloem), certain parts of the plants starve e.g., the roots, and the plants die.]</p>

	Children and pets may re-enter the treated area after it has dried.
---	---

CONTROLS [KILLS]		
Annual yellow sweetclover	Daisy, oxeye	
Aster	Dandelion	
Austrian fieldcress	Dichondra	
Bedstraw	Dogbane	
Beggarticks	Dogfennel	
Betony, Florida	Dollarweed	
Bindweed, field	Elderberry	
Bird vetch	False dandelion	
Bitter wintercress	False flax	
Bittercress, hairy	False sunflower	
Bitterweed	Fiddleneck	
Black-eyed Susan	Florida pusley	
Black medic	Frenchweed	
Black mustard	Galinsoga	
Blackseed plantain	Goathead	
Blessed thistle	Goldenrod	
Bloodflower	Ground ivy	
milkweed	Gumweed	
Blue lettuce	Hairy fleabane	
Blue vervain	Hawkweed	
Bracted plantain	Healall	
Brassbuttons	Heartleaf drymary	
Bristly oxtongue	Heath aster	
Broadleaf dock	Hedge bindweed	
Broadleaf plantain	Hedge mustard	
Broomweed	Hemp	
Buckhorn	Henbit	
Buckhorn plantain	Hoary cress	
Bulbous buttercup	Hoary plantain	
Bull thistle	Hoary vervain	
Bullnettle	Horsenettle	
Burclover	Jimsonweed	
Burdock	Knawel	
Burning nettle	Knotweed	
Burweed	Kochia	
Buttercup	Lambsquarters	
Buttonweed	Lespedeza	
Canada thistle	Mallow	
Carolina geranium	Matchweed	
Carpetweed	Mexicanweed	
Catchweed	Milk vetch	
bedstraw	Morningglory	
Catnip	Mouseear	
Catsear	hawkweed	
Chickweed, common	Mugwort	
Chickweed, mouseear	Musk thistle	
Chicory	Mustard	
Cinquefoil	Narrowleaf plantain	
Clover, crimson	Narrowleaf vetch	
Clover, hop	Nettle	
Clover, red	Orange hawkweed	
Clover, strawberry	Oriental cocklebur	
Clover, sweet	Oxalis	
Clover, white	Parsley-piert	
Cockle	Parsnip	
Cocklebur	Pearlwort	
Common mullein	Pennycress	
Creeping Charlie	Pennywort	
Creeping Jenny	Pepperweed	
Cudweed	Pigweed	
Curly dock	Pineywoods bedstraw	
Daisy, English	Plains coreopsis (tickseed)	
Daisy, fleabane	Plantain	
	Poison ivy	

Continued in next column

CONTROLS LIST continued		
CONTROLS [KILLS]		
Poison oak	Tansy mustard	
Pokeweed	Tanweed	
Poorjoe	Thistle	
Prairie sunflower	Trailing crownvetch	
Prickly lettuce	Tumble mustard	
Prickly sida	Tumble pigweed	
Prostrate knotweed	Velvetleaf	
Prostrate pigweed	Venice mallow	
Prostrate spurge	Virginia buttonweed	
Prostrate vervain	Virginia creeper	
Puncturevine	Virginia pepperweed	
Purslane, common	Water pennywort	
Ragweed	Wavyleaf bullthistle	
Red sorrel	Western clematis	
Redroot pigweed	Western salsify	
Redstem filaree	White mustard	
Rough cinquefoil	Wild aster	
Rough fleabane	Wild buckwheat	
Russian pigweed	Wild carrot	
Russian thistle	Wild four-o'clock	
Scarlet pimpernel	Wild garlic	
Scotch thistle	Wild geranium	
Sheep sorrel	Wild lettuce	
Shepherdspurse	Wild marigold	
Slender plantain	Wild mustard	
Smallflower galinsoga	Wild onion	
Smooth dock	Wild parsnip	
Smooth pigweed	Wild radish	
Sorrel	Wild rape	
Sowthistle	Wild strawberry	
Spanishneedles	Wild sweet potato	
Speedwell	Wild vetch	
Spiny amaranth	Woodsorrel	
Spiny cocklebur	Woolly croton	
Spiny sowthistle	Woolly morningglory	
Spotted catsear	Woolly plantain	
Spotted spurge	Wormseed	
Spurweed	Yarrow	
Stinging nettle	Yellow rocket	
Strawberry, India mock	Yellowflower	
Tall nettle	pepperweed	
Tall vervain	and other broadleaf weeds	

IMPORTANT: Directions for Storage and Disposal

Storage Conditions: Store in original container out of the reach of children, preferably in a locked storage cabinet. Protect from freezing.

- Disposal:**
- Do not reuse empty container.
 - Securely wrap original container in several layers of newspaper and discard in trash.
 - Place empty container in trash.

Important: It is best to use all of the product in accordance with label directions. If it is necessary to dispose of unused product, please follow any applicable state and local guidelines.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS**

CAUTION

- Harmful if swallowed.
- Causes moderate eye irritation.
- Avoid contact with eyes, skin, or clothing.
- When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

After using this product:

- Rinse gloves before removing.
- Wash hands thoroughly with soap and water.
- If product gets on clothing, remove clothing as soon as possible and launder separately before reuse.

FIRST AID	
IN CASE OF ACCIDENTAL...	IMMEDIATELY...
Swallowing	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything to an unconscious person.
Contact with eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
Contact with skin	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 to 20 minutes.• Call a poison control center or doctor for treatment advice.
Inhalation	<ul style="list-style-type: none">• Move person to fresh air.• If not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 877-229-3763 for medical emergency information. Active Ingredients: 7.59% 2,4-D, dimethylamine salt [CAS # 2008-39-1], 3.66% Mecoprop, dimethylamine salt [CAS # 32351-70-5], and 0.84% Dicamba, dimethylamine salt [CAS # 2300-66-5].	

ENVIRONMENTAL HAZARDS

- This product is toxic to aquatic invertebrates. Do not apply directly to lakes, streams, rivers or ponds.
- Drift or runoff may harm aquatic invertebrates and plants.
- Do not contaminate water when disposing of equipment washwaters.
- Do not apply in windy conditions.

NOTICE: Research and testing have determined that the "Directions For Use" are appropriate for the proper use of LAWN HERBICIDE JD CONCENTRATE under expected conditions. The Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.

D - 9443-2 050900

Bayer-Pursell, LLC
1500 Urban Center Drive
Birmingham, AL 35242