

72155-4

12/28/2000


1/6

(FRONT PANEL)

# Lawn Herbicide JD Concentrate

This label contains separate label statements for **Outdoor Residential Pump Sprayer/ Hose-end Sprayer Concentrate** use product and **Outdoor Residential Ready-To-Spray** use product.

- Effective Weed Protection
- Effective Broadleaf Weed Protection
- Effectively Protects Lawns from Broadleaf Weeds
- Kills All Major Broadleaf Weeds
- Kills Weeds - Won't Harm Lawns
- Kills to the Root
- Protects Against All Major Broadleaf Weeds
- Tough Weed Protection
- Weed Protection System
- Systemic
- 1 Quart treats up to 16,000 square feet
- Convenient MEASURE-UP™ Cup
- Kills Broadleaf weeds in St. Augustine & other Southern Lawns
- Kills Over 200 Broadleaf Weeds
- Won't Harm Your Lawn



## ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	7.59 %
Mecoprop, dimethylamine salt	3.66 %
Dicamba, dimethylamine salt	0.84 %
OTHER INGREDIENTS:	87.91 %

Net Contents: 16 fl. oz. (473 mL) or 24 fl. oz. (709 mL) or  
32 fl. oz. (946 mL) or 1 Gal. (3.78 L)

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

See Back Panel [Booklet] for Additional Ingredients and Precautionary Statements



[ ] Information is optional



**(BACK PANEL)**

Lawn Herbicide JD Concentrate kills over 200 types of broadleaf weeds in St. Augustine and other Southern lawns. This special formula kills Dollarweed, Spurge, Oxalis and Dandelion without harming your lawn. Controlling weeds in your lawn has never been easier.

**READ THE LABEL FIRST!**

<b>QUICK FACTS</b> [Marketing claims positioned here]	
<b>KILLS</b>	Over 200 broadleaf weeds such as Dollarweed, Spurge and Oxalis, Dandelion, Clover, Chickweed and other listed weeds
<b>WHERE TO USE</b>	On Lawns (Not for use on Carpetgrass, Dichondra, Floratam variety of St. Augustinegrass or desirable clovers.)
<b>WHEN TO USE</b>	<ul style="list-style-type: none"> <li>• When weeds are actively growing</li> <li>• Do not apply when temperature exceeds 85 °F.</li> </ul>
<b>HOW MUCH TO USE</b>	<b>Northern Lawns: Fescue, Kentucky Bluegrass, and Perennial Ryegrasses</b> <b>and</b> <b>Hardy Southern Lawns: Bermuda and Zoysia grasses</b> 2 fl. oz. per 500 sq. ft. [1 qt treats up to 8,000 sq. ft.] <b>Southern lawns: Bahia, Centipede and St. Augustinegrass grasses</b> 1 fl. oz. per 500 sq. ft. [1 qt treats up to 16,000 sq. ft.]
  For questions or comments, call toll-free 1-877 BayerAL or visit us on-line at <a href="http://www.AdvancedLawn.com">www.AdvancedLawn.com</a>	

**FOR OUTDOOR RESIDENTIAL USE ONLY**

**ACTIVE INGREDIENTS:**

2,4-D, dimethylamine salt ..... 7.59 %  
 Mecoprop, dimethylamine salt ..... 3.66 %  
 Dicamba, dimethylamine salt ..... 0.84 %

**OTHER INGREDIENTS\*:** ..... 87.91 %  
 100.00 %

\*Other ingredients include: Water and surfactants (to keep the active ingredients dispersed in water).

**THIS PRODUCT CONTAINS:**


6.30% 2,4-D acid equivalent  
 3.02% Mecoprop acid equivalent  
 0.69% Dicamba acid equivalent  
 Isomer specific by AOAC Method

EPA Reg. No. 72155-4

EPA Est. No. \_-AA-\_

DIRECTIONS FOR USE	
It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. For best results, read and follow all label directions.	
<b>BEFORE YOU USE</b> Read and follow directions when using.	<ul style="list-style-type: none"> <li>Do not apply near lakes, streams, rivers or ponds.</li> <li>Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Apply when wind is calm.</li> </ul>
<b>HOW TO USE</b>	<ul style="list-style-type: none"> <li>Determine amount of product to use based on grass type, sprayer type and size of area to be treated.</li> <li>Prepare product for use according to How to Apply.</li> <li>Apply to lawn.</li> </ul>
<b>HOW MUCH TO APPLY</b>	<p><b>Northern Lawns: Fescue, Kentucky Bluegrass, and Perennial Rye And</b></p> <p><b>Hardy Southern Lawns: Bermuda and Zoysia grasses</b></p> <p>2 fl. oz. (4 Tbs.) per gallon of water.  Each gallon of solution treats 500 sq. ft.</p> <p><b>Southern Lawns: Bahia, Centipede, and St. Augustinegrass</b></p> <p>1 fl. oz. (2 Tbs.) per gallon of water.  Each gallon of solution treats 500 sq. ft.</p> <p>Apply as a broadcast treatment no more than two times a year.</p> <p><b>TIP:</b> Measure length and width of area to be treated in feet. Multiply these 2 numbers together. This is the area to be treated in square feet (sq. ft.)</p>
<b>HOW TO APPLY</b>	
<b>Pump Style Sprayers</b>	<ol style="list-style-type: none"> <li>Add the appropriate amounts of water and concentrate to the sprayer tank.</li> <li>Close sprayer, shake well and pressurize.</li> <li>Adjust nozzle to a coarse spray pattern.</li> <li>Large Areas – Apply evenly over area.  Individual Weeds – Hold tip of wand about 12 inches from weed, aim at center of weed and spray.</li> <li>Occasionally re-pressurize the sprayer if needed to maintain a good spray pattern.</li> </ol>
Continued in next column	

Directions For Use continued	
<b>Dial Style Hose-End Sprayers</b>	<ol style="list-style-type: none"> <li>Fill sprayer jar with enough concentrate to cover area to be treated. Close sprayer and set dial. <b>TIP:</b> The numbers on Dial-Style Hose-End sprayers are ounces of concentrate per gallon of spray.</li> <li>Attach sprayer to hose and apply.</li> <li>To avoid contact with the treated area, begin by spraying area farthest from faucet walking backwards as you spray.</li> </ol>
<b>Concentrate Plus Water Hose-End Sprayer</b>	<ol style="list-style-type: none"> <li>Pour concentrate into sprayer jar to the required fluid ounce level.</li> <li>Add the appropriate amount of water to the gallons level.</li> <li>Replace sprayer top and shake well before applying.</li> <li>To avoid contact with the treated area, begin by spraying area farthest from faucet walking backwards as you spray.</li> </ol>
<b>FOR BEST RESULTS</b>	<ul style="list-style-type: none"> <li>If the lawn is dry, water 24 hours before using this product.</li> <li>Do not water for 24 hours after application. Do not use this product if rain is likely within 24 hours.</li> <li>Apply after mowing [(mowing encourages weed growth which enhances uptake of the weed killer [herbicide])].</li> <li>Certain hard-to-kill weeds may need to be spot-treated in 14 days.</li> <li>Do not apply to new lawns until the grass has reached a height of at least 2 inches.</li> </ul>
<b>HOW LAWN HERBICIDE JD CONCENTRATE WORKS</b>	<p><b>LAWN HERBICIDE JD CONCENTRATE</b> is a selective [lawn] herbicide - it selectively kills weeds in the lawn but won't harm the lawn itself. It is a blend of proven weed killers designed to get broadleaf weeds such as Dollarweed, Spurge and Oxalis. [This product works on broadleaf weeds by speeding up their growth, the new tissues block the food transport system (phloem), certain parts of the plants starve e.g., the roots, and the plants die.]</p>

	<p>Do not allow people (other than applicator) or pets on treated area during application.</p> <p>People and pets may re-enter the treated area after it has dried.</p>
--	---

CONTROLS [KILLS]		
Annual yellow sweetclover	Daisy, oxeye	
Aster	Dandelion	
Austrian fieldcress	Dichondra	
Bedstraw	Dogbane	
Beggarticks	Dogfennel	
Betony, Florida	Dollarweed	
Bindweed, field	Elderberry	
Bird vetch	False dandelion	
Bitter wintercress	False flax	
Bittercress, hairy	False sunflower	
Bitterweed	Fiddleneck	
Black-eyed Susan	Florida pusley	
Black medic	Frenchweed	
Black mustard	Galinsoga	
Blackseed plantain	Goathead	
Blessed thistle	Goldenrod	
Bloodflower	Ground ivy	
milkweed	Gumweed	
Blue lettuce	Hairy fleabane	
Blue vervain	Hawkweed	
Bracted plantain	Healall	
Brassbuttons	Heartleaf drymary	
Bristly oxtongue	Heath aster	
Broadleaf dock	Hedge bindweed	
Broadleaf plantain	Hedge mustard	
Broomweed	Hemp	
Buckhorn	Henbit	
Buckhorn plantain	Hoary cress	
Bulbous buttercup	Hoary plantain	
Bull thistle	Hoary vervain	
Bullnettle	Horsenettle	
Burclover	Jimsonweed	
Burdock	Knawel	
Burning nettle	Knotweed	
Burweed	Kochia	
Buttercup	Lambsquarters	
Buttonweed	Lespedeza	
Canada thistle	Mallow	
Carolina geranium	Matchweed	
Carpetweed	Mexicanweed	
Catchweed	Milk vetch	
bedstraw	Morningglory	
Catnip	Mouseear	
Catsear	hawkweed	
Chickweed, common	Mugwort	
Chickweed, mouseear	Musk thistle	
Chicory	Mustard	
Cinquefoil	Narrowleaf plantain	
Clover, crimson	Narrowleaf vetch	
Clover, hop	Nettle	
Clover, red	Orange hawkweed	
Clover, strawberry	Oriental cocklebur	
Clover, sweet	Oxalis	
Clover, white	Parsley-piert	
Cockle	Parsnip	
Cocklebur	Pearlwort	
Common mullein	Pennycress	
Creeping Charlie	Pennywort	
Creeping Jenny	Pepperweed	
Cudweed	Pigweed	
Curly dock	Pineywoods	
Daisy, English	bedstraw	
Daisy, fleabane	Plains coreopsis	
	(tickseed)	
	Plantain	
	Poison ivy	

Continued in next column

CONTROLS LIST continued		
CONTROLS [KILLS]		
Poison oak	Tansy mustard	
Pokeweed	Tanweed	
Poorjoe	Thistle	
Prairie sunflower	Trailing crownvetch	
Prickly lettuce	Tumble mustard	
Prickly sida	Tumble pigweed	
Prostrate knotweed	Velvetleaf	
Prostrate pigweed	Venice mallow	
Prostrate spurge	Virginia	
Prostrate vervain	buttonweed	
Puncturevine	Virginia creeper	
Purslane, common	Virginia	
Ragweed	pepperweed	
Red sorrel	Water pennywort	
Redroot pigweed	Wavyleaf bullthistle	
Redstem filaree	Western clematis	
Rough cinquefoil	Western salsify	
Rough fleabane	White mustard	
Russian pigweed	Wild aster	
Russian thistle	Wild buckwheat	
Scarlet pimpernel	Wild carrot	
Scotch thistle	Wild four-o'clock	
Sheep sorrel	Wild garlic	
Shepherdspurse	Wild geranium	
Slender plantain	Wild lettuce	
Smallflower	Wild marigold	
galinsoga	Wild mustard	
Smooth dock	Wild onion	
Smooth pigweed	Wild parsnip	
Sorrel	Wild radish	
Sowthistle	Wild rape	
Spanishneedles	Wild strawberry	
Speedwell	Wild sweet potato	
Spiny amaranth	Wild vetch	
Spiny cocklebur	Woodsorrel	
Spiny sowthistle	Woolly croton	
Spotted catsear	Woolly	
Spotted spurge	morningglory	
Spurweed	Woolly plantain	
Stinging nettle	Wormseed	
Strawberry, India	Yarrow	
mock	Yellow rocket	
Tall nettle	Yellowflower	
Tall vervain	pepperweed	
	and other broadleaf weeds	

**IMPORTANT: Directions for Storage and Disposal**

**Storage Conditions:** Store in original container out of the reach of children, preferably in a locked storage cabinet. Protect from freezing.

- Disposal:**
- Do not reuse empty container.
  - Rinse the empty product container thoroughly and disperse the rinse water on weeds as part of application.
  - Place empty container in trash.

**Important:** It is best to use all of the product in accordance with label directions. If it is necessary to dispose of unused product, please follow any applicable state and local guidelines.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS  
AND DOMESTIC ANIMALS**

**CAUTION**

- Harmful if swallowed.
- Causes moderate eye irritation.
- Avoid contact with eyes, skin, or clothing.
- When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

After using this product:

- Rinse gloves before removing.
- Wash hands thoroughly with soap and water.
- If product gets on clothing, remove clothing as soon as possible and launder separately before reuse.

FIRST AID	
IN CASE OF ACCIDENTAL...	IMMEDIATELY...
Swallowing	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything to an unconscious person.</li> </ul>
Contact with eyes	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
Contact with skin	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
Inhalation	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<p><b>NOTE:</b> When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 877-229-3763 for medical emergency information. Active Ingredients: 7.59% 2,4-D, dimethylamine salt [CAS # 2008-39-1], 3.66% Mecoprop, dimethylamine salt [CAS # 32351-70-5], and 0.84% Dicamba, dimethylamine salt [CAS # 2300-66-5].</p>	

**ENVIRONMENTAL HAZARDS**

- This product is toxic to aquatic invertebrates. Do not apply directly to lakes, streams, rivers or ponds.
- Drift or runoff may harm aquatic invertebrates and plants.
- Do not contaminate water when disposing of equipment washwaters.
- Do not apply in windy conditions.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D and MCPP pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

6/6

**NOTICE:** Research and testing have determined that the "Directions For Use" are appropriate for the proper use of LAWN HERBICIDE JD CONCENTRATE under expected conditions. The Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

**Money Back Guarantee:** If you are not satisfied with this product, we will gladly refund your original purchase price.

**B - 9443a-2     8/30/00**

Bayer-Pursell, LLC  
1500 Urban Center Drive  
Birmingham, AL 35242