

Lawn Herbicide JD Concentrate

This label contains separate label statements for **Outdoor Residential Pump Sprayer/ Hose-end Sprayer Concentrate** use product and **Outdoor Residential Ready-To-Spray** use product.

- Effective Weed Protection
- Effective Broadleaf Weed Protection
- Effectively Protects Lawns from Broadleaf Weeds
- Kills All Major Broadleaf Weeds
- Kills Weeds - Won't Harm Lawns
- Kills to the Root
- Protects Against All Major Broadleaf Weeds
- Tough Weed Protection
- Weed Protection System
- Systemic
- 1 Quart treats up to 16,000 square feet
- Convenient MEASURE-UP™ Cup
- Kills Broadleaf weeds in St. Augustine & other Southern Lawns
- Kills Over 200 Broadleaf Weeds
- Won't Harm Your Lawn
- Rainproof in hours

ACTIVE INGREDIENTS:


2,4-D, dimethylamine salt 7.59 %
Mecoprop, dimethylamine salt 3.66 %
Dicamba, dimethylamine salt 0.84 %
OTHER INGREDIENTS: 87.91 %

Net Contents: 16 fl. oz. (473 mL) or 24 fl. oz. (709 mL) or
32 fl. oz. (946 mL) or 1 Gal. (3.78 L)

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel Booklet for Additional Ingredients and
Precautionary Statements



(FRONT PANEL)

Lawn Herbicide JD Concentrate

(Pump Sprayer/Hose-end Sprayer label)

- Effective Weed Protection
- Effective Broadleaf Weed Protection
- Effectively Protects Lawns from Broadleaf Weeds
- Kills All Major Broadleaf Weeds
- Kills Weeds - Won't Harm Lawns
- Kills to the Root
- Protects Against All Major Broadleaf Weeds
- Tough Weed Protection
- Weed Protection System
- Systemic
- 1 Quart treats up to 16,000 square feet
- Convenient MEASURE-UP™ Cup
- Kills Broadleaf weeds in St. Augustine & other Southern Lawns
- Kills Over 200 Broadleaf Weeds
- Won't Harm Your Lawn
- Rainproof in hours

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	7.59 %
Mecoprop, dimethylamine salt	3.66 %
Dicamba, dimethylamine salt	0.84 %
OTHER INGREDIENTS:	87.91 %

Net Contents: 16 fl. oz. (473 mL) or 24 fl. oz. (709 mL) or
32 fl. oz. (946 mL) or 1 Gal. (3.78 L)

KEEP OUT OF REACH OF CHILDREN



CAUTION

See Back Panel Booklet for Additional Ingredients and
Precautionary Statements

(BACK PANEL)

Lawn Herbicide JD Concentrate kills over 200 types of broadleaf weeds in St. Augustine and other Southern lawns. This special formula kills Dollarweed, Spurge, Oxalis and Dandelion without harming your lawn. Controlling weeds in your lawn has never been easier.

READ THE LABEL FIRST!

QUICK FACTS	
<ul style="list-style-type: none"> • Kills over 200 broadleaf weeds • Won't harm your lawn • Treats up to 16,000 sq. ft. 	
KILLS	Over 200 broadleaf weeds such as Dollarweed, Spurge and Oxalis, Dandelion, Clover, Chickweed and other listed weeds
WHERE TO USE	On Lawns (Do not use on Carpetgrass, Dichondra or Floratam variety of St. Augustinegrass.)
WHEN TO USE	When weeds are actively growing
HOW MUCH TO USE	<p>Northern Lawns: Fescue, Kentucky Bluegrass, and Perennial Ryegrasses and</p> <p>Hardy Southern Lawns: Bermuda, Buffalo and Zoysia grasses 2 fl. oz. per 500 sq. ft.</p> <p>Southern lawns: Bahia, Centipede and St. Augustinegrass grasses 1 fl. oz. per 500 sq. ft.</p>
 	For questions or comments, call toll-free 1-877 BayerAL or visit us on-line at www.BayerAdvanced.com

FOR OUTDOOR RESIDENTIAL USE ONLY

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	7.59 %
Mecoprop. dimethylamine salt	3.66 %
Dicamba. dimethylamine salt	0.84 %
OTHER INGREDIENTS*:	87.91 %
	100.00 %

*Other ingredients include: Water and surfactants (to keep the active ingredients dispersed in water).

THIS PRODUCT CONTAINS:

- 6.30% 2,4-D acid equivalent
- 3.02% Mecoprop acid equivalent
- 0.69% Dicamba acid equivalent
- Isomer specific by AOAC Method

EPA Reg. No. 72155-4

EPA Est. No. _-AA-_

DIRECTIONS FOR USE	
It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. For best results, read and follow all label directions.	
BEFORE YOU USE Read and follow directions when using.	<ul style="list-style-type: none"> Do not apply near lakes, streams, rivers or ponds. Spray when most of the day is expected to remain cooler than 90° F. Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Apply when wind is calm. The maximum number of broadcast applications per treatment site is two per year.
HOW TO USE Shake Well Before Use	<ul style="list-style-type: none"> Determine amount of product to use based on grass type and size of area to be treated. Prepare product for use according to How to Apply. Apply to lawn.
HOW MUCH TO APPLY	<p>Northern Lawns:</p> <ul style="list-style-type: none"> Fescue, Kentucky Bluegrass, and Perennial Rye 2 fl. oz. (4 Tbs.) per gallon of water <p>Hardy Southern Lawns:</p> <ul style="list-style-type: none"> Bermuda, Buffalo and Zoysia grasses 2 fl. oz. (4 Tbs.) per gallon of water <p>Southern Lawns:</p> <ul style="list-style-type: none"> Bahia, Centipede, and St. Augustinegrass 1 fl. oz. (2 Tbs.) per gallon of water <p>For lawns with mixed grass types, use the lowest rate. Each gallon of solution treats 500 sq. ft. TIP: Measure length and width of area to be treated in feet. Multiply these 2 numbers together. This is the area to be treated in square feet (sq. ft.).</p>
HOW TO APPLY	<p>Pump Style Sprayers</p> <ol style="list-style-type: none"> Add the appropriate amounts of water and concentrate to the sprayer tank. Close sprayer, shake well and pressurize. Adjust nozzle to a coarse spray pattern. Large Areas – Apply evenly over area. Individual Weeds – Hold tip of wand about 12 inches from weed, aim at center of weed and spray. Occasionally re-pressurize the sprayer if needed to maintain a good spray pattern.

Continued in next column

Directions For Use continued	
HOW TO APPLY continued	<p>Dial Style Hose-End Sprayers</p> <ol style="list-style-type: none"> Fill sprayer jar with enough concentrate to cover area to be treated. Close sprayer and set dial. TIP: The numbers on Dial-Style Hose-End sprayers are ounces of concentrate per gallon of spray. Attach sprayer to hose and apply. To avoid contact with the treated area, begin by spraying area farthest from faucet walking backwards as you spray. <p>Concentrate Plus Water Hose-End Sprayer</p> <ol style="list-style-type: none"> Pour concentrate into sprayer jar to the required fluid ounce level. Add the appropriate amount of water to the gallons level. Replace sprayer top and shake well before applying. To avoid contact with the treated area, begin by spraying area farthest from faucet walking backwards as you spray.
FOR BEST RESULTS	<ul style="list-style-type: none"> Apply when weeds are young and actively growing. If the lawn is dry, water 24 hours before using this product. Rain or watering 6 to 8 hours after application will not wash away effectiveness. Apply 1 to 3 days after mowing. Certain hard-to-kill weeds may need to be spot-treated in 14 days. Do not apply to new lawns until the grass has reached a height of at least 2 inches. Bare spots may be seeded 3 to 4 weeks after application.
HOW LAWN HERBICIDE JD CONCENTRATE WORKS	<p>LAWN HERBICIDE JD CONCENTRATE is a selective herbicide - it selectively kills weeds in the lawn but won't harm the lawn itself. It is a blend of proven weed killers designed to get broadleaf weeds such as Dollarweed, Spurge and Oxalis.</p>



Do not allow people (other than applicator) or pets on treated area during application.
People and pets may re-enter the treated area after it has dried.

5/11

CONTROLS [KILLS]		
Annual yellow sweetclover	Daisy, oxeye	
Aster	Dandelion	
Austrian fieldcress	Dichondra	
Bedstraw	Dogbane	
Beggarticks	Dogfennel	
Betony, Florida	Dollarweed	
Bindweed, field	Elderberry	
Bird vetch	False dandelion	
Bitter wintercress	False flax	
Bittercress, hairy	False sunflower	
Bitterweed	Fiddleneck	
Black-eyed Susan	Florida pusley	
Black medic	Frenchweed	
Black mustard	Galinsoga	
Blackseed plantain	Goathead	
Blessed thistle	Goldenrod	
Bloodflower	Ground ivy	
milkweed	Gumweed	
Blue lettuce	Hairy fleabane	
Blue vervain	Hawkweed	
Bracted plantain	Healall	
Brassbuttons	Heartleaf drymary	
Bristly oxtongue	Heath aster	
Broadleaf dock	Hedge bindweed	
Broadleaf plantain	Hedge mustard	
Broomweed	Hemp	
Buckhorn	Henbit	
Buckhorn plantain	Hoary cress	
Bulbous buttercup	Hoary plantain	
Bull thistle	Hoary vervain	
Bullnettle	Horsenettle	
Burclover	Jimsonweed	
Burdock	Knawel	
Burning nettle	Knotweed	
Burweed	Kochia	
Buttercup	Lambsquarters	
Buttonweed	Lespedeza	
Canada thistle	Mallow	
Carolina geranium	Matchweed	
Carpetweed	Mexicanweed	
Catchweed	Milk vetch	
bedstraw	Morningglory	
Catnip	Mouseear	
Catsear	hawkweed	
Chickweed, common	Mugwort	
Chickweed, mouseear	Musk thistle	
Chicory	Mustard	
Cinquefoil	Narrowleaf plantain	
Clover, crimson	Narrowleaf vetch	
Clover, hop	Nettle	
Clover, red	Orange hawkweed	
Clover, strawberry	Oriental cocklebur	
Clover, sweet	Oxalis	
Clover, white	Parsley-piert	
Cockle	Parsnip	
Cocklebur	Pearlwort	
Common mullein	Pennycress	
Creeping Charlie	Pennywort	
Creeping Jenny	Pepperweed	
Cudweed	Pigweed	
Curly dock	Pineywoods	
Daisy, English	bedstraw	
Daisy, fleabane	Plains coreopsis	
	(tickseed)	
	Plantain	
	Poison ivy	

Continued in next column

CONTROLS LIST continued		
CONTROLS [KILLS]		
Poison oak	Tansy mustard	
Pokeweed	Tanweed	
Poorjoe	Thistle	
Prairie sunflower	Trailing crownvetch	
Prickly lettuce	Tumble mustard	
Prickly sida	Tumble pigweed	
Prostrate knotweed	Velvetleaf	
Prostrate pigweed	Venice mallow	
Prostrate spurge	Virginia	
Prostrate vervain	buttonweed	
Puncturevine	Virginia creeper	
Purslane, common	Virginia	
Ragweed	pepperweed	
Red sorrel	Water pennywort	
Redroot pigweed	Wavyleaf bullthistle	
Redstem filaree	Western clematis	
Rough cinquefoil	Western salsify	
Rough fleabane	White mustard	
Russian pigweed	Wild aster	
Russian thistle	Wild buckwheat	
Scarlet pimpernel	Wild carrot	
Scotch thistle	Wild four-o'clock	
Sheep sorrel	Wild garlic	
Shepherdspurse	Wild geranium	
Slender plantain	Wild lettuce	
Smallflower	Wild marigold	
galinsoga	Wild mustard	
Smooth dock	Wild onion	
Smooth pigweed	Wild parsnip	
Sorrel	Wild radish	
Sowthistle	Wild rape	
Spanishneedles	Wild strawberry	
Speedwell	Wild sweet potato	
Spiny amaranth	Wild vetch	
Spiny cocklebur	Woodsorrel	
Spiny sowthistle	Wooly croton	
Spotted catsear	Woolly	
Spotted spurge	morningglory	
Spurweed	Woolly plantain	
Stinging nettle	Wormseed	
Strawberry, India	Yarrow	
mock	Yellow rocket	
Tall nettle	Yellowflower	
Tall vervain	pepperweed	
	and other broadleaf weeds	

IMPORTANT: Directions for Storage and Disposal

Storage Conditions:

Store in original container out of the reach of children, preferably in a locked storage cabinet. Protect from freezing.

Disposal:



- Do not reuse empty container.
- Rinse the empty product container thoroughly and disperse the rinse water on weeds as part of application.
- Place empty container in trash.

Important: It is best to use all of the product in accordance with label directions. If it is necessary to dispose of unused product, please follow any applicable state and local guidelines.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

- Harmful if swallowed.
- Causes moderate eye irritation.
- Avoid contact with eyes, skin or clothing.
- When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves.

After using this product:

- Rinse gloves before removing.
- Remove clothing and launder separately before reuse.
- Wash hands thoroughly with soap and water.
- If product gets on clothing, remove clothing as soon as possible and shower.

FIRST AID +

IN CASE OF ACCIDENTAL...	IMMEDIATELY...
Swallowing	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything to an unconscious person.
Contact with eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
Contact with skin	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
Inhalation	<ul style="list-style-type: none"> • Move person to fresh air. • If not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.

NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 877-229-3763 for medical emergency information. Active Ingredients: 7.59% 2,4-D, dimethylamine salt [CAS # 2008-39-1], 3.66% Mecoprop, dimethylamine salt [CAS # 32351-70-5], and 0.84% Dicamba, dimethylamine salt [CAS # 2300-66-5].

ENVIRONMENTAL HAZARDS

- This product is toxic to aquatic invertebrates. Do not apply directly to lakes, streams, rivers or ponds.
- Drift or runoff may harm aquatic invertebrates and non-target plants.
- Do not contaminate water when disposing of equipment washwaters. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D and MCPP pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

NOTICE: Research and testing have determined that the "Directions For Use" are appropriate for the proper use of LAWN HERBICIDE JD CONCENTRATE under expected conditions. The Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.


D-9558-1 062101

Bayer Advanced LLC
1500 Urban Center Drive
Birmingham, AL 35242

Lawn Herbicide JD Concentrate

- Effective Weed Protection
- Effective Broadleaf Weed Protection
- Effectively Protects Lawns from Broadleaf Weeds
- Kills All Major Broadleaf Weeds
- Kills Weeds - Won't Harm Lawns
- Kills to the Root
- Protects Against All Major Broadleaf Weeds
- Tough Weed Protection
- Weed Protection System
- Systemic
- 1 Quart treats up to 16,000 square feet
- Convenient MEASURE-UP™ Cup
- Kills Broadleaf weeds in St. Augustine & other Southern Lawns
- Kills Over 200 Broadleaf Weeds
- Won't Harm Your Lawn
- Rainproof in hours

(Ready - To - Spray label)



ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	7.59 %
Mecoprop, dimethylamine salt	3.66 %
Dicamba, dimethylamine salt	0.84 %

OTHER INGREDIENTS: 87.91 %

Net Contents: 16 fl. oz. (473 mL) or 24 fl. oz. (709 mL) or
32 fl. oz. (946 mL) or 1 Gal. (3.78 L)

KEEP OUT OF REACH OF CHILDREN



CAUTION

See Back Panel Booklet for Additional Ingredients and
Precautionary Statements

(BACK PANEL)

Lawn Herbicide JD Concentrate kills over 200 types of broadleaf weeds in St. Augustine and other Southern lawns. This special formula kills Dollarweed, Spurge, Oxalis and Dandelion without harming your lawn. Controlling weeds in your lawn has never been easier.

READ THE LABEL FIRST!

QUICK FACTS	
<ul style="list-style-type: none"> • Kills over 200 broadleaf weeds • Won't harm your lawn • Treats up to 16,000 sq. ft. 	
KILLS	Over 200 broadleaf weeds such as Dollarweed, Spurge and Oxalis, Dandelion, Clover, Chickweed and other listed weeds
WHERE TO USE	On Lawns (Do not use on Carpetgrass, Dichondra or Floratam variety of St. Augustinegrass.)
WHEN TO USE	When weeds are actively growing
HOW MUCH TO USE	<p>Northern Lawns: Fescue, Kentucky Bluegrass, and Perennial Ryegrasses and Hardy Southern Lawns: Bermuda, Buffalo and Zoysia grasses One qt treats up to 8,000 sq. ft.</p> <p>Southern lawns: Bahia, Centipede and St. Augustinegrass grasses One qt treats up to 16,000 sq. ft.</p> <p>No measuring needed. Dilutes automatically.</p>
 	For questions or comments, call toll-free 1-877-BayerAL or visit us on-line at www.BayerAdvanced.com

FOR OUTDOOR RESIDENTIAL USE ONLY**ACTIVE INGREDIENTS:**

2,4-D, dimethylamine salt	7.59 %
Mecoprop, dimethylamine salt	3.66 %
Dicamba, dimethylamine salt	0.84 %
OTHER INGREDIENTS*:	87.91 %
	100.00 %

*Other ingredients include: Water and surfactants (to keep the active ingredients dispersed in water).

THIS PRODUCT CONTAINS:

6.30% 2,4-D acid equivalent
3.02% Mecoprop acid equivalent
0.69% Dicamba acid equivalent
Isomer specific by AOAC Method

EPA Reg. No. 72155-4


EPA Est. No. _-AA-_

9/11

DIRECTIONS FOR USE	
It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. For best results, read and follow all label directions.	
BEFORE YOU USE Read and follow directions when using.	<ul style="list-style-type: none"> Do not apply near lakes, streams, rivers or ponds. Spray when most of the day is expected to remain cooler than 90° F. Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Apply when wind is calm. The maximum number of broadcast applications per treatment site is two per year.
HOW TO USE Shake Well Before Use	<ul style="list-style-type: none"> Determine amount of product to use based on grass type and size of area to be treated. See How Much To Use section of Quick Facts Box. Prepare and apply product according to the following section, How To Attach and Spray.
HOW TO ATTACH & SPRAY	<p>This product is ready to spray. Just attach the bottle to your garden hose according to the following directions. Water automatically mixes with the concentrate as you spray.</p> <ol style="list-style-type: none"> 1. Connect your garden hose to the bottle. Be sure the blue water control valve on top of the spray nozzle is in the OFF position. 2. Turn on water at faucet. Twist the small, star-shaped, blue product-control button to align the flat portion with the lock tab. Push the button in to open. 3. To begin spraying, point nozzle toward the area to be treated and turn the blue water control valve to the ON position. Tab will be horizontal. 4. To stop spraying, turn the blue water control valve on top of the spray nozzle to OFF. To close, push in the small round button opposite the star-shaped product-control button. Turn off faucet. 5. The unused portion can be saved and used at a later date.

continued in next column


TIPS FOR APPLYING	
HOW TO APPLY	<ol style="list-style-type: none"> 1. Extend hose to the farthest area to be treated and work back toward the faucet to avoid walking over wet, treated grass. 2. Walk at a steady pace while spraying, using an even sweeping motion, slightly overlapping the treated areas.
FOR BEST RESULTS	<ul style="list-style-type: none"> Apply when weeds are young and actively growing. If the lawn is dry, water 24 hours before using this product. Rain or watering 6 to 8 hours after application will not wash away effectiveness. Apply 1 to 3 days after mowing. Certain hard-to-kill weeds may need to be spot-treated in 14 days. Do not apply to new lawns until the grass has reached a height of at least 2 inches. Bare spots may be seeded 3 to 4 weeks after application.
HOW LAWN HERBICIDE JD CONCENTRATE WORKS	<p>LAWN HERBICIDE JD CONCENTRATE is a selective herbicide - it selectively kills weeds in the lawn but won't harm the lawn itself. It is a blend of proven weed killers designed to get broadleaf weeds such as Dollarweed, Spurge and Oxalis.</p>

	<p>Do not allow people (other than applicator) or pets on treated area during application. People and pets may re-enter the treated area after it has dried.</p>
---	--

CONTROLS [KILLS]		
Annual yellow sweetclover	Daisy, oxeye	
Aster	Dandelion	
Austrian fieldcress	Dichondra	
Bedstraw	Dogbane	
Beggarticks	Dogfennel	
Betony, Florida	Dollarweed	
Bindweed, field	Elderberry	
Bird vetch	False dandelion	
Bitter wintercress	False flax	
Bittercress, hairy	False sunflower	
Bitterweed	Fiddleneck	
Black-eyed Susan	Florida pusley	
Black medic	Frenchweed	
Black mustard	Galinsoga	
Blackseed plantain	Goathead	
Blessed thistle	Goldenrod	
Bloodflower	Ground ivy	
milkweed	Gumweed	
Blue lettuce	Hairy fleabane	
Blue vervain	Hawkweed	
Bracted plantain	Healall	
Brassbuttons	Heartleaf drymary	
Bristly ox tongue	Heath aster	
Broadleaf dock	Hedge bindweed	
Broadleaf plantain	Hedge mustard	
Broomweed	Hemp	
Buckhorn	Henbit	
Buckhorn plantain	Hoary cress	
Bulbous buttercup	Hoary plantain	
Bull thistle	Hoary vervain	
Bullnettle	Horsenettle	
Burclover	Jimsonweed	
Burdock	Knawel	
Burning nettle	Knotweed	
Burweed	Kochia	
Buttercup	Lambsquarters	
Buttonweed	Lespedeza	
Canada thistle	Mallow	
Carolina geranium	Matchweed	
Carpetweed	Mexicanweed	
Catchweed	Milk vetch	
bedstraw	Morningglory	
Catnip	Mouseear	
Catsear	hawkweed	
Chickweed, common	Mugwort	
Chickweed, mouseear	Musk thistle	
Chicory	Mustard	
Cinquefoil	Narrowleaf plantain	
Clover, crimson	Narrowleaf vetch	
Clover, hop	Nettle	
Clover, red	Orange hawkweed	
Clover, strawberry	Oriental cocklebur	
Clover, sweet	Oxalis	
Clover, white	Parsley-piert	
Cockle	Parsnip	
Cocklebur	Pearlwort	
Common mullein	Pennycress	
Creeping Charlie	Pennywort	
Creeping Jenny	Pepperweed	
Cudweed	Pigweed	
Curly dock	Pineywoods	
Daisy, English	bedstraw	
Daisy, fleabane	Plains coreopsis	
	(tickseed)	
	Plantain	
	Poison ivy	

Continued in next column

CONTROLS LIST continued		
CONTROLS [KILLS]		
Poison oak	Tansy mustard	
Pokeweed	Tanweed	
Poorjoe	Thistle	
Prairie sunflower	Trailing crownvetch	
Prickly lettuce	Tumble mustard	
Prickly sida	Tumble pigweed	
Prostrate knotweed	Velvetleaf	
Prostrate pigweed	Venice mallow	
Prostrate spurge	Virginia buttonweed	
Prostrate vervain	Virginia creeper	
Puncturevine	Virginia pepperweed	
Purslane, common	Water pennywort	
Ragweed	Wavyleaf bullthistle	
Red sorrel	Western clematis	
Redroot pigweed	Western salsify	
Redstem filaree	White mustard	
Rough cinquefoil	Wild aster	
Rough fleabane	Wild buckwheat	
Russian pigweed	Wild carrot	
Russian thistle	Wild four-o'clock	
Scarlet pimpernel	Wild garlic	
Scotch thistle	Wild geranium	
Sheep sorrel	Wild lettuce	
Shepherdspurse	Wild marigold	
Slender plantain	Wild mustard	
Smallflower	Wild onion	
galinsoga	Wild parsnip	
Smooth dock	Wild radish	
Smooth pigweed	Wild rape	
Sorrel	Wild strawberry	
Sowthistle	Wild sweet potato	
Spanishneedles	Wild vetch	
Speedwell	Woodsorrel	
Spiny amaranth	Wooly croton	
Spiny cocklebur	Woolly morningglory	
Spiny sowthistle	Woolly plantain	
Spotted catsear	Wormseed	
Spotted spurge	Yarrow	
Spurweed	Yellow rocket	
Stinging nettle	Yellowflower	
Strawberry, India	pepperweed	
mock	and other broadleaf	
Tall nettle	weeds	
Tall vervain		

IMPORTANT: Directions for Storage and Disposal	
Storage Conditions:	Store in original container out of the reach of children, preferably in a locked storage cabinet. Protect from freezing.
Disposal:	<div>  <ul style="list-style-type: none"> Do not reuse empty container. Securely wrap original container in several layers of newspaper and discard in trash Place empty container in trash. </div>
Important: It is best to use all of the product in accordance with label directions. If it is necessary to dispose of unused product, please follow any applicable state and local guidelines.	

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

- Harmful if swallowed.
- Causes moderate eye irritation.
- Avoid contact with eyes, skin or clothing.
- When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves.

After using this product:

- Rinse gloves before removing.
- Remove clothing and launder separately before reuse.
- Wash hands thoroughly with soap and water.
- If product gets on clothing, remove clothing as soon as possible and shower.

ENVIRONMENTAL HAZARDS

- This product is toxic to aquatic invertebrates. Do not apply directly to lakes, streams, rivers or ponds.
- Drift or runoff may harm aquatic invertebrates and non-target plants.
- Do not contaminate water when disposing of equipment washwaters.

NOTICE: Research and testing have determined that the "Directions For Use" are appropriate for the proper use of LAWN HERBICIDE JD CONCENTRATE under expected conditions. The Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.

D-9558-2 062101

Bayer Advanced LLC
1500 Urban Center Drive
Birmingham, AL 35242

FIRST AID



**IN CASE OF
ACCIDENTAL...**

IMMEDIATELY...

Swallowing

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything to an unconscious person.

**Contact with
eyes**

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

**Contact with
skin**

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 to 20 minutes.
- Call a poison control center or doctor for treatment advice.

Inhalation

- Move person to fresh air.
- If not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 877-229-3763 for medical emergency information. Active Ingredients: 7.59% 2,4-D, dimethylamine salt [CAS # 2008-39-1], 3.66% Mecoprop, dimethylamine salt [CAS # 32351-70-5], and 0.84% Dicamba, dimethylamine salt [CAS # 2300-66-5].