Lawn Herbicide JD Concentrate

This label contains separate label statements for Outdoor Residential Pump Sprayer/
Hose-end Sprayer Concentrate use product and Outdoor Residential Ready-To-Spray use product.

- Effective Weed Protection
- Effective Broadleaf Weed Protection
- · Effectively Protects Lawns from Broadleaf Weeds
- Kills All Major Broadleaf Weeds
- · Kills Weeds Won't Harm Lawns
- Kills to the Root
- · Protects Against All Major Broadleaf Weeds
- Tough Weed Protection
- Weed Protection System
- Systemic
- 1 Quart treats up to 16,000 square feet
- Convenient MEASURE-UP™ Cup
- Kills Broadleaf weeds in St. Augustine & other Southern Lawns
- Kills Over 200 Eroadleaf Weeds
- Won't Harm You: Lawn
- Rainproof in hours

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	7.59 %
Mecoprop-p, dimethylamine salt	1.83 %
Dicamba, dimethylamine salt	0.84 %
OTHER INGREDIENTS:	39.74 %

Net Contents: 16 fl. oz. (473 mL) or 24 fl. oz. (709 mL) or 32 fl. oz. (946 mL) or 1 Gal. (3.78 L)

CAUTION

See Back Panel Booklet for Additional Ingredients and Precautionary Statements

ACCEPTED

APR 10 2002

Under the Federal Insecticide. Fungicide, and Redenticide Act, as amended, for the pecticide registered under EPA Reg. Phys. 72155-4

Lawn Herbicide JD Concentrate

· Effective Weed Protection

(Pump Sprayer/Hose-end Sprayer label)

- · Effective Broadleaf Weed Protection
- · Effectively Protects Lawns from Broadleaf Weeds
- · Kills All Major Broadleaf Weeds
- · Kills Weeds Won't Harm Lawns
- · Kills to the Root
- Protects Against All Major Broadleaf Weeds
- · Tough Weed Protection
- · Weed Protection System
- Systemic
- · 1 Quart treats up to 16,000 square feet
- Convenient MEASURE-UP™ Cup
- Kills Broadleaf weeds in St. Augustine & other Southern Lawns
- · Kills Over 200 Broadleaf Weeds
- · Won't Harm Your Lawn
- · Rainproof in hours

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	7.59	%
Mecoprop-p, dimethylamine salt		
Dicamba, dimethylamine salt		
OTHER INGREDIENTS:		

Net Contents: 16 fl. oz. (473 mL) or 24 fl. oz. (709 mL) or 32 fl. oz. (946 mL) or 1 Gal. (3.78 L)

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel Booklet for Additional Ingredients and Precautionary Statements

(BACK PANEL)

Lawn Herbicide JD Concentrate kills over 200 types of broadleaf weeds in St. Augustine and other Southern lawns. This special formula kills Dollarweed, Spurge, Oxalis and Dandelion without harming your lawn. Controlling weeds in your lawn has never been easier.

READ THE LABEL FIRST!

QUICK FACTS		
Kills over 200 broadleaf weeds		
Won't harm your lawn		
 Treats up to 16,0 	00 sq. ft.	
KILLS	Over 200 broadleaf weeds such as Dollarweed, Spurge and Oxalis, Dandelion, Clover, Chickweed and other listed weeds	
WHERE TO USE	On Lawns	
	(Do not use on Carpetgrass, Dichondra or Floratam variety of St. Augustinegrass.)	
WHEN TO USE	When weeds are actively growing	
HOW MUCH TO USE	Northern Lawns: Fescue, Kentucky Bluegrass, and Perennial Ryegrasses and	
	Hardy Southern Lawns: Bermuda, Buffalo and Zoysia grasses 2 fl. oz. per 500 sq. ft. Southern lawns: Bahla, Centipede and St. Augustinegrass grasses 1 fl. oz. per 500 sq. ft.	
7	For questions or comments, call toil-free 1-877 BayerAL or visit us on-line at www.BayerAdvanced.com	

FOR OUTDOOR RESIDENTIAL USE ONLY

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	7.59	%
Mecoprop-p, dimethylamine salt	1.83	%
Dicamba, dimethylamine salt	0.84	%
OTHER INGREDIENTS*:	39,74	%

100.00 % *Other ingredients include: Water and surfactants (to keep

the active ingredients dispersed in water).

THIS PRODUCT CONTAINS: 6.30% 2,4-D acid equivalent 1.51% Mecoprop-p acid equivalent 0.69% Dicamba acid equivalent Isomer specific by AOAC Method

EPA Reg. No. 72155-4

EPA Est. No. _-AA-_

DIRECTIONS FO	
inconsistent with its I	-
	and follow all label directions.
BEFORE YOU USE	Do not apply near lakes, streams, rivers or ponds.
Read and follow directions when using.	 Spray when most of the day is expected to remain cooler than 90° F. Avoid drift of spray mist onto
	vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Apply when wind is calm.
	The maximum number of broadcast applications per treatment site is two per year.
HOW TO USE Shake Well Before Use	Determine amount of product to use based on grass type and size of area to be treated.
	Prepare product for use according to How to Apply.
_ ·	Apply to lawn.
HOW MUCH TO	Northern Lawns:
APPLI	Fescue, Kentucky Bluegrass, and Perennial Rye
	2 fl. oz. (4 Tbs.) per gallon of water
	Hardy Southern Lawns:
	Bermuda, Buffalo and Zoysia grasses
	2 fl. oz. (4 Tbs.) per gallon of water
	Southern Lawns:
	Bahia, Centipede, and St. Augustinegrass
	1 fl. oz. (2 Tbs.) per gallon of water
:	For lawns with mixed grass types, use the lowest rate.
	Each gallon of solution treats 500 sq. ft.
	TIP: Measure length and width of area to be treated in feet. Multiply these 2 numbers together. This is the area to be treated in square feet (sq. ft.).
HOW TO APPLY	Pump Style Sprayers
	Add the appropriate amounts of water and concentrate to the sprayer tank.
	Close sprayer, shake well and pressurize.
	Adjust nozzle to a coarse spray pattern.
	Large Areas – Apply evenly over area. Individual Weeds – Hold tip of wand about 12 inches from weed, aim at center of weed and spray.
	 Occasionally re-pressurize the sprayer if needed to maintain a good spray pattern.

Continued in next column

Directions For Use	continued
HOW TO APPLY	Dial Style Hose-End Sprayers
continued	Fill sprayer jar with enough concentrate to cover area to be treated.
	Close sprayer and set dial.
	TIP: The numbers on Dial-Style Hose- End sprayers are ounces of concentrate per gallon of spray.
	Attach sprayer to hose and apply.
;	To avoid contact with the treated area, begin by spraying area farthest from faucet walking backwards as you spray.
İ	Concentrate Plus Water Hose-End Sprayer
	Pour concentrate into sprayer jar to the required fluid ounce level.
	Add the appropriate amount of water to the gallons level.
1	Replace sprayer top and shake well before applying.
	 To avoid contact with the treated area, begin by spraying area farthest from faucet walking backwards as you spray.
FOR BEST RESULTS	 Apply when weeds are young and actively growing.
	If the lawn is dry, water 24 hours before using this product.
	 Rain or watering 6 to 8 hours after application will not wash away effectiveness.
	Apply 1 to 3 days after mowing.
	 Certain hard-to-kill weeds may need to be spot-treated in 14 days.
	 Do not apply to new lawns until the grass has reached a height of at least 2 inches.
	Bare spots may be seeded 3 to 4 weeks after application.
HOW LAWN HERBICIDE JD CONCENTRATE WORKS	LAWN HERBICIDE JD CONCENTRATE is a selective herbicide - it selectively kills weeds in the lawn but won't harm the lawn itself. It is a blend of proven weed killers designed to get broadleaf weeds such as Dollarweed, Spurge and Oxalis.



Do not allow people (other than applicator) or pets on treated area during application.

People and pets may re-enter the treated area after it has dried.

CONTROLS		
[KILLS]	Annual yellow	Daisy, oxeye
[IVIECO]	sweetclover	Dandelion
	Aster	Dichondra
i	Austrian fieldcress	Dogbane
	Bedstraw	Dogfennel
	Beggarticks	Dollarweed
i	Betony, Florida	Elderberry
	Bindweed, field	False dandelion
	Bird vetch	False flax
	Bitter wintercress	False sunflower
	Bittercress, hairy	Fiddleneck
	Bitterweed	Florida pusley
	Black-eyed Susan	Frenchweed
	Black medic	Galinsoga
	Black mustard	Goathead
	Blackseed plantain	Goldenrod
	Blessed thistle	Ground ivy
	Bloodflower	Gumweed
	milkweed	Hairy fleabane
	Blue lettuce	Hawkweed
	Blue vervain	Healall
	Bracted plantain	Heartleaf drymary
	Brassbuttons	Heath aster
	Bristly oxtongue	Hedge bindweed
	Broadleaf dock	Hedge mustard
	Broadleaf plantain	Hemp
	Broomweed	Henbit
	Buckhorn	Hoary cress
	Buckhorn plantain	Hoary plantain
	Bulbous buttercup	Hoary vervain
	Bull thistle	Horsenettle
	Bullnettle	Jimsonweed
	Burclover	Knawel
	Burdock	Knotweed
	Burning nettle	Kochia
	Burweed	Lambsquarters
	Buttercup	Lespedeza
I	Buttonweed	Mallow
	Canada thistle	Matchweed
	Carolina geranium	Mexicanweed
	Carpetweed	Milk vetch
	Catchweed bedstraw	Morningglory
	Catnip	Mouseear hawkweed
l	Catsear	***************************************
	Chickweed.	Mugwort Musk thistle
	common	Mustard
Į.	Chickweed,	Narrowleaf plantain
	mouseear	Narrowleaf vetch
	Chicory	Nettle
:	Cinquefoil	Orange hawkweed
	Clover, crimson	Oriental cocklebur
	Clover, hop	Oxalis
	Clover, red	Parsley-piert
	Clover, strawberry	Parsnip
	Clover, sweet	Pearlwort
	Clover, white	Pennycress
	Cockle	Pennywort
	Cocklebur	Pepperweed
	Common mullein	Pigweed
	Creeping Charlie	Pineywoods
	Creeping Jenny	bedstraw
	Cudweed	Plains coreopsis
	Curly dock	(tickseed)
	Daisy, English	Plantain
	Daisy, fleabane	Poison ivy

Continued in next column

CONTROLS LIST	continued	
CONTROLS		
[KILLS]	Poison oak	Tansy mustard
	Pokeweed	Tanweed
	Poorjoe	Thistle
	Prairie sunflower	Trailing crownvetch
	Prickly lettuce	Tumble mustard
	Prickly sida	Tumble pigweed
	Prostrate knotweed	Velvetleaf
	Prostrate pigweed	Venice mallow
	Prostrate spurge	Virginia
	Prostrate vervain	buttonweed
	Puncturevine	Virginia creeper
	Purslane, common	Virginia
	Ragweed	pepperweed
	Red sorrel	Water pennywort
	Redroot pigweed	Wavyleaf builthistle
	Redstem filaree	Western clematis
	Rough cinquefoil	Western salsify
	Rough fleabane	White mustard
	Russian pigweed	Wild aster
	Russian thistle	Wild buckwheat
	Scarlet pimpernel	Wild carrot
	Scotch thistle	Wild four-o'clock
	Sheep sorrel	Wild garlic
	Shepherdspurse	Wild geranium
	Slender plantain	Wild lettuce
	Smallflower	Wild marigold
	galinsoga	Wild mustard
•	Smooth dock	Wild onion
	Smooth pigweed	Wild parsnip
	Sorrel	Wild radish
	Sowthistle	Wild rape
	Spanishneedles	Wild strawberry
	Speedwell	Wild sweet potato Wild vetch
	Spiny amaranth	Wild Vetch Woodsorrel
	Spiny cocklebur	
	Spiny sowthistle	Wooly croton Woolly
1	Spotted catsear	morningglory
	Spotted spurge	Woolly plantain
	Spurweed	Wormseed
1	Stinging nettle	Yarrow
	Strawberry, India mock	Yellow rocket
	Tail nettle	Yallowflower
	Tall vervain	pepperweed
	Tan Yarram	and other broadleaf
		veeds

IMPORTANT: Directions for Storage and Disposal Store in original container out of the Storage Conditions: reach of children, preferably in a locked

Disposal:



storage cabinet. Protect from freezing. · Do not reuse empty container.

- · Rinse the empty product container thoroughly and disperse the rinse water on weeds as part of application.
- · Place empty container in trash.

Important: It is best to use all of the product in accordance with label directions. If it is necessary to dispose of unused product, please follow any applicable state and local guidelines.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

- · Harmful if swallowed.
- · Causes moderate eve irritation.
- · Avoid contact with eyes, skin or clothing.
- When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves.

After using this product:

- · Rinse gloves before removing.
- · Remove clothing and launder separately before reuse.
- · Wash hands thoroughly with soap and water.
- If product gets on clothing, remove clothing as soon as possible and shower.

FIRST AID	•
IN CASE OF ACCIDENTAL	IMMEDIATELY
Swallowing	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swailow. Do not induce vomiting unless told to do so by a poison control center or doctor
	Do not give anything to an unconscious person.
Contact with eyes	 Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
Contact with skin	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
Inhalation	Move person to fresh air. If not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
NOTE: When calling	g poison control center, have this product

label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 877-229-3763 for medical emergency information. Active Ingredients: 7.59% 2,4-D, dimethylamine salt [CAS # 2008-39-1], 1.83% Mecoprop-p, dimethylamine salt [CAS # 66423-09-4], and 0.84% Dicamba, dimethylamine salt [CAS # 2300-66-5].

ENVIRONMENTAL HAZARDS

- This product is toxic to aquatic invertebrates. Do not apply directly to lakes, streams, rivers or ponds.
- Drift or runoff may harm aquatic invertebrates and non-target plants.
- Do not contaminate water when disposing of equipment washwaters. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D and MCPP pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

NOTICE: Research and testing have determined that the "Directions For Use" are appropriate for the proper use of LAWN HERBICIDE JD CONCENTRATE under expected conditions. The Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.

D-9630-1 101601

Bayer Advanced LLC 1500 Urban Center Drive Birmingham, AL 35242

Lawn Herbicide JD Concentrate

· Effective Weed Protection

(Ready - To - Spray label)

- · Effective Broadleaf Weed Protection
- · Effectively Protects Lawns from Broadleaf Weeds
- · Kills All Major Broadleaf Weeds
- Kills Weeds Won't Harm Lawns
- · Kills to the Root
- · Protects Against All Major Broadleaf Weeds
- Tough Weed Protection
- Weed Protection System
- Systemic
- · 1 Quart treats up to 16,000 square feet
- Convenient MEASURE-UP™ Cup
- Kills Broadleaf weeds in St. Augustine & other Southern Lawns
- Kills Over 200 Broadleaf Weeds
- · Won't Harm Your Lawn
- · Rainproof in hours

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	7.59 %
Mecoprop-p, dimethylamine salt	1.83 %
Dicamba, dimethylamine salt	0.84 %
OTHER INGREDIENTS:	39.74 %

Net Contents: 16 fl. oz. (473 mL) or 24 fl. oz. (709 mL) or 32 fl. oz. (946 mL) or 1 Gal. (3.78 L)

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel Booklet for Additional Ingredients and Precautionary Statements

(BACK PANEL)

Lawn Herbicide JD Concentrate kills over 200 types of broadleaf weeds in St. Augustine and other Southern lawns. This special formula kills Dollarweed, Spurge, Oxalis and Dandelion without harming your lawn. Controlling weeds in your lawn has never been easier.

READ THE LABEL FIRST!

QUICK FACTS		
1 -		
1	Kills over 200 broadleaf weeds	
1	Won't harm your lawn	
Treats up to 16,0		
KILLS	Over 200 broadleaf weeds such as Dollarweed, Spurge and Oxalis, Dandelion, Clover, Chickweed and other listed weeds	
WHERE TO USE	On Lawns	
:	(Do not use on Carpetgrass, Dichondra or Floratam variety of St. Augustinegrass.)	
WHEN TO USE	When weeds are actively growing	
HOW MUCH TO USE	Northern Lawns: Fescue, Kentucky Bluegrass, and Perennial Ryegrasses and Hardy Southern Lawns: Bermuda, Buffalo and Zoysia grasses One qt treats up to 8,000 sq. ft. Southern lawns: Bahia, Centipede and St. Augustinegrass grasses	
	One qt treats up to 16,000 sq. ft.	
	No measuring needed. Dilutes automatically.	
5	For questions or comments, call toll-free 1-877 BayerAL or visit us on-line at www.BayerAdvanced.com	

FOR OUTDOOR RESIDENTIAL USE ONLY

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt 7.59 %	,
Mecoprop-p, dimethylamine salt 1.83 %	,
Dicamba, dimethylamine salt 0.84 %	,
OTHER INGREDIENTS*: 89.74 %	!
100.00 %	

^{*}Other ingredients include: Water and surfactants (to keep the active ingredients dispersed in water).

THIS PRODUCT CONTAINS: 6.30% 2,4-D acid equivalent 1.51% Mecoprop-p acid equivalent 0.69% Dicamba acid equivalent Isomer specific by AOAC Method

EPA Reg. No. 72155-4

EPA Est. No._-AA-_

1	9/11
1	Thu

DIRECTIONS FOR	USE			
It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.				
For best results, read and follow all label directions.				
BEFORE YOU USE Read and follow directions when using.	 Do not apply near lakes, streams, rivers or ponds. Spray when most of the day is expected to remain cooler than 90° F. Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Apply when wind is calm. The maximum number of broadcast applications per treatment site is two per 			
	year.			
HOW TO USE Shake Well	Determine amount of product to use based on grass type and size of area to be treated. See How Much To Use section of Quick Facts Box.			
Before Use	Prepare and apply product according to the following section, How To Attach and Spray.			
HOW TO ATTACH & SPRAY	This product is ready to spray. Just attach the bottle to your garden hose according to the following directions. Water automatically mixes with the concentrate as you spray.			
	Connect your garden hose to the bottle. Be sure the blue water control valve on top of the spray nozzle is in the OFF position.			
	Turn on water at faucet. Twist the small, star-shaped, blue product-control button to align the flat portion with the lock tab. Push the button in to open.			
	To begin spraying, point nozzle toward the area to be treated and turn the blue water control valve to the ON position. Tab will be horizontal.			
	4. To stop spraying, turn the blue water control valve on top of the spray nozzle to OFF. To close, push in the small round button opposite the star-shaped product-control button. Turn off faucet.			
	The unused portion can be saved and used at a later date.			
	_			

continued in next column

TIPS FOR APPLYING		
HOW TO APPLY	Extend hose to the farthest area to be treated and work back toward the faucet to avoid walking over wet, treated grass.	
	Walk at a steady pace while spraying, using an even sweeping motion, slightly overlapping the treated areas.	
FOR BEST RESULTS	Apply when weeds are young and actively growing.	
	If the lawn is dry, water 24 hours before using this product.	
	 Rain or watering 6 to 8 hours after application will not wash away effectiveness. 	
	Apply 1 to 3 days after mowing.	
	Certain hard-to-kill weeds may need to be spot-treated in 14 days.	
	Do not apply to new lawns until the grass has reached a height of at least 2 inches.	
	Bare spots may be seeded 3 to 4 weeks after application.	
HOW LAWN HERBICIDE JD CONCENTRATE WORKS	LAWN HERBICIDE JD CONCENTRATE is a selective herbicide - it selectively kills weeds in the lawn but won't harm the lawn itself. It is a blend of proven weed killers designed to get broadleaf weeds such as Dolfarweed, Spurge and Oxalis.	



Do not allow people (other than applicator) or pets on treated area during application.

People and pets may re-enter the treated area after it has dried.

CONTROLS		
[KILLS]	Annual yellow	Daisy, oxeye
Ç	sweetclover	Dandelicn
	Aster	Dichondra
	Austrian fieldcress	Dogbane
	Bedstraw	Dogfennei
	Beggarticks	Dollarweed
	Betony, Florida	Elderberry
	Bindweed, field	False dandelion
	Bird vetch	False flax
	Bitter wintercress	False sunflower
	Bittercress, hairy	Fiddleneck
	Bitterweed	Florida pusley
	Black-eyed Susan	Frenchweed
	Black medic	Galinsoga
	Black mustard	Goathead
	Blackseed plantain Blessed thistle	Goldenrod
	Bloodflower	Ground ivy Gumweed
	milkweed	Hairy fleabane
	Blue lettuce	Hawkweed
	Blue vervain	Healall
	Bracted plantain	Heartleaf drymary
	Brassbuttons	Heath aster
	Bristly oxtongue	Hedge bindweed
,	Broadleaf dock	Hedge mustard
	Broadleaf plantain	Hemp
	Broomweed	Henbit
	Buckhorn	Hoary creos
	Buckhorn plantain	Hoary plantain
	Bulbous buttercup	Hoary vervain
	Bull thistle	Horsenettla
	Bulinettle	Jimsonweed
	Burclover	Knawel
	Burdock	Knotweed
	Burning nettle	Kochia
	Burweed	Lambsquarters
	Buttercup	Lespedeza
	Buttonweed	Mallow
	Canada thistle	Matchwee :
	Carolina geranium	Mexicanweed
	Carpetweed	Milk vetch
	Catchweed bedstraw	Mouseear
	Catnip	Mouseear hawkweed
	Catsear	Mugwort
	Chickweed,	Musk thistle
	common	Mustard
	Chickweed,	Narrowleaf plantain
	mouseear	Narrowleaf vetch
	Chicory	Nettle
	Cinquefoil	Orange hawkweed
	Clover, crimson	Oriental cocklebur
	Clover, hop	Oxalis
•	Clover, red	Parsley-piert
	Clover, strawberry	Parsnip
	Clover, sweet	Pearlwort
	Clover, white	Pennycress
	Cockle	Pennywort
	Cocklebur	Pepperweed
	Common mullein	Pigweed
	Creeping Charlie	Pineywoods

CONTROLS LIST	continued	
CONTROLS		
[KILLS]	Poison oak	Tansy mustard
•	Pokeweed	Tanweed
	Poorjoe	Thistie
	Prairie sunflower	Trailing crownvetch
	Prickly lettuce	Tumble mustard
	Prickly sida	Tumble pigweed
	Prostrate knotweed	Velvetleaf
!	Prostrate pigweed	Venice mallow
	Prostrate spurge	Virginia buttonweed
	Prostrate vervain	Virginia creeper
	Puncturevine	Virginia pepperweed
	Pursiane, common	Water pennywort
	Ragweed	Wavyleaf bulithistie
	Red sorrel	Western clematis
	Redroot pigweed	Western salsify
	Redstem filaree	White mustard
	Rough cinquefoil	Wild aster
	Rough fleabane	Wild buckwheat
	Russian pigweed	Wild carrot
	Russian thistle	Wild four-o'clock
	Scarlet pimpernel	Wild garlic
	Scotch thistle	Wild geranium
	Sheep sorrel	Wild lettuce
	Shepherdspurse	Wild marigold
	Slender plantain	Wild mustard
	Smailflower	Wild onion
	galinsoga	Wild parsnip
	Smooth dock	Wild radish
	Smooth pigweed	Wild rape
	Sorrel	Wild strawberry
	Sowthistle	Wild sweet potato
	Spanishneedles	Wild vetch
	Speedwell	Woodsorret
	Spiny amaranth	Wooly croton
	Spiny cocklebur	Woolly morningglory
	Spiny sowthistle	Woolly plantain
	Spotted catsear	Wormseed
	Spotted spurge	Yarrow
	Spurweed	Yellow rocket
	Stinging nettle	Yellowflower
i	Strawberry, India mock	pepperweed and other broadleaf
	Tall nettle	weeds
İ	Tall vervain	.,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,
	Tull vervant	

IMPORTANT: Directions for Storage and Disposal Store in original container out of the Storage Conditions:

Disposal:

Plantain

Poison ivy

bedstraw

Plains coreopsis

(tickseed)



reach of children, preferably in a locked storage cabinet. Protect from freezing.

· Do not reuse empty container.

- · Securely wrap original container in several layers of newspaper and discard in trash
- · Place empty container in trash.

Important: It is best to use all of the product in accordance with label directions. If it is necessary to dispose of unused product, please follow any applicable state and local guidelines.

Creeping Jenny

Cudweed

Curly dock

Daisy, English

Daisy, fleabane

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

- · Harmful if swallowed.
- · Causes moderate eye irritation.
- · Avoid contact with eyes, skin or clothing.
- When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves.

After using this product:

- · Rinse gloves before removing.
- · Remove clothing and launder separately before reuse.
- Wash hands thoroughly with soap and water.
- If product gets on clothing, remove clothing as soon as possible and shower.

FIRST AID .		
IN CASE OF ACCIDENTAL	IMMEDIATELY	
Swallowing	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a puison control center or doctor. Do not give anything to an unconscious person. 	
Contact with eyes	 Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. 	
Contact with skin	Take off contaminated clothing. Rinse skiri immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.	
Inhalation	Move person to fresh air. If not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.	

NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 877-229-3763 for medical emergency information. Active Ingredients: 7.59% 2,4-D, dimethylamine salt [CAS # 2008-39-1], 1.83% Mecoprop-p, dimethylamine salt [CAS # 66423-09-4], and 0.84% Dicamba, dimethylamine salt [CAS # 2300-66-5].

ENVIRONMENTAL HAZARDS



- This product is toxic to aquatic invertebrates. Do not apply directly to lakes, streams, rivers or ponds.
- Drift or runoff may harm aquatic invertebrates and non-target plants.
- Do not contaminate water when disposing of equipment washwaters.

NOTICE: Research and testing have determined that the "Directions For Use" are appropriate for the proper use of LAWN HERBICIDE JD CONCENTRATE under expected conditions. The Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.

D-9630-2 101601

Bayer Advanced LLC 1500 Urban Center Drive Birmingham, AL 35242