

# UNITED STATES ENVIRONMENTAL PROTECTION AGENCY Washington, D.C.

SEP 17 2007

OFFICE OF PREVENTION,
PESTICIDES, AND TOXIC SUBSTANCES

Robyn Kneen Bayer Advanced 2 T. W. Alexander Drive Research Triangle Park, N.C. 27709

Dear Ms. Robyn:

SUBJECT:

Label Amendment

Lawn Herbicide JD Concentrate EPA Registration No. 72155-4

Your Application Dated May 31, 2007

The label amendment referred to above, submitted in accordance with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. with the following provisions:

- 1. Under the "Directions for Use" in the "Applications Restrictions" area of the labeling, revise to read as follows:
- "Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- 2. Revise the "EnvironmEntal Hazards" statement to read:

"This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater/."

A stamped copy is enclosed for your records. Please submit one (1) copy of your final printed labeling before you release the product for shipment. This amended labeling supersedes all previously accepted ones.

Sincerely yours,

Joanne I. Miller

Product Manager (23)

Herbicide Branch

Registration Division (75055)

vanne J. Miller

Enclosure

(FRONT PANEL)

# Lawn Herbicide JD Concentrate

This label contains separate label statements for Outdoor Residential Pump Sprayer/ Hose-end Sprayer Concentrate use product and Outdoor Residential Ready-To-Spray use product.

- **Effective Weed Protection**
- Effective Broadleaf Weed Protection
- Effectively Protects Lawns from Broadleaf Weeds
- Kills All Major Broadleaf Weeds
- Kills Weeds Won't Harm Lawns
- Kills to the Root
- Protects Against All Major Broadleaf Weeds
- **Tough Weed Protection**
- Weed Protection System
- Systemic
- 1 Quart treats up to 16,000 square feet
- Convenient MEASURE-UP™ Cup
- Kills Broadleaf weeds in St. Augustine & other Southern Lawns -
- For St. Augustine & other Southern Lawns
- Kills Over 200 Broadleaf Weeds
- Won't Harm Your Lawn
- Rainproof in hours
- 100% Satisfaction Guaranteed\* (or your money back)
- Rainproof in 1 hour!
- Attach to Hose & Spray
- Kills 200+ Weeds

with COMMENTS

In EPA Letter Dated:

Under the Federal Insecticide. Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

72155-4

ACTIVE INGREDIENTS:	
2,4-D, dimethylamine salt	7.59 %
Mecoprop-p, dimethylamine salt	1.83 %
Dicamba, dimethylamine salt	0.84 %
	89.74 %

Net Contents: 16 fl. oz. (473 mL) or 24 fl. oz. (709 mL) or 32 fl. oz. (946 mL) or 1 Gal. (3.78 L)

### **KEEP OUT OF REACH OF CHILDREN** CAUTION

See Back Panel Booklet for Additional Ingredients and Precautionary Statements

### (Pump Sprayer/Hose-end Sprayer label)

(FRONT PANEL)

# Lawn Herbicide JD Concentrate

- Effective Weed Protection
- Effective Broadleaf Weed Protection
- · Effectively Protects Lawns from Broadleaf Weeds
- Kills All Major Broadleaf Weeds
- Kills Weeds Won't Harm Lawns
- Kills to the Root
- Protects Against All Major Broadleaf Weeds
- Tough Weed Protection
- Weed Protection System
- Systemic
- 1 Quart treats up to 16,000 square feet
- Convenient MEASURE-UP™ Cup
- Kills Broadleaf weeds in St. Augustine & other Southern Lawns
- For St. Augustine & other Southern Lawns
- Kills Over 200 Broadleaf Weeds
- Won't Harm Your Lawn
- Rainproof in hours
- 100% Satisfaction Guaranteed\* (or your money back)
- Rainproof in 1 hour!
- Kills 200+ Weeds

#### **ACTIVE INGREDIENTS:**

, . <del>.</del> <del>.</del>	
2,4-D, dimethylamine salt	7.59 %
Mecoprop-p, dimethylamine salt	1.83 %
Dicamba, dimethylamine salt	0.84 %
OTHER INGREDIENTS	89.74 %

Net Contents: 16 fl. oz. (473 mL) or 24 fl. oz. (709 mL) or 32 fl. oz. (946 mL) or 1 Gal. (3.78 L)

# CAUTION

See Back Panel Booklet for Additional Ingredients and Precautionary Statements

#### (BACK PANEL)

#### **OPEN** Resealable Label for Directions & Precautions **PRESS TO RESEAL**

Lawn Herbicide JD Concentrate kills over 200 types of broadleaf weeds in St. Augustine and other Southern lawns.

This special formula kills Dollarweed, Spurge, Oxalis and Dandelion without harming your lawn. Controlling weeds in your lawn has never been easier.

#### **READ THE LABEL FIRST!**

of weeds
. ft.
Over 200 broadleaf weeds such as Dollarweed, Spurge and Oxalis, Dandelion, Clover, Chickweed and other listed weeds
On Lawns
(Do not use on Carpetgrass, Dichondra or Floratam variety of St. Augustinegrass.)
When weeds are actively growing
Northern Lawns: Fescue, Kentucky Bluegrass, and Perennial Ryegrasses
and
Hardy Southern Lawns: Bermuda, Buffalo and Zoysia grasses 2 fl. oz. per 500 sq. ft.
Southern lawns: Bahia, Centipede and St. Augustinegrass grasses  1 fl. oz. per 500 sq. ft.
For questions or comments, call toll-free 1-877 BayerAG (1-877-229-3924) or visit us on-line at <a href="https://www.BayerAdvanced.com">www.BayerAdvanced.com</a>

#### FOR OUTDOOR RESIDENTIAL USE ONLY

$A \cap T \cap$			DIE	ите.
<b>ACTI</b> \	ING	ĸE	DIE	V I J.

2,4-D, dimethylamine salt	7.59 %
Mecoprop-p dimethylamine salt	1.83 %
Dicamba, dimethylamine salt	
· · · · · · · · · · · · · · · · · · ·	89.74 %
OTHER INGREDIENTS :	<u>05.14 %</u>
	100.00 %

Other ingredients include: Water and surfactants (to keep the active ingredients dispersed in water).

#### THIS PRODUCT CONTAINS:

6.30% ........ 2,4-D acid equivalent
1.51% ........ Mecoprop-p acid equivalent
0.69% ........ Dicamba acid equivalent
Isomer specific by AOAC Method

EPA Reg. No. 72155-4

EPA Est. No. \_-AA-\_

DIRECTIONS FOR USE			
It is a violation of Federal L	aw to use this product in a manner inconsistent with its labeling.		
For best results, read and	follow all label directions.		
BEFORE YOU USE	Do not apply near lakes, streams, rivers or ponds.		
Read and follow directions when using.	Spray when most of the day is expected to remain cooler than 90° F.		
	<ul> <li>Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Apply when wind is calm.</li> </ul>		
	The maximum number of broadcast applications per treatment site is two per year.		
HOW TO USE	Determine amount of product to use based on grass type and size of area to be treated.		
Shake Well	Prepare product for use according to How to Apply.		
Before Use	Apply to lawn.		
Continued on next page	3		

#### **DIRECTIONS FOR USE Continued** HOW MUCH TO APPLY Northern Lawns: · Fescue, Kentucky Bluegrass, and Perennial Rye 2 fl. oz. (4 Tbs.) per gallon of water Hardy Southern Lawns: · Bermuda, Buffalo and Zoysia grasses 2 fl. oz. (4 Tbs.) per gallon of water Southern Lawns: · Bahia, Centipede, and St. Augustinegrass 1 fl. oz. (2 Tbs.) per gallon of water For lawns with mixed grass types, use the lowest rate. Each gallon of solution treats 500 sq. ft. TIP: Measure length and width of area to be treated in feet. Multiply these 2 numbers together. This is the area to be treated in square feet (sq. ft.). HOW TO APPLY **Pump Style Sprayers** Add the appropriate amounts of water and concentrate to the sprayer tank. Close sprayer, shake well and pressurize. 3. Adjust nozzle to a coarse spray pattern. 4. Large Areas - Apply evenly over area. Individual Weeds - Hold tip of wand about 12 inches from weed, aim at center of weed and spray. 5. Coccasionally re-pressurize the sprayer if needed to maintain a good spray pattern. **Dial Style Hose-End Sprayers** 1. Fill sprayer jar with enough concentrate to cover area to be treated. Close sprayer and set dial. TIP: The numbers on Dial-Style Hose-End sprayers are ounces of concentrate per gallon of spray. 3. Attach sprayer to hose and apply. 4. To avoid contact with the treated area, begin by spraying area farthest from faucet walking backwards as you spray. Concentrate Plus Water Hose-End Sprayer 1. Pour concentrate into sprayer jar to the required fluid ounce level. 2. Add the appropriate amount of water to the gallons level. 3. Replace sprayer top and shake well before applying. To avoid contact with the treated area, begin by spraying area farthest from faucet walking backwards as you spray. FOR BEST RESULTS Apply when weeds are young and actively growing. If the lawn is dry, water 24 hours before using this product. Rain or watering 6 to 8 hours after application will not wash away effectiveness. Apply 1 to 3 days after mowing. Certain hard-to-kill weeds may need to be spot-treated in 14 days. Do not apply to new lawns until the grass has reached a height of at least 2 inches. Bare spots may be seeded 3 to 4 weeks after application. HOW LAWN HERBICIDE LAWN HERBICIDE JD CONCENTRATE is a selective herbicide - it selectively kills weeds in the lawn but JD CONCENTRATE won't harm the lawn itself. It is a blend of proven weed killers designed to get broadleaf weeds such as WORKS Dollarweed, Spurge and Oxalis.



Do not allow people (other than applicator) or pets on treated area during application. People and pets may re-enter the treated area after it has dried.

CONTROLS	Annual yellow sweetclover	Cockle	Mugwort	Spiny amaranth
[KILLS]	Aster	Cocklebur	Musk thistle	Spiny cocklebur
	Austrian fieldcress	- Common mullein	Mustard	Spiny sowthistle
	Bedstraw .	Creeping Charlie	Narrowleaf plantain	Spotted catsear
	Beggarticks	Creeping Jenny	Narrowleaf vetch	Spotted spurge
	Betony, Florida	Cudweed	Nettle	Spurweed
	Bindweed, field	Curly dock	Orange hawkweed	Stinging nettle
	Bird vetch	Daísy, English	Oriental cocklebur	Strawberry, India mock
	Bitter wintercress	Daisy, fleabane	Oxalis	Tall nettle
	Bittercress, hairy	Daisy, oxeye	Parsley-piert	Tall vervain
	Bitterweed	Dandelion	Parsnip	Tansy mustard
·	Black-eyed Susan	Dichondra	Pearlwort	Tanweed
	Black medic	Dogbane	Pennycress	Thistle
	Black mustard	Dogfennel	Pennywort	Trailing crownvetch
	Blackseed plantain	Dollarweed	Pepperweed ·	Tumble mustard
	Blessed thistle	Elderberry	Pigweed	Tumble pigweed
	Bloodflower milkweed	False dandelion	Pineywoods bedstraw	Velvetleaf
	Blue lettuce	False flax	Plains coreopsis (tickseed)	Venice mallow
	Blue vervain	False sunflower	Plantain	Virginia buttonweed
	Bracted plantain	Fiddleneck	Poison ivy	Virginia creeper
	Brassbuttons	Florida pusley	Poison oak	Virginia pepperweed
	Bristly oxtongue	Frenchweed	Pokeweed	Water pennywort
	Broadleaf dock	Galinsoga	Poorjoe	Wavyleaf bullthistle
	Broadleaf plantain	Goathead	Prairie sunflower	Western clematis
,	Broomweed	Goldenrod	Prickly lettuce	Western salsify
,	Buckhorn	Ground ivy	Prickly sida	White mustard
	Buckhorn plantain	Gumweed .	Prostrate knotweed	Wild aster
	Bulbous buttercup	Hairy fleabane	Prostrate pigweed	Wild buckwheat
	Bull thistle	Hawkweed	· Prostrate spurge	Wild carrot
,	Bullnettle	Healall	Prostrate vervain	Wild four-o'clock
.	Burclover	Heartleaf drymary	Puncturevine	Wild garlic
	Burdock	Heath aster	Purslane, common	Wild geranium
	Burning nettle	Hedge bindweed	Ragweed	Wild lettuce
·	Burweed	Hedge mustard	Red sorrel	Wild marigold
İ	Buttercup	Hemp	Redroot pigweed	Wild mustard
	Buttonweed	Henbit	Redstem filaree	Wild onion
. [	Canada thistle	Hoary cress	Rough cinquefoil	Wild parsnip
	Carolina geranium	Hoary plantain	Rough fleabane	Wild radish
	Carpetweed	Hoary vervain	Russian pigweed	Wild street home
	Catchweed bedstraw	Horsenettle	Russian thistle Scarlet pimpernel	Wild strawberry Wild sweet potato
	Catnip	Jimsonweed Knawel	Scaret plingerner	Wild vetch
	Catsear	Knotweed	Sheep sorrel	Woodsorrel
	Chickweed, common Chickweed, mouseear		•	Wooly croton
,		Kochia	Shepherdspurse	Woolly morningglory
	Chicory	Lambsquarters	Slender plantain Smallflower galinsoga	Woolly plantain
	Cinquefoil	Lespedeza	Smallilower galinsoga Smooth dock	Wormseed
•	Clover, crimson	Mallow	· ·	Yarrow
	Clover, hop	Matchweed	Smooth pigweed	Yarrow Yellow rocket
	Clover, red	Mexicanweed Milk voteb		, , , , , , , , , , , , , , , , , , , ,
•	Clover, strawberry	Milk vetch	Sowthistle	Yellowflower pepperweed and other broadleaf
	Clover, sweet	Morningglory	Spanishneedles	weeds
	Clover, white	Mouseear hawkweed	Speedwell	

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

- · Harmful if swallowed.
- · Causes moderate eye irritation.
- · Avoid contact with eyes, skin or clothing.
- · When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves.

#### After using this product:

- · Rinse gloves before removing.
- · Remove clothing and launder separately before reuse.
- · Wash hands thoroughly with soap and water.
- If product gets on clothing, remove clothing as soon as possible and shower.

♣ FIRST AID		
IN CASE OF ACCIDENTAL	IMMEDIATELY	
IF SWALLOWED	<ul> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>Do not give anything to an unconscious person.</li> </ul>	
IF IN EYES	<ul> <li>Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>	
IF ON SKIN OR CLOTHING	<ul> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>	
IF INHALED	<ul> <li>Move person to fresh air.</li> <li>If not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> </ul>	
	Call a poison control center or doctor for further treatment advice.	

**NOTE:** When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 877-229-3763 for medical emergency information. Active Ingredients: 7.59% 2,4-D, dimethylamine salt [CAS # 2008-39-1], 1.83% Mecoprop-p, dimethylamine salt [CAS # 66423-09-4], and 0.84% Dicamba, dimethylamine salt [CAS # 2300-66-5].

#### **ENVIRONMENTAL HAZARDS**

- · This product is toxic to aquatic invertebrates. Do not apply directly to lakes, streams, rivers or ponds.
- Drift or runoff may harm aquatic invertebrates and non-target plants.
- Do not contaminate water when disposing of equipment washwaters. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D and MCPP pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

Storage Conditions:	Store in original container in a cool dry place, out of the reach of children, preferably in a locked storage		
	cabinet. Protect from freezing.		
Disposal:	If Empty:		
	Do not reuse empty container.		
	Place empty container in trash or offer for recycling if available.		
	If Partly Filled:		
	<ul> <li>If it is necessary to dispose of unused product, please call your local solid waste agency or you may call toll-free 1-877-229-3724 for disposal instructions.</li> </ul>		
•	Never place unused product down any indoor or outdoor drain.		

<sup>\*</sup>Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.



www.bayeradvanced.com
A Business Unit of Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709

(Ready - To - Spray label)
(FRONT PANEL)

# Lawn Herbicide JD Concentrate

- Effective Weed Protection
- Effective Broadleaf Weed Protection
- Effectively Protects Lawns from Broadleaf Weeds
- Kills All Major Broadleaf Weeds
- Kills Weeds Won't Harm Lawns
- Kills to the Root
- Protects Against All Major Broadleaf Weeds
- Tough Weed Protection
- Weed Protection System
- Systemic
- 1 Quart treats up to 16,000 square feet
- Convenient MEASURE-UP™ Cup
- Kills Broadleaf weeds in St. Augustine & other Southern Lawns
- For St. Augustine & other Southern Lawns
- Kills Over 200 Broadleaf Weeds
- Won't Harm Your Lawn
- Rainproof in hours
- 100% Satisfaction Guaranteed\* (or your money back)
- Rainproof in 1 hour!
- Attach to Hose & Spray
- Kills 200+ Weeds

#### **ACTIVE INGREDIENTS:**

NOTIFE MORESIEM OF	
2,4-D, dimethylamine salt	7.59 %
Mecoprop-p, dimethylamine salt	
Dicamba, dimethylamine salt	0.84 %
OTHER INGREDIENTS:	89.74 %

Net Contents: 16 fl. oz. (473 mL) or 24 fl. oz. (709 mL) or 32 fl. oz. (946 mL) or 1 Gal. (3.78 L)

# KEEP OUT OF REACH OF CHILDREN CAUTION

See Back Panel Booklet for Additional Ingredients and Precautionary Statements

#### (BACK PANEL)

#### **OPEN** Resealable Label for Directions & Precautions **PRESS TO RESEAL**

Lawn Herbicide JD Concentrate kills over 200 types of broadleaf weeds in St. Augustine and other Southern lawns.

This special formula kills Dollarweed, Spurge, Oxalis and Dandelion without harming your lawn. Controlling weeds in your lawn has never been easier.

#### **READ THE LABEL FIRST!**

QUICK FACTS	
Kills over 200 broadle	eaf weeds
Won't harm your lawr	n en
<ul> <li>Treats up to 16,000 s</li> </ul>	q. ft.
KILLS	Over 200 broadleaf weeds such as Dollarweed, Spurge and Oxalis, Dandelion, Clover, Chickweed and other listed weeds
WHERE TO USE	On Lawns
	(Do not use on Carpetgrass, Dichondra or Floratam variety of St. Augustinegrass.)
WHEN TO USE	When weeds are actively growing
HOW MUCH TO USE	Northern Lawns: Fescue, Kentucky Bluegrass, and Perennial Ryegrasses
	and
	Hardy Southern Lawns: Bermuda, Buffalo and Zoysia grasses One qt treats up to 8,000 sq. ft.
·	Southern lawns: Bahia, Centipede and St. Augustinegrass grasses One qt treats up to 16,000 sq. ft.
	No measuring needed. Dilutes automatically.
☎ 🖳	For questions or comments, call toll-free 1-877 BayerAG (1-877-229-3924) or visit us on-line at www.BayerAdvanced.com

#### FOR OUTDOOR RESIDENTIAL USE ONLY

#### **ACTIVE INGREDIENTS:**

2,4-D, dimethylamine salt	1.83 %
OTHER INGREDIENTS*:	89.74 %

\* Other ingredients include: Water and surfactants (to keep the active ingredients dispersed in water).

#### THIS PRODUCT CONTAINS:

6.30% ........2,4-D acid equivalent
1.51% ........ Mecoprop-p acid equivalent
0.69% ........ Dicamba acid equivalent
Isomer specific by AOAC Method

EPA Reg. No. 72155-4

EPA Est. No.\_-AA-\_

#### **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For best results, read and follow all label directions.

#### **BEFORE YOU USE**

Read and follow directions when using.

- · Do not apply near lakes, streams, rivers or ponds.
- Spray when most of the day is expected to remain cooler than 90° F.
- Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Apply when wind is calm.
- The maximum number of broadcast applications per treatment site is two per year.

#### **HOW TO USE**

### [For Ready to spray hose end bottle]

#### CONNECT

- 1. Shake well before use.
- 2. Connect sprayer to garden hose.
- 3. Turn on water.



#### **SPRAY**

- Extend hose to farthest area to be treated, point sprayer nozzle in direction to be sprayed.
- Begin spraying <Insert sprayer-specific language here> Hayes HP-2 and Calmar RTS sprayers: Press lock tab while using other hand to turn sprayer valve to ON position. Green Garden RTS sprayer: Turn sprayer valve to ON position.
- 3. Walk at a steady pace while spraying, using an even sweeping motion, and slightly overlap the treated areas. Work back toward the faucet to avoid walking over wet, treated grass.

#### **FINISH**

- 1. To stop spraying, < Insert sprayer-specific language here.> Hayes HP-2 RTS sprayer: Press lock tab while using other hand to turn sprayer valve to OFF position. Calmar & Green Garden RTS sprayers: turn sprayer valve to OFF position.
- 2. Turn off water at faucet.
- 3. Relieve water pressure in hose by turning sprayer valve back to water position until water slows to a drip then turn sprayer valve to OFF position.
- 4. Disconnect sprayer from hose.

#### TIPS FOR APPLYING **HOW TO APPLY** Extend hose to the farthest area to be treated and work back toward the faucet to avoid walking over wet, treated grass. 2. Walk at a steady pace while spraying, using an even sweeping motion, slightly overlapping the treated areas. FOR BEST RESULTS Apply when weeds are young and actively growing. If the lawn is dry, water 24 hours before using this product. Rain or watering 6 to 8 hours after application will not wash away effectiveness. Apply 1 to 3 days after mowing. Certain hard-to-kill weeds may need to be spot-treated in 14 days. Do not apply to new lawns until the grass has reached a height of at least 2 inches. Bare spots may be seeded 3 to 4 weeks after application. **HOW LAWN HERBICIDE** LAWN HERBICIDE JD CONCENTRATE is a selective herbicide - it selectively kills weeds in the lawn JD CONCENTRATE but won't harm the lawn itself. It is a blend of proven weed killers designed to get broadleaf weeds such WORKS as Dollarweed, Spurge and Oxalis.



Do not allow people (other than applicator) or pets on treated area during application.

People and pets may re-enter the treated area after it has dried.

CONTROLS	Annual yellow sweetclover	Cockle	Mugwort	Spiny amaranth
[KILLS]	Aster	Cocklebur	Musk thistle	Spiny cocklebur
	Austrian fieldcress	Common mullein	Mustard	Spiny sowthistle
•	Bedstraw	Creeping Charlie	Narrowleaf plantain	Spotted catsear
	Beggarticks	Creeping Jenny	Narrowleaf vetch	Spotted spurge
	Betony, Florida	Cudweed	Nettle	Spurweed
	Bindweed, field	Curly dock	Orange hawkweed	Stinging nettle
	Bird vetch	Daisy, English	Oriental cocklebur	Strawberry, India mock
	Bitter wintercress	Daisy, fleabane	Oxalis	Tall nettle
	Bittercress, hairy	Daisy, oxeye	Parsley-piert	Tall vervain
	Bitterweed	Dandelion	Parsnip	Tansy mustard
	Black-eyed Susan	Dichondra	Pearlwort	Tanweed
	Black medic	Dogbane	Pennycress	Thistle
	Black mustard	Dogfennel	Pennywort	Trailing crownvetch
	Blackseed plantain	Dollarweed	Pepperweed	Tumble mustard
	Blessed thistle	Elderberry	Pigweed	Tumble pigweed
	Bloodflower milkweed	False dandelion	Pineywoods bedstraw	Velvetleaf
	Blue lettuce	False flax	Plains coreopsis (tickseed)	Venice mallow
	Blue vervain	False sunflower	Plantain	Virginia buttonweed
:	Bracted plantain	Fiddleneck	Poison ivy	Virginia creeper
	Brassbuttons	Florida pusley	Poison oak	Virginia pepperweed
,	Bristly oxtongue	Frenchweed	Pokeweed	Water pennywort
,	Broadleaf dock	Galinsoga	Poorjoe	Wavyleaf bulithistle
	Broadleaf plantain	Goathead	Prairie sunflower	Western clematis
	Broomweed	Goldenrod	Prickly lettuce	Western salsify
	Buckhorn	Ground ivy	Prickly sida	White mustard
	Buckhorn plantain	Gumweed	Prostrate knotweed	Wild aster
	Bulbous buttercup	Hairy fleabane	Prostrate pigweed	Wild buckwheat
·	Bull thistle	Hawkweed	Prostrate spurge	Wild carrot
	Bullnettle	Healall	Prostrate vervain	Wild four-o'clock
	Burclover	Heartleaf drymary	Puncturevine	Wild garlic
·	Burdock	Heath aster	Purslane, common	Wild geranium
	Burning nettle	Hedge bindweed	Ragweed	Wild lettuce
,	Burweed	Hedge mustard	Red sorrel	Wild marigold
	Buttercup	Hemp	Redroot pigweed	Wild mustard
	Buttonweed	Henbit	Redstem filaree	Wild onion
	Canada thistle	Hoary cress	Rough cinquefoil	Wild parsnip
	Carolina geranium	Hoary plantain	Rough fleabane	Wild radish
	Carpetweed	Hoary vervain	Russian pigweed	Wild rape
	Catchweed bedstraw	Horsenettle	Russian thistle	Wild strawberry
	Catnip	Jimsonweed	Scarlet pimpernel	Wild sweet potato
	Catsear	Knawel	Scotch thistle	Wild vetch
	Chickweed, common	Knotweed	Sheep sorrel	Woodsorrel
Ì	Chickweed, mouseear	Kochia	Shepherdspurse	. Wooly croton
	Chicory	Lambsquarters	Slender plantain	Woolly morningglory
Ī.	Cinquefoil	Lespedeza .	Smallflower galinsoga	Wootly plantain
	Clover, crimson	Mallow	Smooth dock	Wormseed
ļ.	Clover, hop	Matchweed	Smooth pigweed	Yarrow
	Clover, red	Mexicanweed	Sorrel	Yellow rocket
	Clover, strawberry	Milk vetch	Sowthistle	Yellowflower pepperweed
	Clover, sweet	Morningglory	Spanishneedles	and other broadleaf
	Clover, white	Mouseear hawkweed	Speedwell	weeds

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

- · Harmful if swallowed.
- · Causes moderate eye irritation.
- · Avoid contact with eyes, skin or clothing.
- · When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves.

#### After using this product:

- · Rinse gloves before removing.
- · Remove clothing and launder separately before reuse.
- · Wash hands thoroughly with soap and water.
- If product gets on clothing, remove clothing as soon as possible and shower.

FIRST AID +				
IN CASE OF ACCIDENTAL	IMMEDIATELY			
IF SWALLOWED	Call a poison control center or doctor immediately for treatment advice.			
	Have person sip a glass of water if able to swallow.			
	Do not induce vomiting unless told to do so by a poison control center or doctor.			
	Do not give anything to an unconscious person.			
IF IN EYES	Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.			
	Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.			
• *	Call a poison control center or doctor for treatment advice.			
IF ON SKIN OR CLOTHING	Take off contaminated clothing.			
	Rinse skin immediately with plenty of water for 15 to 20 minutes.			
	Call a poison control center or doctor for treatment advice.			
IF INHALED	Move person to fresh air.			
	<ul> <li>If not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> </ul>			
	Call a poison control center or doctor for further treatment advice.			

**NOTE:** When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 877-229-3763 for medical emergency information. Active Ingredients: 7.59% 2,4-D, dimethylamine salt [CAS # 2008-39-1], 1.83% Mecoprop-p, dimethylamine salt [CAS # 66423-09-4], and 0.84% Dicamba, dimethylamine salt [CAS # 2300-66-5].

#### **ENVIRONMENTAL HAZARDS**

- This product is toxic to aquatic invertebrates. Do not apply directly to lakes, streams, rivers or ponds.
- · Drift or runoff may harm aquatic invertebrates and non-target plants.
- · Do not contaminate water when disposing of equipment washwaters.

IMPORTANT: Directions for Storage and Disposal			
Storage Conditions:	Store in original container in a cool dry place, out of the reach of children, preferably in a locked stora cabinet. Protect from freezing.		
Disposal:	<ul> <li>If Empty:</li> <li>Do not reuse empty container.</li> <li>Place empty container in trash or offer for recycling if available.</li> <li>If Partiy Filled:</li> <li>If it is necessary to dispose of unused product, please call your local solid waste agency or you may call toll-free 1-877-229-3724 for disposal instructions.</li> <li>Never place unused product down any indoor or outdoor drain.</li> </ul>		

**NOTICE:** Research and testing have determined that the "Directions For Use" are appropriate for the proper use of LAWN HERBICIDE JD CONCENTRATE under expected conditions. The Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

\*Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.



www.bayeradvanced.com
A Business Unit of Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709