

72155-4

8/5/2009

Jacket
10/15



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number: 72155-4
Date of Issuance: AUG 5 2009

NOTICE OF PESTICIDE:
 Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:
Name of Pesticide Product:
Lawn Herbicide JD Concentrate

Name and Address of Registrant (include ZIP Code):
Bayer Advanced
2 T.W. Alexander Drive
Research Triangle Park, NC 27709

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is reregistered in accordance with FIFRA section 4(g)(2)(C) provided you:

1. Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data.
2. Make all the following changes to the product label:
 - a. Per the acute toxicity review, child resistant packaging (CRP) is triggered for this product.
 - b. Change the signal word from "CAUTION" to "DANGER".
 - c. Add a line in the ingredients statement with the heading "Total" and the value "100%" directly below the other ingredient percentages.
 - d. The Hazards to Humans and Domestic Animals statement must be changed to read:
"DANGER: Corrosive. Causes irreversible eye damage. Harmful if swallowed or absorbed through skin. Do not get in eyes or on clothing. Avoid contact with skin. Wear protective eyewear, long-sleeved shirt, long pants, shoes and socks, and chemical-resistant gloves made of any waterproof material. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse."

Signature of Approving Official:

Joanne Miller
Product Manager 23
Herbicide Branch
Registration Division (7505P)

Date:

AUG 5 2009

- e. On page 3, remove the word "special" from the claim "This special formula kills Dollarweed, Spurge, Oxalis and Dandelion without harming your lawn.", as it implies the product possesses unique characteristics due to its composition, which are false and misleading. Also, add the asterisk at the end of the above sentence, referencing the footnote "when used as directed".
- f. Based on toxicity ranking per the acute toxicity review, the First Aid statements must be placed on the label in the following order:
"IF IN EYES
IF SWALLOWED
IF ON SKIN OR CLOTHING
IF INHALED"
- g. A "Note to Physician" must be added to the label, preferably under the First Aid section, with the statement "Probably mucosal damage may contraindicate the use of gastric lavage."

Also, it is recommended that additional text be added that addresses eye irritation concerns. The following are some suggested types of information that could be included, if applicable:

- Technical information on symptomatology;
 - Use of supportive treatments to maintain life functions;
 - Medicine that will counteract the specific physiological effects of the pesticide;
 - Company telephone number to specific medical personnel who can provide specialized medical advice.
- h. Revise the first sentence in the Environmental Hazards section to read "This pesticide is toxic to fish and aquatic invertebrates **and may adversely affect non-target plants.**"
- i. Add the following restriction to the Directions for Use section:
- "Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn or garden area only."

A stamped copy of your label is enclosed for your records. You must submit one (1) copy of the final printed label before you release the product for shipment. Products shipped after 12 months from the date of this Letter or the next printing of the label whichever occurs first, must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Enclosure

Stamped 5-14-09 - 30 of 15

(FRONT PANEL)

COPY

Lawn Herbicide JD Concentrate

This label contains separate label statements for **Outdoor Residential Pump Sprayer/ Hose-end Sprayer Concentrate** use product and **Outdoor Residential Ready-To-Spray** use product.

- Effective Weed Protection
- Effective Broadleaf Weed Protection
- Effectively Protects Lawns from Broadleaf Weeds
- Kills All Major Broadleaf Weeds
- Kills Weeds - Won't Harm Lawns†
- Kills to the Root
- Protects Against All Major Broadleaf Weeds
- Tough Weed Protection
- Weed Protection System
- Systemic
- 1 Quart treats up to 16,000 square feet
- Convenient MEASURE-UP™ Cup
- Kills Broadleaf weeds in St. Augustine & other Southern Lawns
- For St. Augustine & other Southern Lawns
- Kills Over 200 Broadleaf Weeds
- Won't Harm Your Lawn†
- Rainproof in hours
- 100% Satisfaction Guaranteed* (or your money back)
- Rainproof in 1 hour!
- Attach to Hose & Spray
- Kills 200+ Weeds
- [optional advertising claims when product is sold in "low-water" hose-end sprayers, as determined by testing]
- Water [Saver][Saving] Formula
- Water [Saver][Saving] Applicator
- Water [Saver][Saving] Sprayer
- Uses Up To [80][XX]% Less Water*
- [with footnote disclaimer] *vs leading Southern weed hose end sprayers
- won't harm lawns†
- †when used as directed

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	7.59 %
Mecoprop-p, dimethylamine salt	1.83 %
Dicamba, dimethylamine salt	0.84 %

OTHER INGREDIENTS: 89.74 %

Net Contents: 16 fl. oz. (473 mL) or 24 fl. oz. (709 mL) or 32 fl. oz. (946 mL) or 1 Gal. (3.78 L)

KEEP OUT OF REACH OF CHILDREN

CAUTION

ACCEPTED


with **COMMENTS** See Back Panel Booklet for Additional Ingredients and Precautionary Statements

In EPA Letter Dated:

AUG 5 2009

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

72155-4



(Pump Sprayer/Hose-end Sprayer label)

(FRONT PANEL)

Lawn Herbicide JD Concentrate

- Effective Weed Protection
- Effective Broadleaf Weed Protection
- Effectively Protects Lawns from Broadleaf Weeds
- Kills All Major Broadleaf Weeds
- Kills Weeds - Won't Harm Lawns†
- Kills to the Root
- Protects Against All Major Broadleaf Weeds
- Tough Weed Protection
- Weed Protection System
- Systemic
- 1 Quart treats up to 16,000 square feet
- Convenient MEASURE-UP™ Cup
- Kills Broadleaf weeds in St. Augustine & other Southern Lawns
- For St. Augustine & other Southern Lawns
- Kills Over 200 Broadleaf Weeds
- Won't Harm Your Lawn†
- Rainproof in hours
- 100% Satisfaction Guaranteed* (or your money back)
- Rainproof in 1 hour!
- Kills 200+ Weeds
- [optional advertising claims when product is sold in "low-water" hose-end sprayers, as determined by testing]
- Water [Saver][Saving] Formula
- Water [Saver][Saving] Applicator
- Water [Saver][Saving] Sprayer
- Uses Up To [80][XX]% Less Water*
- [with footnote disclaimer] *vs leading Southern weed hose end sprayers
- won't harm lawns†
- †when used as directed

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	7.59 %
Mecoprop-p, dimethylamine salt	1.83 %
Dicamba, dimethylamine salt	0.84 %

OTHER INGREDIENTS: 89.74 %

Net Contents: 16 fl. oz. (473 mL) or 24 fl. oz. (709 mL) or 32 fl. oz. (946 mL) or 1 Gal. (3.78 L)

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Back Panel Booklet for Additional Ingredients and Precautionary Statements

50915

(BACK PANEL)

OPEN Resealable Label for Directions & Precautions **PRESS TO RESEAL**

Lawn Herbicide JD Concentrate kills over 200 types of broadleaf weeds in St. Augustine and other Southern lawns. This special formula kills Dollarweed, Spurge, Oxalis and Dandelion without harming your lawn. Controlling weeds in your lawn has never been easier.

READ THE LABEL FIRST!

QUICK FACTS	
<ul style="list-style-type: none"> • Kills over 200 broadleaf weeds • Won't harm your lawn† • Treats up to 16,000 sq. ft. 	
KILLS	Over 200 broadleaf weeds such as Dollarweed, Spurge and Oxalis, Dandelion, Clover, Chickweed and other listed weeds
WHERE TO USE	On Lawns † (Do not use on Carpetgrass, Dichondra or Floratam variety of St. Augustinegrass.)
WHEN TO USE	When weeds are actively growing
HOW MUCH TO USE	<p>Northern Lawns: Fescue, Kentucky Bluegrass, and Perennial Ryegrasses and Hardy Southern Lawns: Bermuda, Buffalo and Zoysia grasses 2 fl. oz. per 500 sq. ft.</p> <p>Southern lawns: Bahia, Centipede and St. Augustinegrass grasses 1 fl. oz. per 500 sq. ft.</p>
For questions or comments, call toll-free 1-877 BayerAG (1-877-229-3724) or visit us on-line at www.BayerAdvanced.com	

FOR OUTDOOR RESIDENTIAL USE ONLY

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt.....	7.59 %
Mecoprop-p, dimethylamine salt.....	1.83 %
Dicamba, dimethylamine salt.....	0.84 %
OTHER INGREDIENTS*:	89.74 %
	100.00 %

* Other ingredients include: Water and surfactants (to keep the active ingredients dispersed in water).

THIS PRODUCT CONTAINS:

6.30%	2,4-D acid equivalent
1.51%	Mecoprop-p acid equivalent
0.69%	Dicamba acid equivalent
Isomer specific by AOAC Method	

EPA Reg. No. 72155-4

EPA Est. No. _-AA-_

DIRECTIONS FOR USE	
<p>It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. For best results, read and follow all label directions. Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.</p>	
BEFORE YOU USE Read and follow directions when using.	<ul style="list-style-type: none"> • Do not apply near lakes, streams, rivers or ponds. • Spray when most of the day is expected to remain cooler than 90° F. • Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Apply when wind is calm. • The maximum number of broadcast applications per treatment site is two per year.
HOW TO USE Shake Well Before Use	<ul style="list-style-type: none"> • Determine amount of product to use based on grass type and size of area to be treated. • Prepare product for use according to How to Apply. • Apply to lawn.
Continued on next page	

DIRECTIONS FOR USE Continued	
HOW MUCH TO APPLY	<p>Northern Lawns:</p> <ul style="list-style-type: none"> Fescue, Kentucky Bluegrass, and Perennial Rye 2 fl. oz. (4 Tbs.) per gallon of water <p>Hardy Southern Lawns:</p> <ul style="list-style-type: none"> Bermuda, Buffalo and Zoysia grasses 2 fl. oz. (4 Tbs.) per gallon of water <p>Southern Lawns:</p> <ul style="list-style-type: none"> Bahia, Centipede, and St. Augustinegrass 1 fl. oz. (2 Tbs.) per gallon of water <p>For lawns with mixed grass types, use the lowest rate. Each gallon of solution treats 500 sq. ft. TIP: Measure length and width of area to be treated in feet. Multiply these 2 numbers together. This is the area to be treated in square feet (sq. ft.).</p>
HOW TO APPLY	<p>Pump Style Sprayers</p> <ol style="list-style-type: none"> Add the appropriate amounts of water and concentrate to the sprayer tank. Close sprayer, shake well and pressurize. Adjust nozzle to a coarse spray pattern. Large Areas – Apply evenly over area. Individual Weeds – Hold tip of wand about 12 inches from weed, aim at center of weed and spray. Occasionally re-pressurize the sprayer if needed to maintain a good spray pattern. <p>Dial Style Hose-End Sprayers</p> <ol style="list-style-type: none"> Fill sprayer jar with enough concentrate to cover area to be treated. Close sprayer and set dial. TIP: The numbers on Dial-Style Hose-End sprayers are ounces of concentrate per gallon of spray. Attach sprayer to hose and apply. To avoid contact with the treated area, begin by spraying area farthest from faucet walking backwards as you spray. <p>Concentrate Plus Water Hose-End Sprayer</p> <ol style="list-style-type: none"> Pour concentrate into sprayer jar to the required fluid ounce level. Add the appropriate amount of water to the gallons level. Replace sprayer top and shake well before applying. To avoid contact with the treated area, begin by spraying area farthest from faucet walking backwards as you spray.
FOR BEST RESULTS	<ul style="list-style-type: none"> Apply when weeds are young and actively growing. If the lawn is dry, water 24 hours before using this product. Rain or watering 6 to 8 hours after application will not wash away effectiveness. Apply 1 to 3 days after mowing. Certain hard-to-kill weeds may need to be spot-treated in 14 days. Do not apply to new lawns until the grass has reached a height of at least 2 inches. Bare spots may be seeded 3 to 4 weeks after application.
HOW LAWN HERBICIDE JD CONCENTRATE WORKS	<p>LAWN HERBICIDE JD CONCENTRATE is a selective herbicide - it selectively kills weeds in the lawn but won't harm the lawn itself. It is a blend of proven weed killers designed to get broadleaf weeds such as Dollarweed, Spurge and Oxalis.</p>



Do not allow people (other than applicator) or pets on treated area during application.
People and pets may re-enter the treated area after it has dried.

CONTROLS [KILLS]	Annual yellow sweetclover Aster Austrian fieldcress Bedstraw Beggarticks Betony, Florida Bindweed, field Bird vetch Bitter wintercress Bittercress, hairy Bitterweed Black-eyed Susan Black medic Black mustard Blackseed plantain Blessed thistle Bloodflower milkweed Blue lettuce Blue vervain Bracted plantain Brassbuttons Bristly oxtongue Broadleaf dock Broadleaf plantain Broomweed Buckhorn Buckhorn plantain Bulbous buttercup Bull thistle Bullnettle Burclover Burdock Burning nettle Burweed Buttercup Buttonweed Canada thistle Carolina geranium Carpetweed Catchweed bedstraw Catnip Catsear Chickweed, common Chickweed, mouseear Chicory Cinquefoil Clover, crimson Clover, hop Clover, red Clover, strawberry Clover, sweet Clover, white	Cockle Cocklebur Common mullein Creeping Charlie Creeping Jenny Cudweed Curly dock Daisy, English Daisy, fleabane Daisy, oxeye Dandelion Dichondra Dogbane Dogfennel Dollarweed Elderberry False dandelion False flax False sunflower Fiddleneck Florida pusley Frenchweed Galinsoga Goathead Goldenrod Ground ivy Gumweed Hairy fleabane Hawkweed Healall Heartleaf drymary Heath aster Hedge bindweed Hedge mustard Hemp Henbit Hoary cress Hoary plantain Hoary vervain Horsenettle Jimsonweed Knapel Knotweed Kochia Lambsquarters Lespedeza Mallow Matchweed Mexicanweed Milk vetch Morningglory Mouseear hawkweed	Mugwort Musk thistle Mustard Narrowleaf plantain Narrowleaf vetch Nettle Orange hawkweed Oriental cocklebur Oxalis Parsley-piert Parsnip Pearlwort Pennycress Pennywort Pepperweed Pigweed Pineywoods bedstraw Plains coreopsis (tickseed) Plantain Poison ivy Poison oak Pokeweed Poorjoe Prairie sunflower Prickly lettuce Prickly sida Prostrate knotweed Prostrate pigweed Prostrate spurge Prostrate vervain Puncturevine Purslane, common Ragweed Red sorrel Redroot pigweed Redstem filaree Rough cinquefoil Rough fleabane Russian pigweed Russian thistle Scarlet pimpernel Scotch thistle Sheep sorrel Shepherdspurse Slender plantain Smallflower galinsoga Smooth dock Smooth pigweed Sorrel Sowthistle Spanishneedles Speedwell	Spiny amaranth Spiny cocklebur Spiny sowthistle Spotted catsear Spotted spurge Spurweed Stinging nettle Strawberry, India mock Tall nettle Tall vervain Tansy mustard Tanweed Thistle Trailing crownvetch Tumble mustard Tumble pigweed Velvetleaf Venice mallow Virginia buttonweed Virginia creeper Virginia pepperweed Water pennywort Wavyleaf bullthistle Western clematis Western salsify White mustard Wild aster Wild buckwheat Wild carrot Wild four-o'clock Wild garlic Wild geranium Wild lettuce Wild marigold Wild mustard Wild onion Wild parsnip Wild radish Wild rape Wild strawberry Wild sweet potato Wild vetch Woodsorrel Woolly croton Woolly morningglory Woolly plantain Wormseed Yarrow Yellow rocket Yellowflower pepperweed and other broadleaf weeds
-----------------------------------	--	--	---	--

80415

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

- Harmful if swallowed.
- Causes moderate eye irritation.
- Avoid contact with eyes, skin or clothing.
- When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves.

After using this product:

- Rinse gloves before removing.
- Remove clothing and launder separately before reuse.
- Wash hands thoroughly with soap and water.
- If product gets on clothing, remove clothing as soon as possible and shower.

FIRST AID

+	IMMEDIATELY...
IN CASE OF ACCIDENTAL...	
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything to an unconscious person.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
<p>NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 877-229-3763 for medical emergency information. Active Ingredients: 7.59% 2,4-D, dimethylamine salt [CAS # 2008-39-1], 1.83% Mecoprop-p, dimethylamine salt [CAS # 66423-09-4], and 0.84% Dicamba, dimethylamine salt [CAS # 2300-66-5].</p>	


ENVIRONMENTAL HAZARDS

- This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark, except as noted on appropriate labels.
- Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas.
- Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D and MCPP pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

9 of 15

IMPORTANT: Directions for Storage and Disposal	
Storage Conditions:	Store in original container in a cool dry place, out of the reach of children, preferably in a locked storage cabinet. Protect from freezing.
Disposal: 	If Empty: <ul style="list-style-type: none">• Do not reuse empty container.• Place empty container in trash or offer for recycling if available. If Partly Filled: <ul style="list-style-type: none">• If it is necessary to dispose of unused product, please call your local solid waste agency or you may call toll-free 1-877-229-3724 for disposal instructions.• Never place unused product down any indoor or outdoor drain.

***Money Back Guarantee:** If you are not satisfied with this product, we will gladly refund your original purchase price.



A Business Unit of Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709

100815

(Ready - To - Spray label)
(FRONT PANEL)

Lawn Herbicide JD Concentrate

- Effective Weed Protection
- Effective Broadleaf Weed Protection
- Effectively Protects Lawns from Broadleaf Weeds
- Kills All Major Broadleaf Weeds
- Kills Weeds - Won't Harm Lawns†
- Kills to the Root
- Protects Against All Major Broadleaf Weeds
- Tough Weed Protection
- Weed Protection System
- Systemic
- 1 Quart treats up to 16,000 square feet
- Convenient MEASURE-UP™ Cup
- Kills Broadleaf weeds in St. Augustine & other Southern Lawns
- For St. Augustine & other Southern Lawns
- Kills Over 200 Broadleaf Weeds
- Won't Harm Your Lawn†
- Rainproof in hours
- 100% Satisfaction Guaranteed* (or your money back)
- Rainproof in 1 hour!
- Attach to Hose & Spray
- Kills 200+ Weeds
- [optional advertising claims when product is sold in "low-water" hose-end sprayers, as determined by testing]
- Water [Saver][Saving] Formula
- Water [Saver][Saving] Applicator
- Water [Saver][Saving] Sprayer
- Uses Up To [80][XX]% Less Water*
- [with footnote disclaimer] *vs leading Southern weed hose end sprayers
- won't harm lawns†
- †when used as directed

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	7.59 %
Mecoprop-p, dimethylamine salt	1.83 %
Dicamba, dimethylamine salt	0.84 %

OTHER INGREDIENTS: 89.74 %

Net Contents: 16 fl. oz. (473 mL) or 24 fl. oz. (709 mL) or 32 fl. oz. (946 mL) or 1 Gal. (3.78 L)

KEEP OUT OF REACH OF CHILDREN
CAUTION

See Back Panel Booklet for Additional Ingredients and Precautionary Statements

11/09/15


(BACK PANEL)

OPEN Resealable Label for Directions & Precautions **PRESS TO RESEAL**

Lawn Herbicide JD Concentrate kills over 200 types of broadleaf weeds in St. Augustine and other Southern lawns.

This special formula kills Dollarweed, Spurge, Oxalis and Dandelion without harming your lawn. Controlling weeds in your lawn has never been easier.

READ THE LABEL FIRST!

QUICK FACTS	
<ul style="list-style-type: none"> • Kills over 200 broadleaf weeds • Won't harm your lawn† • Treats up to 16,000 sq. ft. 	
KILLS	Over 200 broadleaf weeds such as Dollarweed, Spurge and Oxalis, Dandelion, Clover, Chickweed and other listed weeds
WHERE TO USE	On Lawns † (Do not use on Carpetgrass, Dichondra or Floratam variety of St. Augustinegrass.)
WHEN TO USE	When weeds are actively growing
HOW MUCH TO USE	<p>Northern Lawns: Fescue, Kentucky Bluegrass, and Perennial Ryegrasses and Hardy Southern Lawns: Bermuda, Buffalo and Zoysia grasses One qt treats up to 8,000 sq. ft.</p> <p>Southern lawns: Bahia, Centipede and St. Augustinegrass grasses One qt treats up to 16,000 sq. ft.</p> <p>No measuring needed. Dilutes automatically.</p>
	For questions or comments, call toll-free 1-877 BayerAG (1-877-229-3724) or visit us on-line at www.BayerAdvanced.com

FOR OUTDOOR RESIDENTIAL USE ONLY

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt.....	7.59 %
Mecoprop-p, dimethylamine salt.....	1.83 %
Dicamba, dimethylamine salt.....	0.84 %
OTHER INGREDIENTS*:	89.74 %
	100.00 %

* Other ingredients include: Water and surfactants (to keep the active ingredients dispersed in water).

THIS PRODUCT CONTAINS:

6.30% 2,4-D acid equivalent
1.51% Mecoprop-p acid equivalent
0.69% Dicamba acid equivalent
Isomer specific by AOAC Method

EPA Reg. No. 72155-4

EPA Est. No. -AA-

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For best results, read and follow all label directions.



Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

BEFORE YOU USE
Read and follow directions when using.


- Do not apply near lakes, streams, rivers or ponds.
- Spray when most of the day is expected to remain cooler than 90° F.
- Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Apply when wind is calm.
- The maximum number of broadcast applications per treatment site is two per year.

HOW TO USE
[For Ready to spray hose end bottle]

Shake Well Before Use



- CONNECT**
1. Shake well before use.
 2. Connect sprayer to garden hose.
 3. Turn on water.



- SPRAY**
1. Extend hose to farthest area to be treated, point sprayer nozzle in direction to be sprayed.
 2. Begin spraying <Insert sprayer-specific language here> Hayes HP-2 and Calmar RTS sprayers: Press lock tab while using other hand to turn sprayer valve to ON position. Green Garden RTS sprayer: Turn sprayer valve to ON position.
 3. Walk at a steady pace while spraying, using an even sweeping motion, and slightly overlap the treated areas. Work back toward the faucet to avoid walking over wet, treated grass.

- FINISH**
1. To stop spraying, <Insert sprayer-specific language here.> Hayes HP-2 RTS sprayer: Press lock tab while using other hand to turn sprayer valve to OFF position. Calmar & Green Garden RTS sprayers: turn sprayer valve to OFF position.
 2. Turn off water at faucet.
 3. Relieve water pressure in hose by turning sprayer valve back to ON position until water slows to a drip then turn sprayer valve to OFF position.
 4. Disconnect sprayer from hose.

TIPS FOR APPLYING

HOW TO APPLY

1. Extend hose to the farthest area to be treated and work back toward the faucet to avoid walking over wet, treated grass.
2. Walk at a steady pace while spraying, using an even sweeping motion, slightly overlapping the treated areas.

FOR BEST RESULTS

- Apply when weeds are young and actively growing.
- If the lawn is dry, water 24 hours before using this product.
- Rain or watering 6 to 8 hours after application will not wash away effectiveness.
- Apply 1 to 3 days after mowing.
- Certain hard-to-kill weeds may need to be spot-treated in 14 days.
- Do not apply to new lawns until the grass has reached a height of at least 2 inches.
- Bare spots may be seeded 3 to 4 weeks after application.

HOW LAWN HERBICIDE JD CONCENTRATE WORKS

LAWN HERBICIDE JD CONCENTRATE is a selective herbicide - it selectively kills weeds in the lawn but won't harm the lawn itself. It is a blend of proven weed killers designed to get broadleaf weeds such as Dollarweed, Spurge and Oxalis.



Do not allow people (other than applicator) or pets on treated area during application. People and pets may re-enter the treated area after it has dried.

CONTROLS [KILLS]	Annual yellow sweetclover	Cockle	Mugwort	Spiny amaranth
	Aster	Cocklebur	Musk thistle	Spiny cocklebur
	Austrian fieldcress	Common mullein	Mustard	Spiny sowthistle
	Bedstraw	Creeping Charlie	Narrowleaf plantain	Spotted catsear
	Beggarticks	Creeping Jenny	Narrowleaf vetch	Spotted spurge
	Betony, Florida	Cudweed	Nettle	Spurweed
	Bindweed, field	Curly dock	Orange hawkweed	Stinging nettle
	Bird vetch	Daisy, English	Oriental cocklebur	Strawberry, India mock
	Bitter wintercress	Daisy, fleabane	Oxalis	Tall nettle
	Bittercress, hairy	Daisy, oxeeye	Parsley-piert	Tall vervain
	Bitterweed	Dandelion	Parsnip	Tansy mustard
	Black-eyed Susan	Dichondra	Pearlwort	Tanweed
	Black medic	Dogbane	Pennycress	Thistle
	Black mustard	Dogfennel	Pennywort	Trailing crownvetch
	Blackseed plantain	Dollarweed	Pepperweed	Tumble mustard
	Blessed thistle	Elderberry	Pigweed	Tumble pigweed
	Bloodflower milkweed	False dandelion	Pineywoods bedstraw	Velvetleaf
	Blue lettuce	False flax	Plains coreopsis (tickseed)	Venice mallow
	Blue vervain	False sunflower	Plantain	Virginia buttonweed
	Bracted plantain	Fiddleneck	Poison ivy	Virginia creeper
	Brassbuttons	Florida pusley	Poison oak	Virginia pepperweed
	Bristly oxtongue	Frenchweed	Pokeweed	Water pennywort
	Broadleaf dock	Galinsoga	Poorjoe	Wavyleaf bullthistle
	Broadleaf plantain	Goathead	Prairie sunflower	Western clematis
	Broomweed	Goldenrod	Prickly lettuce	Western salsify
	Buckhorn	Ground ivy	Prickly sida	White mustard
	Buckhorn plantain	Gumweed	Prostrate knotweed	Wild aster
	Bulbous buttercup	Hairy fleabane	Prostrate pigweed	Wild buckwheat
	Bull thistle	Hawkweed	Prostrate spurge	Wild carrot
	Bullnettle	Healall	Prostrate vervain	Wild four-o'clock
	Burclover	Heartleaf drymary	Puncturevine	Wild garlic
	Burdock	Heath aster	Purslane, common	Wild geranium
	Burning nettle	Hedge bindweed	Ragweed	Wild lettuce
	Burweed	Hedge mustard	Red sorrel	Wild marigold
	Buttercup	Hemp	Redroot pigweed	Wild mustard
	Buttonweed	Henbit	Redstem filaree	Wild onion
	Canada thistle	Hoary cress	Rough cinquefoil	Wild parsnip
	Carolina geranium	Hoary plantain	Rough fleabane	Wild radish
	Carpetweed	Hoary vervain	Russian pigweed	Wild rape
	Catchweed bedstraw	Horsenettle	Russian thistle	Wild strawberry
	Catnip	Jimsonweed	Scarlet pimpernel	Wild sweet potato
	Catsear	Knawel	Scotch thistle	Wild vetch
	Chickweed, common	Knotweed	Sheep sorrel	Woodsorrel
	Chickweed, mouseear	Kochia	Shepherdspurse	Woolly croton
	Chicory	Lambsquarters	Slender plantain	Woolly morningglory
	Cinquefoil	Lespedeza	Smallflower galinsoga	Woolly plantain
	Clover, crimson	Mallow	Smooth dock	Wormseed
Clover, hop	Matchweed	Smooth pigweed	Yarrow	
Clover, red	Mexicanweed	Sorrel	Yellow rocket	
Clover, strawberry	Milk vetch	Sowthistle	Yellowflower pepperweed	
Clover, sweet	Morningglory	Spanishneedles	and other broadleaf	
Clover, white	Mouseear hawkweed	Speedwell	weeds	

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

- Harmful if swallowed.
- Causes moderate eye irritation.
- Avoid contact with eyes, skin or clothing.
- When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves.

After using this product:

- Rinse gloves before removing.
- Remove clothing and launder separately before reuse.
- Wash hands thoroughly with soap and water.
- If product gets on clothing, remove clothing as soon as possible and shower.

FIRST AID +

IN CASE OF ACCIDENTAL...

IMMEDIATELY...

IF SWALLOWED

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything to an unconscious person.

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 to 20 minutes.
- Call a poison control center or doctor for treatment advice.

IF INHALED


- Move person to fresh air.
- If not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 877-229-3763 for medical emergency information.
Active Ingredients: 7.59% 2,4-D, dimethylamine salt [CAS # 2008-39-1], 1.83% Mecoprop-p, dimethylamine salt [CAS # 66423-09-4], and 0.84% Dicamba, dimethylamine salt [CAS # 2300-66-5].

ENVIRONMENTAL HAZARDS

- This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark, except as noted on appropriate labels.
- Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas.
- Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

IMPORTANT: Directions for Storage and Disposal.	
Storage Conditions:	Store in original container in a cool dry place, out of the reach of children, preferably in a locked storage cabinet. Protect from freezing.
Disposal:	 <ul style="list-style-type: none"> • If Empty: • Do not reuse empty container. • Place empty container in trash or offer for recycling if available. • If Partly Filled: • If it is necessary to dispose of unused product, please call your local solid waste agency or you may call toll-free 1-877-229-3724 for disposal instructions. • Never place unused product down any indoor or outdoor drain.

NOTICE: Research and testing have determined that the "Directions For Use" are appropriate for the proper use of LAWN HERBICIDE JD CONCENTRATE under expected conditions. The Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

***Money Back Guarantee:** If you are not satisfied with this product, we will gladly refund your original purchase price.



A Business Unit of Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709

Lawn Herbicide JD Concentrate (MASTER) Approved 09/17/07, Notification 06/26/08, Submitted Corrected Label 01/22/09