

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

July 29, 2016

Norma C. Pangilinan, Ph.D. Registration Manager Bayer Advanced P.O. Box 12014 2TW Alexander Drive Research Triangle Park, NC 27709

Subject: Label Amendment – Addition of NY State Restrictions and minor format changes

Product Name: TEBUCONAZOLE 2.9% SE CONCENTRATE SYSTEMIC

FUNGICIDE

EPA Registration Number: 72155-14

Application Date: 04/21/2016 Decision Number: 516635

Dear Ms. Pangilinan:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Page 2 of 2 EPA Reg. No. 72155-14 Decision No. 516635

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Maryam K. Muhammad by phone at 703-347-0301, or via email at Muhammad.maryam@epa.gov.

Sincerely,

Hope Johnson, Product Manager 21 Fungicide Branch Registration Division (7505P) Office of Pesticide Programs

Enclosure

(FRONT PANEL)

Tebuconazole 2.9% SE Concentrate Systemic Fungicide

This label contains two separate label statements for, Outdoor Residential Pump Sprayer/Hose-end Sprayer Concentrate product use and Outdoor Residential Ready-To-Spray* product use.

- [One quart] makes [up to] 42 gallons [diluted] spray
- For control of [listed] [*] diseases on plants
- Controls and prevents [listed] [*] common diseases on [roses,] [flowers,] [trees,] [and] [shrubs]
- Controls all [listed] [*] fungal diseases that harm [roses,] [flowers,] [trees,] [and] [shrubs]
- Rescues plants from diseases listed on label
- Systemic fungicide works through the leaves and stems to protect and cure
- Provides up to 2 weeks protection
- Rainproof Protection won't wash off once it is dry
- Leaves no white residue
- Protects from the inside out
- Controls [listed] [*] diseases and restores beauty to plants
- 100% Satisfaction Guaranteed
- Systemic formula controls and [Prevents] [Cures] [listed] [*] diseases
 - Black Spot
 - Powdery Mildew
 - Rust & Other listed diseases

*For use with garden hose

*listed (Note to PM: The word "listed" can appear as a footnote with the appropriate phrase, *listed pests/diseases)

ACTIVE INGREDIENT:

Tebuconazole	2.9%
OTHER INGREDIENTS:	<u>97.1%</u>
TOTAL:	100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

See Back Panel Booklet for Additional Precautionary Statements

Not for sale, distribution or use in New York State.

ACCEPTED

Jul 29, 2016

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

72155-14

FIRST AID		
If swallowed:	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. 	
	 Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. 	
If on skin or clothing:	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice. 	
If inhaled:	 If breathing is affected, move person to fresh air. If not breathing, call an ambulance and give artificial respiration. Contact doctor or poison control center for further treatment advice. 	

NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 877-229-3763 for medical emergency information.

(Pump Sprayer/Hose-end Sprayer label) (FRONT PANEL)

Tebuconazole 2.9% SE Concentrate Systemic Fungicide

- [One quart] makes [up to] 42 gallons [diluted] spray
- For control of [listed] [*] diseases on plants
- Controls and prevents [listed] [*] common diseases on [roses,] [flowers,] [trees,] [and] [shrubs]
- Controls all [listed] [*] fungal diseases that harm [roses,] [flowers,] [trees,] [and] [shrubs]
- Rescues plants from diseases listed on label
- Systemic fungicide works through the leaves and stems to protect and cure
- Provides up to 2 weeks protection
- Rainproof Protection won't wash off once it is dry
- · Leaves no white residue
- · Protects from the inside out
- Controls [listed] [*] diseases and restores beauty to plants
- 100% Satisfaction Guaranteed
- Systemic formula controls and [Prevents] [Cures] [listed] [*] diseases
 - Black Spot
 - Powdery Mildew
 - Rust & Other listed diseases

*listed (Note to PM: The word "listed" can appear as a footnote with the appropriate phrase, *listed pests/diseases)

ACTIVE INGREDIENT:	
Tebuconazole	2.9%
OTHER INGREDIENTS:	<u>97.1%</u>
TOTAL:	100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

See Back Panel Booklet for Additional Precautionary Statements

Not for sale, distribution or use in New York State.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice. If inhaled: If breathing is affected, move person to fresh air. If not breathing, call an ambulance and give artificial respiration. Contact doctor or poison control center for further treatment advice.	FIRST AID		
 Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice. If inhaled: If breathing is affected, move person to fresh air. If not breathing, call an ambulance and give artificial respiration. 	If swallowed:	 Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. 	
If not breathing, call an ambulance and give artificial respiration.	If on skin or clothing:	Rinse skin immediately with plenty of water for 15 to 20 minutes.	
	If inhaled:	If not breathing, call an ambulance and give artificial respiration.	

NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 877-229-3763 for medical emergency information.

(BACK PANEL)

TEBUCONAZOLE 2.9% SE CONCENTRATE controls, stops and prevents plant diseases. This systemic formula provides rainproof protection against most common plant diseases such as black spot, powdery mildew, and rust. Now it is easier than ever to prevent diseases and have beautiful plants.

Read the label first!

QUICK FACTS

- Protects against [listed] [*] diseases on roses, flowers, trees, and shrubs
- Controls and prevents listed common [listed] [*] diseases on roses, flowers, trees, and shrubs
- Controls [listed] [*] diseases and restores beauty to plants

CONTROLS	Anthracnose, Black Spot, Flower Blight, Leaf Spot, Petal Blight, Powdery Mildew, Rust, Scab, Southern Blight, Boxwood Blight	
WHERE TO USE	Roses and Flowers	
	Azaleas, Camellias, and Rhododendrons	
	Landscape Trees & Shrubs	
	Ground Covers & Vines	
	Houseplants	
WHEN TO USE	To treat an existing disease	
	To prevent and protect against future diseases	
COVERAGE	Makes up to 42 gallons of spray solution (1 Qt.)	
	For questions or comments, call toll-free 1-877-BayerAG	

FOR OUTDOOR RESIDENTIAL USE ONLY

ACTIVE INGREDIENT:

Tebuconazole	2.9%
OTHER INGREDIENTS	<u>97.1%</u>
TOTAL	00.00%

EPA Reg. No. 72155-14 EPA Est. No. _-AA-_

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

- Harmful if swallowed, inhaled, or absorbed through skin.
- Washing hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco will
 reduce exposure.
- · Avoid contact with skin, eyes, or clothing.
- · Removing and washing contaminated clothing before reuse may reduce exposure.
- · Avoid breathing vapor or spray mist.
- When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

ENVIRONMENTAL HAZARDS

- This product is toxic to aquatic invertebrates.
- Drift or runoff may be hazardous to aquatic organisms in water adjacent to treated areas.
- To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters.
 Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

	DIRECTIONS FOR USE			
	aw to use this product in a manner inconsistent with its labeling.			
For best results, read and for	ollow all label directions.			
BEFORE YOU USE	Do not apply near lakes, streams, rivers, or ponds.			
	Do not apply to plants grown for food.			
	Food utensils such as measuring spoons and measuring cups must not be used for food purposes after use with fungicides.			
	Move indoor houseplants outside before treating.			
HOW TO USE	Determine how much spray solution to make up – See "How Much to Use".			
Shake Well Before Use	Make up spray solution by following directions specific to your sprayer – See "Sprayer Types".			
	Wet all plant surfaces until evenly coated with spray, but not dripping.			
HOW MUCH TO USE	Add ¾ fl. oz. (1 ½ TBS) to 1 gallon of water			
SPRAYER TYPES	Pump Style Sprayers			
	Add the appropriate amounts of concentrate and water to the sprayer tank.			
	2. Close sprayer, shake well and pressurize.			
	3. Adjust nozzle to a coarse spray pattern and apply.			
	4. Occasionally re-pressurize the sprayer if needed to maintain a good spray pattern.			
	Dial Style Hose-End Sprayers			
	1. Fill sprayer jar with enough concentrate for the area to be treated. DO NOT ADD WATER.			
	2. Close sprayer and set dial as recommended below:			
	- Ortho Dial 'N Spray Setting: 1 TBSP			
	- Chameleon Hose-End Sprayer Setting: 1 TSP			
	TIP: The numbers on dial indicate amount of concentrate per gallon of spray.			
	3. Hook to garden hose and apply.			
	To avoid contact with the treated area, begin by spraying area farthest from faucet walking backwards as you spray.			
	4. Pour any unused product back into this original container.			
	Concentrate Plus Water Hose-End Sprayer			
	Pour concentrate into sprayer jar to the required fluid ounce level.			
	2. Add the appropriate amount of water to the gallons level.			
	Replace sprayer top and shake well.			

PLANTS	CO	NTF	ROLS PLANT	WHEN TO APPLY	
		4.	Attach sprayer t	o hose and apply.	
		٠.	. topiass spiays	top and onanto from	

PLANTS	CONTROLS PLANT	WHEN TO APPLY			
	DISEASES	To Treat Existing Disease	To Prevent Diseases		
Roses	Black Spot Powdery Mildew Rust	Apply every 14 days for a total of 3 applications beginning at the first sign of disease. Single application rate is	Apply every 14 days during the growing season, starting when leaves first appear.		
Flowers	Leaf Spot Powdery Mildew Rust Southern Blight	0.0014 lb a.i./gal.	Apply no more than 3 times per year, 14 days apart, beginning with Spring bud break. Single application rate is 0.0014 lb a.i./gal.		
Crabapples (Ornamental), Dogwoods and Other Landscape [Ornamental] Trees	Anthracnose Leaf Spot Powdery Mildew Rust Scab		Petal Blight – Apply 2-3 times per week into the flowers as they open and develop color.		
Azaleas, Camellias, Rhododendrons and Other Landscape [Ornamental] Shrubs,	Anthracnose Black Spot Leaf Spot Petal Blight Powdery Mildew				
Ground Covers and Vines	Rust Southern Blight				

Houseplants	Leaf Spot Powdery Mildew		Apply no more than 3 times per year, 14 days apart.
Boxwood	Boxwood Blight	Contact your local Extension Office for disease confirmation and if positive, destroy the infected plant(s) in a manner that will minimize exposure to other boxwoods.	Apply this product 3 times at 14 day intervals when conditions are conducive to disease development (64 – 80°F, high humidity/rain/overhead irrigation). Single application rate is 0.0014 lb a.i./gal.
FOR BEST RESULTS	 Apply at the shorter interval specified under "Directions for Use" when disease or conditions favorable to disease (generally cloudy, wet and warm weather) are present. Prevent damage to your plants caused by disease by starting a spray schedule when diseases are likely in your area. For more information on what you can do to control plant diseases, contact your local County Extension Service (usually found in the White Pages of the phone book under the name of your county.) 		
HOW [IT] [TEBUCONAZOLE 2.9% SE CONCENTRATE] WORKS		R-way action: it kills existing fungi, forms a poto the plant to keep on protecting regardles	
RESTRICTIONS	Do not exceed 3 a	pplications per year.	



Do not allow people and pets to enter the treated area until sprays have dried.

Do not contaminate wat	ter, food, or feed by storage or disposal.		
Pesticide Storage:	Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing.		
Container Handling:	Nonrefillable Container. Do not reuse or refill this container.		
	If Empty:		
	Place empty container in trash or offer for recycling, if available.		
	If Partly Filled:		
	 If it is necessary to dispose of unused product, please call your local solid waste agency or you may call toll-free 1-877-229-3724 for disposal instructions. 		
	Never place unused product down any indoor or outdoor drain.		

NOTICE: Research and testing have determined that the "Directions For Use" are appropriate for the proper use of TEBUCONAZOLE 2.9% SE CONCENTRATE SYSTEMIC FUNGICIDE under expected conditions. The Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.

Net Contents: 16 fl. oz. (473 mL) or 32 fl. oz. (946 mL)

(Ready-to-Spray label*) (FRONT PANEL)

Tebuconazole 2.9% SE Concentrate Systemic Fungicide

- [One quart] makes [up to] 42 gallons [diluted] spray
- For control of [listed] [*] diseases on plants
- Controls and prevents [listed] [*] common diseases on roses, flowers, trees, and shrubs
- Controls all [listed] [*] fungal diseases that harm roses, flowers, trees, and shrubs
- Rescues plants from diseases listed on label
- · Systemic fungicide works through the leaves and stems to protect and cure
- Provides up to 2 weeks protection
- · Rainproof Protection won't wash off once it is dry
- Leaves no white residue
- · Protects from the inside out
- Controls [listed] [*] diseases and restores beauty to plants
- 100% Satisfaction Guaranteed
- Systemic formula controls and [Prevents] [Cures] [listed] [*] diseases
 - Black Spot
 - Powdery Mildew
 - Rust & Other listed diseases

TOTAL: 100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

See Back Panel Booklet for Additional Precautionary Statements

Not for sale, distribution or use in New York State.

*For use with garden hose

FIRST AID		
If swallowed:	Call a poison control center or doctor immediately for treatment advice.	
	Have person sip a glass of water if able to swallow.	
	Do not induce vomiting unless told to do so by a poison control center or doctor.	
	Do not give anything by mouth to an unconscious person.	
If on skin or clothing:	Take off contaminated clothing.	
	Rinse skin immediately with plenty of water for 15 to 20 minutes.	
	Call a poison control center or doctor for treatment advice.	
If inhaled:	If breathing is affected, move person to fresh air.	
	If not breathing, call an ambulance and give artificial respiration.	
	Contact doctor or poison control center for further treatment advice.	

NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 877-229-3763 for medical emergency information.

(BACK PANEL)

TEBUCONAZOLE 2.9% SE CONCENTRATE controls, stops and prevents plant diseases. This systemic formula provides rainproof protection against most common plant diseases such as black spot, powdery mildew, and rust. Now it is easier than ever to prevent diseases and have beautiful plants.

Read the label first!

QUICK FACTS

- · Protects against listed diseases on roses, flowers, trees, and shrubs
- Controls and prevents listed common diseases on roses, flowers, trees, and shrubs
- · Controls listed diseases and restores beauty to plants
- Ready-to-Spray ... No Mixing, No Measuring

, , ,	3
CONTROLS	Anthracnose, Black Spot, Flower Blight, Leaf Spot, Petal Blight, Powdery Mildew, Rust, Scab, Southern Blight, Boxwood Blight
WHERE TO USE	Roses and Flowers
	Azaleas, Camellias and Rhododendrons
	Landscape Trees & Shrubs
	Ground Covers & Vines
	Houseplants
WHEN TO USE	To treat an existing disease
	To prevent and protect against future diseases
COVERAGE	No measuring or mixing needed. Dilutes automatically as you spray.
	Makes up to 42 gallons of spray solution (1 qt.)
	For questions or comments, call toll-free 1-877-BayerAG.

FOR OUTDOOR RESIDENTIAL USE ONLY

ACTIVE INGREDIENT:

Tebuconazole	2.9%
OTHER INGREDIENTS	<u>97.1%</u>
TOTAL	100.00%

EPA Reg. No. 72155-14 EPA Est. No. _-AA-_

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

- · Harmful if swallowed, inhaled or absorbed through skin.
- Washing hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco will
 reduce exposure.
- · Avoid contact with skin, eyes, or clothing.
- Removing and washing contaminated clothing before reuse may reduce exposure.
- · Avoid breathing vapor or spray mist.
- · When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

ENVIRONMENTAL HAZARDS

- This product is toxic to aquatic invertebrates.
- Drift or runoff may be hazardous to aquatic organisms in water adjacent to treated areas.
- To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

DIRECTIONS FOR USE					
It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. For best results, read and follow all label directions.					
·	T				
BEFORE YOU USE		lakes, streams, rivers, or ponds.			
		nts grown for food.			
	Food utensils such after use with fung	as measuring spoons and measuring cups icides.	s must not be used for food purposes		
	Move indoor house	eplants outside before treating.			
HOW TO USE Shake Well Before Use		spray. Just attach this bottle to your hose the concentrate as you spray.	according to directions. Water		
HOW TO ATTACH AND SPRAY	Connect your gard nozzle is in the OF	len hose to the bottle. Be sure the blue water F position.	er control valve on top of the spray		
		aucet. Twist the small star-shaped, blue prock tab. Push the button in to open.	oduct-control button to align the flat		
	3. To begin spraying	point nozzle toward the area to be treated ab will be horizontal.	and turn the blue water control valve to		
	4. To stop spraying,				
	5. Relieve water pres	ssure by turning blue water control valve to ter control valve back to OFF. Disconnect g	the ON position until water slows to a		
	-	n can be saved and used at a later date.	garden need nem betae.		
	Tips for Applying				
		e farthest area to be treated and work back	toward the faucet to avoid walking into		
	1	ace while spraying, using an even sweepin	ng motion, slightly overlapping the		
PLANTS	CONTROLS PLANT WHEN TO APPLY				
	DISEASES	To Treat Existing Disease	To Prevent Diseases		
Roses	Black Spot Powdery Mildew Rust	Apply every 14 days for a total of 3 applications beginning at the first sign of disease. Single application rate is	Apply every 14 days during the growing season, starting when leaves first appear.		
Flowers	Leaf Spot Powdery Mildew Rust Southern Blight	0.0014 lb a.i./gal.	Apply no more than 3 times per year, 14 days apart, beginning with Spring bud break. Single application rate is 0.0014 lb a.i./gal.		
Crabapples (Ornamental), Dogwoods and Other Landscape [Ornamental] Trees	Anthracnose Leaf Spot Powdery Mildew Rust Scab		Petal Blight – Apply 2-3 times per week into the flowers as they open and develop color.		
Azaleas, Camellias, Rhododendrons and Other Landscape [Ornamental] Shrubs, Ground Covers and Vines	Anthracnose Black Spot Leaf Spot Petal Blight Powdery Mildew Rust Southern Blight				
Houseplants	Leaf Spot Powdery Mildew		Apply no more than 3 times per year, 14 days apart.		
Boxwood	Boxwood Blight	Contact your local Extension Office for disease confirmation and if positive, destroy the infected plant(s) in a manner that will minimize exposure to other boxwoods.	Apply this product 3 times at 14 day intervals when conditions are conducive to disease development (64 – 80°F, high humidity/rain/overhead irrigation). Single application rate is 0.0014 lb a.i./gal.		

FOR BEST RESULTS	 Apply at the shorter interval specified under "Directions for Use" when disease or conditions favorable to disease (generally cloudy, wet and warm weather) are present. Prevent damage to your plants caused by disease by starting a spray schedule when diseases are likely in your area. For more information on what you can do to control plant diseases, contact your local County Extension Service (usually found in the White Pages of the phone book under the name of your county.) 	
HOW TEBUCONAZOLE 2.9% SE CONCENTRATE WORKS	This product provides 3-way action: it kills existing fungi, forms a protective barrier on the outside of the plant and is absorbed into the plant to keep on protecting regardless of weather conditions (up to 2 weeks).	
RESTRICTIONS	Do not exceed 3 applications per year.	

₹≒	Do not allow people and pets to enter the treated area until sprays have dried.
	STORAGE AND DISPOSAL
Do not contaminate wat	ter, food, or feed by storage or disposal.
Pesticide Storage:	Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing.
Container Handling:	Nonrefillable Container. Do not reuse or refill this container. If Empty: Place empty container in trash or offer for recycling, if available.
	If Partly Filled:
	 If it is necessary to dispose of unused product, please call your local solid waste agency or you may call toll-free 1-877-229-3724 for disposal instructions.
	Never place unused product down any indoor or outdoor drain.

NOTICE: Research and testing have determined that the "Directions For Use" are appropriate for the proper use of TEBUCONAZOLE 2.9% SE CONCENTRATE SYSTEMIC FUNGICIDE under expected conditions. The Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.

Net Contents: 16 fl. oz. (473 mL) or 32 fl. oz. (946 mL)

Produced for



A Business Unit of Bayer CropScience LP P.O. Box 12014, 2 T.W. Alexander Drive Research Triangle Park, North Carolina 27709

Tebuconazole 2.9% SE Concentrate (PENDING) 10/09/2015, 03/30/2016, 04/06/2016, 04/11/2016, 04/15/2016, 07/26/2016, 07/28/2016