

ACCEPTED

Beta-cyfluthrin 0.0015% Ready-to-Use Insecticide

For use on: Flowers, Shrubs, Groundcovers, Lawns, Around Homes, and Vegetables

GUARANTEED SATISFACTION

READY TO USE

KILLS INSECTS

MULTIPURPOSE INSECTICIDE

RAINPROOF within 1 hour of application

DOUBLE PROTECTION FORMULA

- Fast Acting...Kills On Contact
- Kills on contact plus protects for 2 weeks (14 days)
- Long Lasting...Keeps On Killing For Up To 14 Days

FAST ACTING LONG LASTING

- Kills outdoor pests on contact
- Kills over 60 different outdoor pests
- Lasts up to 2 weeks (14 days)
- · For use on flower and vegetable gardens, shrubs, lawns, and around homes.
- Just One Application Quickly Kills Insects
- Keeps On Killing For Up To 2 weeks (14 Days)
- Use on tomatoes up to day of harvest
- Contains Tempo Ultra®

ACTIVE INGREDIENT:

Net Contents: 24 fl oz

STOP - Read This Entire Label Before Use **KEEP OUT OF REACH OF CHILDREN**

CAUTION

See back panel / booklet for additional precautionary statements.

(BACK PANEL)

This product kills caterpillars, cutworms and other nuisance pests on contact and keeps protecting for 2 weeks (14 days) in a single application. No mixing and measuring hassles. No messy cleanup. Controlling annoying outdoor insects around your home and in your garden has never been easier.

QUICK FACTS

- Ready to Use...No Mixing, No Measuring
- Quickly Stops Harmful Insects Before They Destroy Plants
- . May be used up to day of harvest on tomatoes
- Effective Against Broad Range of Listed Pests
- Quickly Stops Harmful Insects Before They Do Damage
- Kills Household Nuisance Pest Outdoors

KILLS	(ART: aphid, cabbage looper, whitefly)						
•	Annual Bluegrass weevil	Elm leaf beetles	Potato aphids				
	(hyperodes weevil)	Flea beetles	Potato leafhoppers				
	Ants	Flies	Sawflies				
	Aphids	Garden webworms	Scale (crawler stages)				
	Armyworms	Grasshoppers	Seed weevils (adults)				
	Aster leafhoppers	Gypsy moth larvae	Sod webworms				
	Bagworms	Headclipper weevils (adults)	Sowbugs				
	Banded sunflower moths	Japanese beetles (adults)	Spiders				
	Boxelder bugs	Lace bugs	Spittlebugs Stem weevil (adults) Stink bugs Sunflower beetles Sunflower moths Tomato fruitworms Tomato homworms Tomato pinworms Thrips Wasps Webworms Weevils (including root weevils) Western flower thrips Whiteflies				
	Budmoths	Leaf beetles					
	Cabbage Loopers (adults)	Leaf-feeding beetles					
	Carrot Weevils	Leaf-feeding caterpillars					
	Caterpillars	Leafhoppers (including Glassy-					
	Celery Leaftiers	Winged Sharpshooter)					
	Chinch bugs Clover mites Cockroaches Colorado potato beetles Corn borers Corn earworms Corn rootworms (adults) Crickets Cutworms	Leafminers Mealybugs Midges Mosquitoes Nuisance beetles Pepper weevils (adults) Pillbugs Pine Shoot Moths Pine Tip Moths Plant bugs					
WHERE TO USE WHEN TO USE							
				Earwigs			
				Flower and vegetable gardens,	Flower and vegetable gardens, shrubs, ground covers, lawns, and around homes When insects or damage first appear For severe infestations, repeat every 7 to 14 days as necessary.		
				When insects or damage first a			
				For severe infestations, repeat			
				Questions or comments, call to	II-free (877) 229-3724		
	70. 💾	www.BayerAdvanced.com					

ACTIVE INGREDIENT:

Beta-cyfluthrin	0.0015%
OTHER INGREDIENTS:	99.9985%
	100.000 β-

Beta-cyfluthrin is the chemical name for Tempo Ultra®

FOR OUTDOOR RESIDENTIAL USE ONLY

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

- Harmful if swallowed or inhaled.
- · Causes eye irritation.
- Avoid contact with skin, eyes, or clothing.
- Avoid breathing spray mist.
- · Wash thoroughly with soap and water after handling.

I	◆ First Aid
If Swallowed	Call a poison control center or doctor immediately for treatment advice.
	Have person sip a glass of water if able to swallow.
	Do not induce vomiting unless told to do so by a poison control center or doctor.
	Do not give anything by mouth to an unconscious person.
If in eyes	Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
	Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes.
	Call a poison control center or doctor for treatment advice.
If Inhaled	Move person to fresh air.
	 If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
	Call a poison control center or doctor for further treatment advice.

NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's affice, bring this product label to show medical personnel. You may call toll-free (877) 229-3763 for medical emergency information.

Active Ingredient: 0.0015% beta-cyfluthrin [CAS # 68359-37-5]

Note to Physician: No specific antidote is available. Treat the patient symptomatically.

Environmental Hazards

- · This pesticide is toxic to fish. Do not apply directly to water.
- This pesticide is highly toxic to bees. Do not apply Beta-cyfluthrin 0.0015% Ready-to-Use Insecticide when bees are active.
- · Apply this product only as specified on this label.

DIRECTIONS FO	OR USE	
It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For best results, read and follow all label directions.		
BEFORE YOU USE Read and follow these directions when using:	 Apply the product to flowering plants during early morning or late evening when bees are not present. Do not spray the product into fish pools, ponds, streams, or lakes. Do not use in small, enclosed areas where spray can be inhaled. 	
WHEN TO USE	When insects or damage first appear.	
HOW TO USE Shake well before using.	To apply Beta-cyfluthrin 0.0015% Ready-to-Use Insecticide: • Adjust nozzle to desired spray pattern. • Hold sprayer about 12 inches from foliage. • Spray leaves, stems, and tender new shoots. • To kill insects that may hide, spray the underside of the leaves. Apply until leaves are evenly coated with spray, but not dripping. (ART: spraying undersides of leaves)	
HOW OFTEN TO USE	Kills insects on contact and keeps on killing for weeks. For severe infestations, repeat every 7 to 14 days as necessary.	

PEST			
Aphids	Grasshoppers	Leaf-feeding beetles	Scale
Bagworms	Gypsy moth larvae	Leafhoppers (including	(crawler stages)
Boxelder bugs	Japanese beetle (adults)	Glassy-winged Sharpshooter)	Spittlebugs
Caterpillars	Lace bugs	Mealybugs	Thrips
Clover mites	Leaf-feeding caterpillars	Pillbugs	Webworms
Elm leaf beetles	and the second of the second o	Sawflies	Weevils (including root weevils)
Limited beeties			Whiteflies
HOW TO APPLY			

Spray foliage, tender new shoots and the underside of leaves thoroughly.

To kill insects that hide, spray the underside of the leaves. Apply until leaves are evenly coated with spray droplets, but not dripping.

(ART)

VEGETABLES			200
CARROTS, RADISHES,	SWEET CORN, TOMATOES		
 No Waiting. 			
 You may spray u 	ip to day of harvest.		
 Do not spray mo 	re than 5 times per season.		
PEST			
Aphids	Colorado potato beetles	Grasshoppers	Thrips
Armyworms	Corn borers	Leafhoppers (including	Tomato fruitworms
Aster leafhoppers	Corn earworms	Glassy-winged Sharpshooter)	Tomato hornworms
Cabbage loopers	Corn rootworm (adults)	Leafminers	Tomato pinworms
Carrot weevils (adult)	Cutworms	Potato Aphids	Weevils (adult)
Chinch bugs	Flea Beetles	Stink bugs	Western flower thrips

PEPPERS			
 Do not spray more 	t 7 days before harvesting. than 6 times per season.		
PEST			<u> </u>
Armyworms	Corn borers	Leafhoppers (including	Potato leafhoppers
Cabbage loopers	Corn earworms	Glassy-winged Sharpshooter)	Thrips
Celery leaftiers	Garden webworms	Leafminers	Weevils (adult)
•		Pepper weevils (adult)	
SUNFLOWERS		turi (s fra	
 After spraying, wa 	t 30 days before harvesting.		
	than 3 times per season.	 	***
PEST		· · · ·	·
Banded sunflower moths	Grasshoppers	Midges	Sunflower beetles
Budmoths	Headclipper weevils (adult)	Seed weevils (adult)	Sunflower moths
Cutworms	Leafhoppers (including Glassy-winged Sharpshooter)	Stem weevils (adult)	Weevils (adult)

FOR BEST RESULTS:

Do not water the foliage within one hour after applying. To avoid serious damage to plants, eliminate pests early in the season before they multiply.

HOUSEHOLD NUISANCE PESTS OUTDOORS			
Ants	Crickets	Midges	Sowbugs
Boxelder bugs	Earwigs	Mosquitoes	Spiders
Clover mites	Flies	Nuisance beetles	Wasps
Cockroaches		Pillbugs	

Spray around doors, window frames and other places where insects may enter house. Spray around foundations and porches. Spray on ant hills and ant trails.

'ART)

PEST Ants Crickets Earwigs Japanese beetles (adults) Armyworms Cutworms Grasshoppers Sod webworms (lawn moths) Chinch bugs HOW TO APPLY For localized infestations, spray to penetrate the lawn and reach the soil. Do not mow for one day after application. For large areas, use Beta-cyfluthrin 0.38% Concentrate Insecticide.

₹₩

Children and pets may re-enter the treated area after spray has dried.

Storage Conditions:	Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing.
Disposal:	If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Important: It is best to use all of the product in accordance with label directions. If it is necessary to dispose of unused product, please follow any applicable state or local guidelines.

NOTICE

Research and testing have determined that the "Directions for Use" are appropriate for the proper use of Beta-cyfluthrin 0.0015% Ready-to-Use Insecticide under expected conditions. The Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

ን-9806 110804

Bayer Advanced LLC, A Business Unit of CropScience LP 95 Chestnut Ridge Road Montvale, NJ 07645