


[FRONT PANEL]

Imidacloprid 0.012% +Beta-cyfluthrin 0.0015% Ready-to-Use Insecticide

KILLS INSECTS

For use on: Flowers, Roses, Shrubs

- Just one application quickly kills insects
- Protects up to 30 days
- Rainproof Within 1 Hour of Application
- Exclusive [insert trademark]™ Formula (Technology)
- Next Generation (of)Insect Control
- More powerful (effective)
- Long-Lasting Protection
- Kills surface pests in (24) hours
- Leading(cutting edge) insect control
- Leading (Edge) (insect control) Technology
- Fast Acting Formula
- 2 way (action) Formula
- Powerful Fast Acting (2-Way) Formula



MERIT + TEMPO Ready-to-Use Insecticide is a highly effective insecticide designed to kill insect pests on

- flowers
- roses
- house plants
- ground covers
- shrubs

Contains Merit® and Tempo Ultra®

ACTIVE INGREDIENTS:

Beta-cyfluthrin	0.0015 %
Imidacloprid.....	0.012 %

OTHER INGREDIENTS:

.....	99.9865 %
-------	-----------


Net Contents: 24 fl oz

284

STOP - Read This Entire Label Before Use
KEEP OUT OF REACH OF CHILDREN
 FOR RESIDENTIAL USE ONLY

CAUTION

See back panel booklet for ingredients and precautionary statements
 [BACK PANEL]

QUICK FACTS																																													
<ul style="list-style-type: none"> • Kills Insects on Contact • Keeps on Killing for Up to 30 Days • Ready to Use -- No Mixing, No Measuring 																																													
KILLS	<p align="center">(ART: Aphid, Japanese Beetle, Leafminer, Whitefly)</p> <table border="0"> <tr> <td>Adelgids</td> <td>Flies</td> <td>Mealybugs</td> <td>Rose Slugs</td> </tr> <tr> <td>Ants</td> <td>Fungus gnats</td> <td>Midges</td> <td>Sawflies</td> </tr> <tr> <td>Aphids</td> <td>Grasshoppers</td> <td>Mosquitoes</td> <td>Scale insects (crawler stages)</td> </tr> <tr> <td>Bagworms</td> <td>Gypsy moth larvae</td> <td>Pillbugs</td> <td>Sowbugs</td> </tr> <tr> <td>Boxelder bugs</td> <td>Japanese beetles (adult)</td> <td>Pine Shoot moths</td> <td>Spiders</td> </tr> <tr> <td>Caterpillars</td> <td>Lace bugs</td> <td>Pine Tip moths</td> <td>Spittlebugs</td> </tr> <tr> <td>Clover mites</td> <td>Leaf-feeding beetles</td> <td>Plant bugs</td> <td>Thrips</td> </tr> <tr> <td>Cockroaches</td> <td>Leaf-feeding caterpillars</td> <td>Psyllids</td> <td>Wasps</td> </tr> <tr> <td>Crickets</td> <td>Leafhoppers (including Glassy-Winged Sharpshooter)</td> <td>Rose Midges</td> <td>Webworms</td> </tr> <tr> <td>Earwigs</td> <td>Leafminers</td> <td></td> <td>Weevils (including root weevils)</td> </tr> <tr> <td>Elm leaf beetles</td> <td></td> <td></td> <td>Whiteflies</td> </tr> </table>	Adelgids	Flies	Mealybugs	Rose Slugs	Ants	Fungus gnats	Midges	Sawflies	Aphids	Grasshoppers	Mosquitoes	Scale insects (crawler stages)	Bagworms	Gypsy moth larvae	Pillbugs	Sowbugs	Boxelder bugs	Japanese beetles (adult)	Pine Shoot moths	Spiders	Caterpillars	Lace bugs	Pine Tip moths	Spittlebugs	Clover mites	Leaf-feeding beetles	Plant bugs	Thrips	Cockroaches	Leaf-feeding caterpillars	Psyllids	Wasps	Crickets	Leafhoppers (including Glassy-Winged Sharpshooter)	Rose Midges	Webworms	Earwigs	Leafminers		Weevils (including root weevils)	Elm leaf beetles			Whiteflies
Adelgids	Flies	Mealybugs	Rose Slugs																																										
Ants	Fungus gnats	Midges	Sawflies																																										
Aphids	Grasshoppers	Mosquitoes	Scale insects (crawler stages)																																										
Bagworms	Gypsy moth larvae	Pillbugs	Sowbugs																																										
Boxelder bugs	Japanese beetles (adult)	Pine Shoot moths	Spiders																																										
Caterpillars	Lace bugs	Pine Tip moths	Spittlebugs																																										
Clover mites	Leaf-feeding beetles	Plant bugs	Thrips																																										
Cockroaches	Leaf-feeding caterpillars	Psyllids	Wasps																																										
Crickets	Leafhoppers (including Glassy-Winged Sharpshooter)	Rose Midges	Webworms																																										
Earwigs	Leafminers		Weevils (including root weevils)																																										
Elm leaf beetles			Whiteflies																																										
COVERAGE	Treats up to 34 medium-sized plants																																												
WHERE TO USE	Flowers, roses and shrubs																																												
WHEN TO USE	When insects or damage first appear																																												
	Questions or comments, call toll-free (877) 229-3724 www.BayerAdvanced.com																																												

ACTIVE INGREDIENTS:

Beta-cyfluthrin 0.0015 %
 Imidacloprid 0.012 %

OTHER INGREDIENTS:* 99.9865 %
 100.000 %

* Other ingredients include: Water, glycerin (stabilizer), organic solvent (less than 0.1%, to dissolve the active ingredient for mixing) and surfactants (to keep the active ingredient dispersed in water).

Beta-cyfluthrin is the chemical name for Tempo Ultra® and Imidacloprid is the chemical name for Merit®.

EPA Reg. No. 72155-xxx

EPA Est. No. XXXX-AA-1

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

- Harmful if swallowed or inhaled.
- Causes eye irritation.
- Avoid contact with skin, eyes, or clothing.
- Avoid breathing spray mist.
- Wash thoroughly with soap and water after handling.

First Aid

IN THE CASE OF ACCIDENTAL...

IMMEDIATELY...

Swallowing

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

Contact with eyes

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes.
- Call a poison control center or doctor for treatment advice.

Inhalation

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouthy-to-mouth if possible.
- Call a poison control center or doctor for further treatment advice

NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. For medical emergency information, you may call toll-free 1-877-229-3763.

Active Ingredients: 0.0015% Beta-cyfluthrin [CAS # 68359-37-5], 0.012% Imidacloprid [CAS # 138261-41-3]

Environmental Hazards

- This pesticide is highly toxic to aquatic invertebrates and fish. Do not apply directly to water.
- This pesticide is highly toxic to bees. Do not apply Imidacloprid 0.012%+Beta-cyfluthrin 0.0015% Ready-to-Use Insecticide when bees are active.
- Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For best results, be sure to read and follow all label directions.

BEFORE YOU USE

Read and follow these directions when using.

- Apply the product to flowering plants during early morning or late evening, when bees are not present.
- Do not spray the product into fish pools, ponds, streams, or lakes.
- Remove animal feeding dishes prior to application.
- Do not use in small, enclosed areas where spray may be inhaled.
- Do not apply to fruits or vegetables used for food.

WHEN TO USE

When insects or damage first appear
To avoid serious damage, eliminate pests early in the season before they multiply.

HOW TO USE

- Adjust nozzle to desired spray pattern.
- Hold sprayer about 12 inches from foliage.


4 8 4

Shake well before using.	<ul style="list-style-type: none"> Spray leaves, stems and tender new shoots. (ART: of steps above) To kill insects that hide, spray the underside of the leaves. Apply until leaves are evenly coated with spray, but not dripping. (ART: spraying undersides of leaves)
HOW OFTEN	Kills insects on contact and keeps on killing for up to 30 days. For severe infestations, repeat every 7 to 14 days as necessary.

FOR BEST RESULTS

Do not wet the foliage within one hour after applying. To avoid serious damage to plants, eliminate pests early in the season before they multiply.

	Children and pets may re-enter after spray has dried.
---	---

IMPORTANT: Directions for Storage and Disposal	
Storage Conditions:	Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing.
Disposal: 	<ul style="list-style-type: none"> Do not reuse empty spray bottle. Rinse the empty product container thoroughly. Place container in trash.
<i>Important:</i> It is best to use all of the product in accordance with label directions. If it is necessary to dispose of unused product, please follow any applicable state or local guidelines.	

NOTICE

Research and testing have determined that the "Directions for Use" are appropriate for the proper use of Imidacloprid 0.012%+Beta-cyfluthrin 0.0015% Ready-to-Use Insecticide under expected conditions. The Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

D-9771 100803

Bayer Advanced LLC,
 A Business Unit of CropScience LP
 95 Chestnut Ridge Road
 Montvale, NJ 07645