

U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (H7505C) 401 "M" St., S.W. Washington, D.C. 10460

EPA Reg. Number 72155-60 base of lasuance:

MAR 1 8 2005

NOTICE OF PESTICIDE:

X Registration Reregistration

(under FIFRA, as amended)

Term of Issuance: Conditional

Name of Pesticide Product:

Trimec® + Prodiamine Granular Herbicide Plus Lawn Fertilizer

Name and Address of Registrant (include ZIP Code):

Bayer Advanced 95 Chestnut Ridge Road Montvale, NJ 07645

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insectitide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been towered by others.

This product is registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Add the phrase "EPA Registration No.72155-60" the label before you release the product for shipment.

COMMENTS CONTINUED ON PAGE 2 OF THIS NOTICE OF REGISTRATION

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product under the enclosed stamped copy of the label constitutes acceptance of these conditions.

Enclosure

Joanne I. Miller Product Manager (23) Herbicide Branch Registration Division (7505C)

Figurature of Amprovince trivial:

Joanne D. Miller

1,8001

MAR 1 & 2005

Page 2 EPA Reg. No. 72155-60 Comments Continued:

- 2. Submit the storage stability data (EPA Guidelines Number 830.6317) and Corrosion Characteristic data (EPA Guidelines Number 830.6320) within one year from the date of this Notice of Registration.
- 3. Add to the front panel of the labeling: either "Not for Commercial Use" or "For Homeowner Use Only".
- 4. On the front panel, delete the claims:

"Contains Trimec®"

"Does not stain concrete"

"Two Weed Killers in One!"

"Stops Listed Weeds From Invading Your Lawn"

- 5. On page 5 delete the claim: "and other broadleaf weeds". Note: weeds must be identified for this pesticide product.
- 6. Submit one (1) copy of the final printed labeling before you release this product for shipment.
- 7. Submit and/or cite all data required for the registration of this product when the Agency requires all registrants of similar products to submit data; and submit acceptable responses required for reregistration of this product under FIFRA, section 4.

A Stamped copy of the label is enclosed for your records.

Trimec® + Prodiamine Granular Herbicide Plus Lawn Fertilizer

- READY-TO-Spread Granules
- · Kills Dandelion, Chickweed, Clover and Many Common Lawn Weeds
- Controls Broadleaf Weeds
- Controls over 200 Broadleaf Weeds
- Contains Trimec*
- Kills Weeds
- Easy-to-Use Granules
- Two Weed Killers in One!
 - o Prevents Crabgrass
 - Kills Broadleaf weeds
 - Kills the Weeds Feeds the Lawn
- Won't Harm Lawn (when used as directed)
- Prevents Crabgrass Before It Starts
- Effective Weed Prevention for up to 6 Weeks
- Prevents Crabgrass
- Stops Crabgrass Before It Starts
- Does not stain concrete
- Controls Listed Weed Grasses and Broadleaf Weeds
- Stops Listed Weeds From Invading Your Lawn

ACCEPTED with COMMENTS In EPA Letter Dated:

MAR 1 8 2005 Under the Federal Insecticide. Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

72155-60

| ACTIVE INGREDIENTS: | |
|---------------------------|--------|
| 2,4-D (CAS # 94-75-7) | 0.69% |
| | |
| | |
| Dicamba (CAS # 1918-00-9) | 0.06% |
| | 98.82% |

Net Weight: 16 or 48 Pounds

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional ingredients and precautionary statements

*Registered Trademark of PBI/Gordon Corporation

[BACK PANEL]

4/8

This product provides weed control by killing unsightly weeds, such as dandelion, chickweed, & clover, and stopping [preventing] crabgrass, foxtail, spurge, and oxalis. At the same time its exclusive Triple Action formula feeds continuously at every level for up to X months. And it's guaranteed not to burn. The result is a greener, thicker lawn.

READ THE LABEL FIRST! (Read the Label First Icon)

| Quick Facts | | | | |
|--|---|--|--|--|
| Feeds the Lawn and | ds the Lawn in One Easy Step Prevents Crabgrass in One Easy Step eds , and feeds the lawn in one easy step. sum | | | |
| FEEDS | X-X-X Provides Essential Nutrients to Grass, Roots, and Soil (for x months) | | | |
| KILLS | Dandelion, Chickweed, Clovers, Crabgrass, Foxtail, Spurge, Oxalis and Other Listed Broadleaf and Weed grasses. | | | |
| COVERAGE | Treats 5,000 sq. ft. (16 lbs.) | | | |
| | Treats 15,000 sq. ft. (48 lbs.) | | | |
| VHERE TO USE | On lawns (Do not use on Bentgrass, Carpetgrass, Dichondra or St. Augustinegrass [including Floratam]) | | | |
| WHEN TO USE | Early Spring (through Summer) | | | |
| HOW MUCH TO USE | See spreader settings | | | |
| T | Questions or comments, call toll-free 1-877 BayerAL (877) 229-3725 Or visit us on-line at www.BayerAdvanced.com | | | |

FOR OUTDOOR RESIDENTIAL USE ONLY

^DA Reg. No. 72155-XXX

EPA Est. No. XXXX-AA-XX

Information regarding the contents and levels of metals in this product is available on the internet at http://www.regulatory-info-states.com.

When applied as directed, this product meets the guidelines for metals adopted by the Association of American Plant Food Control Officials.

5/8

DIRECTIONS FOR USE

" is a violation of Federal law to use this product in a manner inconsistent with its labeling.

BEFORE YOU USE

Read and follow these


directions when using:

- Apply when grass is moist early in the morning when dew is still present or after a brief watering –
 and wind is calm.
- Avoid applying when temperatures are consistently above 90°F.
- Use care when applying near desirable plants such as flowers, vegetables, shrubs and trees as contact with this weed killer may damage them.
- Follow these recommendations for use on new or refurbished lawns.
 - o **Seedina**
 - Do not apply to overseeded turf for 60 days after seeding or until the second mowing, whichever is longer.
 - Do not overseed for 6 weeks after using this product.
 - Sodding Do not apply to newly set sod.
 - Sprigging or Plugging Stolen rooting of newly sprigged or plugged grass may be temporarily slowed.

Apply before the typical crabgrass germination date shown for your area.

Approx. Date

- 1 After May 30
- 2 After May 10
- 3 After April 20
- 4 After March 20
- 5 Jan. 1 to March 20



HOW TO USE

Adjust spreader to the proper setting (see Spreader Settings).

To be sure you cover the entire area uniformly, first treat the border of the lawn then fill in the center. Make parallel passes, walking at a steady pace to ensure even distribution of the granules.

Do NOT apply if rain is expected within 24 hours.

Do NOT water for 24 hours after application.

Do NOT apply more than twice a year.

FOR BEST RESULTS

- Best results are obtained when this product is used as part of the (Bayer Advanced Lawn care program).
- Mow lawn to normal height 1 2 days before applying this product to expose low growing weeds and stimulate uptake of the weed killer.
- If lawn is dry, water 1 2 days before applying this product.
- Make sure lawn receives a thorough watering 2 days after application, either from rain and/or sprinkling.
- A second application may be necessary in areas where there was a lot of crabgrass the year before or where there is a long growing season. Make second application 6 – 8 weeks later. Use the shorter interval when conditions are hot or wet.

HOW Trimec + Prodiamine Granular Herbicide Plus Lawn Fertilizer Works

This product controls existing weeds and controls new weeds in your lawn. Weeds begin twisting & curling in just days and die within 4 weeks.



Do not allow people (other than the applicator) or pets on treatment area during application. Do not enter treatment area until dust has settled.

Kills: Annual Bluegrass (from seed) Annual Yellow Sweetclover strian Fieldcress Barnyardgrass (from seed) **Bedstraw Beggarticks** Betony, Florida Bindweed, Field **Bird Vetch Bitter Wintercress** Bittercress, Hairy **Bitterweed** Black-eyed Susan Black Medic **Black Mustard** Blackseed Plantain **Blessed Thistle** Bloodflower Milkweed **Blue Lettuce** 3lue Vervain **Bracted Plantain Brassbuttons Bristly Oxtongue Broadleaf Dock Broadleaf Plantain Broomweed** Buckhorn **Buckhom Plantain Bulbous Buttercup Bull Thistle** Illnettle urclover Burdock **Burning Nettle** Burweed Buttercup **Buttonweed** Canada Thistle Carolina Geranium Carpetweed Catchweed Bedstraw Catnip Catsear Chickweed, Common Chickweed, Mouseear Chicory Cinquefoil Clover Clover, Crimson Clover, Hop Clover, Red Clover, Strawberry Clover, Sweet Clover, White Cockle Cocklebur Common Mullein Crabgrass, Large & Smooth (from seed) **Preeping Charlie** reeping Jenny

Cudweed

Curly Dock

Daisy, English

Daisy, Fleabane Daisy, Oxeye **Dandelion** Dichondra Dogbane **Doafennel** Dollarweed Dock Elderberry False Dandelion False Flax False Sunflower **Fiddleneck** Florida Pusley Frenchweed Galinsoga Goathead Goldenrod Ground lvy Gumweed Hairy Fleabane Hawkweed Healall Heartleaf Drymary **Heath Aster** Hedge Bindweed Hedge Mustard Hemp Henbit **Hoary Cress** Hoary Plantain Hoary Vervain Horsenettle Jimsonweed Knawel Knotweed Kochia Lambsquarters Lespedeza Mallow Matchweed Mexicanweed Milk Vetch Morningglory Mouseear Hawkweed Mugwort **Musk Thistle** Mustard Narrowleaf Plantain Narrowleaf Vetch Nettle Orange Hawkweed **Oriental Cocklebur Oxalis** Panicum (from seed) Parsley-piert Parsnip Pearlwort Pennycress Pennywork Peppergrass Pepperweed Pigweed Pineywoods Bedstraw

Plains Coreopsis (Tickseed) Plantain Poison Ivy Poison Oak Pokeweed **Poorjoe** Prairie Sunflower **Prickly Lettuce** Prickly Sida Prostrate Knotweed **Prostrate Pigweed** Prostrate Spurge Prostrate Vervain **Puncturevine** Foxtails, Annual (from seed) Purslane Purslane, Common Ragweed Red Sorrel Redroot Pigweed Goosegrass (from seed) Redstem Filaree Rescuegrass (from seed) Rough Cinquefoil Rough Fleabane Russian Pigweed Russian Thistle Scarlet Pimpernel Scotch Thistle Sheep Sorrel Shepherdspurse Signalgrass (from seed) Slender Plantain Smallflower Galinsoga Smooth Dock Smooth Pigweed Sorrel Sowthistle Johnsongrass (from seed) Spanishneedles Speedwell Spiny Amaranth Spiny Cocklebur Spiny Sowthistle **Spotted Catsear** Spurge Spotted Spurge Spurweed Stinging Nettle Strawberry, India Mock Tall Nettle Tall Vervain **Tansy Mustard** Tanweed Thistle Trailing Crownvetch Tumble Mustard **Tumble Pigweed** Velvetleaf Venice Mallow Virginia Buttonweed Virginia Creeper Virginia Pepperweed Water Pennywort Wavyleaf Bullthistle Western Clematis Western Salsify White Mustard Wild Aster Wild Buckwheat

Wild Carrot
Wild Four-o'clock
Wild Garlic
Wild Geranium
Wild Lettuce
I Marigold
Mustard
Wild Onion
Wild Parsnip

Wild Radish
Wild Rape
Wild Strawberry
Wild Sweet Potato
Wild Vetch
Woodsorrel
Wooly Croton
Wooly Morningglory
Wooly Plantain

Wormseed Yarrow Yellow Rocket Yellowflower Pepperweed and other broadleaf weeds

| all our toll-free Consumer Service hotline 1-877-B | ayerAL for help. |
|--|--|
| /1,000 sq. ft. | |
| Setting | Distance Between Passes (ft.) |
| (Spreader settings to be determined.) | To be determined |
| sq. ft. | |
| Setting | |
| (Spreader settings to be determined.) | |
| | (Spreader settings to be determined.) sq. ft. Setting |

Settings are approximate. The condition of your spreader, the speed at which you walk and uneven terrain may affect the rate of application. Calibrate your spreader according to manufacturer's directions before each application.

TIP 1,000 sq. ft. = 20 x 50 ft.

Do not use a presider that applies are

Do not use a spreader that applies granules in narrow, concentrated or uneven bands. Be sure the spreader works properly and spreads granules evenly.

| IMPORTANT: Directions for Storage and Disposal | | | | |
|--|---|---|--|--|
| TORAGE CONDITIONS: | | Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. | | |
| DISPOSAL | Î | If Empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain. | | |

NOTICE

Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. The Buyer assumes responsibility for tack of performance or safety if not used according to the directions.

Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

CAUTION

- Harmful if absorbed through skin, inhaled, or swallowed.
- Causes moderate eye irritation.
- Avoid contact with skin, eyes, or clothing.
- Wear protective eyewear.
- Avoid breathing dust.
- When using this product, wear long-sleeved shirt, long pants, socks and shoes.
- Washing hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco will reduce exposure.
 - After using this product remove ciothing and launder separately before reuse.
- Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

| First Aid + | | | | |
|------------------------|---|--|--|--|
| If on skin or clothing | Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advise. | | | |
| lf Inhaled | Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration preferable mouth to mouth, if possible. Contact poison control center or doctor for further treatment advice. | | | |
| If swallowed | Call a poison control center or doctor immediately for treatment advice Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. | | | |
| If in eyes | Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advise. | | | |

NOTE: Have the product container or label with you when calling a poison control center, doctor, or going for treatment. You may call toll-free 1-877-229-3763 for medical emergency information. Active Ingredients: 2,4-D (CAS # 94-75-7), Prodiamine (CAS # 29091-21-2), Mecoprop-p (CAS # 16484-77-8) and Dicamba (CAS # 1918-00-9).

Environmental Hazards

- Drift or runoff may adversely affect nontarget plants.
- Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites.
 Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark.
 Do not contaminate water when disposing of equipment washwaters.
- This product has low solubility in water. At the limit of solubility, this product is not toxic to fish. However, at concentrations substantially above the level of water solubility, it may be toxic to fish.

D - 9791 030905

Bayer Advanced A business unit of Bayer CropScience, LP 95 Chestnut Ridge Road Montvale, NJ 07645