72155 -73

08/09/2010



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

August 9, 2010

Charles W. Boyd, Registrations Manager Bayer Advanced 2 T. W. Alexander Drive Research Triangle Park, NC 27709

Subject: Label Amendment—addition of alternate statements in accordance with EPA's pyrethroid label call-in Product Name: Imidacloprid + Tau-fluvalinate + Tebuconazole Concentrate EPA Reg. No.: 72155-73 Application Dated: June 11, 2010 Decision Number: 436276

Dear Mr. Boyd:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable with the following label revision provisions:

1. Revise the Storage and Disposal statements by adding the following statement: "Do not contaminate food, water, or feed by storage or disposal."

A copy of the label stamped "accepted with comments" is enclosed for your records. You must submit two copies of the final printed label revised in accordance with the comments listed above before the product is released for shipment under this label. If you have any questions, you may contact Clayton Myers at (703) 347-8874 or <u>myers.clayton@epa.gov</u>.

Sincerely,



Richard Gebken, Product Manager Insecticide Branch Registration Division (7505P)

Enclosure Stamped Label

Page 1 of 1

[FRONT PANEL]

Imidacloprid + Tau-fluvalinate + Tebuconazole

Concentrate Insecticide, Miticide, and Fungicide

This label contains two separate label statements for, Outdoor Residential Pump Sprayer/Hose-end Sprayer Concentrate product use and Outdoor Residential Ready-To-Spray product use.

- For use on: Roses, Flowers, Houseplants, Ground Covers, Vines, Ornamentals, Shrubs, and Trees
- Just one application quickly kills insects
- Complete plant protection Protects plants from roots to leaves/branches (top to bottom)
- Protects plants from the inside out
- Up to 30 day insect protection (guard, defense, shield)
- Protects against insects for up to 30 days
- Keeps killing insects for up to 30 days
- Continuous killing (power) for up to 30 days
- Rainproof Protection
- Protection that won't wash off
- Fast Acting ... Kills On Contact
- Kills insects on contact
- Long Lasting ... Keeps On Killing For up to 4 Weeks (30 days)
- Residual action, keeps killing for up to 4 weeks
- Protects 24/7, around the clock protection (guard, killing) for up to 4 weeks
- Contains Merit®
- Proprietary Merit® inside
- Systemic formula works through the leaves and stems to protect and cure
- A systemic protectant and curative
- A curative and preventative
- Cures and protects against listed insects, mites and diseases on roses, flowers, houseplants, ground covers, vines, ornamentals, shrubs, and trees
- Cures and prevents listed insect, mite and disease problems on roses, flowers, houseplants, ground covers, vines, ornamentals, shrubs, and trees
- 3 products in One: Insect Protection, Spider Mite Protection, and Disease Protection
- One application, three benefits: Insect Protection, Spider Mite Protection, and Disease Protection
- All-in-One, three products: insect, disease, and spider mite protection
- Systemic Protectant Formula gets (moves) inside the plant to protect and cure

- Prevents, stops, and cures listed pests on roses, flowers, houseplants, ground covers, vines, ornamentals, shrubs, and trees
- Controls (Kills, Eliminates) listed pests that harm roses, flowers, houseplants, ground covers, vines, ornamentals, shrubs, and trees
- Rescues (guards) plants from pests
- Cures and Prevents Black Spot, Powdery Mildew and Rust Disease(s)
- Kills (Knocks Out) Insects and Mites on Contact
- Controls black spot, powdery mildew, (and) rust, (and more)
- Controls (Kills) aphids, caterpillars, (and) mites, (and more)
- Premium (Superior) care 3 in 1 formula controls insect, mites, and diseases
- 4 in 1 formula kills insects on contact, controls mites, controls diseases, and keeps on protecting for (up to 4) weeks (provides long lasting protection)
- Total (Complete, All-in-one) Protection Formula Kills insects, mites and disease on contact, and keeps on protecting for (up to 4) weeks
- Exclusive (Proprietary) Formula Three way control controls insects, mites, and diseases
- Systemic formula, controls insects, mites and diseases
- Three benefits in one package
- One application with three benefits: controls insects, mites and diseases
- Systemic three way action (control)
- Fast, easy and efficient
- The Rose & Flower enhancer (enforcer)
- Merit® inside
- Maintains and restores beauty
- Three garden products in one
- From the creators of Merit®
- Proven three way control
- The Rose and Flower defender (guardian, protector)
- 100% Satisfaction Guaranteed* (or your money back)
- Kills aphids, spider mites, Japanese beetles & more

ACTIVE INGREDIENTS:		
Imidacloprid		
Tau-fluvalinate Tebuconazole		
OTHER INGREDIENTS:		
Net Contents: 32 fl. oz. (946 ml)	000000	0 0 0
STOP - Read This Entire Label Before Use	6 6 6 6 6 6 6 0	o
KEEP OUT OF REACH OF CHILDREN		0 0 6 6 6 6 6 6 6 6
CAUTION	00 0000 00	0 0 0 0 0 0 0 0 0 0
See back panel booklet for additional precautionary statements		U
WINCOMMENTS into, distribution and or use in Nassau, Suffolk, Kings, or Queens co	ounties of New Y	ork
In EPA Letter Dated:		
Under the Federal Insecticide, Fungicide		00001
pesticide Registered under EPA Reg. No:		

 $\left(\begin{array}{c} \cdot \end{array} \right)$

Imidacloprid + Tau-fluvalinate + Tebuconazole

Concentrate Insecticide, Miticide, and Fungicide

(Pump Sprayer/Hose-end Sprayer label)

Marketing claims as on page 1

ACTIVE INGREDIENTS:

Imidacloprid	
Tau-fluvalinate	
Tebuconazole	
OTHER INGREDIENTS:	

Net Contents: 32 fl. oz. (946 ml)

STOP - Read This Entire Label Before Use


KEEP OUT OF REACH OF CHILDREN

For Outdoor Residential Use Only

CAUTION

See back panel booklet for additional precautionary statements

Not for sale, sale into, distribution and or use in Nassau, Suffolk, Kings, or Queens counties of New York



[BACK PANEL]

[This product's exclusive triple protection formula provides long lasting control against insects, diseases and mites in one easy step.]

This product's exclusive 3-way systemic formula protects against insects, mites, and diseases in one easy step (Bayer Advanced's proprietary 4-in-1 formula kills insects on contact, controls mites, controls diseases, and keeps on protecting for (up to 4) weeks (provides long lasting protection)). This systemic concentrate formula provides long lasting insect, mite, and disease protection on roses, flowers, houseplants, ground covers, vines, ornamentals, shrubs, and trees. Growing beautiful trees, shrubs, flowers, and roses has never been easier! Better Science, Better Results.

QUICK FACTS	· · · · · · · · · · · · · · · · · · ·	
 Effective Against a Broad Range of Insects, Mites, and Diseases Keeps on Killing Insects for Up to 30 Days Easy, Accurate Measuring & Mixing 		 Apply with most common garden sprayers Kills insects on contact Highly effective against mites One quart covers up to 192 roses
INSECTS and MITES CONTROLLED	Aphids, caterpillars, Japanese beetles, leafminers, spider mites, whiteflies, and other listed pests.	
DISEASES CONTROLLED	Anthracnose, black spot, leaf spot, petal blight, powdery mildew, rust, and scab.	
COVERAGE	Makes (Creates, Delivers) up to 12 gallons of spray	
WHERE TO USE	Roses, flowers, houseplants, ground covers, vines, ornamentals, shrubs, and trees For use on non-edible plants only	
WHEN TO USE	To prevent insect, mite, or disease infestations or control them when they first appear.	
HOW MUCH TO USE	Dilute 2.67 oz (5¼ Tablespoons) concentrate in 1 gallon of water	
	Questions or comments, call toll-free (877) 229-3724 1-877-Bayer-AG	

ACTIVE INGREDIENTS:

Imidacloprid	
Tau-fluvalinate	
Tebuconazole	
OTHER INGREDIENTS:	


Imidacloprid is the chemical name for Merit® Insecticide.

EPA Reg. No. 72155-73

EPA Est. No. XXXX-AA-1

116

FOR OUTDOOR RESIDENTIAL USE ONLY



PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

- Harmful if swallowed or absorbed through skin.
- · Avoid contact with skin, eyes, or clothing.
- · Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.
- · Remove and wash contaminated clothing before reuse.

First Aid		
If Swallowed • Call a poison control center or doctor immediately for treatment advice		
	Have person sip a glass of water if able to swallow.	
	Do not induce vomiting unless told to do so by a poison control center or doctor.	
	Do not give anything to an unconscious person.	
If On Skin or Clothing • Take off contaminated clothing.		
	Rinse skin immediately with plenty of water for 15-20 minutes.	
	Call a poison control center or doctor for treatment advice.	
	tainer or label with you when calling a poison control center or doctor, or going for treatment. For medical nay call toll-free 1-877-229-3763.	

• Active Ingredients: Imidacloprid [CAS #138261-41-3], Tau-fluvalinate [CAS #102851-06-9], Tebuconazole [CAS #107534-96-3]

Environmental Hazards

- This product is toxic to fish and aquatic organisms.
- Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark.
- To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.
- This pesticide is toxic to bees.
- Do not apply this product when bees are active.
- Apply this product only as specified on this label.
- Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For best results, read and follow all label directions.

BEFORE YOU USE	Apply the product to flowering plants during early morning or late evening, when bees are not present.
	Do not spray the product into fish pools, ponds, streams, or lakes.
Read and follow these	Remove animal feeding dishes prior to application.
directions when using:	Do not use in small, enclosed areas where spray may be inhaled.
	For houseplants, move them outside, treat, let dry, and bring back indoors.
	Do not apply to fruits or vegetables used for food.
	Shake well before using.
WHEN TO USE	When insects, mites, or disease first appear.
	• To avoid serious damage, eliminate pests and diseases early in the season before they multiply.
MIXING & MEASURING	SEE SPRAYER TYPES

HOW TO SPRAY	Adjust nozzle on your sprayer to desired spray pattern.
	Hold sprayer about 12 inches from foliage.
	Spray leaves, stems and tender new shoots.
	(ART: of steps above)
	 Apply until both sides of the leaves are evenly coated with spray, but not dripping.
	(ART: spraying undersides of leaves)
HOW OFTEN	Kills insects on contact and keeps on killing for up to 30 days.
	For diseases and severe infestations, repeat every 7 to 14 days as necessary.
HOW IT WORKS	This product provides 3 way action: kills existing pests, forms protective barrier on outside of plant and is absorbed into the plant regardless of weather conditions. (This product provides 4 way action: kills insects on contact, controls mites, controls diseases, and keeps on protecting for (up to 4) weeks (provides long lasting protection)).

 \bigcap

٠

•

INSECTS AND MITES	(ART: Aphid, Japanese Beetle,			
CONTROLLED	Leafminer, Lace bugs)			
	Adelgids Aphids Armyworms Bagworms Beetles Boxelder bugs Budworms Cankerworms Cankerworms Caterpillars Chinch bugs Clover mites Crickets Cucumber beetles Cutworms	Earwigs Elm leaf beetles Firebrats Flea beetles Fungus gnats Grasshoppers Gypsy moth larvae Japanese beetles (adult) Lace bugs Leaf-feeding beetles Leaf-feeding caterpillars Leafhoppers (including Glassy-Winged Sharpshooter)	Leafminers (including Birch Leafminers and Holly Leafminers) Loopers Mealybugs Midges Mites Pillbugs Plant bugs Psyllids Rose Midges Rose Slugs Sawflies Scale insects (crawler stages)	Silverfish Sowbugs Spider Mites Spittlebugs Stink Bugs Thrips Two-Spotted Spider Mites Viburnum leaf beetle Webworms Weevils (including adult root weevils) Whiteflies

DISEASES CONTROLLED	CONTROLS	PLANTS
	Black Spot Powdery Mildew Rust	Roses
	Leaf Spot Powdery Mildew Rust	Flowers
	Anthracnose Leaf Spot Powdery Mildew Rust Scab	Crabapples (Ornamental), Dogwoods and Other Small Landscape [Ornamental] Trees
	Anthracnose Black Spot Leaf Spot Petal Blight Powdery Mildew Rust	Azaleas, Camellias, Rhododendrons, and Other Landscape [Ornamental] Shrubs, Ground Covers, and Vines
	Leaf Spot Powdery Mildew	Houseplants

To prevent diseases: apply at least 3 times, every 7-14 days apart during the growing season, starting when new leaves first appear.

FOR BEST RESULTS

Do not wet the foliage within one hour after applying. To avoid serious damage to plants, eliminate pests early in the season before they multiply.

\$₩	Children and pets may re-enter after spray has dried.

Restrictions

- Do not water the treated area to the point of run-off
- Do not make applications during rain.
- All outdoor applications must be limited to spot or crack-and-crevice treatment only, except for the following permitted uses:
 - (1) Treatment to soil or vegetation around structures;
 - (2) Applications to lawns, turf, and other vegetation;
- Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

SPRAYER TY	PES	
(ART: ICON OF	For Pump Style Tank Sprayer:	
PUMP TYPE)	To make one gallon of spray:	
	1. Pour 2.67 fl. oz. (5¼ Tablespoons) of the liquid concentrate.	
	2. Dilute in 1 gallon of water.	
	3. Close sprayer and shake well before applying.	
(ART: ICON	For Dial-Style Hose-End Sprayers:	
DIAL TYPE)	Dilution rate: 2.67 fl. oz. per gallon (2 tablespoons = 1 fl. oz.)	
	1. Fill sprayer jar with enough concentrate to treat the desired plants.	
	2. Close sprayer and set dial.	
	3. Attach sprayer to hose and apply.	
	OR	
	Fill sprayer jar with enough concentrate to cover plants to be treated and follow sprayer label directions for applying.	
(ART: ICON OF	For Concentrate-Plus-Water Hose-End Sprayers:	
WATER TYPE)	Dilution rate: 2.67 fl. oz. per gallon (2 tablespoons = 1 fl. oz.)	
	1. Pour concentrate into sprayer jar to the required fluid ounce level.	
	2. Add the appropriate amount of water to the gallons level.	
	3. Replace sprayer top and shake well before applying.	
	OR	
	Pour concentrate into sprayer jar to the required fluid ounce level and follow sprayer label directions for applying.	

Important – Directions for Storage and Disposal		
StorageStore in original container in a cool, dry place, out of the reach of chConditionspreferably a locked storage cabinet.		
Disposal ÎÎÎ	 If empty: Non-refillable container. Do not refill or reuse container. Place empty container in trash or offer for recycling if available. If partly filled: 	
	 Call your local solid waste agency or toll free 1-877-229-3724 for disposal instructions. Never place unused product down any indoor or outdoor drain. 	



NOTICE

Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. To the extent consistent with applicable law, the Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

*Money Back Guaranteed: If you are not satisfied with this product, we will gladly refund your original purchase price.



.

A Business Unit of Bayer CropScience, L.P. 2 T.W. Alexander Drive Research Triangle Park, NC 27709 [FRONT PANEL]

Imidacloprid + Tau-fluvalinate + Tebuconazole

Concentrate Insecticide, Miticide, and Fungicide

(Ready-to-Spray label)

Marketing claims as on page 1

Additional claims for Ready-to-Spray only:

- Pre-mixed and ready-to-spray (ready-to-go)
- Ready to spray ... No mixing, and no measuring

ACTIVE INGREDIENTS:

Imidacloprid	
Tau-fluvalinate	
Tebuconazole	
OTHER INGREDIENTS:	

Net Contents: 32 fl. oz. (946 ml)

STOP - Read This Entire Label Before Use KEEP OUT OF REACH OF CHILDREN FOR OUTDOOR RESIDENTIAL USE ONLY

CAUTION

See back panel booklet for additional precautionary statements

Not for sale, sale into, distribution and or use in Nassau, Suffolk, Kings, or Queens counties of New York

10/14

[BACK PANEL]

[This product's exclusive triple protection formula provides long lasting control against insects, diseases and mites in one easy step.]

This product's exclusive 3-way systemic formula protects against insects, mites, and diseases in one easy step (proprietary 4-in-1 formula kills insects on contact, controls mites, controls diseases, and keeps on protecting for (up to 4) weeks (provides long lasting protection)). This ready-to-spray formula provides long lasting insect, mite, and disease protection on roses, flowers, houseplants, ground covers, vines, ornamentals, shrubs, and trees. Growing beautiful trees, shrubs, flowers, and roses has never been easier! Better Science, Better Results.

QUICK FACTS		
 Effective Against a Bro Diseases Keeps on Killing Insect 30 days of Insect Cont 	. ,	 Ready-to-Spray formula no mixing, no measuring Kills insects on contact Highly effective against mites One quart covers up to 192 roses
INSECTS and MITES CONTROLLED	Aphids, caterpillars, Japanese beetles,	leafminers, spider mites, whiteflies, and other listed pests.
DISEASES CONTROLLED	Anthracnose, black spot, leaf spot, petal blight, powdery mildew, rust, and scab.	
COVERAGE	Treats (Protects) up to 192 roses	
WHERE TO USE	Roses, flowers, houseplants, ground covers, vines, ornamentals, shrubs, and trees For use on non-edible plants only	
WHEN TO USE	To prevent insect, mite, or disease infestations or control them when they first appear.	
	Questions or comments, call toll-free (877) 229-3724 1-877-Bayer-AG

ACTIVE INGREDIENTS:

Imidacloprid	
Tau-fluvalinate	
Tebuconazole	
OTHER INGREDIENTS:	

Imidacloprid is the chemical name for Merit® Insecticide.

EPA Reg. No. 72155-73

EPA Est. No. XXXX-AA-1

FOR OUTDOOR RESIDENTIAL USE ONLY

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

- · Harmful if swallowed or absorbed through skin.
- · Avoid contact with skin, eyes, or clothing.
- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.
- Remove and wash contaminated clothing before reuse.

First Aid		
If Swallowed	Call a poison control center or doctor immediately for treatment advice	
	Have person sip a glass of water if able to swallow.	
	Do not induce vomiting unless told to do so by a poison control center or doctor.	
	Do not give anything to an unconscious person.	
If On Skin or Clothing	Take off contaminated clothing.	
	Rinse skin immediately with plenty of water for 15-20 minutes.	
	Call a poison control center or doctor for treatment advice.	

NOTE: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergency information, you may call toll-free 1-877-229-3763.

• Active Ingredients: Imidacloprid [CAS #138261-41-3], Tau-fluvalinate [CAS #102851-06-9], Tebuconazole [CAS #107534-96-3]

Environmental Hazards

- This product is toxic to fish and aquatic organisms.
- Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark.
- To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.
- This pesticide is toxic to bees.
- Do not apply this product when bees are active.
- Apply this product only as specified on this label.
- · Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

DIRECTIONS FOR USE		
It is a violation of Federal	law to use this product in a manner inconsistent with its labeling.	
For best results, read and	d follow all label directions.	
BEFORE YOU USE	Apply the product to flowering plants during early morning or late evening, when bees are not present.	
	Do not spray the product into fish pools, ponds, streams, or lakes.	
Read and follow these	Remove animal feeding dishes prior to application.	
directions when using:	Do not use in small, enclosed areas where spray may be inhaled.	
	For houseplants, move them outside, treat, let dry, and bring back indoors.	
	Do not apply to fruits or vegetables used for food.	
	Shake well before using.	
WHEN TO USE	When insects, mites, or disease first appear.	
	• To avoid serious damage, eliminate pests and diseases early in the season before they multiply.	
HOW TO USE	This product is ready to spray. Just attach this bottle to your hose according to directions. Water	
Shake Well Before Use	automatically mixes with the concentrate as you spray.	

HOW TO ATTACH AND SPRAY	 Connect your garden hose to the bottle. Be sure the blue water control valve on top of the spray nozzle is in the OFF position. 	
ON	2. To begin spraying, tum on water at faucet. Point nozzle towards the area to be treated and Press lock tab while using other hand to turn sprayer valve to on position.	
	3. To stop spraying, turn the water control valve on top of the spray nozzle to OFF. Turn off faucet.	
	 Relieve water pressure by turning water control valve to the ON position until water slows to a drip. Turn water control valve back to OFF. Disconnect garden hose from bottle. 	
	5. The unused portion can be saved and used at a later date.	
OFF	Tips for Applying	
	 Extend hose to the farthest plants to be treated and work back toward the faucet to avoid contacting wet, treated plants. 	
	 Use an even sweeping motion, ensuring thorough coverage of both the top and bottom leaf surfaces. 	
[HOW TO ATTACH AND SPRAY]	 [Connect your garden hose to the bottle. Be sure the (color inserted here) water control value on top of the spray nozzle is in the OFF position. 	
	2. Turn on water at faucet. Twist the small, star-shaped, (color inserted here) product-control button to align the flat portion with the lock tab. Push the button in to open.	
	 To begin spraying, point nozzle towards the area to be treated and turn the (color) water control valve to ON position. Tab will be horizontal. 	
	 To stop spraying, turn the (color) water control valve on top of the spray nozzle to OFF. To close, push in the small round button opposite the star-shaped product-control button. Turn off faucet.] 	
	 Relieve water pressure by turning water control valve to the ON position until water slows to a drip Turn water control valve back to OFF. Disconnect garden hose from bottle. 	
	6. The unused portion can be saved and used at a later date.	
	Tips for Applying	
	 Extend hose to the farthest plants to be treated and work back toward the faucet to avoid contacting wet, treated plants. 	
	 Use an even sweeping motion, ensuring thorough coverage of both the top and bottom leaf surfaces. 	
	[Alternative Sprayer Directions]	
HOW OFTEN	Kills insects on contact and keeps on killing for up to 30 days.	
	For diseases and severe infestations, repeat every 7 to 14 days as necessary.	
HOW IT WORKS	This product provides 3 way action: kills existing pests, forms protective barrier on outside of plant and is absorbed into the plant regardless of weather conditions. (This product provides 4 way action: kills insects on contact, controls mites, controls diseases, and keeps on protecting for (up to 4) weeks (provides long lasting protection)).	
a a a		

(

٠

,

.

(

INSECTS AND MITES	(ART: Aphid, Japanese Beetle,			
CONTROLLED	Leafminer, Lace bugs)			
	Adelgids Aphids Armyworms Bagworms Beetles Boxelder bugs Budworms Cankerworms Cankerworms Caterpillars Chinch bugs Clover mites Crickets Cucumber beetles Cutworms	Earwigs Elm leaf beetles Firebrats Flea beetles Fungus gnats Grasshoppers Gypsy moth larvae Japanese beetles (adult) Lace bugs Leaf-feeding beetles Leaf-feeding caterpillars Leafhoppers (including Glassy-Winged Sharpshooter)	Leafminers (including Birch Leafminers and Holly Leafminers) Loopers Mealybugs Midges Mites Pillbugs Plant bugs Psyllids Rose Midges Rose Slugs Sawflies Scale insects (crawler stages)	Silverfish Sowbugs Spider Mites Spittlebugs Stink Bugs Thrips Two-Spotted Spider Mites Viburnum leaf beetle Webworms Weevils (including adult root weevils) Whiteflies

DISEASES CONTROLLED	CONTROLS	PLANTS
	Black Spot	Roses
	Powdery Mildew	
	Rust	
	Leaf Spot	Flowers
	Powdery Mildew	
	Rust	
	Anthracnose	Crabapples (Ornamental), Dogwoods and
	Leaf Spot	Other Small Landscape [Ornamental] Trees
	Powdery Mildew	
	Rust	
	Scab	
	Anthracnose	Azaleas, Camellias, Rhododendrons, and
	Black Spot	Other Landscape [Ornamental] Shrubs,
	Leaf Spot	Ground Covers, and Vines
	Petal Blight	
	Powdery Mildew	
	Rust	
	Leaf Spot	Houseplants
	Powdery Mildew	
	ase: apply at least 3 times, every 7-14 days, begin	
To prevent diseases:	apply at least 3 times, every 7-14 days, during th	e growing season, starting when new leaves first appear.

FOR BEST RESULTS

j

Do not wet the foliage within one hour after applying. To avoid serious damage to plants, eliminate pests early in the season before they multiply.

Children and pets may re-enter after spray has dried.

Restrictions

- · Do not water the treated area to the point of run-off
- Do not make applications during rain.
- All outdoor applications must be limited to spot or crack-and-crevice treatment only, except for the following permitted uses:
 (1) Treatment to soil or vegetation around structures;
- (2) Applications to lawns, turf, and other vegetation;
- Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

Important – Directions for Storage and Disposal	
Storage Conditions	Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet.
Disposal	If empty:
Ŵ	 Non-refillable container. Do not refill or reuse container. Place empty container in trash or offer for recycling if available.
	If partly filled:
	 Call your local solid waste agency or toll free 1-877-229-3724 for disposal instructions.
	 Never place unused product down any indoor or outdoor drain.

NOTICE

Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. To the extent consistent with applicable law, the Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

*Money Back Guaranteed: If you are not satisfied with this product, we will gladly refund your original purchase price.



A Business Unit of Bayer CropScience, L.P. 2 T.W. Alexander Drive Research Triangle Park, NC 27709

Imidacloprid + Tau-fluvalinate + Tebuconazole Concentrate (MASTER) Approved 11/02/09, Notification 05/26/10