

72155-77

05/18/2007

114



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number:
72155-77

Date of Issuance:
MAY 18 2007

Term of Issuance: **Conditional**

Name of Pesticide Product:
Lawn Herbicide 3F Concentrate

NOTICE OF PESTICIDE:
 Registration
 Reregistration
(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Bayer Advanced
2 T W Alexander Drive
Research Triangle Park, NC 27709

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) and (B) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit such data.
2. Submit within one year of the date of this registration notice, storage stability (830.6317) and corrosion characteristics (830.6320) studies.
3. Make the following label changes:
 - a. Add an Application Restriction statement to read "Do not apply this product in a way that will contact any person, pet either directly or through drift. Keep people and pets out of the area during application."
 - b. Add the following statement to Environmental Hazards "Do not apply to water. Do not contaminate water when disposing of equipment washwaters or rinsate." Also in this same section, delete the statement that reads "Do not apply directly to lakes, streams, rivers or ponds."

Signature of Approving Official:
Joanne I. Miller
Product Manager 23
Herbicide Branch
Registration Division (7505P)

Date:
MAY 18 2007

- c. Make the **"If in Eyes"** First Aid statement first.
 - d. Revise the EPA Registration Number to read, "EPA Reg. No. **72155-77**".
4. Submit one copy of the revised final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505P)

3/14

(Front Panel)

Bayer Advanced Lawn Herbicide 3F Concentrate

This label contains two separate label statements for Outdoor Residential Pump sprayer/Hose-end Sprayer Concentrate product use and Outdoor Residential Ready-To-Spray product use.

- Kills Lawn Weeds plus Crabgrass [in One Step]
- Kills Over 200 Lawn Weeds Plus Crabgrass in One Easy Step
- Kills Crabgrass plus Broadleaf Weeds
- Kills Listed Weeds
- Kills Hard-to-Control Weeds
- Systemic [Herbicide] [Activity]
- Kills Both Broadleaf and Grassy Weeds
- Kills Dandelion, Crabgrass, Clover, plus 200 Other listed Weeds
- Rainproof in [1] hour[s]
- New!
- Two-in-One Product that Kills Crabgrass & Broadleaf Weeds
- Two Types of Weeds; One Solution
- Guaranteed* Not to harm your lawn
- All-In-One Weed Killer [for lawns]
- Starts working on contact
- Kills even tough weeds – roots & all
- Kills 200+ Weeds
- Unique Formula
- Fast results
- Kills lawn weeds plus crabgrass
- Gets the root
- Won't Harm [your] lawn [s]
- Will not harm your lawn
- Kills tough weeds
- Kills the weeds, not your lawn
- Kills to the root
- Kills weeds, not lawns
- One [easy] step
- Two products in one
- Kills all major broadleaf weeds
- Weed protection system
- Visible results
- Kills weeds - Fast
- Kills both Crabgrass & Broadleaf weeds
- Proprietary Formula that Kills Both Crabgrass and Broadleaf Weeds
- Effectively Kills Both Broadleaf and Weed grasses
- Save time and energy
- Always ready to spray
- Easy to use
- Fast coverage
- Easy & Fast coverage
- Targets weeds
- Kills weeds the first time, every time
- Guaranteed* not to harm lawns
- Guaranteed* Results
- 100% Satisfaction Guaranteed* (or your money back)
- For use on Kentucky bluegrass, perennial ryegrass, fescue (except creeping fescue), centipedegrass, St. Augustinegrass (except Floratam), zoysiagrass
- Tough weed protection
- 4-way weed control action
- Exclusive 4-Way Weed control
- Proprietary [unique] [advanced] formula
- Attach to hose and spray

Active Ingredients:

2,4-D, Triisopropanolamine salt.....	4.91%
Mecoprop-p, Potassium salt.....	1.55%
Dicamba, Potassium salt	0.77%
Fenoxaprop-p-ethyl†.....	0.41%
Other Ingredients	92.36 %
	100.00%

†Protected by U.S. Patent No. 5,254,527
Net 32 fl oz. (946 mL)

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See back panel for additional precautionary statements

Editorial Note – [Bracketed text] is optional language

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

MAY 18 2007
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

72155-77

4/14

(Pump Sprayer/Hose-end Sprayer label)

(FRONT PANEL)

Bayer Advanced Lawn Herbicide 3F Concentrate

[Editorial Note – Marketing Claims from page 1]

Active Ingredients:

2,4-D, Triisopropanolamine salt.....	4.91%
Mecoprop-p, Potassium salt.....	1.55%
Dicamba, Potassium salt	0.77%
Fenoxaprop-p-ethyl†.....	0.41%
Other Ingredients	92.36 %
	100.00%

†Protected by U.S. Patent No. 5,254,527

Net 32 fl oz. (946 mL)

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional precautionary statements


5/14

(Back Panel)

**OPEN Resealable Label for Directions & Precautions
PRESS TO RESEAL**

Bayer Advanced Lawn Herbicide 3F Concentrate's exclusive and unique formula combines 4 proven weed killers to kill over 200 types of lawn weeds plus crabgrass in one easy step. And it's specially formulated so it won't harm your lawn when used as directed. Rainproof in 1 hour. Controlling weeds plus crabgrass in your lawn has never been easier!



QUICK FACTS	
Marketing claims to be selected from the front page	
KILLS	Broadleaf Weeds: Over 200 types including dandelions, clovers, chickweeds Grassy Weeds: Crabgrass, goosegrass, foxtail, sandbur, and many others
WHERE TO USE	On lawns without harming lawn grasses – Centipedegrass, Fescue (except Creeping Fescue), Kentucky Bluegrass, Perennial Ryegrass, St. Augustinegrass (except Floratam), Zoysiagrass (DO NOT USE on Bahiagrass, Bermudagrass, Buffalograss, Carpetgrass, Creeping Fescue, Floratam variety of St. Augustinegrass, Dichondra or desirable clovers.)
[COVERAGE]	[This bottle treats 6,400 sq ft.]
	For questions or comments, call toll-free 1-877 BayerAG (1-877-229-3724) or visit our website at www.BayerAdvanced.com

FOR OUTDOOR RESIDENTIAL USE ONLY

Active Ingredients:	
2,4-D, Triisopropanolamine salt	4.91%
Mecoprop-p, Potassium salt	1.55%
Dicamba, Potassium salt	0.77%
Fenoxaprop-p-ethyl [†]	0.41%
Other Ingredients	92.36 %
	100.00%

THIS PRODUCT CONTAINS: 2.64% 2,4-D acid equivalent, 1.32% Mecoprop-p acid equivalent, 0.66% Dicamba acid equivalent. Isomer Specific by AOAC Methods.

EPA Reg. No. 72155 – XX EPA Est. No. XXX-AA-XX

6/14

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

HOW MUCH TO USE
 Measure area to be sprayed. Calculate 'square feet' by multiplying length times width. [This bottle will treat 6,400 sq. ft.]
 The maximum application rate to turf is 1.0 pound of 2,4-D acid equivalent per acre per application per site.
 The maximum number of broadcast applications per treatment site is 2 per year.

HOW TO USE
Shake Well Before Use

PUMP-UP TANK SPRAYERS (Best for spot spraying individual weeds)
 1. Add appropriate amounts of water & product to sprayer tank – 3 fl. oz. (6 Tbs) product in 1 gallon of water treats 500 sq. ft.
 2. Close sprayer, shake well & pressurize. Adjust nozzle to a coarse spray.
 Large Areas – Apply evenly over area.
 Individual Weeds – Hold tip of wand about 12 inches from weed, aim at center of weed and spray.
 3. Occasionally re-pressurize sprayer to maintain a good spray pattern.

DIAL STYLE HOSE-END SPRAYERS (Best for spraying large areas of the lawn)
 1. Fill sprayer jar with enough product for area to be treated – 3 fl. oz. (6 Tbs) product treats 500 sq. ft. **DO NOT ADD WATER!**
 2. Close sprayer, set dial and apply.
 3. Pour any unused product back into this original container.

CONCENTRATE-PLUS-WATER HOSE-END SPRAYERS (Best for spraying large areas of the lawn)
 1. Pour product into sprayer jar to the required fluid ounce level – 3 fl oz (6 Tbs) treats 500 sq. ft.
 2. Add the appropriate amount of water to the gallons level.
 3. Replace sprayer top, shake well and apply.

[FOR BEST RESULTS]

- [Spray when weeds are [small [young] and] actively growing.]
- [Spray when daily high temperature is below 90°F.]
- [Rain or watering 1 hour after application will not wash away effectiveness.]
- [Apply when wind is calm [to avoid spray drift onto desirable plants (fruits, vegetables, flowers, groundcovers, shrubs & trees)]]
- [Reapply in 2 weeks if necessary.]
- [Do not apply while grass is emerging from winter dormancy.]
- [Newly seeded lawns may be sprayed after the fourth mowing.]
- [Bare spots may be seeded 4 weeks after application.]
- [Do not mow for 1 to 2 days before & after spraying [Uncut weeds have more surface area to hold more weed killer. They are also more likely to be hit by the spray. Lawn clippings can get in the way of the herbicide hitting the weed. Weed killers need time to move throughout the weed – mowing may remove the treated parts].]
- [Leave grass clippings on the lawn from the first 3 mowings after using this product. Do not use as compost or mulch.]
- [If lawn is dry, water 1 to 2 days before using this product. [Weeds take up herbicides best when not under stress.]]

HOW IT WORKS
 BAYER ADVANCED LAWN HERBICIDE 3F CONCENTRATE contains four proven weed killers that target lawn weeds and crabgrass. The herbicides enter the lawn weeds through their leaves and move inside the weeds, even down into the roots. And lawns will not be harmed when used as directed.

Kills over 200 weeds including:

Black medic Carpetweeds Chickweeds Clovers Crabgrass Creeping Charlie Cudweed Curly dock Dandelion English daisy False dandelion Filaree	Florida pusley Foxtails Ground ivy Heartleaf drymary Henbit Kikuyugrass Oxalis Pennywort Plantains Purslane Red sorrel Sheep sorrel Spurges	Spurweed Thistles Toadflax Wild carrot Wild garlic Wild geranium Wild onion & many other broadleaf & grass weeds (See appendix for a complete list of the weeds controlled)
---	---	---

7/14



Do not allow people (other than applicator) or pets on treatment area during application and until spray has dried.

PRECAUTIONARY STATEMENTS

CAUTION

Hazards to Humans and Domestic Animals

- Harmful if swallowed
- Causes moderate eye irritation.
- Avoid contact with eyes, skin, or clothing.

After using this product:

- If product gets on clothing, remove clothing as soon as possible and launder clothing separately before reuse.
- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

FIRST AID +

If Swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in Eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for further treatment advice.


NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 1-877-229-3763 for medical emergency information.

Active Ingredients: 4.91% 2,4-D Triisopropanolamine salt (CAS # 32341-80-3), 1.55% Mecoprop-p Potassium salt single isomer (CAS # 66423-5-0), 0.77% Dicamba, Potassium salt (CAS # 10007-85-9), and 0.41% Fenoxaprop-p-ethyl (CAS # 71283-80-2).

ENVIRONMENTAL HAZARDS

- This product is toxic to fish & aquatic invertebrates. Do not apply directly to lakes, streams, rivers or ponds.
- Drift or runoff may harm fish, aquatic invertebrates and non-target plants.

8/14

Important – Directions for Storage and Disposal	
Storage Conditions	Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing
Disposal 	If empty: Do not reuse container. Place empty container in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or toll free 1-877-229-3724 for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE: Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. The Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

***Money Back Guarantee:** If you are not satisfied with this product, we will gladly refund your original purchase price.



www.bayeradvanced.com

A Business Group of Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709

Lawn Herbicide 3F Concentrate (PENDING) 10/20/06

9/14

(Ready-To- Spray label)
(FRONT PANEL)

Bayer Advanced Lawn Herbicide 3F Concentrate

This label contains two separate label statements for, Outdoor Residential Pump Sprayer/Hose-end Sprayer Concentrate product use and Outdoor Residential Ready-To-Spray product use.

[Editorial Note – Marketing Claims from page 1]

Active Ingredients:

2,4-D, Triisopropanolamine salt.....	4.91%
Mecoprop-p, Potassium salt.....	1.55%
Dicamba, Potassium salt	0.77%
Fenoxaprop-p-ethyl†	0.41%
Other Ingredients	92.36 %
	100.00%

Net 32 fl oz. (946 mL)

KEEP OUT OF REACH OF CHILDREN
CAUTION

See back panel for additional precautionary statements

†Protected by U.S. Patent No. 5,254,527


10/14

(Back Panel)

**OPEN Resealable Label for Directions & Precautions
PRESS TO RESEAL**

Bayer Advanced Lawn Herbicide 3F Concentrate's exclusive and unique formula combines 4 proven weed killers to kill over 200 types of lawn weeds plus crabgrass in one easy step. And it's specially formulated so it won't harm your lawn when used as directed. Rainproof in 1 hour. Controlling weeds plus crabgrass in your lawn has never been easier!



QUICK FACTS	
Marketing claims to be selected from the front page	
KILLS	Broadleaf Weeds: Over 200 types including dandelions, clovers, chickweeds Grassy Weeds: Crabgrass, goosegrass, foxtail, sandbur, and many others
WHERE TO USE	On lawns without harming lawn grasses – Centipedegrass, Fescue (except Creeping Fescue), Kentucky Bluegrass, Perennial Ryegrass, St. Augustinegrass (except Floratam), Zoysiagrass (DO NOT USE on Bahiagrass, Bermudagrass, Buffalograss, Carpetgrass, Creeping Fescue, Floratam variety of St. Augustinegrass, Dichondra or desirable clovers.)
[COVERAGE]	[This bottle treats 6,400 sq ft.]
	For questions or comments, call toll-free 1-877 BayerAG (1-877-229-3724) or visit our website at www.BayerAdvanced.com

FOR OUTDOOR RESIDENTIAL USE ONLY

Active Ingredients:	
2,4-D, Triisopropanolamine salt.....	4.91%
Mecoprop-p, Potassium salt.....	1.55%
Dicamba, Potassium salt	0.77%
Fenoxaprop-p-ethyl [†]	0.41%
Other Ingredients	92.36 %
	100.00%

THIS PRODUCT CONTAINS: 2.64% 2,4-D acid equivalent, 1.32% Mecoprop-p acid equivalent, 0.66% Dicamba acid equivalent. Isomer Specific by AOAC Methods

EPA Reg. No. 72155 – XX


EPA Est. No. XXX-AA-XX

11/14

DIRECTIONS FOR USE


It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

<p>HOW MUCH TO USE</p>	<p>Measure area to be sprayed. Calculate 'square feet' by multiplying length times width. This bottle will treat 6,400 sq. ft. The maximum application rate to turf is 1.0 pound of 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.</p>
<p>HOW TO USE Shake Well Before Use</p>	<p>CONNECT</p> <ol style="list-style-type: none"> 1. Connect sprayer to garden hose. 2. Turn on water. <p>SPRAY</p> <ol style="list-style-type: none"> 1. Extend hose to farthest area to be treated, point nozzle in direction to be sprayed. 2. Turn on sprayer -- < Insert sprayer-specific language here > <i>Calmar RTS sprayer:</i> press lock tab while using other hand to turn knob counter-clockwise to ON position. <i>Green Garden RTS sprayer:</i> turn sprayer valve to ON position. <i>Hayes HP-2 RTS sprayer:</i> press lock tab while using other hand to turn knob clockwise to ON position. 3. Walk at a steady pace while spraying, using an even sweeping motion, and slightly overlap the treated areas. Work back toward the faucet to avoid walking over wet, treated grass. <p>FINISH</p> <ol style="list-style-type: none"> 1. Turn off sprayer -- < Insert sprayer-specific language here > <i>Calmar RTS sprayer:</i> turn knob clockwise to OFF position. <i>Green Garden RTS sprayer:</i> [Nothing required]. <i>Hayes HP-2 RTS sprayer:</i> turn knob counter-clockwise to OFF position. 2. Turn off water at faucet. 3. Relieve pressure in hose by turning sprayer knob back to ON position until water slows to a drip then turn sprayer knob to OFF position. 4. Disconnect sprayer from hose.
<p>[FOR BEST RESULTS]</p>	<ul style="list-style-type: none"> • [Spray when weeds are [small [young] and] actively growing.] • [Spray when daily high temperature is below 90°F.] • [Rain or watering 1 hour after application will not wash away effectiveness.] • [Apply when wind is calm [to avoid spray drift onto desirable plants (fruits, vegetables, flowers, groundcovers, shrubs & trees)]] • [Reapply in 2 weeks if necessary.] • [Do not apply while grass is emerging from winter dormancy.] • [Newly seeded lawns may be sprayed after the fourth mowing.] • [Bare spots may be seeded 4 weeks after application.] • [Do not mow for 1 to 2 days before & after spraying [Uncut weeds have more surface area to hold more weed killer. They are also more likely to be hit by the spray. Lawn clippings can get in the way of the herbicide hitting the weed. Weed killers need time to move throughout the weed – mowing may remove the treated parts.].] • [Leave grass clippings on the lawn from the first 3 mowings after using this product. Do not use as compost or mulch.] • [If lawn is dry, water 1 to 2 days before using this product. [Weeds take up herbicides best when not under stress.]]
<p>HOW IT WORKS</p>	<p>BAYER ADVANCED LAWN HERBICIDE 3F CONCENTRATE contains four proven weed killers that target lawn weeds and crabgrass. The herbicides enter the lawn weeds through their leaves and move inside the weeds, even down into the roots. And lawns will not be harmed when used as directed.</p>



12/14

Kills over 200 weeds including:		
Black medic Carpetweeds Chickweeds Clovers Crabgrass Creeping Charlie Cudweed Curly dock Dandelion English daisy False dandelion Filaree	Florida pusley Foxtails Ground ivy Heartleaf drymary Henbit Kikuyugrass Oxalis Pennywort Plantains Purslane Red sorrel Sheep sorrel Spurges	Spurweed Thistles Toadflax Wild carrot Wild garlic Wild geranium Wild onion & many other broadleaf & grass weeds (See appendix for a complete list of the weeds controlled)

	Do not allow people (other than applicator) or pets on treatment area during application and until spray has dried.
---	---

PRECAUTIONARY STATEMENTS

CAUTION

Hazards to Humans and Domestic Animals

- Harmful if swallowed
- Causes moderate eye irritation.
- Avoid contact with eyes, skin, or clothing.

After using this product:

- If product gets on clothing, remove clothing as soon as possible and launder clothing separately before reuse.
- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

FIRST AID +

If Swallowed	<ul style="list-style-type: none"> ▪ Call a poison control center or doctor immediately for treatment advice. ▪ Have person sip a glass of water if able to swallow. ▪ Do not induce vomiting unless told to do so by a poison control center or doctor. ▪ Do not give anything by mouth to an unconscious person.
If in Eyes	<ul style="list-style-type: none"> ▪ Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. ▪ Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. ▪ Call a poison control center or doctor for further treatment advice.


NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 1-877-229-3763 for medical emergency information.
 Active Ingredients: 4.91% 2,4-D Triisopropanolamine salt (CAS # 32341-80-3), 1.55% Mecoprop-p Potassium salt single isomer (CAS # 66423-5-0), 0.77% Dicamba, Potassium salt (CAS # 10007-85-9), and 0.41% Fenoxaprop-p-ethyl (CAS # 71283-80-2).

13/14

ENVIRONMENTAL HAZARDS

- This product is toxic to fish & aquatic invertebrates. Do not apply directly to lakes, streams, rivers or ponds.
- Drift or runoff may harm fish, aquatic invertebrates and non-target plants.

Important – Directions for Storage and Disposal

Storage Conditions	Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing
Disposal 	If empty: Do not reuse container. Place empty container in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or toll free 1-877-229-3724 for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE: Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. The Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

***Money Back Guarantee:** If you are not satisfied with this product, we will gladly refund your original purchase price.



www.bayeradvanced.com

A Business Group of Bayer CropScience LP
 P.O. Box 12014, 2 T.W. Alexander Drive
 Research Triangle Park, North Carolina 27709

Lawn Herbicide 3F RTS (PENDING) 10/20/06

Appendix

WEEDS CONTROLLED			
Annual Yellow Sweetclover	Cocklebur	Mouseear Hawkweed	Speedwell
Aster	Common Mullein	Mugwort	Spiny Amaranth
Austrian Fieldcress	Crabgrass , Large/Hairy	Musk Thistle	Spiny Cocklebur
Barnyardgrass	Crabgrass, Small/Smooth	Mustard	Spiny Sowthistle
Bedstraw	Creeping Charlie	Narrowleaf Plantain	Spotted Catsear
Beggarticks	Creeping Jenny	Narrowleaf Vetch	Spotted Spurge
Betony, Florida	Cudweed	Nettle	Sprangletop
Bindweed, Field	Curly Dock	Orange Hawkweed	Spurweed
Bird Vetch	Daisy, English	Oriental Cocklebur	Stinging Nettle
Bitter Wintercress	Daisy, Fleabane	Oxalis	Strawberry, India Mock
Bittercress, Hairy	Daisy, Oxeye	Panicum	Tall Nettle
Bitterweed	Dandelion	Parsley-piert	Tall Vervain
Black-eyed Susan	Dichondra	Parsnip	Tansy Mustard
Black Medic	Dogbane	Pearlwort	Tansy Ragwort
Black Mustard	Dogfennel	Pennycress	Tanweed
Blackseed Plantain	Dollarweed	Pennywort	Thistle
Blessed Thistle	Elderberry	Pepperweed	Trailing Crownvetch
Bloodflower Milkweed	False Dandelion	Pigweed	Tumble Mustard
Blue Lettuce	False Flax	Pineywoods Bedstraw	Tumble Pigweed
Blue Vervain	False Sunflower	Plains Coreopsis (Tickseed)	Velvetleaf
Bracted Plantain	Fiddleneck	Plantain	Venice Mallow
Brassbuttons	Florida Pusley	Poison Ivy	Virginia Buttonweed
Bristly Oxtongue	Foxtail	Poison Oak	Virginia Creeper
Broadleaf Dock	Frenchweed	Pokeweed	Virginia Pepperweed
Broadleaf Plantain	Galinsoga	Poorjoe	Water Pennywort
Broomweed	Goathead	Prairie Sunflower	Wavyleaf Bullthistle
Buckhorn	Goldenrod	Prickly Lettuce	Western Clematis
Buckhorn Plantain	Goosegrass	Prickly Sida	Western Salsify
Bulbous Buttercup	Ground Ivy	Prostrate Knotweed	White Mustard
Bull Thistle	Gumweed	Prostrate Pigweed	Wild Aster
Bullnettle	Hairy Fleabane	Prostrate Spurge	Wild Buckwheat
Burclover	Hawkweed	Prostrate Vervain	Wild Carrot
Burdock	Healall	Puncturevine	Wild Four-o'clock
Burning Nettle	Heartleaf Drymary	Purslane, Common	Wild Garlic
Burweed	Heath Aster	Ragweed	Wild Geranium
Buttercup	Hedge Bindweed	Red Sorrel	Wild Lettuce
Buttonweed	Hedge Mustard	Redroot Pigweed	Wild Marigold
Canada Thistle	Hemp	Redstem Filaree	Wild Mustard
Carolina Geranium	Henbit	Rough Cinquefoil	Wild Onion
Carpetweed	Hoary Cress	Rough Fleabane	Wild Parsnip
Catchweed Bedstraw	Hoary Plantain	Russian Pigweed	Wild Radish
Catnip	Hoary Vervain	Russian Thistle	Wild Rape
Catsear	Horsenettle	Sandbur	Wild Strawberry
Chickweed, Common	Japanese Stiltgrass	Scarlet Pimpernel	Wild Sweet Potato
Chickweed, Mouseear	Jimsonweed	Scotch Thistle	Wild Vetch
Chicory	Johnsongrass (seedling)	Sheep Sorrel	Woodsorrel
Cinquefoil	Knawel	Shepherdspurse	Wooly Croton
Clover, Crimson	Knotweed	Silver Crabgrass	Wooly Morningglory
Clover, Hop	Kochia	Slender Plantain	Wooly Plantain
Clover, Red	Lambsquarters	Smallflower Galinsoga	Wormseed
Clover, Strawberry	Lespedeza	Smooth Dock	Yarrow
Clover, Sweet	Mallow	Smooth Pigweed	Yellow Rocket
Clover, White	Matchweed	Sorrel	Yellowflower Pepperweed
Cockle	Mexicanweed	Sowthistle	and other broadleaf weeds
	Milk Vetch	Spanishneedles	
	Morningglory		