10-19-2009



U.S. ENVIRONMENTAL PROTECTION **AGENCY**

Office of Pesticide Programs Registration Division (7505P) Ariel Rios Building 1200 Pennsylvania Ave., NW Washington, D.C. 20460

NOTICE OF PESTICIDE:

Registration X Reregistration (under FIFRA, as amended)

EPA Reg. Numbe	1
72155-77	

Date of Issuance:

OCT 19 2009

Term of Issuance:

Name of Pesticide Product:

Bayer Advanced Lawn Herbicide 3F Concentrate

Name and Address of Registrant (include ZIP Code): Bayer Advanced, A Business Unit of Bayer CropScience

P.O. Box 12014, 2 T.W. Alexander Drive

Research Triangle Park, NC 27709

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is reregistered in accordance with FIFRA provided that you:

1) Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data.

Signature of Approving Official:	Date:
	OCT 1 9 2009
Joanne Miller	
Product Manager 23	
Herbicide Branch	·
Registration Division (7505P)	

Page 2 EPA Reg. 72155-77

- 2) Per the acute toxicity review, the text "wear long-sleeved shirt, long pants, and shoes and socks" must be added to the Hazards to Humans and Domestic Animals statements currently on the label.
- 3) Per the 2,4-D, Dicamba and MCPP-p REDs label tables, the following text must be added to the Environmental Hazard statements currently on the label:

"This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater."

- 4) Per the MCPP-p RED, a 30-day retreatment interval must be added to the label.
- 5) Per the MCPP-p RED, the following text must be added to the directions for use:

"Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays and oceans. Do not apply when windy. To prevent product run-off, do not over water to the point of runoff or apply when raining or when raining or when rain is expected that day."

- 6) The claim "NEW" may only remain on the label for 6 months from the date of this letter. To the Storage and Disposal section change "Storage Conditions" to "Pesticide Storage" and "Disposal" to "Pesticide Disposal".
- 7) To the Notice section, add "to the extent consistent with applicable law" in front of "the buyer assumes responsibility".

A stamped copy of the label is enclosed for your records. You must submit one copy of the final printed label before you release the product for shipment. Products shipped after 12 months from the date of this letter must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA. Your release for shipment of the product constitutes acceptance of these conditions. This label supersedes all other previously accepted labels. If you have any questions please call Erik Kraft at 703-308-9358 or email at Kraft.Erik@epa.gov.

3-14

Bayer Advanced Lawn Herbicide 3F Concentrate

This label contains two separate label statements for Outdoor Residential Pump sprayer/Hose-end Sprayer Concentrate product use and Outdoor Residential Ready-To-Spray product use.

- · Kills Lawn Weeds plus Crabgrass [in One Step]
- Kills Over 200 Lawn Weeds Plus Crabgrass in One Easy Step
- Kills Crabgrass plus Broadleaf Weeds
- Kills Listed Weeds
- Kills Hard-to-Control Weeds
- Systemic [Herbicide] [Activity]
- Kills Both Broadleaf and Grassy Weeds
- Kills Dandelion, Crabgrass, Clover, plus 200 Other listed Weeds
- Rainproof in [1] hour[s]
- New!
- Two-in-One Product that Kills Crabgrass & Broadleaf Weeds
- Two Types of Weeds; One Solution
- Guaranteed* Not to harm your lawn
- All-In-One Weed Killer [for lawns]
- · Starts working on contact
- Kills even tough weeds roots & all
- Kills 200+ Weeds
- Unique Formula
- Fast results
- Kills lawn weeds plus crabgrass
- Gets the root
- Won't Harm [your] lawn [s]
- Will not harm your lawn
- · Kills tough weeds
- Kills the weeds, not your lawn
- Kills to the root
- Kills weeds, not lawns

- One [easy] step
- Two products in one
- · Kills all major broadleaf weeds
- Weed protection system
- Visible results
- Kills weeds Fast
- Kills both Crabgrass & Broadleaf weeds
- Proprietary Formula that Kills Both Crabgrass and Broadleaf Weeds
- Effectively Kills Both Broadleaf and Weed grasses
- Save time and energy
- Always ready to spray
- · Easy to use
- Fast coverage
- Easy & Fast coverage
- Targets weeds
- · Kills weeds the first time, every time
- · Guaranteed* not to harm lawns
- Guaranteed* Results
- 100% Satisfaction Guaranteed* (or your money back)
- For use on Kentucky bluegrass, perennial ryegrass, fescue (except creeping fescue), centipedegrass, St. Augustinegrass (except Floratam), zoysiagrass
- Tough weed protection
- 4-way weed control action
- Exclusive 4-Way Weed control
- Proprietary [unique] [advanced] formula
- Attach to hose and spray

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

ACCEPTED

with COMMENTS

72155-77

†Protected by U.S. Patent No. 5,254,527 Net 32 fl oz. (946 mL)

CAUTION

See back panel for additional precautionary statements

Editorial Note - [Bracketed text] is optional language

(Pump Sprayer/Hose-end Sprayer label)

4/4

(FRONT PANEL)

Bayer Advanced Lawn Herbicide 3F Concentrate

[Editorial Note - Marketing Claims from page 1]

Active Ingredients:	
2,4-D, Triisopropanolamine salt	4.91%
Mecoprop-p, Potassium salt	
Dicamba, Potassium salt	0.77%
Fenoxaprop-p-ethyl [†]	0.41%
Other Ingredients	
	100.00%

†Protected by U.S. Patent No. 5,254,527

Net 32 fl oz. (946 mL)

CAUTION

See back panel for additional precautionary statements

OPEN Resealable Label for Directions & Precautions PRESS TO RESEAL

Bayer Advanced Lawn Herbicide 3F Concentrate's exclusive and unique formula combines 4 proven weed killers to kill over 200 types of lawn weeds plus crabgrass in one easy step. And it's specially formulated so it won't harm your lawn when used as directed. Rainproof in 1 hour. Controlling weeds plus crabgrass in your lawn has never been easier!



QUICK FACTS		
Marketing claims to be selected from the front page		
KILLS Broadleaf Weeds: Over 200 types including dandelions, clovers, chickweeds Grassy Weeds: Crabgrass, goosegrass, foxtail, sandbur, and many others		
WHERE TO USE	On lawns without harming lawn grasses – Centipedegrass, Fescue (except Creeping Fescue), Kentucky Bluegrass, Perennial Ryegrass, St. Augustinegrass (except Floratam), Zoysiagrass	
10 03E	(DO NOT USE on Bahiagrass, Bermudagrass, Buffalograss, Carpetgrass, Creeping Fescue, Floratam variety of St. Augustinegrass, Dichondra or desirable clovers.)	
[COVERAGE]	[This bottle treats 6,400 sq ft.]	
	For questions or comments, call toll-free 1-877 BayerAG (1-877-229-3724) or visit our website at www.BayerAdvanced.com	

FOR OUTDOOR RESIDENTIAL USE ONLY

Active Ingredients:	
2,4-D, Triisopropanolamine salt	4.91%
Mecoprop-p, Potassium salt	1.55%
Dicamba, Potassium salt	
Fenoxaprop-p-ethyl [†]	0.41%
Other Ingredients	
	100.00%

THIS PRODUCT CONTAINS: 2.64% 2,4-D acid equivalent, 1.32% Mecoprop-p acid equivalent, 0.66% Dicamba acid equivalent. Isomer Specific by AOAC Methods.

EPA Reg. No. 72155 - 77

EPA Est. No. XXX-AA-XX

DIRECTIONS FOR	IISE
	aw to use this product in a manner inconsistent with its labeling.
HOW MUCH TO USE	Measure area to be sprayed. Calculate 'square feet' by multiplying length times width. [This bottle will treat 6,400 sq. ft.] The maximum application rate to turf is 1.0 pound of 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.
HOW TO USE Shake Well Before Use	 PUMP-UP TANK SPRAYERS (Best for spot spraying individual weeds) Add appropriate amounts of water & product to sprayer tank – 3 fl. oz. (6 Tbs) product in 1 gallon of water treats 500 sq. ft. Close sprayer, shake well & pressurize. Adjust nozzle to a coarse spray. Large Areas – Apply evenly over area. Individual Weeds – Hold tip of wand about 12 inches from weed, aim at center of weed and spray. Occasionally re-pressurize sprayer to maintain a good spray pattern.
	 DIAL STYLE HOSE-END SPRAYERS (Best for spraying large areas of the lawn) Fill sprayer jar with enough product for area to be treated – 3 fl. oz. (6 Tbs) product treats 500 sq. ft. DO NOT ADD WATER! Close sprayer, set dial and apply. Pour any unused product back into this original container. CONCENTRATE-PLUS-WATER HOSE-END SPRAYERS (Best for spraying large areas of the lawn) Pour product into sprayer jar to the required fluid ounce level – 3 fl oz (6 Tbs) treats 500 sq. ft. Add the appropriate amount of water to the gallons level. Replace sprayer top, shake well and apply.
[FOR BEST RESULTS]	 [Spray when weeds are [small [young] and] actively growing.] [Spray when daily high temperature is below 90°F.] [Rain or watering 1 hour after application will not wash away effectiveness.] [Apply when wind is calm [to avoid spray drift onto desirable plants (fruits, vegetables, flowers, groundcovers, shrubs & trees)]] [Reapply in 2 weeks if necessary.] [Do not apply while grass is emerging from winter dormancy.] [Newly seeded lawns may be sprayed after the fourth mowing.] [Bare spots may be seeded 4 weeks after application.] [Do not mow for 1 to 2 days before & after spraying [Uncut weeds have more surface area to hold more weed killer. They are also more likely to be hit by the spray. Lawn clippings can get in the way of the herbicide hitting the weed. Weed killers need time to move throughout the weed – mowing may
HOW IT WORKS	 remove the treated parts.].] [Leave grass clippings on the lawn from the first 3 mowings after using this product. Do not use as compost or mulch.] [If lawn is dry, water 1 to 2 days before using this product. [Weeds take up herbicides best when not under stress.]] BAYER ADVANCED LAWN HERBICIDE 3F CONCENTRATE contains four proven weed killers that target lawn weeds and crabgrass. The herbicides enter the lawn weeds through their leaves and move inside the weeds, even down into the roots. And lawns will not be harmed when used as directed.

Black medic	Florida pusley	Spurweed
Carpetweeds	Foxtails	Thistles
Chickweeds	Ground ivy	Toadflax
Clovers	Heartleaf drymary	Wild carrot
Crabgrass	Henbit	Wild garlic
Creeping Charlie	Kikuyugrass	Wild geranium
Cudweed	Oxalis	Wild onion
Curly dock	Pennywort	& many other broadleaf & grass weeds
Dandelion	Plantains	(See appendix for a complete list of the
English daisy	Purslane	weeds controlled)
False dandelion	Red sorrel	
Filaree	Sheep sorrel	
•	Spurges	



Do not apply this product in a way that will contact any person, pet either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until spray has dried.

PRECAUTIONARY STATEMENTS

CAUTION

Hazards to Humans and Domestic Animals

- Harmful if swallowed
- · Causes moderate eye irritation.
- · Avoid contact with eyes, skin, or clothing.

After using this product:


- If product gets on clothing, remove clothing as soon as possible and launder clothing separately before reuse.
- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

IRST AID +	
If in Eyes	 Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for further treatment advice.
If Swallowed	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 1-877-229-3763 for medical emergency information. Active Ingredients: 4.91% 2,4-D Triisopropanolamine salt (CAS # 32341-80-3), 1.55% Mecoprop-p Potassium salt single isomer (CAS # 66423-5-0), 0.77% Dicamba, Potassium salt (CAS # 10007-85-9), and 0.41% Fenoxaprop-p-ethyl (CAS # 71283-80-2).

ENVIRONMENTAL HAZARDS

- This product is toxic to fish & aquatic invertebrates.
- Drift or runoff may harm fish, aquatic invertebrates and non-target plants.
- To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters.
 Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.



NOTICE: Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. The Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

*Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.



A Business Group of Bayer CropScience LP P.O. Box 12014, 2 T.W. Alexander Drive Research Triangle Park, North Carolina 27709

Lawn Herbicide 3F Concentrate (PENDING) 01/28/09

(Ready-To- Spray label) (FRONT PANEL)

Bayer Advanced Lawn Herbicide 3F Concentrate

This label contains two separate label statements for, Outdoor Residential Pump Sprayer/Hose-end Sprayer Concentrate product use and Outdoor Residential Ready-To-Spray product use.

[Editorial Note - Marketing Claims from page 1]

Active Ingredients:		·
2,4-D, Triisopropanolamine salt		4.91%
Mecoprop-p, Potassium salt		
Dicamba, Potassium salt		
Fenoxaprop-p-ethyl [†]	•	
Other Ingredients		
-	·	100.00%

Net 32 fl oz. (946 mL)

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for additional precautionary statements

†Protected by U.S. Patent No. 5,254,527

OPEN Resealable Label for Directions & Precautions PRESS TO RESEAL

Bayer Advanced Lawn Herbicide 3F Concentrate's exclusive and unique formula combines 4 proven weed killers to kill over 200 types of lawn weeds plus crabgrass in one easy step. And it's specially formulated so it won't harm your lawn when used as directed. Rainproof in 1 hour. Controlling weeds plus crabgrass in your lawn has never been easier!



QUICK FACTS	,
Marketing claims to be selected from the front page	
KILLS	Broadleaf Weeds: Over 200 types including dandelions, clovers, chickweeds Grassy Weeds: Crabgrass, goosegrass, foxtail, sandbur, and many others
WHERE TO USE	On lawns without harming lawn grasses – Centipedegrass, Fescue (except Creeping Fescue), Kentucky Bluegrass, Perennial Ryegrass, St. Augustinegrass (except Floratam), Zoysiagrass
10 032	(DO NOT USE on Bahiagrass, Bermudagrass, Buffalograss, Carpetgrass, Creeping Fescue, Floratam variety of St. Augustinegrass, Dichondra or desirable clovers.)
[COVERAGE]	[This bottle treats 6,400 sq ft.]
☎ 🖳	For questions or comments, call toll-free 1-877 BayerAG (1-877-229-3724) or visit our website at www.BayerAdvanced.com

FOR OUTDOOR RESIDENTIAL USE ONLY

Active Ingredients:	
2,4-D, Triisopropanolamine salt	4.91%
Mecoprop-p, Potassium salt	1.55%
Dicamba, Potassium salt	0.77%
Fenoxaprop-p-ethyl [†]	0.41%
Other Ingredients	
	100.00%

THIS PRODUCT CONTAINS: 2.64% 2,4-D acid equivalent, 1.32% Mecoprop-p acid equivalent, 0.66% Dicamba acid equivalent. Isomer Specific by AOAC Methods

EPA Reg. No. 72155 -77

EPA Est. No. XXX-AA-XX

DIRECTIONS FOR	USE
It is a violation of Federal I	aw to use this product in a manner inconsistent with its labeling.
HOW MUCH TO USE	Measure area to be sprayed. Calculate 'square feet' by multiplying length times width. This bottle will treat 6,400 sq. ft. The maximum application rate to turf is 1.0 pound of 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast aplications per treatment site is 2 per year.
HOW TO USE	CONNECT 1. Connect sprayer to garden hose. 2. Turn on water.
Shake Well Before Use	 Extend hose to farthest area to be treated, point nozzle in direction to be sprayed. Turn on sprayer - < Insert sprayer-specific language here > Calmar RTS sprayer: press lock tab while using other hand to turn knob counter-clockwise to ON position. Green Garden RTS sprayer: turn sprayer valve to ON position. Hayes HP-2 RTS sprayer: press lock tab while using other hand to turn knob clockwise to ON position. Walk at a steady pace while spraying, using an even sweeping motion, and slightly overlap the treated areas. Work back toward the faucet to avoid walking over wet, treated grass.
	 Turn off sprayer - < Insert sprayer-specific language here > Calmar RTS sprayer: turn knob clockwise to OFF position. Green Garden RTS sprayer: [Nothing required]. Hayes HP-2 RTS sprayer: turn knob counter-clockwise to OFF position. Turn off water at faucet. Relieve pressure in hose by turning sprayer knob back to ON position until water slows to a drip then turn sprayer knob to OFF position. Disconnect sprayer from hose.
[FOR BEST RESULTS]	 [Spray when weeds are [small [young] and] actively growing.] [Spray when daily high temperature is below 90°F.] [Rain or watering 1 hour after application will not wash away effectiveness.] [Apply when wind is calm [to avoid spray drift onto desirable plants (fruits, vegetables, flowers, groundcovers, shrubs & trees)]] [Reapply in 2 weeks if necessary.]
	 [Do not apply while grass is emerging from winter dormancy.] [Newly seeded lawns may be sprayed after the fourth mowing.] [Bare spots may be seeded 4 weeks after application.] [Do not mow for 1 to 2 days before & after spraying [Uncut weeds have more surface area to hold more weed killer. They are also more likely to be hit by the spray. Lawn clippings can get in the way of the herbicide hitting the weed. Weed killers need time to move throughout the weed – mowing may remove the treated parts.].] [Leave grass clippings on the lawn from the first 3 mowings after using this product. Do not use as compost or mulch.] [If lawn is dry, water 1 to 2 days before using this product. [Weeds take up herbicides best when not under stress.]]
HOW IT WORKS	BAYER ADVANCED LAWN HERBICIDE 3F CONCENTRATE contains four proven weed killers that target lawn weeds and crabgrass. The herbicides enter the lawn weeds through their leaves and move inside the weeds, even down into the roots. And lawns will not be harmed when used as directed.

Kills over 200 weeds including:

Carpetweeds
Chickweeds
Clovers
Crabgrass
Creeping Charlie
Cudweed
Curly dock
Dandelion
English daisy

False dandelion

Filaree

Florida pusley Foxtails Ground ivy Heartleaf drymary Henbit

Heartleaf dryr Henbit Kikuyugrass Oxalis Pennywort Plantains Purslane Red sorrel Sheep sorrel Spurges Spurweed Thistles Toadflax Wild carrot Wild garlic

Wild geranium
Wild onion
& many other broadleaf & grass weeds
(See appendix for a complete list of the

weeds controlled)



Do not apply this product in a way that will contact any person, pet either directly or through drift. Keep people and pets out of the area during application. Do not allow people (other than applicator) or pets to enter the treated area until spray has dried.

PRECAUTIONARY STATEMENTS

CAUTION

Hazards to Humans and Domestic Animals

- Harmful if swallowed
- · Causes moderate eye irritation.
- · Avoid contact with eyes, skin, or clothing.

After using this product:

- If product gets on clothing, remove clothing as soon as possible and launder clothing separately before reuse.
- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

FIRST AID +	
If in Eyes	 Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for further treatment advice.
If Swallowed	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 1-877-229-3763 for medical emergency information. Active Ingredients: 4.91% 2,4-D Triisopropanolamine salt (CAS # 32341-80-3), 1.55% Mecoprop-p Potassium salt single isomer (CAS # 66423-5-0), 0.77% Dicamba, Potassium salt (CAS # 10007-85-9), and 0.41% Fenoxaprop-p-ethyl (CAS # 71283-80-2).

ENVIRONMENTAL HAZARDS

- This product is toxic to fish & aquatic invertebrates.
- Drift or runoff may harm fish, aquatic invertebrates and non-target plants.
- To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters.
 Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

Important – Directions for Storage and Disposal			
Storage Conditions	Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing		
Disposal —	If empty: Non-refillable container. Do not refill or reuse container. Place empty container in trash or offer for recycling if available.		
	If partly filled: Call your local solid waste agency or toll free 1-877-229-3724 for disposal instructions. Never place unused product down any indoor or outdoor drain.		

NOTICE: Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. The Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

*Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.



A Business Group of Bayer CropScience LP P.O. Box 12014, 2 T.W. Alexander Drive Research Triangle Park, North Carolina 27709

Lawn Herbicide 3F RTS (PENDING) 01/28/09

WEEDS CONTROLLED

Annual Yellow Sweetclover Aster Austrian Fieldcress Barnyardgrass Bedstraw Beggarticks Betony, Florida Bindweed, Field Bird Vetch Bitter Wintercress Bittercress, Hairy

Bittercress, Hairy Bitterweed Black-eyed Susan Black Medic Black Mustard Blackseed Plantain Blessed Thistle Bloodflower Milkwee

Bloodflower Milkweed
Blue Lettuce
Blue Vervain
Bracted Plantain
Brassbuttons
Bristly Oxtongue
Broadleaf Dock
Broadleaf Plantain
Broomweed
Buckhorn
Buckhorn Plantain
Bulbous Buttercup

Bull Thistle
Bullnettle
Burclover
Burdock
Burning Nettle
Burweed
Buttercup
Buttonweed
Canada Thistle
Carolina Geranium

Carpetweed Catchweed Bedstraw

Catnip Catsear

Chickweed, Common Chickweed, Mouseear

Chicory
Cinquefoil
Clover, Crimson
Clover, Hop
Clover, Red
Clover, Strawberry
Clover, Sweet
Clover, White
Cocklé

Cocklebur

Common Mullein Crabgrass , Large/Hairy Crabgrass, Small/Smooth

Crabgrass, Small Creeping Charlie Creeping Jenny Cudweed Curly Dock Daisy, English Daisy, Fleabane Daisy, Oxeye Dandelion Dichondra Dogbane Dogfennel

Dollarweed
Elderberry
False Dandelion
False Flax
False Sunflower
Fiddleneck
Florida Pusley
Foxtail
Frenchweed

Galinsoga
Goathead
Goldenrod
Goosegrass
Ground Ivy
Gumweed
Hairy Fleabane
Hawkweed

Healall Heartleaf Drymary Heath Aster Hedge Bindweed Hedge Mustard

Hedge Mustard Hemp Henbit Hoary Cress Hoary Plantain Hoary Vervain Horsenettle

Japanese Stiltgrass

Jimsonweed Johnsongrass (seedling)

Johnsongrass (se Knawel Knotweed Kochia Lambsquarters Lespedeza Mallow Matchweed Mexicanweed Milk Vetch Morningglory Mouseear Hawkweed

Mugwort Musk Thistle Mustard

Narrowleaf Plantain Narrowleaf Vetch

Nettle

Orange Hawkweed Oriental Cocklebur

Oxalis
Panicum
Parsley-piert
Parsnip
Pearlwort
Pennycress
Pennywort
Pepperweed
Pigweed

Pineywoods Bedstraw
Plains Coreopsis (Tickseed)

Plantain Poison Ivy Poison Oak Pokeweed Poorjoe

Prairie Sunflower Prickly Lettuce Prickly Sida Prostrate Knotweed Prostrate Pigweed Prostrate Spurge Prostrate Vervain Puncturevine

Purslane, Common Ragweed Red Sorrel Redroot Pigweed Redstem Filaree Rough Cinquefoil Rough Fleabane Russian Pigweed Russian Thistle Sandbur Scarlet Pimpernel

Scotch Thistle
Sheep Sorrel
Shepherdspurse
Silver Crabgrass
Slender Plantain
Smallflower Galinsoga
Smooth Dock
Smooth Pigweed

Sorrel Sowthistle Spanishneedles Speedwell
Spiny Amaranth
Spiny Cocklebur
Spiny Sowthistle
Spotted Catsear
Spotted Spurge
Sprangletop
Spurweed
Stinging Nettle
Strawberry, India Mock

Tall Nettle
Tall Vervain
Tansy Mustard
Tansy Ragwort
Tanweed

Thistle **Trailing Crownvetch** Tumble Mustard Tumble Pigweed Velvetleaf Venice Mallow Virginia Buttonweed Virginia Creeper Virginia Pepperweed Water Pennywort Wavyleaf Bullthistle Western Clematis Western Salsify White Mustard Wild Aster Wild Buckwheat

Wild Four-o'clock
Wild Garlic
Wild Geranium
Wild Lettuce
Wild Marigold
Wild Mustard
Wild Onion
Wild Parsnip
Wild Radish
Wild Rape
Wild Strawberry
Wild Sweet Potato
Wild Vetch

Wild Carrot

Woodsorrel Wooly Croton Wooly Morningglory Wooly Plantain Wormseed Yarrow

Yarrow Yellow Rocket

Yellowflower Pepperweed and other broadleaf weeds