


72155-87

04/30/2009

1/16

	U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505P) Ariel Rios Building 1200 Pennsylvania Ave., NW Washington, D.C. 20460		EPA Reg. Number: 72155-87	Date of Issuance: APR 30 2009
	NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration (under FIFRA, as amended)		Term of Issuance: Conditional	Name of Pesticide Product: Lawn 3iP Herbicide Concentrate
Name and Address of Registrant (include ZIP Code): Bayer Advanced 2 T.W. Alexander Drive RTP, NC 27709				
Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.				
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:</p> <ol style="list-style-type: none"> 1. Submit and/or cite all data required for the registration/reregistration review of this product when the Agency requires all registrants of similar products to submit data. 2. Make the following label changes: <ol style="list-style-type: none"> A. Add the phrase "EPA Registration No.72155-87" to the label and the Establishment Number before you release the product for shipment. B. Add the "Net Contents" to the labeling prior to releasing this product for shipment. C. Delete the words "exclusive" and "unique" that describes the formula on pages 3 and 8 				
COMMENTS CONTINUED ON PAGE 2 OF THIS NOTICE OF REGISTRATION.				
Signature of Approving Official: Joanne I. Miller Product Manager 23 Herbicide Branch Registration Division (7505P)			Date: APR 30 2009	

page 2
EPA Reg. No. 72155-87
Comments Continued:

and the claim "has never been easier" also on pages 3 and 8.

3. Submit the data listed below:

Submit the following data required for the registration of this pesticide product within 1 year from the date of this Notice of Registration:

<u>EPA Guideline Data Number</u>	<u>Guideline Descriptor</u>
830.6317	Storage Stability Study
830.6320	Corrosion Characteristics

Observations should be made at 0, 3, 6, 9 and 12 month intervals and submit the results electronically along with a hard copy.

- 4. Submit to the EPA Analytical Laboratory, 701 Mapes Road, Fort Meade, MD 20755-5350 a validated enforcement analytical method (EPA Guidelines 830.1800), within 6 months from the date of this Notice of Registration.
- 5. Submit one (1) copy of the final printed labeling before you release this product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product under the enclosed stamped copy of the label constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Enclosure

3/16

(FRONT PANEL)

Lawn 3iP Herbicide Concentrate/Ready-to-Spray

[ABN: Season Long Weed Control for Lawns Concentrate/Ready-to-Use or Southern Season Long Weed Control for Lawns Concentrate/ Ready-to-Use]

This label contains two separate label statements for, **Pump Sprayer/Hose-end Sprayer Concentrate** product use and **Pre-packed Hose-end Ready-to-Spray** product use.

▪ *Editorial Note – Marketing Claims Positioned Here*

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt ¹	4.73%
Dicamba, potassium salt ²	0.52%
Mecoprop-p, potassium salt ³	1.10%
Isoxaben ⁴	2.63%
OTHER INGREDIENTS:	91.02%
TOTAL	100.00%

[Active ingredient statement]:

THIS PRODUCT CONTAINS:

0.337 lb 2,4-D acid equivalent per gal or 3.94%
 0.038 lb Dicamba acid equivalent per gal or 0.44%
 0.080 lb Mecoprop-p acid equivalent per gal or 0.94%
 0.225 lb Isoxaben per gal or 2.63%
 Isomer Specific by AOAC Method.

¹CAS No. 2008-39-1
²CAS No. 10007-85-9
³CAS No. 66423-05-0
⁴CAS No. 82558-50-7

Note: If the design, format, or small size of the container labels makes it impractical to present the entire ingredient statement on the front panel, then the sub statement will be placed on the back panel.

Net Contents: 32 fl oz (946 mL)

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See back panel for additional precautionary statements.

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

APR 30 2000
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
is needed, for the pesticide
registered under EPA Reg. No.

72155-87

Editorial Note – [Bracketed text] is optional language

4
16

(Pump Sprayer/Hose-end Sprayer Concentrate Label)

(FRONT PANEL)

Lawn 3iP Herbicide Concentrate

[ABN: Season Long Weed Control for Lawns Concentrate or Southern Season Long Weed Control for Lawns Concentrate]

▪ Editorial Note – Marketing Claims Positioned Here

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt ¹	4.73%
Dicamba, potassium salt ²	0.52%
Mecoprop-p, potassium salt ³	1.10%
Isoxaben ⁴	2.63%
OTHER INGREDIENTS:	<u>91.02%</u>
TOTAL	100.00%

[Active ingredient statement]:

THIS PRODUCT CONTAINS:

0.337 lb 2,4-D acid equivalent per gal or 3.94%
 0.038 lb Dicamba acid equivalent per gal or 0.44%
 0.080 lb Mecoprop-p acid equivalent per gal or 0.94%
 0.225 lb Isoxaben per gal or 2.63%

¹CAS No. 2008-39-1
²CAS No. 10007-85-9
³CAS No. 66423-05-0
⁴CAS No. 82558-50-7

Isomer Specific by AOAC Method.

Note: If the design, format, or small size of the container labels makes it impractical to present the entire ingredient statement on the front panel, then the sub statement will be placed on the back panel.

Net Contents: 32 fl oz (946 mL)

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional precautionary statements.



Editorial Note – [Bracketed text] is optional language

(Back Panel)

OPEN Resealable Label for Directions & Precautions **PRESS TO RESEAL**

[The days of multiple applications to kill weeds are over!] Just one application of Lawn 3iP Herbicide Concentrate's [exclusive] [unique], [2-in-1], [3-in-1] [4-in-1] formula kills & prevents lawn weeds up to 6 months! [It delivers Season Long Weed Control that's simple and easy to use] [and it will not harm your lawn when used as directed]. [PLUS, when applied in Early Spring, prevents Crabgrass all season long]. Ideal for large areas – treats up to 10,000 square feet. [Rainproof in one hour.] Controlling weeds in your lawn [up to [6][X] months] has never been easier!



QUICK FACTS	
<i>Editorial Note – Marketing Claims Positioned Here</i>	
KILLS & PREVENTS	Kills over 200 broadleaf weeds such as chickweed, dandelion, dollarweed, henbit, clover & spurge Prevents common broadleaf weeds & grass weeds including crabgrass & annual bluegrass.
WHERE TO USE	ON ESTABLISHED LAWNS without harming lawn grasses †(Do not use on carpetgrass, Floratam variety of St. Augustinegrass, dichondra or desirable clovers.)
COVERAGE	This bottle treats 5,000 – 10,000 sq ft
 	For questions or comments, call toll-free 1-877 BayerAG (1-877-229-3924) or visit us on-line at www.BayerAdvanced.com

FOR OUTDOOR RESIDENTIAL USE ONLY

[Active ingredient statement]:




THIS PRODUCT CONTAINS:

- 0.337 lb 2,4-D acid equivalent per gal or 3.94%
- 0.038 lb Dicamba acid equivalent per gal or 0.44%
- 0.080 lb Mecoprop-p acid equivalent per gal or 0.94%
- 0.225 lb Isoxaben per gal or 2.63%

Isomer Specific by AOAC Method.


EPA Reg. No. 72155-XX

EPA Est. No.


DIRECTIONS FOR USE	
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.	
BEFORE YOU USE	<ul style="list-style-type: none"> • Use this product only on established lawns. • Apply when the air is calm to avoid spray drift onto desirable plants.
HOW MUCH TO USE	<ul style="list-style-type: none"> • Measure area to be sprayed. Calculate 'square feet' by multiplying length times width. • For best results, use spray solution immediately. If it is necessary to store, shake vigorously before use. Do not store for more than 3 days. <p>Bermudagrass, Buffalograss, Fescue, Kentucky Bluegrass, Perennial Ryegrass, Zoysiagrass 3.2 fl. oz. (6½ TBS) in 1 - 2 gallons of water treats 500 sq. ft.</p> <p>Bahiagrass, Centipedegrass, St. Augustinegrass (not for use on Floratam variety) 1.6 fl. oz. (3 ¼ TBS) in 1 - 2 gallons of water treats 500 sq. ft.</p> <p>Food utensils, such as tablespoons and measuring cups, should not be used for food purposes after use with any pesticide.</p> <p>The maximum application rate of this product is 6.4 fl. oz. per 1,000 sq. ft. per application per site. The maximum number of broadcast applications per treatment site is 2 per year.</p>
HOW TO USE	<ul style="list-style-type: none"> • Shake bottle well before use. • To prevent new weeds from growing, you must spray the entire area you want protected, not just the emerged weeds. • Wait 2 days after applying, then activate residual barrier by watering with ¼ - ½" of irrigation. <p><i>Best for spraying individual weeds & small areas.</i></p> <p>Add appropriate</p> <ol style="list-style-type: none"> 1. Add appropriate amounts of water & product to sprayer tank. 2. Close sprayer, shake well and pressurize. Adjust nozzle to a coarse spray. Individual weeds – Hold tip of wand about 12" from weed, aim at center of weed & spray. Also spray the area underneath & around the weed to prevent new weeds from growing. Small areas – apply evenly over area. 3. Occasionally re-pressurize sprayer as needed to maintain a good spray pattern. <p><i>Best for spraying large areas such as the entire lawn.</i></p> <ol style="list-style-type: none"> 1. Fill sprayer jar with enough product for the area to be treated. DO NOT ADD WATER. 2. Close sprayer, set dial, hook to garden hose and apply. To avoid contact with the treated area, begin by spraying area farthest from faucet walking backwards as you spray. 3. Pour any unused product back into this original container. <p><i>Best for spraying large areas such as the entire lawn.</i></p> <ol style="list-style-type: none"> 1. Pour product into sprayer jar to the required 'Fluid Ounce' level. 2. Add the appropriate amount of water to the 'Gallons' level. 3. Close sprayer, hook to garden hose, shake well & apply. To avoid contact with the treated area, begin by spraying area farthest from faucet walking backwards as you spray.
PUMP STYLE TANK SPRAYERS 	
DIAL-STYLE HOSE-END SPRAYERS 	
CONCENTRATE PLUS WATER HOSE-END SPRAYERS 	

<p>[FOR BEST RESULTS]</p>	<p>[WHEN TO APPLY]</p> <ul style="list-style-type: none"> • [Spray when <ul style="list-style-type: none"> - weeds are small and actively growing - temperatures are between 50 - 90°F] • [Do not apply while grass is emerging from dormancy.] • [This product also prevents crabgrass when applied early spring before the fourth mowing or before the temperatures are regularly in the 80's.] • [This product also prevents annual bluegrass (Poa annua) when applied August – December.] • [Certain hard-to-kill weeds may be retreated in 2 weeks if necessary.] [Spray individual weeds using this product in a tank sprayer or with Lawn 3iP Herbicide Ready-to-Use.] • [Heavy raking will disturb the weed preventative barrier and reduce the effectiveness of this product.] • [Newly seeded lawn may be sprayed after fourth mowing.] • [Bare spots may be seeded 60 days after application.] <p>[MOWING]</p> <ul style="list-style-type: none"> • [Do not mow for 1 to 2 days after spraying.] • [Clippings from the first three mowings should be left on the treated area. Do not use the clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.] <p>[WATERING]</p> <ul style="list-style-type: none"> • [If lawn is dry, water 1 to 2 days before using this product.] • [Rain or watering 1 hour after application will not wash away effectiveness.]
<p>HOW IT WORKS</p>	<p>This product contains a proven blend of three weed killers to kill existing weeds Plus a residual herbicide to keep them from coming back for up to 6 months. Weeds begin to wilt & die within [24] hours [and are completely dead in 14 – 28 days]. And, it won't harm your lawn when used as directed.</p>

<p>KILLS & PREVENTS</p> <p>[Weeds presented here to be chosen from Appendix 1.]</p> <p>and other broadleaf & grass weeds</p>

	<p>Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.</p>
---	--

<p>PRECAUTIONARY STATEMENTS</p>
<p style="text-align: center;">CAUTION</p> <p style="text-align: center;">Hazards to Humans and Domestic Animals</p>
<ul style="list-style-type: none"> • Harmful if absorbed through skin or if swallowed. • Causes moderate eye irritation. • Avoid contact with skin, eyes, or clothing. • Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. • Remove and wash contaminated clothing before reuse.

FIRST AID 	
If on Skin	<ul style="list-style-type: none"> ▫ Take off contaminated clothing. ▫ Rinse skin immediately with plenty of water for 15 – 20 minutes. ▫ Call a poison control center or doctor for further treatment advice.
If Swallowed	<ul style="list-style-type: none"> ▫ Call a poison control center or doctor immediately for treatment advice. ▫ Have person sip a glass of water if able to swallow. ▫ Do not induce vomiting unless told to do so by a poison control center or doctor. ▫ Do not give anything by mouth to an unconscious person.
If in Eyes	<ul style="list-style-type: none"> ▫ Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. ▫ Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. ▫ Call a poison control center or doctor for further treatment advice.
<p>NOTE: Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may call toll-free 1-877-229-3763 for medical emergency information.</p> <p>Active Ingredients: 4.73% 2,4-D, dimethylamine salt (CAS # 2008-39-1), 0.52% Dicamba, potassium salt (CAS # 10007-85-9), 1.10% Mecoprop-p, potassium salt (CAS # 66423-05-0) and 2.63% Isoxaben (CAS # 82558-50-7).</p>	

ENVIRONMENTAL HAZARDS
<ul style="list-style-type: none"> • This product is toxic to fish and aquatic invertebrates. Do not apply directly to water. • Drift or runoff maybe hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate. • To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

Important – Directions for Storage and Disposal	
Storage Conditions	Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing.
Disposal	<p>If empty: Non-refillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.</p> <p>If partly filled: Call your local solid waste agency or toll free 1-877-229-3724 for disposal instructions. Never place unused product down any indoor or outdoor drain.</p>



NOTICE: Research and testing have determined that the “Directions for Use” are appropriate for the proper use of this product under expected conditions. To the extent consistent with applicable law, the Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

***Money Back Guarantee:** If you are not satisfied with this product, we will gladly refund your original purchase price.

9
16

(Pre-packed Hose-end Ready-to-Spray Label)

(FRONT PANEL)

Lawn 3iP Herbicide Ready-to-Spray

[ABN: Season Long Weed Control for Lawns Ready-to-Use or Southern Season Long Weed Control for Lawns Ready-to-Use]

▪ *Editorial Note – Marketing Claims Positioned Here*

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt ¹	4.73%
Dicamba, potassium salt ²	0.52%
Mecoprop-p, potassium salt ³	1.10%
Isoxaben ⁴	2.63%
OTHER INGREDIENTS:	91.02%
TOTAL	100.00%

[Active ingredient statement]:

THIS PRODUCT CONTAINS:

0.337 lb 2,4-D acid equivalent per gal or 3.94%
 0.038 lb Dicamba acid equivalent per gal or 0.44%
 0.080 lb Mecoprop-p acid equivalent per gal or 0.94%
 0.225 lb Isoxaben per gal or 2.63%
 Isomer Specific by AOAC Method.

¹CAS No. 2008-39-1
²CAS No. 10007-85-9
³CAS No. 66423-05-0
⁴CAS No. 82558-50-7

Note: If the design, format, or small size of the container labels makes it impractical to present the entire ingredient statement on the front panel, then the sub statement will be placed on the back panel.

Net Contents: 32 fl oz (946 mL)

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional precautionary statements.

Editorial Note – [Bracketed text] is optional language

29
16

(Back Panel)


OPEN Resealable Label for Directions & Precautions **PRESS TO RESEAL**

[The days of multiple applications to kill weeds are over!] Just one application of Lawn 3iP Herbicide Ready-to-Spray's [exclusive] [unique], [2-in-1], [3-in-1] [4-in-1] formula kills & prevents lawn weeds up to 6 months! [It delivers Season Long Weed Control that's simple and easy to use] [and it will not harm your lawn when used as directed]. [PLUS, when applied in Early Spring, prevents Crabgrass all season long]. [Just Hook to Your Hose and Go -] Convenient sprayer top connects to garden hose and automatically dispenses the weed killer. Ideal for large areas – treats up to 10,000 square feet. [Rainproof in one hour.] Controlling weeds in your lawn [up to [6][X] months] has never been easier!



QUICK FACTS

Editorial Note – Marketing Claims Positioned Here

KILLS & PREVENTS	Kills over 200 broadleaf weed such as chickweed, dandelion, dollarweed, henbit, clover & spurge Prevents common broadleaf weeds & grass weeds including crabgrass & annual bluegrass.
WHERE TO USE	ON ESTABLISHED LAWNS without harming lawn grasses †(Do not use on carpetgrass, Floratam variety of St. Augustinegrass, dichondra or desirable clovers.)
COVERAGE	This bottle treats 5,000 – 10,000 sq ft. [Just hook to garden hose and go.]
	For questions or comments, call toll-free 1-877 BayerAG (1-877-229-3924) or visit us on-line at www.BayerAdvanced.com

FOR OUTDOOR RESIDENTIAL USE ONLY

[Active ingredient statement]:

THIS PRODUCT CONTAINS:

- 0.337 lb 2,4-D acid equivalent per gal or 3.94%
- 0.038 lb Dicamba acid equivalent per gal or 0.44%
- 0.080 lb Mecoprop-p acid equivalent per gal or 0.94%
- 0.225 lb Isoxaben per gal or 2.63%
- Isomer Specific by AOAC Method

EPA Reg. No. 72155-XX

EPA Est. No.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

BEFORE YOU USE

- Use this product only on established lawns.
- Apply when the air is calm to avoid spray drift onto desirable plants.

HOW MUCH TO USE

Measure area to be sprayed. Calculate 'square feet' by multiplying length times width.

Bermudagrass, Buffalograss, Fescue, Kentucky Bluegrass, Perennial Ryegrass, Zoysiagrass – this bottle (32 fl. oz.) treats 5,000 sq. ft.

Bahiagrass, Centipedegrass, St. Augustinegrass (not for use on Floratam variety) – this bottle (32 fl. oz.) treats 10,000 sq. ft.


The maximum application rate of this product is 32 fl. oz. per 5,000 sq. ft. per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

HOW TO USE

Shake Well Before Use

- To prevent new weeds from growing, you must spray the entire area you want protected, not just the emerged weeds.
- Wait 2 days after applying then activate residual barrier by watering with ¼ - ½" of irrigation.


Editorial Note – Either the Single Rate Sprayer or the Dual Rate Sprayer directions for use will appear on the commercial package but not both.



[For single rate sprayer]


CONNECT

1. Shake well before use.
2. Connect sprayer to garden hose.
3. Turn on water.



SPRAY

1. Extend hose to farthest area to be treated, point sprayer nozzle in direction to be sprayed.
2. Begin spraying – Push & hold lock tab away from knob, use other hand to turn knob [counter]-clockwise to ON position.
3. Walk at a steady pace while spraying, using an even sweeping motion, and slightly overlap the treated areas. Work back toward the faucet to avoid walking over wet, treated grass.



FINISH

1. To stop spraying, push & hold lock tab away from knob and use other hand to turn knob [counter]-clockwise to OFF position.
2. Turn off water at faucet.
3. Relieve water pressure in hose by turning sprayer to [ON] [WATER] position until water slows to a drip then turn sprayer OFF.
4. Disconnect sprayer from hose.

[For dual rate sprayer]





1. Connect your garden hose to the bottle. Be sure the blue water control valve on top of the spray nozzle is in the OFF position. Turn on water at faucet.



2. To apply high rate (treats 5,000 sq. ft.): Pull the small star-shaped blue product-control button out until it stops, Twist the blue product-control button so that the 'L' on the product-control button is at the top of the control valve AND the product control valve is aligned with the flat portion of the lock tab. Push the button in to open. Proceed to step #3.


To apply low rate (treats 10,000 sq. ft.): Pull the small star-shaped, blue product-control button out until it stops. Twist the product-control button so that the 'G' on the product-control button is at the top of the control valve AND the product control valve is aligned with the flat portion of the lock tab. Push the button in to open (The button will only go half in for this setting). Proceed to step #3.

	 <p>3. Extend hose to farthest area to be treated, point sprayer nozzle in direction to be sprayed. To begin spraying, point nozzle toward the area to be treated and turn the blue water control valve to the ON position. Tab will be horizontal. Walk at a steady pace while spraying, using an even, sweeping motion, and slightly overlap the treated areas. Work back toward the faucet to avoid walking over wet, treated grass.</p>  <p>4. To stop spraying, turn the blue water control valve on top of the spray nozzle to OFF. To close, push in the small round button opposite the star-shaped product-control button. Turn off faucet,</p> <p>5. Relieve water pressure by turning blue water control valve to the ON position until water slows to a drip. Turn blue water control valve back to OFF. Disconnect garden hose from bottle.</p>
--	--

<p>[FOR BEST RESULTS]</p>	<p>[WHEN TO APPLY]</p> <ul style="list-style-type: none"> • [Spray when <ul style="list-style-type: none"> - weeds are small and actively growing - temperatures are between 50 - 90°F] • [Do not apply while grass is emerging from dormancy.] • [This product also prevents crabgrass when applied early spring before the fourth mowing or before the temperatures are regularly in the 80's.] • [This product also prevents annual bluegrass (Poa annua) when applied August – December.] • [Certain hard-to-kill weeds may be retreated in 2 weeks if necessary.] [Spray individual weeds using this product in a tank sprayer or with Lawn 3IP Herbicide Ready-to-Use.] • [Heavy raking will disturb the weed preventative barrier and reduce the effectiveness of this product.] • [Newly seeded lawn may be sprayed after fourth mowing.] • [Bare spots may be seeded 60 days after application.] <p>[MOWING]</p> <ul style="list-style-type: none"> • [Do not mow for 1 to 2 days after spraying.] • [Clippings from the first three mowings should be left on the treated area. Do not use the clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.] <p>[WATERING]</p> <ul style="list-style-type: none"> • [If lawn is dry, water 1 to 2 days before using this product.] • [Rain or watering 1 hour after application will not wash away effectiveness.]
----------------------------------	---

<p>HOW IT WORKS</p>	<p>This product contains a proven blend of three weed killers to kill existing weeds Plus a residual herbicide to keep them from coming back for up to 6 months. Weeds begin to wilt & die within [24] hours and are completely dead in 14 – 28 days. And, it won't harm your lawn when used as directed.</p>
----------------------------	---

<p>KILLS & PREVENTS</p>	
<p>[Weeds presented here to be chosen from Appendix 1.]</p> <p>and other broadleaf & grass weeds</p>	


	<p>Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.</p>
---	--

13
16

PRECAUTIONARY STATEMENTS	
CAUTION	
Hazards to Humans and Domestic Animals	
<ul style="list-style-type: none"> • Harmful if absorbed through skin or if swallowed. • Causes moderate eye irritation. • Avoid contact with skin, eyes, or clothing. • Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. • Remove and wash contaminated clothing before reuse. 	

FIRST AID +	
If on Skin	<ul style="list-style-type: none"> ▪ Take off contaminated clothing. ▪ Rinse skin immediately with plenty of water for 15 – 20 minutes. ▪ Call a poison control center or doctor for further treatment advice.
If Swallowed	<ul style="list-style-type: none"> ▪ Call a poison control center or doctor immediately for treatment advice. ▪ Have person sip a glass of water if able to swallow. ▪ Do not induce vomiting unless told to do so by a poison control center or doctor. ▪ Do not give anything by mouth to an unconscious person.
If in Eyes	<ul style="list-style-type: none"> ▪ Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. ▪ Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. ▪ Call a poison control center or doctor for further treatment advice.
<p>NOTE: Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may call toll-free 1-877-229-3763 for medical emergency information.</p> <p>Active Ingredients: 4.73% 2,4-D, dimethylamine salt (CAS # 2008-39-1), 0.52% Dicamba, potassium salt (CAS # 10007-85-9), 1.10% Mecoprop-p, potassium salt (CAS # 66423-05-0) and 2.63% Isoxaben (CAS # 82558-50-7)</p>	

ENVIRONMENTAL HAZARDS	
<ul style="list-style-type: none"> • This product is toxic to fish and aquatic invertebrates. Do not apply directly to water. • Drift or runoff maybe hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate. • To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. 	

Important – Directions for Storage and Disposal	
Storage Conditions	Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing.
Disposal 	<p>If empty: Non-refillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.</p> <p>If partly filled: Call your local solid waste agency or toll free 1-877-229-3724 for disposal instructions. Never place unused product down any indoor or outdoor drain.</p>

NOTICE: Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. To the extent consistent with applicable law, the Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

14
16

***Money Back Guarantee:** If you are not satisfied with this product, we will gladly refund your original purchase price.



**A Business Unit of Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709**

Lawn 3iP (PENDING) 05/08/08, Resubmitted 04/01/09, Resubmitted 04/13/09

Appendix 1

KILLS			
Annual Bluegrass (<i>Poa annua</i>)	Cocklebur	Morningglory	Spiny Cocklebur
Annual Yellow Sweetclover	Common Groundsel	Mouseear Hawkweed	Spiny Sowthistle
Aster	Common Mullein	Mugwort	Spotted Catsear
Austrian Fieldcress	Crabgrass	Musk Thistle	Spotted Spurge
Barnyardgrass	Creeping Charlie	Mustard	Spurweed
Bedstraw	Creeping Jenny	Narrowleaf Plantain	Stinging Nettle
Beggarticks	Cudweed	Narrowleaf Vetch	Strawberry, India Mock
Betony, Florida	Curly Dock	Nettle	Tall Nettle
Bindweed, Field	Daisy, English	Orange Hawkweed	Tall Vervain
Bird Vetch	Daisy, Fleabane	Oriental Cocklebur	Tansy Mustard
Bitter Wintercress	Daisy, Oxeye	Oxalis Panicle Willowweed	Tanweed
Bittercress, Hairy	Dandelion	Parsley-piert	Telegraphplant
Bitterweed	Desert Rockpurslane	Parsnip	Texas Panicum
Black-eyed Susan	Dichondra	Pearlwort	Thistle
Blackleaved Fleabane	Dogbane	Pennsylvania Smartweed	Trailing Crownvetch
Black Nightshade	Dogfennel	Pennycress	Tumble Mustard
Black Medic	Dollarweed	Pennywort	Tumble Pigweed
Black Mustard	Elderberry	Pepperweed	Velvetleaf
Blackseed Plantain	Fall Panicum	Pigweed	Venice Mallow
Blessed Thistle	False Dandelion	Pineappleweed	Virginia Buttonweed
Bloodflower Milkweed	False Flax	Pineywoods Bedstraw	Virginia Creeper
Blue Lettuce	False Sunflower	Plains Coreopsis (Tickseed)	Virginia Pepperweed
Blue Vervain	Fiddleneck	Plantain	Water Pennywort
Bracted Plantain	Florida Pusley	Poison Ivy	Wavyleaf Bullthistle
Brassbuttons	Frenchweed	Poison Oak	Western Clematis
Bristly Oxtongue	Galinsoga	Pokeweed	Western Salsify
Broadleaf Dock	Goathead	Poorjoe	White Mustard
Broadleaf Plantain	Goldenrod	Prairie Sunflower	Wild Aster
Broomweed	Ground Ivy	Prickly Lettuce	Wild Buckwheat
Buckhorn	Gumweed	Prickly Sida	Wild Carrot
Buckhorn Plantain	Hairy Fleabane	Prostrate Knotweed	Wild Celery
Bulbous Buttercup	Hawkweed	Prostrate Pigweed	Wild Four-o'clock
Bull Thistle	Healall	Prostrate Spurge	Wild Garlic
Bullnettle	Heartleaf Drymary	Prostrate Vervain	Wild Geranium
Burclover	Heath Aster	Puncturevine	Wild Lettuce
Burdock	Hedge Bindweed	Purslane, Common	Wild Marigold
Burning Nettle	Hedge Mustard	Ragweed	Wild Mustard
Burweed	Hemp	Red Sorrel	Wild Oat
Buttercup	Henbit	Redroot Pigweed	Wild Onion
Buttonweed	Hoary Cress	Redstem Filaree Rough	Wild Parsnip
Canada Thistle	Hoary Plantain	Cinquefoil	Wild Radish
Carolina Geranium	Hoary Vervain	Rough Fleabane	Wild Rape
Carpetweed	Horsenettle	Russian Pigweed	Wild Strawberry
Catchweed Bedstraw	Horseweed	Russian Thistle	Wild Sweet Potato
Catnip	Italian Ryegrass	Scarlet Pimpernel	Wild Vetch
Catsear	Jimsonweed	Scotch Thistle	Witchgrass
Chickweed, Common	Junglerice	Sheep Sorrel	Woodsorrel
Chickweed, Mouseear	Knawel	Shepherdspurse	Wooly Croton
Chicory	Knotweed	Sibara	Wooly Morningglory
Cinquefoil	Kochia	Slender Plantain	Wooly Plantain
Clover	Lambsquarters	Smallflower Galinsoga	Wormseed
Clover, Crimson	Lanceleaf Groundcherry	Smooth Dock	Yarrow
Clover, Hop	Lespedeza	Smooth Pigweed	Yellow Foxtail
Clover, Red	London Rocket	Sorrel	Yellow Rocket
Clover, Strawberry	Mallow	Southwest Cupgrass	Yellowflower Pepperweed
Clover, Sweet	Matchweed	Sowthistle	and other broadleaf weeds
Clover, White	Mexicanweed	Spanishneedles	Kills & Prevents
Cockle	Milk Vetch	Speedwell	Prevents only
		Spiny Amaranth	

16
16

Appendix 2

Optional Marketing Claims:

- One quart covers [up to] 5,000 [10,000] square feet.
- 1 quart treats [up to] 5,000 [10,000] sq. ft.
- Treats 5,000 [10,000] sq. ft.
- Keep from freezing.
- Kills [Controls] even the tough weeds...roots and all!
- Kills Broadleaf Weeds
- Kills Hard-to-Control Weeds
- Systemic [Herbicide] [Activity]
- Kills Dandelion, Clover, plus 200 Other listed Weeds
- Kills the root.
- Water based.
- Kills [Controls] [200+] broadleaf weeds [to the root].
- Mixes with Water - Concentrate
- Attach to [garden] hose and spray
- Hook to Hose and Go
- Proven performance
- Simple, [Convenient], [Easy] to use
- Two-in-One Product that Kills Weeds and Stops them from coming back
- 3-in-1 Product: Kills Weeds, Prevents New Weeds [up to 6 Months] & Prevents Crabgrass
- New [Formulation]!
- [Brand] New [Formula]
- Kills 200+ Weeds (to the root)
- Rainproof in 1 Hour
- Starts Working Immediately
- All Weather Formula (rainproof in 1 hour)
- 100% Satisfaction Guaranteed* (or your money back)
- Season Long Weed Control [with 1 Application]
- 1 Application Provides Season Long Weed Control
- Kills & Prevents Weeds Up to 6 Months
- Kills Weeds & Prevents New Weeds [from Coming Back] [up to 6 months]
- Protects Your Lawn From Weeds up to 6 Months
- Keeps weeds out [of your lawn] up to 6 months
- Prevents [New] Weeds Up to 6 Months
- [Exclusive] [Unique], [Proprietary] [2-in-1] [3-in-1] [4-in-1] Formula
- Prevents [New] Weeds up to 6 Months
- Weed Free Lawn [Up to 6 Months]
- Crabgrass Free Lawn [Up to 6 Months]
- Visible Results in 24 Hours
- Prevents Crabgrass [up to 6 months]
- Season Long Crabgrass Control
- PLUS Prevents Crabgrass!
- Prevents Crabgrass All Season
- Kills Over 200 Lawn Weeds in One Easy Step
- Kills the weeds not the lawn†
- Won't Harm your Lawn†
- Won't Harm Lawns† (when used as directed)
- Kills [Current] Weeds to the Root
- Kills [Current] [Existing] Weeds [Up to 6 Months]
- Prevents Grasses Up to 6 Months
- Season Long Weed Control
- Season Long Weed Control [Killer]
- Extended [Weed] Control [Killer]
- [Up to] 6 Month Weed Control
- Use on Bermudagrass, Centipede, St. Augustine and Zoysia lawns.
- Won't harm listed lawns [when used as directed] †