

72155-104

04-26-2011

1/19



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

Date of Issuance:

72155-104

4-26-11

Term of Issuance: Unconditional

Name of Pesticide Product:

Lawn 3FL2 Herbicide
Concentrate/Ready-to-Spray

NOTICE OF PESTICIDE:

Registration
 Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Bayer Advanced / Division of Bayer CropScience LP
P.O. box 12014
2 T.W. Alexander Drive
Research Triangle Park, NC 27709

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration /registration review of your product when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Kable Bo Davis
Product Manager 25
Herbicide Branch, Registration Division (7505P)

Date:

4-26-11

2. Submit one copy of the final printed label for the record before you release the product for shipment.
3. Submit the results of the one year storage stability and corrosion characteristics studies within 18 months of the date of this notice.
4. Add the Correct EPA Reg. # (72155-104) and EPA Est. # to the label.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records. If you have any questions please contact Erik Kraft at 703-308-9358 or kraft.erik@epa.gov.


3
19

Lawn 3FL2 Herbicide Concentrate/Ready-to-Spray

[ABN: Southern Season Long Weed Control for Lawns¹ Concentrate/Ready-to-Spray]

This label contains four separate label statements as follows:

- A) Pump Sprayer/Hose-end Sprayer Concentrate
- B) Pre-packed Hose-end Ready-to-Spray



M-393633-01-1

E-SUBMISSION

194

(FRONT PANEL)

(Part A: Pump Sprayer/Hose-end Sprayer Concentrate Label)

Lawn 3FL2 Herbicide Concentrate

[ABN: Southern Season Long Weed Control for Lawns¹ Concentrate]

▪ *Editorial Note – Marketing Claims Positioned Here [see Appendix 2]*

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	3.45%
Dicamba, potassium salt.....	0.379%
Mecoprop-p potassium salt.....	0.812%
Indaziflam.....	0.242%
OTHER INGREDIENTS:	95.117%
TOTAL	100.00%

THIS PRODUCT CONTAINS:

- 0.249 lb 2,4-D acid equivalent per gal or 2.86%
- 0.028 lb Dicamba acid equivalent per gal or 0.324%
- 0.060 lb Mecoprop-p equivalent per gal or 0.689%
- 0.021 lb Indaziflam per gal or 0.242%
- Isomer Specific by AOAC Method.

Note: If the design, format, or small size of the container labels makes it impractical to present the entire ingredient statement on the front panel, then the sub statement will be placed on the back panel.

Net Contents : 40 fl. oz. (1.18 L), 32 fl. oz. (946 mL), 24 fl. oz. (709 mL), 2.25 fl. oz. (66 mL), or other sizes within this range.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional precautionary statements.

Editorial Note – [Bracketed text] is optional language

5
19

(Back Panel)

OPEN Resealable Label for Directions & Precautions **PRESS TO RESEAL**

[The days of multiple applications to kill weeds are over!]. Just one application of Lawn 3FL Herbicide Concentrate's [patented] [exclusive] [proprietary] [4-in-1] [3-in-1] [2-in-1] formula kills over 200 weeds & prevents new lawn weeds for up to 6 months!. [It delivers Season Long Weed Control [up to 6 months] that's simple and easy to use] [and it will not harm your lawn when used as directed]...[No need [to re-apply every couple of weeks] [for frequent re-application]]. [PLUS, when applied in Early Spring, prevents... Crabgrass all season long]. Ideal for large areas – treats [up to] 7,100 square feet. [Packaging-related note: Adjust sq. ft. coverage proportionately for sizes other than 32 fl. oz.] [Rainproof in [1][4][24] hour.] Controlling [listed] weeds [and preventing crabgrass] in your lawn up to 6 months has never been easier!



QUICK FACTS	
<i>Editorial Note – Marketing Claims Positioned Here [see Appendix 2]</i>	
KILLS & PREVENTS For up to 6 Months with one application	[Kills over 200 broadleaf weeds such as Dollarweed, Chickweed, Dandelion, Henbit, Clover & Spurge] [Prevents [XX] [listed] grassy & broadleaf weeds such as Crabgrass, Annual Bluegrass, Goosegrass, White Clover, Dandelion, Plantain, Henbit and Black Medic] <i>Choose from marketing claims</i>
WHEN TO [APPLY] [USE]	[Apply anytime – spring or fall] <i>or choose from other applicable marketing claim</i>
WHERE TO USE	ON ESTABLISHED LAWNS of the turf types listed below: Bermudagrass, Centipedegrass, St. Augustinegrass, and Zoysia DO NOT use on improved varieties of St. Augustinegrass (such as Floratam), Fescue, Bentgrass, Carpetgrass, Kentucky Bluegrass, Kikuyugrass, Ryegrass or Dichondra lawns
HOW TO USE	To prevent new weeds from growing, spray entire area you want protected. Wait at least [24 hours] [1 day] after applying, then activate preventative weed barrier by watering. See Directions for Use for further details.
COVERAGE	This bottle treats 5,000 – 7,100 sq ft. [(7,100 sq ft for centipede and St. Augustine [grasses] [lawns]), [(5,000 sq ft. for Bermudagrass and Zoysia Lawns)] [Packaging-related note: Adjust sq. ft. coverage proportionately for sizes other than 32 fl. oz.]
[OPTIONAL GRAPHIC IMAGES:]	WAIT
	For questions or comments, call toll-free 1-877 BayerAG (1-877-229-3924) [or visit our website at www.bayeradvanced.com]

ACTIVE INGREDIENTS:	
2,4-D, dimethylamine salt	3.45%
Dicamba, potassium salt.....	0.379%
Mecoprop-p potassium salt.....	0.812%
Indaziflam*	0.242%
OTHER INGREDIENTS:	95.117%
TOTAL	100.00%

* Protected by patent number US 6,069,114

THIS PRODUCT CONTAINS:
 0.249 lb 2,4-D acid equivalent per gal or 2.86%
 0.028 lb Dicamba acid equivalent per gal or 0.324%
 0.060 lb Mecoprop-p equivalent per gal or 0.689%
 0.021 lb Indaziflam per gal or 0.242%
 Isomer Specific by AOAC Method.

EPA Reg. No. 72155-XX
 EPA Est. No.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

BEFORE YOU USE
Protect from Freezing

- Use this product only on established lawns – new lawns must be at least 4 [or X] months old if sodded, 2 [or X] months old if sprigged & 12 [or X] months old if seeded.
- Do not overlap treated areas or make repeat applications within 3 months as over application could damage the lawn.
- Do not apply to recently aerated or verticutted turf until it has recovered.
- Apply when the air is calm to avoid spray drift onto desirable plants.
- This product may affect young seedlings downslope of treated area after heavy rainfall. Do not apply if heavy rainfall is expected within the next 48 hours.

HOW MUCH TO USE

- Measure area to be sprayed. Calculate 'square feet' by multiplying length times width.
 - Use spray solution within 3 days of preparation.
- Bermudagrass, Zoysiagrass:
3.2 fl. oz. (6½ TBS) in [1] [1-2] [2] gallons of water treats 500 sq. ft. [Packaging-related note: Adjust sq. ft. coverage proportionately for sizes other than 32 fl. oz.]
- Centipedegrass, St. Augustinegrass (not for use on improved varieties, such as Floratam), [Kentucky bluegrass] or lawns containing one of more of these grasses:
2 ¼ fl. oz. (4 ½ TBS) in [1] [1-2] [2] gallons of water treats 500 sq. ft. [or 4.5 fl. oz. (9 TBS) in [1] [1-2] [2] gallons of water treats 1,000 sq. ft.] [Packaging-related note: Adjust sq. ft. coverage proportionately for sizes other than 32 fl. oz.]
- Food utensils, such as tablespoons and measuring cups, should not be used for food purposes after use with any pesticide.
- The maximum application rate of this product is 6.4 fl. oz. per 1,000 sq. ft. per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

HOW TO USE

- Shake bottle well before use.
- To prevent new weeds from growing, you must spray the entire area you want protected, not just the emerged weeds.
- [Wait [1 day] [24 hours] after applying then] activate preventative weed barrier by watering with ¼ - ½" of irrigation.

PUMP STYLE TANK SPRAYERS



- Best for spraying individual weeds & small areas.*
1. Add appropriate amounts of water & product to sprayer tank.
[For 2.25 fl. oz. container: Pour entire contents into spray tank and fill with water to [1] [2] gallon mark].
 2. Close sprayer, shake well and pressurize. Adjust nozzle to a coarse spray.
Individual weeds – Hold tip of wand about 12 inches from weed, aim at center of weed and spray. Also spray the area underneath & around the weed to prevent new weeds from growing.
Small areas – Apply evenly over area.
 3. Occasionally re-pressurize sprayer to maintain a good spray pattern.


DIAL-STYLE HOSE-END SPRAYERS




- Best for spraying large areas such as the entire lawn.*
1. Fill sprayer jar with enough product for the area to be treated. DO NOT ADD WATER.
 2. Close sprayer, set dial, hook to garden hose and apply. To avoid contact with the treated area, begin by spraying area farthest from faucet walking backwards as you spray.
 3. Pour any unused product back into this original container.

CONCENTRATE PLUS WATER HOSE-END SPRAYERS

- Best for spraying large areas such as the entire lawn.*
1. Pour product into sprayer jar to the required 'Fluid Ounce' level.
 2. Add the appropriate amount of water to the 'Gallons' level.
 3. Close sprayer, hook to garden hose, shake well, & apply. To avoid contact with the treated area, begin by spraying area farthest from faucet walking backwards as you spray.

	
<p>[FOR BEST RESULTS]</p>	<p>[WHEN TO APPLY]</p> <ul style="list-style-type: none"> • [Spray when: - weeds are small and actively growing - temperatures are between 50 – [85][90]°F.] • [Do not apply while grass is emerging from dormancy.] • [This product also prevents crabgrass & goosegrass [other listed weeds as appropriate] when applied early spring before the fourth mowing] • [This product also prevents annual bluegrass (<i>Poa annua</i>) when applied August – November.] • [Certain hard-to-kill weeds may be retreated in 30 days if necessary. Spray individual weeds using this product or any other Bayer Advanced lawn weed herbicide] • [Heavy raking will disturb the weed preventative barrier and reduce the effectiveness of this product.] • [Lawns may be seeded 8 months (St. Augustinegrass and Centipedegrass) or 12 months (Bermudagrass and Zoysia) after applying this product. Lawns may be sodded 4 [or X] [months] and sprigged 2 [or X] months after applying this product.] <p>[MOWING]</p> <ul style="list-style-type: none"> • [Do not mow for 1 to 2 days after spraying.] • [Clippings from the first three mowings should be left on the treated area. Do not use the clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.] <p>[WATERING]</p> <ul style="list-style-type: none"> • [If lawn is dry, water 1 to 2 days before using this product.] • [Rain or watering [1][4][24] hour after application will not wash away effectiveness.] <p>[CLEAN-UP]</p> <ul style="list-style-type: none"> • [Rinse sprayer 3 times after use with clean water. Spray rinse water onto treatment area.]
<p>HOW IT WORKS</p>	<p>This product contains a proven blend of 3 weed killers [in one] to kill existing weeds plus a residual herbicide to keep them from coming back for up to 6 months. Weeds begin to wilt & die within [24] hours [and are completely dead in [14 – 28] days] [and new weeds will not grow up to 6 months]. And, it won't harm your lawn when used as directed.</p>

<p>KILLS & PREVENTS [or KILLS (a "1" indicates Kills & Prevents)]</p>	
<p>[Weeds presented here to be chosen from Appendix 1.]</p> <p>[and many other weeds]</p>	


	<p>Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.</p>
---	--

PRECAUTIONARY STATEMENTS	
CAUTION	
Hazards to Humans and Domestic Animals	
<ul style="list-style-type: none"> • Harmful absorbed through skin. • Harmful if swallowed. • Avoid contact with skin. • Causes moderate eye irritation. • Remove and wash contaminated clothing before reuse. • When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves. <p>After using this product:</p> <ul style="list-style-type: none"> • Rinse gloves before removing and wash hands. • If product gets on skin, wash thoroughly with soap & water. • If product gets on clothing, remove clothing as soon as possible and launder clothing separately before reuse. • Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. 	

FIRST AID	
If Swallowed	<ul style="list-style-type: none"> ▪ Call a poison control center or doctor immediately for treatment advice. ▪ Have person sip a glass of water if able to swallow. ▪ Do not induce vomiting unless told to do so by a poison control center or doctor. ▪ Do not give anything by mouth to an unconscious person.
If on Skin	<ul style="list-style-type: none"> ▪ Take off contaminated clothing. ▪ Rinse skin immediately with plenty of water for 15 – 20 minutes. ▪ Call a poison control center or doctor for further treatment advice.
If in Eyes	<ul style="list-style-type: none"> ▪ Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. ▪ Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. ▪ Call a poison control center or doctor for further treatment advice.
<p>NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 1-877-229-3763 for medical emergency information.</p> <p>Active Ingredients: 3.45% 2,4-D, dimethylamine salt (CAS # 2008-39-1), 0.379% Dicamba, potassium salt (CAS # 10007-85-9), 0.812% Mecoprop-p, potassium salt (CAS # 66423-05-0) and 0.242% Indaziflam (CAS # 950782-86-2).</p>	

ENVIRONMENTAL HAZARDS	
<ul style="list-style-type: none"> • This product is toxic to fish, aquatic invertebrates, and plants. Drift or runoff maybe hazardous to aquatic organisms in water adjacent to treated areas. • To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. • This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow. • This pesticide may impact water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. 	

9
19

Important – Directions for Storage and Disposal	
Pesticide Storage Conditions	Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing.
Pesticide Disposal 	If empty: Non-refillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or toll free 1-877-229-3724 for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE: Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. To the extent consistent with applicable law, the Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

***Money Back Guarantee:** If you are not satisfied with this product, we will gladly refund your original purchase price.

Produced For



A Business Group of Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709

10
5

(FRONT PANEL)

(Part B: Pre-packed Hose-end Ready-to-Spray Label)

Lawn 3FL2 Herbicide Ready-to-Spray

[ABN: Southern Season Long Weed Control for Lawns¹ Ready-to-Spray]

▪ *Editorial Note – Marketing Claims Positioned Here [see Appendix 2]*

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	3.45%
Dicamba, potassium salt.....	0.379%
Mecoprop-p potassium salt.....	0.812%
Indaziflam.....	0.242%
OTHER INGREDIENTS:.....	95.117%
TOTAL	100.00%

THIS PRODUCT CONTAINS:

- 0.249 lb 2,4-D acid equivalent per gal or 2.86%
- 0.028 lb Dicamba acid equivalent per gal or 0.324%
- 0.060 lb Mecoprop-p equivalent per gal or 0.689%
- 0.021 lb Indaziflam per gal or 0.242%
- Isomer Specific by AOAC Method.

Note: If the design, format, or small size of the container labels makes it impractical to present the entire ingredient statement on the front panel, then the sub statement will be placed on the back panel.

Net Contents : 40 fl. oz. (1.18 L), 32 fl. oz. (946 mL), 24 fl. oz. (709 mL), 2.25 fl. oz. (66 mL), or other sizes within this range.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional precautionary statements.

Editorial Note – [Bracketed text] is optional language

11
1
9



(Back Panel)

OPEN Resealable Label for Directions & Precautions PRESS TO RESEAL

[The days of multiple applications to kill weeds is over!]. Just one application of Lawn 3FL Herbicide Ready-to-Spray's [patented] [exclusive] [proprietary] [4-in-1] [3-in-1] [2-in-1] formula kills over 200 weeds & prevents new lawn weeds for up to 6 months! [It delivers Season Long Weed Control [up to 6 months] that's simple and easy to use] [and it will not harm your lawn when used as directed]. [No need-[to-re-apply-every-couple-of-weeks]-[for frequent-re-application]-] [PLUS, when-applied-in Early Spring, prevents Crabgrass all season long]. [Just Hook to Your Hose and Go -]. Convenient sprayer top connects to garden hose and automatically dispenses the weed killer. Ideal for large areas – treats [up to] 7,100 square feet. [Packaging-related note: Adjust sq. ft. coverage proportionately for sizes other than 32 fl. oz.] [Rainproof in [1][4][24] hour.] Controlling [listed] weeds [and preventing crabgrass] in your lawn up to 6 months has never been easier!



QUICK FACTS

<i>Editorial Note – Marketing Claims Positioned Here [see Appendix 2]</i>	
KILLS & PREVENTS For up to 6 Months with one application	[Kills over 200 broadleaf weeds such as Dollarweed, Chickweed, Dandelion, Henbit, Clover & Spurge] [Prevents XX [listed] grassy & broadleaf weeds such as Crabgrass, Annual Bluegrass, Goosegrass, White Clover, Dandelion, Plantain, Henbit and Black Medic] <i>Choose from marketing claims</i>
WHEN TO [APPLY] [USE]	[Apply anytime – spring or fall] or choose from other applicable marketing claim
WHERE TO USE	ON ESTABLISHED LAWNS of the turf types listed below: Bermudagrass, Centipedegrass, St. Augustinegrass, and Zoysia DO NOT use on improved varieties of St. Augustinegrass (such as Floratam), Fescue, Bentgrass, Carpetgrass, Kentucky Bluegrass, Kikuyugrass, Ryegrass or Dichondra lawns
HOW TO USE	To prevent new weeds from growing, spray entire area you want protected. Wait [X] [days] [hours] after applying, then activate preventative weed barrier by watering.] See Directions for Use for further details.
COVERAGE	This bottle treats 5,000 – 7,100 sq ft. [Just hook to garden hose & go.] [(7,100 sq ft for centipede and St. Augustine [grasses] [lawns]), [(5,000 sq ft. for Bermudagrass and Zoysia Lawns)] [Packaging-related note: Adjust sq. ft. coverage proportionately for sizes other than 32 fl. oz.]
[OPTIONAL GRAPHIC IMAGES:]	
	For questions or comments, call toll-free 1-877 BayerAG (1-877-229-3924) [or visit our website at www.bayeradvanced.com]

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	4.89%
Dicamba, potassium salt.....	0.54%
Mecoprop-p potassium salt.....	1.15%
Indaziflam*.....	0.25%

OTHER INGREDIENTS:	93.17%
TOTAL	100.00%

* Protected by patent number US 6,069,114

THIS PRODUCT CONTAINS:









- 0.249 lb 2,4-D acid equivalent per gal or 2.86%
- 0.028 lb Dicamba acid equivalent per gal or 0.324%
- 0.060 lb Mecoprop-p equivalent per gal or 0.689%
- 0.021 lb Indaziflam per gal or 0.242%

EPA Reg. No. 72155-XX
EPA Est. No.

12
19

Isomer Specific by AOAC Method.


DIRECTIONS FOR USE	
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.	
BEFORE YOU USE	<ul style="list-style-type: none">• Use this product only on established lawns – new lawns must be at least 4 [or X] months old if sodded, 2 [or X] months old if sprigged & 12 [or X] months old if seeded.
Protect from Freezing	<ul style="list-style-type: none">• Do not overlap treated areas or make repeat applications within 3 months as over application could damage the lawn.• Do not apply to recently aerated or verticutted turf until it has recovered.• Apply when the air is calm to avoid spray drift onto desirable plants.• This product may affect young seedlings downslope of treated area after heavy rainfall. Do not apply if heavy rainfall is expected within the next 48 hours.
HOW MUCH TO USE	<p>Measure area to be sprayed. Calculate 'square feet' by multiplying length times width. [In order to gauge how quickly you should walk [to cover] [5,000 sq. ft.] [7,100 sq. ft.], this bottle will empty in approximately [XX] minutes.] <i>[Packaging-related note: Adjust sq. ft. coverage proportionately for sizes other than 32 fl. oz.]</i></p> <p>Bermudagrass, Zoysiagrass – this bottle treats 5,000 sq ft. <i>[Packaging-related note: Adjust sq. ft. coverage proportionately for sizes other than 32 fl. oz.]</i></p> <p>Centipedegrass, St. Augustinegrass (not for use on improved varieties, such as Floratam), Kentucky bluegrass – this bottle treats 7,100 sq. ft. <i>[Packaging-related note: Adjust sq. ft. coverage proportionately for sizes other than 32 fl. oz.]</i></p> <p>The maximum application rate of this product is 32 fl. oz. per 5,000 sq. ft. per application per site. The maximum number of broadcast applications per treatment site is 2 per year.</p>

<p>HOW TO USE</p> <p>Shake Well Before Use</p>	<ul style="list-style-type: none"> To prevent new weeds from growing, you must spray the entire area you want protected, not just the emerged weeds. [Wait [X] [days] [hours] after applying then activate preventative weed barrier by watering with ¼ - ½" of irrigation.] <p><i>Editorial Note – Either the Single Rate Sprayer or the Dual Rate Sprayer directions for use will appear on the commercial package but not both.</i></p>
<p>[Optional graphic images]:</p>  	<p>[For single rate sprayer]</p> <p>CONNECT</p> <ol style="list-style-type: none"> Shake well before use. Connect sprayer to garden hose. Turn on water.  <p>SPRAY</p> <ol style="list-style-type: none"> Extend hose to farthest area to be treated, point sprayer nozzle in direction to be sprayed. Begin spraying – Push & hold lock tab away from knob, use other hand to turn knob [counter-]clockwise to ON position. Walk at a steady pace while spraying, using an even sweeping motion, and slightly overlap the treated areas. Work back toward the faucet to avoid walking over wet, treated grass. <p>FINISH</p> <ol style="list-style-type: none"> To stop spraying, push & hold lock tab away from knob and use other hand to turn knob [counter-]clockwise to OFF position. Turn off water at faucet. Relieve water pressure in hose by turning sprayer to [ON] [WATER] position until water slows to a drip then turn sprayer OFF. Disconnect sprayer from hose. <p>[For dual rate sprayer]</p>  <ol style="list-style-type: none"> Connect your garden hose to the bottle. Be sure the blue water control valve on top of the spray nozzle is in the OFF position. Turn on water at faucet.  <ol style="list-style-type: none"> To apply high rate (treats 5,000 sq. ft.) [<i>Packaging-related note: Adjust sq. ft. coverage proportionately for sizes other than 32 fl. oz.</i>]: Pull the small star-shaped blue product-control button out until it stops, Twist the blue product-control button so that the 'L' on the product-control button is at the top of the control valve AND the product control valve is aligned with the flat portion of the lock tab. Push the button in to open. Proceed to step #3.  <p>To apply low rate (treats 7,100 sq. ft.) [<i>Packaging-related note: Adjust sq. ft. coverage proportionately for sizes other than 32 fl. oz.</i>]: Pull the small star-shaped, blue product-control button out until it stops. Twist the product-control button so that the 'G' on the product-control button is at the top of the control valve AND the product control valve is aligned with the flat portion of the lock tab. Push the button in to open (The button will only go half way in for this setting). Proceed to step #3.</p>  <ol style="list-style-type: none"> Extend hose to farthest area to be treated, point sprayer nozzle in direction to be sprayed. To begin spraying, point nozzle toward the area to be treated and turn the blue water control valve to the ON position. Tab will be horizontal. Walk at a steady pace while spraying, using an even sweeping motion, and slightly overlap the treated areas. Work back toward the faucet to avoid walking over wet, treated grass.  <ol style="list-style-type: none"> To stop spraying, turn the blue water control valve on top of the spray nozzle to OFF. To close, push in the small round button opposite the star-shaped product-control button. Turn off faucet, <ol style="list-style-type: none"> Relieve water pressure by turning blue water control valve to the ON position until water slows to a drip. Turn blue water control valve back to OFF. Disconnect garden hose from bottle.

14
19

[FOR BEST RESULTS]	<p>[WHEN TO APPLY]</p> <ul style="list-style-type: none"> • [Spray when: - weeds are small and actively growing - temperatures are between 50 – [85][90]°F.] • [Do not apply while grass is emerging from dormancy.] • [This product also prevents crabgrass & goosegrass [other listed weeds as appropriate] when applied early spring before the fourth mowing] • [This product also prevents annual bluegrass (<i>Poa annua</i>) when applied August – November.]
	<p>[MOWING]</p> <ul style="list-style-type: none"> • [Certain hard-to-kill weeds may be retreated in 30 days if necessary. Spray individual weeds using this product or any other Bayer Advanced lawn weed herbicide] • [Heavy raking will disturb the weed preventative barrier and reduce the effectiveness of this product.] • [Lawns may be seeded 8 months (St. Augustinegrass and Centipedegrass) or 12 months (Bermudagrass and Zoysia) after applying this product. Lawns may be sodded 4 [or X] [months] and sprigged 2 [or X] months after applying this product.] <p>[WATERING]</p> <ul style="list-style-type: none"> • [If lawn is dry, water 1 to 2 days before using this product.] • [Rain or watering [1][4][24] hour after application will not wash away effectiveness.]
HOW IT WORKS	<p>This product contains a proven blend of [three weed killers] [3 weed killers in one] to kill existing weeds plus a residual herbicide to keep them from coming back for up to 6 months. Weeds begin to wilt & die within [24] hours [and are completely dead in [14 – 28] days] [and new weeds will not grow for up to 6 months]. And, it won't harm your lawn when used as directed.</p>

KILLS & PREVENTS [or KILLS (a "1" indicates Kills & Prevents)]	
<p>[Weeds presented here to be chosen from Appendix 1.]</p> <p>[and many other weeds]</p>	

	<p>Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.</p>
---	--


15
19

PRECAUTIONARY STATEMENTS	
CAUTION	
Hazards to Humans and Domestic Animals	
<ul style="list-style-type: none"> • Harmful if swallowed or absorbed through skin. • Causes moderate eye irritation. • Avoid contact with eyes, skin, or clothing. • Remove and wash contaminated clothing before reuse. • When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves. <p>After using this product:</p> <ul style="list-style-type: none"> • Rinse gloves before removing and wash hands. • If product gets on skin, wash thoroughly with soap & water. • If product gets on clothing, remove clothing as soon as possible and launder clothing separately before reuse. • Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. 	

FIRST AID	
If Swallowed	<ul style="list-style-type: none"> ▪ Call a poison control center or doctor immediately for treatment advice. ▪ Have person sip a glass of water if able to swallow. ▪ Do not induce vomiting unless told to do so by a poison control center or doctor. ▪ Do not give anything by mouth to an unconscious person.
If on Skin	<ul style="list-style-type: none"> ▪ Take off contaminated clothing. ▪ Rinse skin immediately with plenty of water for 15 – 20 minutes. ▪ Call a poison control center or doctor for further treatment advice.
If in Eyes	<ul style="list-style-type: none"> ▪ Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. ▪ Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. ▪ Call a poison control center or doctor for further treatment advice.
<p>NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 1-877-229-3763 for medical emergency information.</p> <p>Active Ingredients: 3.45% 2,4-D, dimethylamine salt (CAS # 2008-39-1), 0.379% Dicamba, potassium salt (CAS # 10007-85-9), 0.812% Mecoprop-p, potassium salt (CAS # 66423-05-0) and 0.242% Indaziflam (CAS # 950782-86-2).</p>	

16
19

ENVIRONMENTAL HAZARDS	
<ul style="list-style-type: none"> • This product is toxic to fish, oysters, aquatic invertebrates and non-target plants. Drift or runoff maybe hazardous to aquatic organisms in water adjacent to treated areas. • To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. • This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater. 	

Important – Directions for Storage and Disposal	
Pesticide Storage Conditions	Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing.
Pesticide Disposal 	If empty: Non-refillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or toll free 1-877-229-3724 for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE: Research and testing have determined that the “Directions for Use” are appropriate for the proper use of this product under expected conditions. To the extent consistent with applicable law, the Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

***Money Back Guarantee:** If you are not satisfied with this product, we will gladly refund your original purchase price.

Produced For



A Business Group of Bayer CropScience LP
 P.O. Box 12014, 2 T.W. Alexander Drive
 Research Triangle Park, North Carolina 27709

Lawn 3FL2 Herbicide Concentrate/Ready-to-Spray (Pending) 11/03/10

Appendix 1: KILLS & PREVENTS [or KILLS (a "1" indicates Kills & Prevents)]			
Annual Bluegrass (<i>Poa annua</i>) ²	Creeping Charlie	Morningglory	Spanishneedles
Annual Sedge ²	Crabgrass ²	Mouseear Hawkweed	Speedwell ¹
Annual Yellow Sweetclover	Creeping Jenny	Mouse Barley ²	Spiny Amaranth
Aster	Cudweed ¹	Mugwort	Spiny Cocklebur
Austrian Fieldcress	Curly Dock ¹	Musk Thistle	Spiny Sowthistle
Barnyardgrass ²	Cutleaf Eveningprimrose ²	Mustard	Spotted Catsear ¹
Bedstraw	Daisy, English	Narrowleaf Plantain	Spotted-Spurge
Beggarticks	Daisy, Fleabane	Narrowleaf Vetch	Spurweed
Betony, Florida	Daisy, Oxeye	Nettle	Stinging Nettle
Bindweed, Field	Dandelion ¹	Orange Hawkweed	Strawberry, India Mock
Bird Vetch	Dichondra	Oriental Cocklebur	Sunflower, Common ²
Bitter Wintercress	Dogbane	Oxalis ¹	Swinecress ²
Bittercress, Hairy	Dogfennel	Panicum Willowweed ²	Tall Nettle
Bitterweed	Dollarweed	Parsley-piert	Tall Vervain
Black-eyed Susan	Doveweed ²	Parsnip	Tansy Mustard
Black Medic ¹	Eclipta ²	Pearlwort	Tansy Ragwort
Black Mustard ¹	Elderberry	Pennycress	Tanweed
Black Nightshade, American ²	False Chamomile ²	Pennywort	Thistle ¹
Blackseed Plantain	False Dandelion	Pepperweed	Trailing Crownvetch
Blessed Thistle	False Flax	Phyllanthus, Longstalked ²	Tropic Ageratum ²
Bloodflower Milkweed	False Sunflower	Pigweed ¹	Tufted Lovegrass ²
Blue Lettuce	Fiddleneck	Pineywoods Bedstraw	Tumble Mustard
Blue Vervain	Fleabane, Blackleafed ²	Plains Coreopsis (Tickseed)	Tumble Pigweed
Bracted Plantain	Florida Pusley ¹	Plantain ¹	Twincress ²
Brassbuttons ¹	Foxtail ²	Poison Ivy	Velvetleaf
Bristly Oxtongue	Foxtail Brome ²	Poison Oak	Venice Mallow
Broadleaf Dock	Frenchweed	Pokeweed	Vetch, Purple ²
Broadleaf Plantain	Galinsoga	Poorjoe	Virginia Buttonweed
Broomweed	Goathead	Prairie Sunflower	Virginia Creeper
Buckhorn	Goldenrod	Prickly Lettuce	Virginia Pepperweed
Buckhorn Plantain ¹	Goosegrass ²	Prickly Sida	Water Pennywort
Bulbous Buttercup	Ground Ivy	Prostrate Knotweed ¹	Wavyleaf Bullthistle
Bull Thistle	Guineagrass ²	Prostrate Pigweed ¹	Western Clematis
Bullnettle	Gumweed	Prostrate Spurge ¹	Western Salsify
Burclover ¹	Hairy Fleabane ¹	Prostrate Vervain	White Mustard
Burdock	Hairy Nightshade ²	Puncturevine	Wild Aster
Burning Nettle	Hawkweed	Purslane, Common ¹	Wild Buckwheat ¹
Burweed ¹	Healall	Ragweed	Wild Carrot ¹
Buttercup	Heartleaf Drymary	Red Sorrel ¹	Wild Four-o'clock
Buttonweed	Heath Aster	Red Tasselflower ²	Wild Garlic
Canada Thistle ¹	Hedge Bindweed	Redmaids ²	Wild Geranium
Carolina Geranium	Hedge Mustard	Redroot Pigweed ¹	Wild Lettuce
Carpetweed ¹	Hemp	Redstem Filaree	Wild Marigold
Catchweed Bedstraw	Henbit ¹	Redstem Fleabane/Storksbill ²	Wild Mustard ¹
Catnip	Hoary Cress	Rice Flatsedge ²	Wild Onion
Catsear	Hoary Plantain	Rough Cinquefoil	Wild Parsnip
Cheat ²	Hoary Vervain	Rough Fleabane	Wild Radish
Chickweed, Common ¹	Horseweed/ Marestail ²	Russian Pigweed	Wild Rape
Chickweed, Mouseear ¹	Horsenettle	Russian Thistle	Wild Strawberry
Chicory	Italian Ryegrass ²	Ryegrass ²	Wild Sweet Potato
Cinquefoil	Jimsonweed	Sandbur	Wild Vetch
Clover, Crimson	Knawel	Scarlet Pimpernel	Woodsorrel ¹
Clover, Hop	Knotweed ¹	Scotch Thistle	Wooly Croton
Clover, Red	Kochia ¹	Sesbania, Hemp ²	Wooly Morningglory
Clover, Strawberry	Lambsquarters ¹	Sheep Sorrel	Wooly Plantain
Clover, Sweet	Lespedeza	Shepherdspurse ¹	Wormseed
Clover, White ¹	London Rocket ²	Slender Plantain	Yarrow
Cockle	Mallow ¹	Smallflower Galinsoga	Yellow Gromwell ²
Cocklebur	Matchweed	Smooth Dock	Yellow Rocket, common ¹
Common Evening Primrose ²	Mexicanweed	Smooth Pigweed	Yellowflower Pepperweed
Common Groundsel ²	Milk Vetch	Sorrel	[and other broadleaf and grassy weeds]
Common Mullein		Sowthistle ¹	

¹ Kills & Prevents [add footnote to specific weeds as appropriate]

18
—
19

² Prevents only [add footnote to specific weeds as appropriate]

Appendix 2

Optional Marketing Claims:

- New [Formulation]!
- [Brand] New [Formula]!
- Breakthrough Formulation
- Breakthrough Convenience
- [Patented] [Exclusive][Proprietary] [4-in-1] [3-in-1] [2-in-1] Formula
- Proprietary Formula that Kills Broadleaf Weeds
- Systemic [Herbicide] [Activity]
- Kills to the root.
- Kills [Current] Weeds to the Root
- Kills [Controls] even the tough weeds ... to the root!
- Kills 200+ Weeds [to the root]
- Kills [Controls] [200+] broadleaf weeds [to the root]
- Kills Dandelion, Clover, plus 200 Other listed Weeds
- Kills Over 200 Lawn Weeds in One Easy Step
- Kills Lawn Weeds in One Step
- Kills Broadleaf Weeds
- Kills Both Broadleaf and Grassy Weeds
- Kills Listed Weeds
- Kills Hard-to-Control Weeds
- Effectively Kills Both Broadleaf and [Weed][Crab] grass[es]
- Kills the weeds not the lawn† (when used as directed)
- Won't Harm your Lawn† (when used as directed)
- Won't Harm Lawns† (when used as directed)
- [Patented] [Exclusive] [Unique] [Proprietary] 2-in-1 Product that Kills [Lawn] Weeds and [Stops] [Prevents] them [from] coming back
- [Patented] [Exclusive] [Unique] [Proprietary] 3-in-1 Product: Kills Weeds, Prevents New Weeds [up to 6 Months] & Prevents Crabgrass
- Season Long Weed Control [with 1 Application]
- Season Long [Lawn] Weed & [Crab] Grass Control
- Season Long [Lawn] Weed Control [Killer]
- Season Long Crabgrass Control
- 1 Application Provides Season Long [Lawn] [Weed] Control
- One Application = Multiple applications of traditional products
- One application provides up to 6 months [weed] [crabgrass] control
- [Up to] 6 Month Weed Control [with one application]
- Kills Weeds [& Prevents New Weeds] [from Coming Back] [up to 6 months]
- Kills [Current] [Existing] Weeds [up to 6 Months]
- Kills [& Prevents] Weeds [up to 6 Months]
- Keeps weeds out [of your lawn] [up to 6 months]
- Prevents [New] Weeds up to 6 Months [with one application]
- Prevents [Listed] Grasses [up to 6 Months]
- Protects Your Lawn From [Grassy & Broadleaf] Weeds Up To 6 Months
- Tough Weed Protection [for up to 6 months]
- Weeds don't return [for up to 6 months]
- Weed Free Lawn [Up to 6 Months]
- Extended [Weed] Control [Killer]
- Lawn Weed Control is Easier than Ever Before
- Simple, [Convenient], [Easy] to use
- PLUS Prevents Crabgrass!
- Prevents Crabgrass All Season
- Prevents Crabgrass [up to 6 months]
- Crabgrass Free Lawn [Up to 6 Months]
- [Now] [Plus] Prevents [New] Weeds [Plus] [Crabgrass] [Grassy Weeds] [Poa Annua] [Annual Bluegrass] [Goosegrass] [up to 6 months]
- [Also] Kills [Prevents] Dollaweed [up to 6 months]
- Prevents Annual Bluegrass [Poa Annua]
- Kills Dollarweed [and other common southern lawn weeds]
- Starts Working Immediately
- Visible Results in 24 Hours
- Proven Performance
- Rainproof in [1][4][24] Hour
- All Weather Formula (rainproof in [1][4][24] hour)
- Guaranteed* Not to Harm Lawns† (when used as directed)
- 100% Satisfaction Guaranteed* (or your money back)
- Water based.
- Fast [& Easy] Coverage
- Simple, [Convenient], [Easy] to use
- Mixes with Water – Concentrate
- Water saving applicator
- Attach to [garden] hose and spray
- Just hook to garden hose & go
- Hook to hose and go
- Always Ready to Spray
- One quart covers [up to] 5,000 [7,100] square feet.
- 1 quart treats [up to] 5,000 [7,100] sq. ft.
- *[Packaging-related not for claims]*
- Treats [up to] 5,000 [7,100] sq. ft.
- Now 25% More [Free versus 32 oz size]
- 25% (33%) More [FREE] versus XX oz size
- Apply anytime spring or fall
- † (When used as directed. [DO NOT use on] [improved varieties of St. Augustinegrass (such as Floratam)], [Fescue], [Bentgrass], [Carpetgrass], [Kentucky Bluegrass], [Kikuyugrass], [Ryegrass] or [Dichondra] lawns
- *See weed list for [complete list of] ["Kills and "Prevents"] weeds.
- [New] [exclusive] [XX oz] [special value] [size]