| (- | | 1 | - |
|--|---|--|---|
| STATED STATED. IS NOT THE STATED STATED STATED IN THE STATED STATED IN THE STATE STATED IN THE STATE STATE STATES IN THE STATE STATES IN THE STATE STATES IN THE STATES INTO A STATES IN | U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505P) 1200 Pennsylvania Ave., N.W. Washington, D.C. 20460 | EPA Reg. Number: 72155-113 | Date of Issuance: 4/21/15 |
| | NOTICE OF PESTICIDE: | Term of Issuance: | |
| | X Registration | Unconditional | |
| | (under FIFRA, as amended) | Name of Pesticide Prod | uct: |
| | | 3FL1B HERBIC | IDE GRANULE |
| Name and Address of R Bayer CropScien PO Box 12014 2 TW Alexander RTP, NC 27709 | | | |
| | g differing in substance from that accepted in connection with this registrati or to use of the label in commerce. In any correspondence on this product a | | |
| under the Federa Registration is in Agency. In order time suspend or o name in connecti registrant a right This product is u 1. Submit an product v 2. The data and 830.6 | nformation furnished by the registrant, the above n I Insecticide, Fungicide and Rodenticide Act. a no way to be construed as an endorsement or recor- to protect health and the environment, the Admin cancel the registration of a pesticide in accordance ion with the registration of a product under this Ac- to exclusive use of the name or to its use if it has be nconditionally registered in accordance with FIFR nd/or cite all data required for registration/reregistra- when the Agency requires all registrants of similar requirements for storage stability and corrosion ch 6320) are not satisfied. A one year study is required to a months from the date of registration to provide | ommendation of th istrator, on his mo with the Act. The t is not to be const been covered by ot A section 3(c)(5) p ration/registration products to submit aracteristics (Guid d to satisfy these d | is product by the tion, may at any acceptance of any rued as giving the hers. provided that you: review of your t such data. |
| Signature of Approving | Official: | Date: | |
| Kathryn Montag | V. W Jon Juguo ue, Product Manager 23 h, Registration Division (7505P) | 4/21/15 | ; |

- 3. A revised version of MRID 49415109 must be submitted which includes the expanded rationale for the request to waive pH data (Guideline 830.7000).
- 4. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 72155-113."
- 5. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 03/24/2015
- Alternate CSF 1 dated 03/25/2015
- Alternate CSF 2 dated 03/26/2015

If you have any questions, you may contact Dominic Schuler at 703-347-0260 or via email at <u>schuler.dominic@epa.gov</u>.

Sincerely,

astryn V. Wontaguo

Kathryn V. Montague, Product Manager 23 Herbicide Branch Registration Division (7505P) Office of Pesticide Programs

3FL1B Herbicide Granule

ABN: Bayer Advanced Extended Control Weed [and] [&] Feed II; Bayer Advanced Season Weed [and] [&] Feed II; Bayer Advanced Season Long Weed Control Plus Fertilizer II; Bayer Advanced 3-in-[1] [One] Weed [and] [&] Feed II; 3 in 1 Weed & Feed for Southern Lawns II

| Editorial Note – Marketing Claims Positioned Here | |
|---|----------|
| ACTIVE INGREDIENTS: | |
| Dicamba acid | 0.073% |
| Penoxsulam | 0.02.40/ |
| Indaziflam | 0.022% |
| OTHER INGREDIENTS: | |
| TOTAL: | 100.000% |

[GUARANTEED ANALYSIS X-X-X]

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for additional precautionary statements.

Editorial Note - [Bracketed text] is optional language



and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 72155-113

(Back Panel)

Tired of killing weeds in your lawn – even after you've put down your traditional weed and feed granules? Just one application of [3FL1B Herbicide Granule] kills lawn weeds and prevents new ones, including Crabgrass, for up to six months. Plus it greens, feeds and strengthens your lawn for a thick, healthy turf. [Product Name] [It] delivers [season long] [extended] weed control that is simple and easy to use and it will not harm your lawn [when used as directed.] No need for frequent re-application [plus when applied in early spring, it prevents Crabgrass] [and Goosegrass] [and when applied late summer, Annual Bluegrass] [all season long.] Ideal for large areas –[treats 5,000] [square feet] [sq. ft.] with the 15.0 lb size [or] [XXXX square feet] [sq ft] with the [XXIb size.] [Up to 6 months weed control – plus feeding – has never been easier]. [Spend your summer enjoying your lawn – not maintaining it!] [Killing and preventing weeds in your lawn for up to 6 months plus getting a healthy turf has never been easier.]



| [QUICK FACTS] | | | |
|--|--|--|--|
| Editorial Note – Marketing Claims Positioned Here [see Appendix] | | | |
| [KILLS & PREVENTS] | Kills broadleaf weeds including [any applicable weed from weed list] | | |
| [For up to 6 Months] | Prevents grassy & broadleaf weeds such as[any applicable weed from weed list] | | |
| [PLUS FEEDS] | Feeds and strengthens the lawn. Contains slow release fertilizer for long lasting feeding. | | |
| [WHEN TO] | [Apply] Anytime [in] [early] spring [or fall]. | | |
| [APPLY] [USE] | | | |
| [WHERE | Use only on established lawns of the turf types listed below: | | |
| TO USE] | Bermudagrass, Centipedegrass,[all varieties of] St. Augustinegrass [including Floratam], Zoysia [and Kentucky Bluegrass] | | |
| | DO NOT use on Ormond variety of Bermudagrass, Fescue, Bentgrass, Carpetgrass, Kikuyugrass, Ryegrass or Dichondra lawns | | |
| [COVERAGE] | This bag treats 5,000 sq ft [15 lb size] [or X,XXX sq ft [for XX lb size]] | | |
| (? ,) | Questions or comments, call toll-free (877) 229-3724 www.BayerAdvanced.com | | |
| [🕿 🛄] | | | |

ACTIVE INGREDIENTS:

| Dicamba acid | 0.073% |
|--------------------|----------|
| Penoxsulam | 0.034% |
| Indaziflam. | 0.022% |
| OTHER INGREDIENTS: | |
| TOTAL: | 100.000% |
| | |

[FERTILIZER ANALYSIS X-X-X]

| [Total Nitrogen (N) [Available Phosphate (P2O5) | |
|--|---|
| [Soluble Potash (K2O) | - |
| [Derived from commercially available fertilizer] | |

EPA Reg. No. 72155-xxx EPA Est. No.

| ➡ FIRST AID | | |
|------------------------|--|--|
| If swallowed | Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. | |
| If on skin or clothing | Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 minutes. Call a poison control center or doctor for further treatment advice. | |
| If in eyes | Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for further treatment advice. | |

NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 1-877-229-3763 for medical emergency information.

Active Ingredients: 0.088% Dicamba, (CAS # 1918-00-9), 0.041% Penoxsulam (CAS # 219714-96-2) and 0.027% Indaziflam (CAS # 950782-86-2).

PRECAUTIONARY STATEMENTS

J CAUTION HAZARDS TO HUMANS AND DOMESTIC ANIMALS

- Harmful if swallowed or absorbed through skin.
- Causes moderate eye irritation.
- Avoid contact with eyes, skin, or clothing.
- Remove and wash contaminated clothing before reuse.
- When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

After using this product:

- Rinse gloves before removing and wash hands.
- If product gets on skin, wash thoroughly with soap & water.
- If product gets on clothing, remove clothing as soon as possible and launder clothing separately before reuse.
- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

ENVIRONMENTAL HAZARDS

- This product is toxic to fish, aquatic invertebrates, and plants. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean water mark. Do not contaminate water when disposing of rinsate or washwater. This product may impact water through spray drift or runoff. Follow directions for use to avoid spray drift and runoff. A level well maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams and springs will reduce the potential of this product entering water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.
- Ground Water Advisory: This pesticide has properties and characteristics associated with chemicals detected in
 groundwater. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the
 water table is shallow.
- Surface Water Advisory: This pesticide may impact water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application.

| | DIRECTIONS FOR USE | |
|------------------------|---|--|
| It is a violation of F | ederal law to use this product in a manner inconsistent with its labeling. | |
| BEFORE YOU USE | Use this product only on established Southern lawns – new lawns must be at least 3 months old if sodded 16 months old if sprigged & 16 months old if seeded. Do not overlap treated areas or make repeat applications within 90 days as over application could damage the lawn. Apply only to actively growing turf. Do not apply to recently aerated or verticutted turf until it has recovered. Apply when the air is calm to avoid drift onto desirable plants. | |
| HOW TO USE | Use rotary or drop type spreader to apply granules evenly over the lawn. To be sure you cover lawn areas uniformly, first treat the border of the lawn then fill in the center. Make parallel passes, walking at a steady pace to ensure even distribution of the granules. Sweep, brush, or blow any accidentally applied material off driveways, sidewalks etc. back on to the lawn. Wait 24 hours after applying then activate residual barrier by lightly watering in [with 1/10 - ¼ inch of irrigation]. Do not make more than two applications per site in any 12 month period. Heavy raking will disturb the weed preventative barrier and reduce the effectiveness of this product. | |
| FOR BEST RESULTS | WHEN TO APPLY Apply when weeds are small and actively growing temperatures are between 50–90°F. May be applied to dry or (for best results) wet lawns. Do not apply while turfgrass is emerging from dormancy. Do not apply to centipedegrass after October 1. To prevent Crabgrass and Goosegrass apply early spring before the temperatures are regularly in the lower 80° F range. To prevent Annual Bluegrass apply August - December Lawns may be seeded 12 months, sodded 6 months , and sprigged 12 months after applying this product. [For optimal dollarweed control, apply this product mid-fall.] | |
| | [MOWING] If the lawns needs mowing, do it 1 to 2 days before (for best results) or 1 day after applying this product. Clippings from the first three mowings should be left on the treated area. Do not use the clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens. | |
| | [WATERING] If lawn is dry, water 1 to 2 days before using this product. Do not water for at least 24 hours after application or apply if rain is expected within 24 hours. | |
| HOW IT WORKS | [Insert Product Name] contains a proven blend of weed killers to kill existing weeds plus a residual herbicide to keep them from coming back for up to 6 months. This product starts working immediately to stop weed growth until weeds are dead [within weeks] [4 to 10 weeks]. Plus, it feeds your lawn for quick greening and long lasting feeding. | |
| | Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled. | |

ı**† 1**1

| Beggarweed, Creeping Beggarweed, Dwarf Betony, FloridaAgeratum, Tropic Barley, MouseMulberry Weed Mustard, BlackBurnweed, American Chickweed, Common Chickweed, Common Chickweed, Mouseear Nightshade, Hairy Bluegrass, Annual Borwe, Foxtail Burweed, Lawn Cokfeburg, Common Chickweed, Virginia* Clover, Red Cockfebur, Common Chickweed, Common Chickweed, Virginia* Cokfeburg, Common Chickweed, Common California Burlover Chamomile, False Chamomile, False Chamomile, False Coaldenrod Hawkweed Healial (Purple Cudweed) Crabgrass, Large (Hairy Crabgrass, Large (Hairy Crabgrass, Large (Hairy Crabgrass, Large (Hairy Crabgrass, Spronth Cudweed, Linear Leaf Notweed Kylinga, Green* Lespedeza, Common Kyllinga, Green* Lespedeza, Common Kyllinga, Green* Lespedeza, Common Kyllinga, Green* Kutsedge, Yellow*Ageratum, Tropic Barnyardgrass Burweed, Hairy Parbenium Crabgrass, Large (Hairy Crabgrass, Smooth Cudweed) Southern Brassbuttons Southern Bras | KILLS | PREVENTS | | KILLS AND PREVENTS |
|---|---|--|---|--|
| Buttercup, Hairy Buttonweed, Virginia*Black Nightshade, Hairy Bluegrass, AnnualNightshade, Hairy PartheniumDandelion, Common Dock, CurlyClover, Hop Clover, RedBrome, Foxtail Burweed, LawnPhyllanthus, Longstalked Pigweed, RedrootEcliptaCoffeeweed, (Common Coffeeweed, (Common Chickory)Charnomile, False Chanomile, FalsePlantain, Paleseed PuncturevineKnotweed, Prostrate Hantain, PaleseedKnotweed, Prostrate Kyllinga, Annual*Daisy)Crabgrass, Henry Crabgrass, Henry Dollarweed (Pennywort)Crabgrass, Large (Hairy Crabgrass, Smooth Crabgrass, SmoothRedmaids Sedge, AnnualPlantain, Buckhorn Purslane, Common Pusley, FloridaBadelion, Cate Lash(Purple Cudweed) DoveweedSedge, Annual Sedge, AnnualShepardspurse Spurge, SpottedKnotweed Kylinga, Green* Lespedeza, Common Flatsedge, Kice Nutsedge, Veillo*Evening Primrose, Cutleaf Fleabane, Hairy Foxtail, GreenSouthern Brassbuttons Swinecress Tassel Flower Vetoh, PurpleWoodsorrel, YellowVelvetleaf Vetoh, Purple Pepperweed, Virginia Pepperweed, Virginia Pepperweed, Virginia Pepperweed, Virginia Pepperweed, Virginia Pepperweed, Virginia Pepperweed, Virginia Pepperweed, Virginia Pepperweed, Virginia Pilantain, Broadleaf Spreted Catsear Spreted Catsear Spreted Catsear Spreted Catsear Spurweed Mallow, Little ThistleMarestailMatoweed Purbate PurbateKyllinga, Fragrant MarestailWallow, Little MarestailWeiden Blae Purbate VeidealPurstere Veided Catsear SpurweedKyllinga, Fragrant L | Beggarweed, Dwarf | Barley, Mouse | Mustard, Black Mustard, Wild | |
| SpurweedMallow, LittleThistleMarestail | Beggarweed, Creeping Beggarweed, Dwarf Betony, Florida Burdock Buttercup, Hairy Buttonweed, Virginia* Clover, Hop Clover, Red Cocklebur, Common Coffeeweed, (Common Chickory) Daisy, English (Lawn Daisy) Dandelion, False Dollarweed (Pennywort) Goldenrod Hawkweed Heallal Ivy, Ground (Creeping Charlie) Knotweed Kyllinga, Green* Lespedeza, Common Mallow, Roundleaf Matchweed Nutsedge, Purple* Nutsedge, Yellow* Parsley-piert Peppergrass Pepperweed, Virginia Plantain, Broadleaf Smartweed Sorrel, Sheep Speedwell, Common | Ageratum, Tropic Barley, Mouse Barnyardgrass Bittercress Black Nightshade, Hairy Bluegrass, Annual Brome, Foxtail Buckwheat, Wild Burweed, Lawn California Burclover Chamomile, False Cheatgrass Crabgrass, Henry Crabgrass, Large (Hairy Crabgrass, Large (Hairy Crabgrass, Large (Hairy Crabgrass) Crabgrass, Smooth Cudweed, Linear Leaf (Purple Cudweed) Dandelion, Cat's Ear Doveweed Evening Primrose, Common Evening Primrose, Cutleaf Flatsedge, Rice Fleabane, Hairy Foxtail, Giant Foxtail, Green Foxtail, Green Foxtail, Green Foxtail, Green Foxtail, Yellow Filaree, Redstem Goosegrass Groundsel, Common Guineagrass Guineagrass Kochia Kyllinga, Fragrant | Mulberry Weed Mustard, Black Mustard, Wild Nightshade, American Black Nightshade, Hairy Parthenium Phyllanthus, Longstalked Pigweed, Prostrate Pigweed, Redroot Plantain, Paleseed Puncturevine Redmaids Rocket, London Ryegrass, Italian Ryegrass, Perennial Sandbur Sedge, Annual Sesbania, Hemp Sida, Prickly (Teaweed) Southern Brassbuttons Sowthistle, Annual Speedwell, Corn Spurge, Prostrate Sunflower, Common Swinecress Tassel Flower Velvetleaf Vetch, Purple Wild Carrot Willowherb | Burnweed, American Chickweed, Common Chickweed, Mouseear Clover, White Dandelion, Common Dock, Curly Eclipta Henbit Knotweed, Prostrate Kyllinga, Annual* Lambsquarters Medic, Black Plantain, Buckhorn Purslane, Common Pusley, Florida Ragweed, Common Shepardspurse Spurge, Spotted Thistle, Canada |
| Wild Violet Woodsorrel, Creeping | Spotted Catsear Spurweed Thistle Vetch, Common Wild Violet | Lovegrass, Tufted Mallow, Little | | |

* Suppression

SPREADER SETTINGS

If your spreader is not listed, please call our Consumer Service Hotline toll-free (877) 229-3724 for help.

| SPREADER | SETTING to deliver 3 lbs per 1,000 sq ft |
|---|---|
| [Spreader models to be determined.] | [Spreader settings to be determined.] |
| rate of application. Calibrate your spreader according to Tip: 1,000 sq ft = 20 ft x 50 ft | r, the speed at which you walk and uneven terrain may affect the manufacturer's directions. |
| | |

| STORAGE AND DISPOSAL | | | |
|---|--|--|--|
| Do not contaminate water, for | Do not contaminate water, food, or feed by storage and disposal | | |
| Pesticide Storage | Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing. | | |
| Pesticide Disposal and Container Handling [ÎÎÎ] | If empty: Non-refillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or toll free 1-877-229-3724 for disposal instructions. Never place unused product down any indoor or outdoor drain. | | |
| ſ ⁽ | | | |

NOTICE: Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. To the extent consistent with applicable law, the Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

*Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price. For the full Spanish translation go to www.BayerAdvanced.com

Net Contents

Produced For



A Division of Bayer CropScience LP P.O. Box 12014, 2 T.W. Alexander Drive Research Triangle Park, North Carolina 27709

3FL1B Herbicide Granule (Pending) 12/08/2014, 12/12/2014, 12/15/2014, 04/01/2015, 04/10/205, 4/13/2015, 04/14/2015, 04/15/2015, 04/20/2015

Appendix -

Optional Advertising Claims:

- New [Formulation]!
- [Brand] New [Formula]!
- Breakthrough Formulation
- Breakthrough Convenience
- [Exclusive] [Proprietary] [3-in-1] [2-in-1] Formula*
- Proprietary Formula that Kills Broadleaf Weeds
- Systemic [Herbicide] [Activity]
- Kills to the root.
- Kills [Current] Weeds to the Root
- Kills [Controls] even the tough weeds...to the root!
- Kills 50+ Weeds [to the root]
- Kills [Controls] [48] broadleaf weeds [to the root]
- Kills Dollarweed, Clover, Dandelion, plus 50 Other listed Weeds
- Kills Over 50 Lawn Weeds in One Easy Step
- Kills Lawn Weeds in One Step
- Kills Broadleaf weeds
- Kills Both Broadleaf and Grassy Weeds
- Kills Listed Weeds
- Kills Hard-to-Control Weeds
- [Effectively] Kills Both Broadleaf and Weed [Crab] Grass[es]
- Kills the weeds not the lawn† (when used as directed)
- Won't Harm your Lawn† (when used as directed)
- Won't Harm Lawns† (when used as directed)
- [Exclusive] [Unique] [Proprietary] [2-in-1[One]] [3-in-1][One] [Season Long] [Extended Control] [Weed [&][and] Feed] Product that Kills [Lawn] Weeds and [Stops] [Prevents] them [from] coming back
- [Exclusive] [Unique] [Proprietary] [3-in-1] [Season Long]
 [Extended Control] Product: Kills Weeds, Prevents New Weeds
 [up to 6 Months] & Prevents Crabgrass
- Season Long Weed Control [(up to 6 months)] [with 1 Application]
- Season Long [(up to 6 months)] [Lawn] Weed & [Crab] Grass
 Control
- Season Long [(up to 6 months)] [Lawn] Weed Control [Killer]
- Season Long Crabgrass Control [(up to 6 months)]
- 1 Application Provides Season Long [(up to 6 Months)] [Lawn] [Weed] Control
- One application provides up to 6 months [weed] [crabgrass] control
- [Up to] 6 Month Weed Control [with one application]
- Kills Weeds [& Prevents New Weeds] [from Coming Back] [(up to 6 months)]
- Kills [Current] [Existing] Weeds Up to 6 Months
- Kills [& Prevents] Weeds Up to 6 Months
- Keeps weeds out [of your lawn] up to 6 months
- Prevents [New] Weeds up to 6 Months [with one application]
- Prevents [listed] Grasses Up to 6 Months
- Protects your lawn from weeds up to 6 months
- Tough weed protection [for up to 6 months]
- Weeds don't return for up to 6 months
- Weed Free Lawn up to 6 Months
- Extended [Weed] Control [Killer]
- Lawn Weed Control is Easier than Ever Before
- Simple, [Convenient], [Easy] to use
- PLUS Prevents Crabgrass!
- Prevents Crabgrass All Season [(up to 6 months)]
- Prevents Crabgrass [(up to 6 months)]
- Crabgrass Free Lawn Up to 6 Months
- Prevents Crabgrass, [Goosegrass] [and Annual Bluegrass (poa

annua)] [(up to 6 months)]

- [Now] [Plus] Prevents [New] Weeds [Plus] [Crabgrass] [Grassy Weeds] [Poa Annua] [Annual Bluegrass] [up to 6 months]
- Prevents Annual Bluegrass [poa annua]
- Kills Dollarweed [and other common southern lawn weeds]
- Starts Working Immediately
- Guaranteed* not to harm lawns† (when used as directed)
- 100% Satisfaction Guaranteed* (or your money back)
- Easy-to-Spread Granules
- [12.5 lbs][25 lbs] [XX lbs] treats [up to] [5,000] [10,000] [X,XXX] sq ft
- Treats 5,000 [10,000] [X,XXX] sq ft
- Xx% More [FREE] versus XX lb size
- Apply anytime spring or fall
- 3 in 1
- 2 in 1
- All -in- [1] [One}
- Season Long Weed Control
- Season Long
- Extended Control Weed & Feed
- Weed [and] [&] Feed
- Extended Control
- Kills [Broadleaf] weeds [including] [insert any weed from weed list as applicable]
- Prevents [Broadleaf] [and] [Grassy] Weeds [including] [insert any weed from weed list as applicable]
- Prevents Crabgrass
- Kills or Prevents Dollarweed, Clover, Dandelion [and] [&] Crabgrass
- Controls Dollarweed, Clover, [Dandelion] [and][&] Crabgrass
- Say good-bye to Crabgrass [for] [up to] [six] [6] [months]
- Feeds your lawn
- † [For use on] [all] Southern [Turf] [Lawns] including all varieties of St. Augustinegrass [including Floratam], [Bermudagrass], [Centipedegrass], [Zoysia], [and Kentucky Bluegrass]
- Up to [6] [Six] Months of Control
- Kills existing weeds [and] [&] prevents new [weeds] [ones] for up to [six] [6] months
- [New] [XX lbs] Bonus Size
- [XX%] [Bonus size] More than [XX] Size
- Now covers [XXXX][square] [sq] [feet] [ft]
- Long [lasting] residual control
- Controls more grassy weeds than [the competition] [other competitive products]
- Green[s] lawns with superior weed control
- Kills & prevents weeds for up to 6 months
- Kills existing weeds
- Kills [XX]+ types of broadleaf weeds, including Dollarweed and Clover
- Prevents [XX]+ types of broadleaf and grassy weeds, including Crabgrass

 † (When used as directed. [DO NOT use on] Ormond variery of bermudagrass, [Fescue], [Bentgrass], [Carpetgrass], [Kikuyugrass], [Ryegrass] or [Dichondra] lawns

*3-in-1 refers to 3 types of herbicide active ingredients; and 2-in-1 refers to pre-emergent and post-emergent weed control in one product.

- Prevents new weeds up to 6 months
- Kills existing weeds like Dollarweed & Clover
- Prevents new weeds like Crabgrass up to 6 months
- Lawns love it.
- Your lawn needs love too.
- Your yard needs you.
- The total lawn solution.
- A whole new way to weed.
- Everything your lawn needs in one bag.
- The best yard on the block.
- Make your neighbors jealous.
- Feed without weeds.
- Improved feeding, without weeding.
- Lawn solutions made simple.
- Set your lawn apart.
- Concentrated Power Longer Lasting Results
- [Lush, green healthy lawns are better able to choke out weeds, purify [clean] the air, and prevent runoff and soil erosion. Healthy, well-fed lawns are also able to use water more efficiently than unfed lawns because they have more roots. Use Bayer Advanced 3 in 1 Weed & Feed for Southern lawns to [keep your lawn looking its best.] [keep your lawn [weed-free], healthy and green.]]
- [Getting a healthy Southern Turf has never been easier.]



Fertilizer Claims

- Feeds, [Greens] and Strengthens [your lawn][Southern turf][Southern Lawns]
- Plus Feeds
- Builds thick, green turf from the roots up
- Strengthens your lawn
- Builds strong roots
- Builds a thick lush green lawn
- Even feeding for a thicker, greener lawn
- Slow Release Fertilizer/Feeding
- Long Lasting Feeding
- Feeds & strengthens all types of southern lawns
- Feeds southern lawns.
- Fertilizes, greens and strengthens all types of southern turf
- Feeds and strengthens southern lawns
- Greener. Thicker. Stronger.

[For placement under Fertilizer Guarantee Section] 3FL1B Herbicide Granule X-X-X

Guaranteed Analysis

Derived from:

Net Weight:

F1559

Information regarding the contents and levels of metals in this product are available on the Internet at [http://www.regulatory-info-ba.com] [http://www.aapfco.org/metals.htm] .

[The following optional statements are added in order to comply with the Florida (fertilizer) Urban Turf Rule.]

[Florida Applications: X annual feedings [X lbs N/1000 sq ft] with the following exceptions: On centipedegrass do not exceed X feedings [X lbs N/1000 sq ft] in the north and X feedings [X lbs N/1000 sq ft] in the central and south. On zoysiagrass do not exceed X feedings [X lbs N/1000 sq ft] in the north.]

[North Florida extends north from Cedar Key to Daytona Beach.] [Central Florida extends across Florida from Cedar Key to Daytona Beach down to south of Sarasota County. South Florida includes all remaining counties from Sarasota to Coco Beach South]

[In Florida, X feedings may exceed the Nitrogen limits on certain grass types in some [geographic] areas.] [Check with your local Cooperative Extension Agency to obtain specific information on local turf best management practices [for feeding your lawn.]

[Check with your county or city government to determine if there are local regulations for fertilizer use.]

[Florida Annual Fertilizer Applications]

[On centipede lawns, do not exceed rate of X lbs. of Nitrogen per 1,000 sq. ft. in the north and X lbs. of Nitrogen per 1,000 sq. ft. in the central and south.]

[North Florida is north of Ocala. Central Florida is defined as south of Ocala to a line extending from Vero Beach to Tampa. South Florida includes the remaining southern portion of the state.]