

72159-2

( 05/18/2009 )

1/16

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460



MAY 18 2009

OFFICE OF  
PREVENTION, PESTICIDES  
AND TOXIC SUBSTANCES

Agrisel USA Inc.  
c/o Frank Sobotka, Ph.D  
IPM Resources LLC  
660 Newtown-Yardley Rd., Suite 105  
Newtown, PA 18940

Dear Dr. Sobotka,

Subject: Amendment to revise errors in rates to match parent product  
ImidaPro 2SC Insecticide  
EPA Registration No. 72159-2  
Your email submission dated: April 23, 2009

The labeling referred to above submitted in connection with the Federal Insecticide, Fungicide and Rodenticide Act, as amended is acceptable.

Submit one copy of your final printed labeling before you release the product for shipment. A stamped copy of the labeling is enclosed for your records. If you have any questions regarding this label, please contact Autumn Metzger at (703) 305-5314 or metzger.autumn@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Venus Eagle".

Venus Eagle  
Product Manager 01  
Insecticide-Rodenticide Branch  
Registration Division (7505P)

2/16

[ Agrisel logo]

# ImidaPro™ 2SC

## Systemic Insecticide

For Foliar and systemic insect control in turf grass (including sod farms), landscape ornamentals, listed fruit and nut trees, interior plantscapes, nursery and greenhouse grown ornamental and vegetable plants.

**ACTIVE INGREDIENT:**

Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2 imidazolidinimine.....21.4%

**OTHER INGREDIENTS:**.....78.6%

**Total:**.....100.0%

Contains 2 pounds of imidacloprid per gallon.

**SHAKE WELL BEFORE USING**

EPA Reg. No.: 72159-2

EPA Est. No.: 37429-GA-01

### KEEP OUT OF REACH OF CHILDREN CAUTION

#### FIRST AID (neonicotinoid)

<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advise.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<p>In case of emergency call toll free the Emergency Response Telephone No.: To be Added. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.</p>	
<p>Note To Physician (neonicotinoid): No specific antidote is available. Treat the patient symptomatically.</p>	

Manufactured For:  
AGRISEL USA INC  
PO Box 3528  
Suwanee, GA 30024  
1-877-247-4735

Net Contents: TBA  
(fl.oz./qt./gal.)

ACCEPTED

MAY 18 2009

Under the Federal Insecticide, Fungicide,  
and Rodenticide Act, as amended, for the  
pesticide registered under:

EPA. Reg. No: 72159-2

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

### CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing.

### Applicators and Other Handlers Must Wear:

- Long-sleeved shirt and long pants.
- Chemical-resistant gloves made of any waterproof material such as, barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton.
- Shoes plus socks.

### ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**Important:** When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "Applicators and Other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

<p><b>USER SAFETY RECOMMENDATIONS:</b></p> <p>User should:</p> <ul style="list-style-type: none"> <li>• Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.</li> <li>• Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.</li> <li>• Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.</li> </ul>
---

### ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is pre-sent or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on the foliage of blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

**Exception:** If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

**Coveralls**

- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton
- Shoes plus socks

### Application to Ornamentals and Vegetable Plants (Including: nurseries, greenhouses and interior landscapes)

#### Application Information

ImidaPro 2SC Systemic Insecticide is for insect control on ornamental and vegetable plants in nurseries, greenhouses and interior landscapes. ImidaPro 2SC Systemic Insecticide is a systemic product and will be translocated upward within the plant. The addition of a nitrogen fertilizer, where applicable, into the solution may enhance the uptake of the active ingredient. Application can be made by foliar application or soil applications; including soil injection, drenches, chemigation and broadcast sprays. When making foliar applications on hard to wet foliage such as holly, pine, or ivy, the addition of a spreader/sticker is recommended. If concentrate or mist type spray equipment is used, an equivalent amount of product must be used on the area sprayed, as would be used in a dilute application.

**Resistance:** Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area. Consult your Cooperative Extension Service for resistance management strategies and recommended pest management practices for your area. For resistance management purposes, a foliar application of any chloronicotinyl insecticide following a IMIDAPRO 2SC SYSTEMIC INSECTICIDE soil application in the same crop is not recommended.

**Incorporation:** Incorporation of IMIDAPRO 2SC SYSTEMIC INSECTICIDE can be achieved by cultivation, irrigation, rainfall, mechanical placement, soil injection, drenching, and broadcast sprays.

**Woody Perennials:** Onset of protection is slower than in herbaceous species. A delay of 2 or more weeks should be expected. Longer delays may be expected with larger plants. Therefore, make application well in advance of expected insect activity.

**Bark Media:** Media with 30 to 50% or more bark content may confer a shorter period of protection when treated with IMIDAPRO 2SC SYSTEMIC INSECTICIDE.

**Tank Mixes:** IMIDAPRO 2SC SYSTEMIC INSECTICIDE has been found to be compatible with commonly used liquid fertilizers, fungicides and insecticides. Check physical compatibility using the correct proportion of products in a small jar test if local experience is unavailable.

#### Application Through Irrigation Systems

ImidaPro 2SC Systemic Insecticide may be applied at rates specified on this label either alone or in tank mixture with other pesticides and chemicals registered for application through irrigation systems. The normal dilution ratio is 1:10 to 1:200, depending on the system. Always meter the product into the irrigation water during the first part of the irrigation cycle. The product may be mixed separately prior to injection. Agitation may be necessary if the mixture is allowed to stand more than 24 hours. Remove scale, pesticide residue and other foreign matter from the tank and entire irrigation system. Apply ImidaPro 2SC Systemic Insecticide only through microirrigation (individual spaghetti tube), drip irrigation, overhead irrigation, ebb and flood, or hand-held or motorized calibrated irrigation equipment.

Do not apply this product through any other type of irrigation system. Crop injury or lack of effectiveness can result from nonuniform distribution of treated water.

If you have any questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or a person who is under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

**SAFETY DEVICES FOR IRRIGATION SYSTEMS CONNECTED TO PUBLIC WATER SUPPLIES:**

If the source of water for your irrigation system is a public water supply, follow the instructions below.

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or over-flow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system inter-lock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

**SAFETY DEVICES FOR IRRIGATION SYSTEMS NOT CONNECTED TO A PUBLIC WATER SUPPLY:**

1. The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system inter-lock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where the pesticide distribution is adversely affected.
6. Systems must use a metering pump such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

**DRENCH AND IRRIGATION APPLICATIONS**

For use only on ornamental and vegetable plants in greenhouses, nurseries and interior plantscapes using soil drenches, micro-irrigation, drip irrigation, overhead irrigation, ebb and flood irrigation, or hand-held or motorized calibrated irrigation equipment.

Pest	Use Pattern		Dosage - IMIDAPRO 2SC SYSTEMIC INSECTICIDE		Remarks
			Container Size (inches)	No. pots treated with 1.7 fl oz (50 ml)	
Adelgids Aphids Armored scale (suppression) Fungus gnats <sup>1</sup> (larvae only) Japanese Beetle (adults) Lacebugs Leaf beetles (including elm and viburnum leaf beetles) Leaf hoppers (including glassy-winged sharpshooter) Leafminers Mealybugs Psyllids Root mealybugs <sup>2</sup> Root Weevil Complex (Such as Black Vine Weevil, ApopkaWeevil, Citrus Root Weevil <sup>3</sup> ) Soft Scale Thrips (suppression) <sup>4</sup> White Grub larvae (such as Japanese Beetle, Masked Chafers, European Chafer, Oriental Beetle, Asiatic Garden Beetle) Whiteflies	Plants in containers	Herbaceous Species - including Vegetable Plants <sup>5</sup> (one or two plants per pot)	2	3000	Evenly distribute one 1.7 fl oz (50 ml) of ImidaPro 2SC Systemic Insecticide in the stated number of pots, using sufficient water volume to wet potting medium without loss of liquid through leaching. Apply according to label directions.  Follow application with moderate irrigation. Irrigate carefully during the next 10 days in order to avoid loss of active ingredient due to leaching.
		3	2000		
		4	1500		
		5	1200		
		6	1000		
		7	850		
		8	750		
		9	675		
		10	600		
		11	550		
		12	500		
			Woody Perennial Species	2	
		3	1350		
		4	1000		
		5	800		
		6	650		
		7	550		
		8	500		
		9	450		
		10	400		
		11	350		
		12	300		
		Use the above Woody perennial Species rates			
		Herbaceous Species - including Vegetable Plants <sup>5</sup> (three or more plants per pot)			
	Ornamental and vegetable Plants <sup>5</sup> grown in flats benches, or beds	1.7 fl oz (50 ml) per 3000 sq ft		Mix required amount in sufficient water to uniformly and accurately cover the area being treated. Do not use less than 2 gallons of mixture per 1000 sq. ft. Apply as a broadcast treatment and incorporate into the medium before planting or apply after plants are established. If application is made to established plants, optimum control will be attained if areas are lightly irrigated after application. Allow no leaching or run out for 10 days after application.	
	Containerized Plants		No. Pots Treated with 1.7 fl oz (50 ml)	Apply in sufficient water to wet the potting medium. For optimum control, make applications prior to egg hatch of the target pest. Irrigate moderately after application to move the active ingredient into the root zone.	
	Plants	Container Size			
		1 gallon	340 to 244		
		2 gallon	280 to 210		
		3 gallon	220 to 165		
		5 gallon	160 to 110		
		7 gallon	100 to 75		
		10 gallon	60 to 45		
		15 gallon	40 to 30		
		20 gallon	20 to 15		

<p>White Grub larvae (such as Japanese Beetle, Masked Chafers, European Chafer, Oriental Beetle, Asiatic Garden Beetle)</p>	<p>Field and Forest Nurseries</p>	<p>Apply as a uniform band on either side of row using a band six (6) inches wider than the actual root ball diameter to be dug. Do not allow bands in adjacent rows to overlap. Use 1.7 fl oz (50 ml) per 1000 ft. of row or 3000 sq. ft. For grub control in areas of turf, apply as a broadcast application using 1.35 to 1.7 fl oz (40 to 50 ml) per 3000 sq. ft.</p>	<p>Vegetation in the area to be treated should be mowed to a height of 3 inches or less prior to application. Mowing to the lowest possible height will insure greater consistency of control. Apply May through July. For optimum control, treatment should be followed by rainfall or irrigation. Do not use less than 2 gallons of spray volume per 1000 square feet.</p>
---	-----------------------------------	---	--

**APPLICATIONS FOR NURSERY, GREENHOUSE AND INTERIORESCAPE PLANTS**

<p>Adelgids Aphids Armored scales (suppression) Black vine weevil larvae Eucalyptus longhorned borers Flatheaded borers (excluding emerald ashborer, but including bronze birch and alder borers)</p>	<p>Japanese beetles (adults) Lacebugs Leaf beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassy-winged sharpshooter) Leafminers Mealybugs</p>	<p>Pine Tip moth larvae Psyllids Royal palm bugs Sawfly larvae Soft scales Thrips (suppression) White grub larvae Whiteflies</p>
---	--	--

<p><b>Trees</b></p>	<p>0.1 to 0.2 fl oz (3 to 6 ml) per inch of cumulative trunk diameter</p>
---------------------	---

**Soil Injection:** GRID SYSTEM: Space holes on 2.5 foot centers, in a grid pattern, extending to the drip line of the tree. CIRCLE SYSTEM: Apply in holes evenly spaced in circles, (use more than one circle dependent upon the size of the tree) beneath the drip line of the tree extending in from that line. BASAL SYSTEM: Space injection holes evenly around the base of the tree trunk no more than 6 to 12 inches out from the base. Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. For optimum control, keep the treated area moist for 7 to 10 days. Do not use less than 4 holes per tree.  
**No Soil Injection Application Allowed in Nassau or Suffolk Counties of New York.**  
**Soil Drench:** Uniformly apply the dosage in no less than 10 gallons of water per 1000 square feet as a drench around the base of the tree, directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone.  
**For Control of Specified Borers:** Application to trees already heavily infested may not prevent the eventual loss of the trees due to existing pest damage and tree stress.

<p><b>Shrubs</b></p>	<p>0.1 to 0.2 fl oz (3 to 6 ml) per foot of shrub height</p>
----------------------	--

**Soil Injection:** Apply to individual plants using dosage indicated. Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. Keep the treated area moist for 7 to 10 days. Do not use less than 4 holes per shrub.  
**No Soil Injection Application Allowed in Nassau or Suffolk Counties of New York.**  
**Soil Drench:** Uniformly apply the dosage in no less than 10 gallons of water per 1000 square feet as a drench around the base of the tree, directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone.

<p><b>Flowers and Ground Covers</b></p>	<p>0.45 to 0.6 fl oz (13 to 17 ml) per 1000 sq ft</p>
---	---

Apply as a broadcast treatment and incorporate into the soil before planting or apply after plants are established. If application is made to established plants, optimum control will be attained if area is irrigated thoroughly after application

- <sup>1</sup> Fungus Gnat larvae in the soil will be controlled by drench or incorporation. No adult Fungus Gnat control. Other foliar insect control is achieved by the uptake of ImidaPro 2SC Systemic Insecticide from a healthy root system translocating the active ingredient up into the plant.
- <sup>2</sup> Root Mealybug control will require a thorough drenching of containerized media. Coverage is essential for control while minimizing the amount of leachate. **Rate: 1.7 fl oz (50 ml) in 150 gallons of water.**
- <sup>3</sup> Citrus Root Weevil: For use on non-bearing citrus nursery stock.
- <sup>4</sup> Thrips suppression on foliage only. Thrips in buds and flowers will not be suppressed.
- <sup>5</sup> **Restriction:** For use on vegetable plants intended for resale only including: Broccoli, Chinese Broccoli, Broccoli Raab, Brussels Sprouts, Cabbage, Chinese Cabbage, Cauliflower, Collards, Eggplant, Ground Cherry, Kale, Kohirabi, Lettuce, Mustard Greens, Pepinos, Peppers, Potatoes, Rape Greens, Sorghum, Sugarbeets, Tomatillo, and Tomato.

**Application to Grassy Areas in Nurseries Only:** ImidaPro 2SC Systemic Insecticide can be used for the control of the following soil inhabiting pests of grassy areas of nurseries: Northern and Southern masked chafers, *Cyclocephala borealis*, *C. immaculata*, and/or *C. lurida*; Asiatic garden beetle, *Maladera castanea*; European chafer, *Rhizotrogus majalis*; Green June beetle, *Cotinis nitida*; May or June beetle, *Phylophaga* spp.; Japanese beetle, *Popillia japonica*; Oriental beetle, *Anomala orientalis*; Billbugs, *Spherophorus* spp.; Annual bluegrass weevil, *Hyperodes* spp.; Black turfgrass ataenius, *Ataenius spretulus* and *Aphodius* spp. and mole crickets, *Scapteriscus* spp. ImidaPro 2SC Systemic Insecticide can also be used for suppression of cutworms and hairy chinchbugs. ImidaPro 2SC Systemic Insecticide can be used as directed on nursery grass in sites such as under or around field or container grown plants, on roadways or other grassy areas in or around nurseries.

The active ingredient in ImidaPro 2SC Systemic Insecticide has sufficient residual activity so that applications can be made preceding the egg laying activity of the target pests. High levels of control can be achieved when applications are made preceding or during the egg laying period. The need for an application can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. Optimum control will be achieved when applications are made prior to egg hatch of the target pests, followed by sufficient irrigation or rainfall to move the active ingredient through the thatch.

Do not make applications when grassy areas are water-logged or the soil is saturated with water. Adequate distribution of the active ingredient cannot be achieved when these conditions exist. The treated grassy area must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile. Application cannot exceed a total of 1.6 pint (0.4 lb of active ingredient) per acre per year. Refer to the "Application in Turf Grass" section for application rates.

**Application Equipment for Use on Grassy Areas in Nurseries:** Apply ImidaPro 2SC Systemic Insecticide in sufficient water to provide adequate distribution in the treated area. The use of accurately calibrated equipment normally used for soil application of insecticides is required. Use equipment which will produce a uniform, coarse droplet spray, using a low pressure setting to eliminate off-target drift. Check calibration periodically to ensure that equipment is working properly.



## ImidaPro 2SC Systemic Insecticide Ebb & Flood Application

**ImidaPro 2SC Systemic Insecticide** may be applied through Ebb and Flood applications. To assure accurate uptake it is recommended that prior to treatment, a minimum of 10 plants be brought up to a known field capacity and allowed to dry out for one or two days. Re-wet these plants to determine how much water on average each plant will absorb to bring it back at field capacity. Use the volume absorbed per plant (keeping pot sizes uniform) multiplied by the number of pots being treated. Add to this volume a required minimum to flood your smallest treatment area. This should minimize the return back to the storage tank. Reuse the returned volume with subsequent irrigation or nutrients on the same plants.

ImidaPro 2SC Systemic Insecticide EBB & FLOOD APPLICATIONS		
Adelgids Aphids Armored scales (suppression) Fungus Gnats (larvae only) <sup>1</sup> Japanese Beetles (adults) Lacebugs Leaf beetles (including elm and viburnum leaf beetles)	Leafhoppers (including glassy-winged sharpshooter) Leafminers Mealybugs Psyllids Root mealybugs <sup>2</sup> Root Weevil Complex: (such as Apopka Weevil, Black Vine Weevil, Citrus Root Weevil <sup>3</sup> )	Soft Scales Thrips (suppression) <sup>4</sup> Whiteflies White Grub Larvae: (such as Japanese Beetle, Masked Chafers, European Chafer, Oriental Beetle, Asiatic Garden Beetle)
Pot sizes (inches)	Herbaceous species including vegetable plants <sup>5</sup> (1 or 2 plants per pot)	Woody perennials, Herbaceous species including vegetable plants <sup>5</sup> (3 or more plants per pot)
	ML/100 plants	ML/100 plants
2	1.6	2.5
3	2.5	3.7
4	3.3	5.0
5	4.2	6.3
6	5.0	7.7
7	5.9	9.1
8	6.6	10.0
9	7.4	11.1
10	8.3	12.5
11	9.0	14.3
12	10.0	16.7
<p><sup>1</sup> Fungus gnat larvae in the soil will be controlled by drench or incorporation. No adult Fungus Gnat control. Other foliar insect control is achieved by the uptake of IMIDAPRO 2SC SYSTEMIC INSECTICIDE from a healthy root system translocating the active ingredient up into the plant.</p> <p><sup>2</sup> Root Mealybug control will require a thorough drenching of containerized media. Coverage is essential for control while minimizing the amount of leachate. <b>Rate: 1.7 fl oz (50 mL) in 150 gallons of water.</b></p> <p><sup>3</sup> Citrus Root Weevil: For use on non-bearing citrus nursery stock.</p> <p><sup>4</sup> Thrips suppression on foliage only. Thrips in buds and flowers will not be suppressed.</p> <p><sup>5</sup> Restriction: For use on vegetable plants intended for resale only including: Broccoli, Chinese Broccoli, Broccoli Raab, Brussels Sprouts, Cabbage, Chinese Cabbage, Cauliflower, Collards, Eggplant, Ground Cherry, Kale, Kohlrabi, Lettuce, Mustard Greens, Pepinos, Peppers, Potatoes, Rape Greens, Sorghum, Sugarbeets, Tomatillo, and Tomato.</p>		

## RESTRICTIONS

Do not graze treated areas or use clippings from treated areas for feed or forage.

Do not apply to soils that are water-logged or saturated, which will not allow the penetration of the insecticide into the root zone of the plants

Do not allow leachate run out for the first 10 days after application, in order to retain the product and facilitate full plant uptake of the active ingredient.

For outdoor ornamentals grown in beds or turf, applications of ImidaPro 2SC Systemic Insecticide cannot exceed a total of 1.6 Pints (0.4 lb of active ingredient) per acre per year.

On plants with a production cycle of less than one year, application is not to exceed a frequency of more than once each 16 weeks for a particular plant. On stock plants and woody crops with a production cycle of greater than one year, application may not exceed once a year.

**Food Crops:** Treated areas may be replanted with any crop specified on an imidacloprid label, or with any crop for which a tolerance exists for the active ingredient.

For crops not listed on an imidacloprid label a 12 month plant-back interval must be observed.

## APPLICATION TO TURF GRASS

ImidaPro 2SC Systemic Insecticide can be used for the control of the following soil inhabiting pests of grassey areas of nurseries: Northern & Southern masked chafers, *Cyclocephala borealis*, *C. immaculate*, and/or *C. lurida*; Asiatic garden beetle, *Maladera castanea*; European chafer, *Rhizotrogus majalis*; Green June beetle, *Cotinis nitida*; May or June beetle, *Phyllophaga* spp.; Japanese beetle, *Popillia japonica*; Oriental beetle, *Anomala orientalis*; Billbugs, *Sphenophorus* spp.; Annual bluegrass weevil, *Listronotus* spp.; Black turf grass ataenius, *Ataenius spretulus* and *Aphodius* spp; European crane fly, *Tipula paludosa*; and mole crickets, *Scapteriscus* spp.. ImidaPro 2SC Systemic Insecticide can also be used for suppression of cutworms and chinch bugs. ImidaPro 2SC Systemic Insecticide can be used as directed on turfgrass in sites home lawns, business and office complexes, shopping complexes, multi-family residential complexes, golf courses, airports, cemeteries, parks, playgrounds, athletic fields and sod farms.

The active ingredient in ImidaPro 2SC Systemic Insecticide has sufficient residual activity so that applications can be made preceding the egg laying activity of the target pests. High levels of control can be achieved when applications are made preceding or during the egg laying period. The need for an application can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. Optimum control will be achieved when applications are made prior to egg hatch of the target pests, followed by sufficient irrigation or rainfall to move the active ingredient through the thatch.

Do not make applications when turf grass areas are waterlogged or the soil is saturated with water. Adequate distribution of the active ingredient cannot be achieved when these conditions exist. The treated turf area must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile. Applications cannot exceed a total of 1.6 pints (0.4 lb of active ingredient) per acre per year.

### APPLICATION EQUIPMENT FOR USE ON TURF GRASS

Apply ImidaPro 2SC Systemic Insecticide in sufficient water to provide adequate distribution in the treated area. The use of accurately calibrated equipment normally used for the application of turf grass insecticides is required. Use equipment that will produce a uniform, coarse droplet spray, using a low pressure setting to eliminate off target drift. Check calibration periodically to ensure that equipment is working properly. **Do not apply through any irrigation system.**

TURF GRASS APPLICATIONS			
CROP	PEST	DOSAGE - ImidaPro 2SC Systemic Insecticide	REMARKS
Turf Grasses	Larvae of: Annual bluegrass weevil Asiatic garden beetle Billbugs Black turf grass ataenius Cutworm (suppression) European chafer European crane fly Green June beetle Japanese beetle Northern masked chafer Oriental beetle <i>Phyllophaga</i> spp. Southern masked chafer	1.25 to 1.6 pt per acre or 0.46 to 0.6 fl oz (14 to 17 mL) per 1000 sq ft	For optimum control of grubs, billbugs and annual bluegrass weevil, and European crane fly make application prior to egg hatch of the target pest. Be sure to read "APPLICATION EQUIPMENT" Section of this label.
	Chinchbugs (suppression) Mole crickets	1.6 pt per acre or 0.6 fl oz (17 mL) per 1000 sq ft	For suppression of chinchbugs, make application prior to or during the hatching of the first instar nymphs. For control of mole crickets make application prior to or during the peak egg hatch period. When adults or large nymphs are present and actively tunneling, ImidaPro 2SC Systemic Insecticide application should be accompanied by a remedial insecticide. Follow label instructions for other insecticides when tank-mixing.
<p>Consult your local turf, state Agricultural Experiment Station, or State Extension Service Specialists for more specific information regarding timing of application.</p> <p><b>RESTRICTIONS:</b> If there is no rainfall within 24 hours, follow treatment with irrigation to move the active ingredient through the thatch. Do not apply more than 1.6 pt (0.4 lb of active ingredient) per acre per year. Avoid mowing turf or lawn area until after sufficient irrigation or rainfall has occurred so that uniformity of application will not be affected.</p>			

APPLICATIONS – ORNAMENTAL TREES, SHRUBS, FLOWERS AND GROUNDCOVERS For industrial, commercial buildings and residential planting areas.			
CROP	PEST	DOSAGE - ImidaPro 2SC Systemic Insecticide	REMARKS
Trees Shrubs Evergreens Flowers Foliage Plants Groundcovers Interior Plantscapes	Adelgids Aphids Japanese beetles Lace bugs Leaf beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassy-winged sharpshooter) Mealybugs Psyllids Sawfly larvae Thrips (suppression) Whiteflies	1.5 fl oz (45 mL) per 100 gal of water	Foliar Applications: Start treatments prior to establishment of high pest populations and reapply on as needed basis.
	White grub larvae (such as Japanese beetle larvae, Chafers, <i>Phyllophaga</i> sp Asiatic garden beetle, Oriental beetle)	0.46 to 0.6 fl oz (14 to 17 mL) per 1000 sq ft	<b>Broadcast Applications:</b> Mix required amount of product in sufficient water to uniformly and accurately cover the area being treated. Do not use less than 2 gallons of water per 1000 sq ft. For optimum control, irrigate thoroughly to incorporate ImidaPro 2SC Systemic Insecticide into the upper soil profile. Refer to use direction specific for FLOWERS and GROUND COVERS concerning additional use directions.

**APPLICATIONS – ORNAMENTAL TREES, SHRUBS, FLOWERS AND GROUNDCOVERS**  
**For use only in and around industrial and commercial buildings and residential areas, and state, national, and private wooded and forested areas for the insect pests listed below.**

CROP	PEST	DOSAGE - ImidaPro 2SC Systemic Insecticide	REMARKS
Trees	Adelgids Aphids Armored scales (suppression) Black vine weevil larvae Eucalyptus longhorned borer Flatheaded borers(except emerald ash borer, but, including bronze birch and alder borer) Japanese beetles Lace bugs Leaf beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassy-winged sharpshooter) Leafminers Mealybugs Pine tip moth larvae Psyllids Royal palm bugs Sawfly larvae Soft scales Thrips (suppression) White grub larvae Whiteflies	0.1 to 0.2 fl oz (3 to 6 mL) per inch of trunk diameter (D.B.H.)	<p><b>Soil Injection:</b> GRID SYSTEM: Space holes on 2.5 foot centers, in a grid pattern, extending to the drip line of the tree. CIRCLE SYSTEM: Apply in holes evenly spaced in circles, (use more than one circle dependent upon the size of the tree) beneath the drip line of the tree extending in from that line. BASAL SYSTEM: Space injection holes evenly around the base of the tree trunk no more than 6 to 12 inches out from the base.</p> <p>Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. For optimum control, keep the treated area moist for 7 to 10 days. Do not use less than 4 holes per tree.</p> <p><b>RESTRICTION: No Soil Injection Applications Allowed in Nassau or Suffolk Counties of New York.</b></p> <p><b>Soil Drench:</b> Uniformly apply the dosage in no less than 10 gallons of water per 1000 square feet as a drench around the base of the tree, directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone.</p> <p><b>For Control of Specified Borers:</b> Application to trees already heavily infested may not prevent the eventual loss of the trees due to existing</p>
Shrubs		0.1 to 0.2 fl oz (3 to 6 mL) per foot of shrub height	<p><b>Soil Injection:</b> Apply to individual plants using dosage indicated.</p> <p>Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. Keep the treated area moist for 7 to 10 days. Do not use less than 4 holes per shrub.</p> <p><b>RESTRICTION: No Soil Injection Applications Allowed in Nassau or Suffolk Counties of New York.</b></p> <p><b>Soil Drench:</b> Uniformly apply the dosage in no less than 10 gallons of water per 1000 square feet as a drench around the base of the tree, directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone.</p>
Flowers and Groundcovers		0.46 to 0.6 fl oz (14 to 17 mL) per 1000 sq ft	Apply as a broadcast treatment and incorporate into the soil before planting or apply after plants are established. If application is made to established plants, optimum control will be attained if area is irrigated thoroughly after application.

FRUIT TREE APPLICATIONS For use only in residential areas.			
CROP	PEST	RATE PER APPLICATION	
Pome Fruits Apple Crabapple Loquat Mayhew Pear Pear (oriental) Quince	Aphids (except Woolly apple aphid) Leafhoppers (including glassy-winged sharpshooter) Leafminer Mealybugs* San Jose scale*	1.5 fl oz (45 mL) per 100 gal of water	6.0 fl oz/A <sup>1</sup>
<p>Apply specified dosage as foliar spray as needed after petal-fall is complete.</p> <p>For control of rosy apple aphid, apply prior to leafrolling caused by the pest.</p> <p>For first generation leafminer control, make first application as soon as petal-fall is complete. Greatest leafminer control will result from the earliest possible application. For second and succeeding generations of leafminer, optimal control is obtained from applications made early in the adult flight against egg and early instar larvae. A second application may be required 10 days later if severe pressure continues or if generations are overlapping. A single application may result in suppression only. ImidaPro 2SC Systemic Insecticide will not control late stage larvae.</p> <p>For San Jose Scale, time applications to the crawler stage. Treat each generation.</p> <p>For late season (preharvest) control of leafhopper species, apply ImidaPro 2SC Systemic Insecticide while most leafhoppers are in the nymphal stage.</p> <p>For optimal control of mealybug, insure good spray coverage of the trunk and scaffolding limbs or other resting sites of the mealybug.</p> <p>Do not apply more than 6.0 fluid ounces per acre in a single application.</p> <p>Do not make more than 5 applications per year.</p> <p>Do not apply more that 30 fluid ounces of ImidaPro 2SC Systemic Insecticide per acre per year.</p> <p>Allow 10 or more days between applications. Allow at least 7 days between last application and harvest.</p> <p>* Not for use in California for control on pears.</p>			
Pecans*	Yellow pecan aphid Black margined aphid Pecan leaf phylloxera Pecan spittlebug Pecan stem phylloxera	1.5 fl oz (45 mL) per 100 gal of water	6.0 fl oz/A <sup>1</sup>
<p>Make foliar applications as pests begin to build before populations become extreme. Two applications at a 10 to 14 day interval may be required to achieve control. Scout and retreat if needed.</p> <p>Thorough uniform coverage of foliage is necessary for optimal control. Addition of an organosilicone-based spray adjuvant at a rate not to exceed the adjuvant manufacturer's specified use rate may improve coverage.</p> <p>Do not make more than 3 applications per year.</p> <p>Do not apply more than a total of 18.0 fluid ounces of ImidaPro 2SC Systemic Insecticide per acre per year.</p> <p>Allow 10 or more days between applications.</p> <p><sup>1</sup>The amount of ImidaPro 2SC Systemic Insecticide required per acre will depend on tree size and volume of foliage present. The rate per acre is based on a standard of 400 gallons of dilute spray solution per acre for large trees.</p> <p>* Use on pecans not permitted in California unless directed by specific supplemental labeling.</p>			

VINE APPLICATIONS				
For industrial, commercial buildings and residential planting areas.				
CROP	PEST		RATE PER APPLICATION	
Grapes	Leafhoppers (including glassy-winged sharpshooter)	Mealybugs	1.5 fl oz (45 mL) per 100 gal of water	3.0 fl oz/A (90 mUA)
Apply specified dosage as a foliar spray using 200 gallons of water per acre. Do not apply more than a total of 6.0 fl ounces of ImidaPro 2SC Systemic Insecticide per acre per year. Allow at least 14 days between applications. Applications may be applied up to and including day of harvest.				

**RESTRICTIONS**

Do not graze treated areas or use clippings from treated areas for feed or forage. Avoid runoff or puddling of irrigation water following application. Keep children and pets off treated area until dry. Avoid application of ImidaPro 2SC Systemic Insecticide to areas that are water logged or saturated, which will not allow penetration into the root zone of the plant. Treated areas may be replanted with any crop specified on an imidacloprid label, or with any crop for which a tolerance exists for the active ingredient. For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12-month plant-back interval must be observed.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

**Pesticide Storage:** Store in a tightly closed container in a cool, dry place.

**Pesticide Disposal:** Pesticide spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Container Disposal (Nonrefillable container 5 gallons or less):** Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Offer for recycling, if available.

**Residue Removal:** Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Refillable container.** Refill this container with pesticide only. Do not reuse this container for any other purpose. Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. Use standard industry practices for cleaning refillable containers.

**Spills:** For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire or other emergency, call **CHEMTREC Day or Night, DOMESTIC NORTH AMERICA 1-800-424-9300.**

**CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of AGRISEL USA INCORPORATED or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold AGRISEL USA INCORPORATED and Seller harmless for any claims relating to such factors.

AGRISEL USA INCORPORATED warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or AGRISEL USA INCORPORATED, and Buyer and User assume the risk of any such use. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW AGRISEL USA INCORPORATED MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

In no event shall AGRISEL USA INCORPORATED or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF AGRISEL USA INCORPORATED AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF AGRISEL USA INCORPORATED OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

AGRISEL USA INCORPORATED and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of AGRISEL USA INCORPORATED.

Registered: [DATE TBA]

**Manufactured For:  
AGRISEL USA INC  
PO Box 3528  
Suwanee, GA 30024  
1-877-247-4735**