


Kaput

For control of Deer Mice,
White-Footed Mice,
and House Mice



MOUSE BLOCKS

NET WT. 10 LBS. (4.54 kg)

ACTIVE INGREDIENT: Warfarin	
3-(α -Acetonylbenzyl-4-hydroxycoumarin).....	00.025%
INERT INGREDIENTS	99.975%
TOTAL	100.000%

EPA Reg. No. 72500-7
EPA Est. No. 72500-CO-1



ABOUT
105
BLOCKS

ACCEPTED
00 28 2004

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended by the FIFRA Act of 1992
Registered under EPA Reg. No. 72500-7

SYNTHETIC

CAUTION

72500-7

10/28/2004

1/2

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS**

CAUTION: Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN: This product reduces the clotting ability of the blood and may cause hemorrhaging. If poisoning occurs, intramuscular and oral administration of Vitamin K1 are indicated, as in poisoning from an overdose of bis-hydroxycoumarin. For human cases, Vitamin K1 is antidotal at doses of 10-20 mg total (not mg/kg). For animal cases, Vitamin K1 is antidotal at 2-5 mg/kg. Repeated doses may need to be given up to two weeks (based upon monitoring of prothrombin times).

ENVIRONMENTAL HAZARDS

Do not apply this product directly to water, or to areas where surface water is present or the intertidal areas below the mean high water mark.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container in a dry place inaccessible to children and pets.

DISPOSAL: Do not reuse empty container. Securely wrap container and unused bait in several layers of newspaper and discard in trash.

NOTICE OF WARRANTY

Manufacturer and Seller make no warranty of merchantability, fitness for any purpose, or otherwise, express or implied, concerning this product or its use, which extend beyond the statements on this label.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children to this product or use it for any purpose other than to control deer mice, house mice and white-footed mice. To help prevent accidents:

1. Store product not in use in locations out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or areas prone to vandalism.
3. Dispose of product container, and unused, spoiled or unconsumed bait as specified on this label.

USE RESTRICTIONS: This product may only be used to control deer mice (*Peromyscus maniculatus*), white-footed mice (*P. leucopus*), and house mice (*Mus musculus*). This product may be used in and around homes, agricultural and industrial buildings, and similar manmade structures. If applied in secured, tamper-resistant bait stations, this product also may be used in parks, woodlots, yards and lots surrounding

residential buildings, and non-crop areas near agricultural buildings. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not use this product within the occupied habitats of threatened or endangered *Peromyscus spp.* mice.

SELECTION OF TREATMENT AREAS:

Determine where mice will most likely find and consume the bait. Generally, these areas are along walls, along building foundations, in cellars and crawl spaces, and other locations where mice or signs of mice are seen in or near homes, cabins, outbuildings, and similar structures. When controlling white-footed mice, the bases of trees located in permitted use sites (see **USE RESTRICTIONS**) may provide suitable baiting points. When bait is applied, remove as much as possible of the foods that the mice have been eating.

APPLICATION DIRECTIONS:

All applications, including indoor applications, must be made in tamper-resistant bait stations unless children, pets, domestic animals, and non-target wildlife are reliably denied access to the treated area by other means. Apply one or two bait blocks per placement. Use two blocks only at points of extremely high mouse activity. Space placements at intervals of 8 to 12 feet in infested area. Maintain an uninterrupted supply of fresh bait for at least 30 days or until signs of mouse activity cease to appear. Replace contaminated or spoiled bait immediately. To discourage re-infestation, eliminate food, water, and harborage as much as possible and attempt to locate and seal openings through which deer mice or white-footed mice enter buildings. If re-infestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.



LTD. CORP.

New Solutions to Old Problems

120 Commerce Dr., Unit 4 Fort Collins, CO 80524
(970) 482-1330 www.scimetricsltd.com



8 53215 00-04 8