

72500-13

4-2-2008

1/12



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505C)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

72500-13

Date of Issuance:

APR 2 2008

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Novel Commensal Rodent Pellet #2

Name and Address of Registrant (include ZIP Code):

Scimetrics Ltd. Corp
c/o RegWest Company, LLC
30856 Rocky Road
Greeley, CO 80631-9375

R23 5-826989
1160

Attention: Ms. Kim Davis

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you

A) submit one (1) copy of final printed labeling to us before you release the product for shipment.

DBP
Daniel B. Peacock, Biologist
Insecticide-Rodenticide Branch
Registration Division (7505C)

Date:

APR 2 2008

B) within 15 months of registration, submit the following data in electronic and paper formats:

1. Storage Stability (830.6317)
2. Corrosion Characteristics (830.6320)

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed. If you have any questions, please contact Dan Peacock by phone (703-305-5407), fax (703-305-6596), or E-Mail (peacock.dan@epa.gov).

- Enclosures:
1. Stamped Label
 2. Minimum Type Size Requirements for Final Printed Labels

Dan Peacock, Flash Drive, 2gb, P:\Documents\Word, WP, Excel, etc\Warfarin - Imidacloprid\72500-13, Notice of Registration, 3-27-2008, ver 2.doc

3/12

Novel Commensal Rodent Pellet #2

Master Label

Master Label Includes:

Sub-Label A: Box/Pail Label

Sub-Label B: Place Pack Label (Unit Packaging)

Sub-Label C: Alternate Place Pack Label (When Sold Individually)

Kills Norway Rats, Roof Rats, House Mice and Voles

First dead rats, mice and voles will appear 4 to 5 days after feeding begins

Rodent Control Pellets

A Bait Containing a Rodenticide to Kill Norway Rats, Roof Rats, House Mice and Voles and an Insecticide to Aid in the Reduction of Flea Populations on Listed Rodent Hosts

Active Ingredients:

Imidacloprid (CAS Number 138261-41-3) 0.020%

Warfarin (CAS Number 81-81-2) 0.025%

Other Ingredients 99.955%

Total 100.000%

Keep Out of Reach of Children

CAUTION

Harmful if swallowed.

See back [side] panel for First Aid and additional Precautionary Statements.

EPA Reg. No. 72500-13

EPA Est. 72500-CO-1

Net Weight: 2 oz. (56 grams)

{Unit Packaging:}

12 oz. (6 packs x 2 oz.)

[9.38 lbs. (75 packs x 2 oz.)] in EPA Letter Dated:

[12.5 lbs. (100 packs x 2 oz.)]

[22.5 lbs. (180 packs x 2 oz.)]

[25 lbs. (200 packs x 2 oz.)]

**ACCEPTED
with COMMENTS**

APR 2 2008

Under the Federal Insecticide,

Fungicide, and Rodenticide Act

amended, for the pesticide

registered under EPA Reg. No.

Manufactured by:

Scimetrics Ltd. Corp.

P.O. Box 1045

Wellington, CO 80549

72500-13

New Solutions to Old Problems

[] Denotes optional/alternate language

{ } Denotes language that does not appear on the market label

4/12

Sub-Label A: Box/Pail Label

{Front Panel}

Novel Commensal Rodent Pellet #2
Place Packs

Kills Norway Rats, Roof Rats, House Mice and Voles
First dead rats, mice and voles will appear 4 to 5 days after feeding begins
Rodent Control Pellets

A Bait Containing a Rodenticide to Kill Norway Rats, Roof Rats, House Mice and Voles and an
Insecticide to Aid in the Reduction of Flea Populations on Listed Rodent Hosts

Active Ingredients:

Imidacloprid (CAS Number 138261-41-3)	0.020%
Warfarin (CAS Number 81-81-2)	0.025%
Other Ingredients	<u>99.955%</u>
Total	100.000%

Keep Out of Reach of Children
CAUTION

Harmful if swallowed.

See back panel for additional precautionary statements.

EPA Reg. No. 72500-13

EPA Est. 72500-CO-1

Net Weight: _____

{Unit Packaging:}

[12 oz. (6 packs x 2 oz.)]

[9.38 lbs. (75 packs x 2 oz.)]

[12.5 lbs. (100 packs x 2 oz.)]

[22.5 lbs. (180 packs x 2 oz.)]

[25 lbs. (200 packs x 2 oz.)]

Manufactured by:

Scimetrics Ltd. Corp

P.O. Box 1045

Wellington, CO 80549

New Solutions to Old Problems

{Back Panel}

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not use this product to control any pests, by any application method, or at any use sites, that are not specified on this label.

Use of this product may aid in the reduction of fleas, including by not necessarily limited to *Xenopsylla cheopis*, that commonly infest listed rodent hosts, if ingesting sufficient quantities of this bait.

5/12

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets or other non-target animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, OR in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and children under six year of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container and unused or unconsumed bait as specified by this label.

RATS AND MICE:

Use Restrictions: This product may be used to control Norway rats, roof rats and house mice at the sites in "Urban Areas" and "Non-urban Areas" that are indicated below. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come into contact with food. Wear gloves when handling bait or animal carcasses. Do not broadcast bait.

Urban Areas: To control Norway rats, roof rats and house mice this product may be used in and around homes, and industrial, commercial and public buildings; in transport vehicles (ship, trains and aircraft) and in and around related port or terminal buildings. Applications "around" buildings must be made in tamper-resistant bait stations placed along walls or other external parts of buildings (e.g. doorways, ramps, loading docks) where rats or mice might seek to gain entrance.

Non-urban Areas: To control Norway rats, roof rats and house mice this product may be used in and around homes and agricultural buildings. All outdoor placements must be in tamper-resistant bait stations.

Selection of Treatment Areas: Determine areas where rodents will most likely find and consume bait. Generally, these areas will be along walls, by gnawed openings, in corners and concealed places, between floors and walls, beside burrow openings and other locations where signs of rodents have been seen. Remove as much alternative food as possible.

Application Directions:

{Only 12-oz size:}

The amount of bait in this 6-pack package may be sufficient to kill 3 to 6 rats.

{All outer packages:}

Norway Rats and Roof Rats: Apply 2 to 8 [2 go 6 for 6-pack] place packs per bait placement. Space bait placements at intervals of 15 to 30 feet in the infested area, at locations consistent with the requirements of this label. Maintain an uninterrupted supply of fresh bait for 10 days or until there no longer are fresh signs of rat activity or signs that rats are feeding on the bait.

House Mice: Open pack and apply 1/4 to 1/2 ounce of bait per typical placement. Larger placement – up to one unopened 2-ounce pack – may be made at points of extremely high mouse activity. Space bait placements at intervals of 8 to 12 feet in the infested area, at locations consistent with the requirements on this label. Maintain an uninterrupted supply of fresh bait for 15 days or until there no longer are fresh signs of house mouse activity or signs that house mice are feeding on the bait.

Follow-up (Rats & Mice): Immediately replace contaminated or spoiled bait. Collect and properly dispose of all visible dead animals and leftover bait. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation occurs, repeat treatment. Where there is a continuous source of reinfestation, establish permanent bait stations and replenish bait as needed.

6/12

VOLES:

Use Restrictions: This product may be used to control meadow voles (*Microtus pennsylvanicus*), pine voles (*M. pinetus*) and montane voles (*M. montanus*) around (within 15 feet of) homes, and industrial, commercial, agricultural and public buildings. Do not use this product within the habitat of threatened or endangered *Microtus* spp. voles, threatened or endangered *Peromyscus* spp. mice, or other threatened or endangered rodents. Do not expose bait on bare ground. Unless children, pets, birds and other wildlife are excluded from the treated areas, all application for controlling voles must be made in tamper-resistant bait stations. All bait applications must be made by hand. Wear gloves when handling bait or animal carcasses. Do not broadcast bait.

Selection of Treatment Areas: Determine areas near buildings where voles or signs of vole activity (girdling or plants, consumption of bulbs, burrow openings and surface runways) have been seen. These areas generally will be in flower beds or under shrubs, bushes or ground cover.

Application Directions: Open pack and load one ounce of bait (half of a pack) into each bait station. Situate bait stations near runways, burrow openings or plants that voles have been damaging. If, and only if, children, pets, domestic animals and wildlife reliably are excluded from the treated area, bait may be applied in on-ounce amounts directly into vole runways or burrow openings. Cover treated runways or holes with a shingle or board, or ground vegetation, if present. Check placements daily. Remove any fouled or spoiled bait, and replenish if vole activity and interest in bait persist. Maintain an uninterrupted supply of fresh bait for 15 days or until there no longer are fresh signs of vole activity and feeding on bait. Collect and properly dispose of any carcasses and leftover bait.

{The following will be used for the 4 and 12 oz. sizes;}

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container in a cool, dry place inaccessible to children and pets. **Disposal: If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

{The following will be used for the 9.38 through 25 lbs. sizes;}

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container in a cool, dry place inaccessible to children and pets. **Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Disposal:** Non-Refillable Container. Do not reuse or refill this container. **{Plastic:}** Triple rinse (or equivalent) then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or incineration; or if allowed by state and local authorities, by burning; if burned, stay out of smoke. **{Paper:}** Dispose of empty paper box in a sanitary landfill or by incineration; or if allowed by state and local authorities, by burning; if burned stay out of smoke.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Keep away from humans, domestic animals and pets. Harmful if swallowed because this material may reduce the clotting ability of the blood and cause bleeding. Causes moderate eye irritation. Do not get in eyes or on skin or clothing. Wash arms, hands and face with soap and water after applying and before eating, drinking, chewing gum or using tobacco. Applicators must wear long-sleeved shirt, long pants, socks, shoes and gloves.

First Aid

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

7/12

If swallowed, call a poison control center, doctor or 1-800-858-7378 immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. **If in eyes**, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Notice to Physician or Veterinarian

This product reduces the clotting ability of the blood and may cause bleeding. If ingested, administer Vitamin K-1 intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.

Environmental Hazards

This product is highly toxic to aquatic invertebrates and toxic to mammals and birds. Do not apply this product directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwaters.

Warranty: To the extent consistent with applicable law, Manufacturer and Seller make no warranty of merchantability, fitness for any purpose, or otherwise express or implied, concerning this product or its use which extend beyond the statements on this label.

[] Denotes optional/alternate language

{ } Denotes language that does not appear on the market label

8/12

Sub-Label B: Place Pack Label (Unit Packaging)

{Front Panel}

Novel Commensal Rodent Pellet #2
Place Pack

Kills Norway Rats, Roof Rats, House Mice and Voles
First dead rats, mice and voles will appear 4 to 5 days after feeding begins
Rodent Control Pellets

A Bait Containing a Rodenticide to Kill Norway Rats, Roof Rats, House Mice and Voles and an
Insecticide to Aid in the Reduction of Flea Populations on Listed Rodent Hosts

Active Ingredients:

Imidacloprid (CAS Number 138261-41-3) 0.020%
Warfarin (CAS Number 81-81-2) 0.025%

Other Ingredients 99.955%
Total 100.000%

Keep Out of Reach of Children
CAUTION

Harmful if swallowed.

See back panel for additional precautionary statements.

EPA Reg. No. 72500-13

EPA Est. 72500-CO-1

Net Weight: 2 oz. (56 grams)

Manufactured by:

Scimetrics Ltd. Corp

P.O. Box 1045

Wellington, CO 80549

New Solutions to Old Problems

{Back Panel}

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read the entire label on the outer package before using this product. It is illegal to sell these place packs individually.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container in a cool, dry place inaccessible to children and pets. **Disposal: If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

9/12

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Keep away from humans, domestic animals and pets. Harmful if swallowed because this material may reduce the clotting ability of the blood and cause bleeding. Causes moderate eye irritation. Do not get in eyes or on skin or clothing. Wash arms, hands and face with soap and water after applying and before eating, drinking, chewing gum or using tobacco. Applicators must wear long-sleeved shirt, long pants, socks, shoes and gloves.

First Aid

Have the product container or label with you when calling a poison control center or doctor or going for treatment. **If swallowed**, call a poison control center, doctor or 1-800-858-7378 immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. **If in eyes**, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Notice to Physician or Veterinarian

This product reduces the clotting ability of the blood and may cause bleeding. If ingested, administer Vitamin K-1 intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.

Environmental Hazards

This product is highly toxic to aquatic invertebrates and toxic to mammals and birds. Do not apply this product directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwaters.

Warranty: To the extent consistent with applicable law, Manufacturer and Seller make no warranty of merchantability, fitness for any purpose, or otherwise express or implied, concerning this product or its use which extend beyond the statements on this label.

[] Denotes optional/alternate language

{ } Denotes language that does not appear on the market label

10/12

Sub-Label C: Alternate Place Pack Label (When Sold Individually)

{Front Panel}

Novel Commensal Rodent Pellet #2
Place Packs

Kills House Mice and Voles

First dead house mice and voles will appear 4 to 5 days after feeding begins

House Mouse and Vole Control Pellets

A Bait Containing a Rodenticide to Kill House Mice and Voles and an
Insecticide to Aid in the Reduction of Flea Populations on Listed Rodent Hosts

Active Ingredients:

Imidacloprid (CAS Number 138261-41-3)	0.020%
Warfarin (CAS Number 81-81-2)	0.025%
Other Ingredients	<u>99.955%</u>
Total	100.000%

Keep Out of Reach of Children

CAUTION

Harmful if swallowed.

See back panel for additional precautionary statements.

EPA Reg. No. 72500-13

EPA Est. 72500-CO-1

Net Weight: 2 oz. (56 grams)

Manufactured by:

Scimetrics Ltd. Corp

P.O. Box 1045

Wellington, CO 80549

New Solutions to Old Problems

{Back Panel}

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not use this product to control any pests, by any application method, or at any use sites, that are not specified on this label.

Use of this product may aid in the reduction of fleas, including but not necessarily limited to *Xenopsylla cheopsis*, that commonly infest listed rodent hosts, if ingesting sufficient quantities of this bait.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets or other non-target animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.

11/12

2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, OR in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and children under six year of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooved livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container and unused or unconsumed bait as specified by this label.

HOUSE MICE:

Use Restrictions: This product may be used to control house mice at the sites in "Urban Areas" and "Non-urban Areas" that are indicated below. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come into contact with food. Wear gloves when handling bait or animal carcasses. Do not broadcast bait.

Urban Areas: To control house mice this product may be used in and around homes, and industrial, commercial and public buildings; in transport vehicles (ship, trains and aircraft) and in and around related port or terminal buildings. Applications "around" buildings must be made in tamper-resistant bait stations placed along walls or other external parts of buildings (e.g. doorways, ramps, loading docks) where rats or mice might seek to gain entrance.

Non-urban Areas: To control house mice this product may be used in and around homes and agricultural buildings.

Selection of Treatment Areas: Determine areas where house mice will most likely find and consume bait. Generally, these areas will be along walls, by gnawed openings, in corners and concealed places, between floors and walls, beside burrow openings and other locations where signs of house mice have been seen. Remove as much alternative food as possible.

Application Directions:

Open pack and apply 1/4 to 1/2 ounce of bait per typical placement. Larger placement – up to one unopened 2-ounce pack – may be made at points of extremely high mouse activity. Space bait placements at intervals of 8 to 12 feet in the infested area, at locations consistent with the requirements on this label. Maintain an uninterrupted supply of fresh bait for 15 days or until there no longer are fresh signs of house mouse activity or signs that house mice are feeding on the bait.

Follow-up: Immediately replace contaminated or spoiled bait. Collect and properly dispose of all visible dead animals and leftover bait. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation occurs, repeat treatment. Where there is a continuous source of reinfestation, establish permanent bait stations and replenish bait as needed.

VOLES:

Use Restrictions: This product may be used to control meadow voles (*Microtus pennsylvanicus*), pine voles (*M. pinetis*) and montane voles (*M. montanus*) around (within 15 feet of) homes, and industrial, commercial, agricultural and public buildings. Do not use this product within the habitat of threatened or endangered *Microtus* spp. voles, threatened or endangered *Peromyscus* spp. mice, or other threatened or endangered rodents. Do not expose bait on bare ground. Unless children, pets, birds and other wildlife are excluded from the treated areas, all application for controlling voles must be made in tamper-resistant bait stations. All bait applications must be made by hand. Wear gloves when handling bait or animal carcasses. Do not broadcast bait.

Selection of Treatment Areas: Determine areas near buildings where voles or signs of vole activity (girdling or plants, consumption of bulbs, burrow openings and surface runways) have been seen. These areas generally will be in flower beds or under shrubs, bushes or ground cover.

Application Directions: Open pack and load one ounce of bait (half of a pack) into each bait station. Situate bait stations near runways, burrow openings or plants that voles have been damaging. If, and only if, children, pets,

12/12

domestic animals and wildlife reliably are excluded from the treated area, bait may be applied in on-ounce amounts directly into vole runways or burrow openings. Cover treated runways or holes with a shingle or board, or ground vegetation, if present. Check placements daily. Remove any fouled or spoiled bait, and replenish if vole activity and interest in bait persist. Maintain an uninterrupted supply of fresh bait for 15 days or until there no longer are fresh signs of vole activity and feeding on bait. Collect and properly dispose of any carcasses and leftover bait.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container in a cool, dry place inaccessible to children and pets. **Disposal: If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Keep away from humans, domestic animals and pets. Harmful if swallowed because this material may reduce the clotting ability of the blood and cause bleeding. Causes moderate eye irritation. Do not get in eyes or on skin or clothing. Wash arms, hands and face with soap and water after applying and before eating, drinking, chewing gum or using tobacco. Applicators must wear long-sleeved shirt, long pants, socks, shoes and gloves.

First Aid

Have the product container or label with you when calling a poison control center or doctor or going for treatment. **If swallowed,** call a poison control center, doctor or 1-800-858-7378 immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. **If in eyes,** hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Notice to Physician or Veterinarian

This product reduces the clotting ability of the blood and may cause bleeding. If ingested, administer Vitamin K-1 intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.

Environmental Hazards

This product is highly toxic to aquatic invertebrates and toxic to mammals and birds. Do not apply this product directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwaters.

Warranty: To the extent consistent with applicable law, Manufacturer and Seller make no warranty of merchantability, fitness for any purpose, or otherwise express or implied, concerning this product or its use which extend beyond the statements on this label.

[] Denotes optional/alternate language

{ } Denotes language that does not appear on the market label