

72500-13

7.25.2008

1/17



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Ms. Kim Davis CC, APC Manager
For Scimetrics Ltd. Corporation
Reg West Company, LLC
30856 Rocky Road
Greely, CO 80631-9375

JUL 25 2008

SUBJECT: Application for Pesticide Notification (PRN 98-10)
Request General Label Change (Product Sizes)
EPA Reg. 72500-13
Application Dated June 27, 2008

Dear Registrant:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 06/27/08 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

A handwritten signature in cursive script, appearing to read "Linda Arrington".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060.

United States Environmental Protection Agency 1200 Pennsylvania Avenue, N.W. Washington, DC 20460-0001	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number <h2 style="text-align: center;">Notification</h2>
--	--	--

Application for Pesticide – Section I

1. Company/Product Number EPA Reg. No. 72500-13	2. EPA Product Manager John Hebert	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/ Product (Name) Novel Commensal Rodent Pellet #2	PM # 7	
5. Name and Address of Applicant (Include ZIP Code) Scimetrics Ltd. Corporation c/o RegWest Company, LLC 30856 Rocky Road Greeley, CO 80631-9375 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i) my product is similar or identical in composition and labeling to: <h2 style="text-align: center;">NOTIFICATION</h2> EPA Reg. No. <u>JUL 25 2008</u> Product Name _____	

Section – II

<input type="checkbox"/> Amendment – Explain below. <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification – Explain below.	<input checked="" type="checkbox"/> Final printed labels in response to Agency letter dated 04/02/2008 <input type="checkbox"/> "Me Too" Application <input type="checkbox"/> Other – Explain below.
--	---

Explanation: Use additional page(s) if necessary. (For Section I and Section II.)

**Submit notification of additional container sizes (with final printed labeling) per PR Notice 98-10.
The additional container sizes are: 1 lb. (8 packs x 2 oz.) and 3.125 lbs. (25 packs x 2 oz.)**

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR § 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to the EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR § 152.46 this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section – III

1. Material This Product Will Be Packaged In:			
Child Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify): _____
*Certification must be submitted		If "Yes" Unit Packaging Wt. No. Per Container	If "Yes" Packaging Wt. No. Per Container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 	
5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling Accompanying Product			
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Stenciled <input type="checkbox"/> Paper glued <input type="checkbox"/> Other _____			

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application).		
Name Kim Davis, CC, APC, Manager	Title Consultant/Agent	Telephone No. (Include Area Code) (970) 353-0611 kim@regwest.com
Certification		6. Date Application Received (Stamped)
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		
2. Signature 	3. Title Consultant/Agent	
4. Typed Name Kim Davis, CC, APC, Manager	5. Date June 27, 2008	

FedEx

FTN 8646 7118 7588



June 27, 2008

Mr. John Hebert, Product Manager 7
Document Processing Desk – **NOTIF/FPL**
Office of Pesticide Programs – **7504P**
U.S. Environmental Protection Agency
2777 South Crystal Drive
One Potomac Yard, Room S-4900
Arlington, VA 22202

Dear Mr. Hebert:

Subject: Scimetrics Ltd. Corp.
Novel Commensal Rodent Pellet #2
EPA Reg. No. 72500-13

On behalf of **Scimetrics Ltd. Corporation**, RegWest Company, LLC is submitting the attached documents as notification of additional product sizes (with final printed labeling) for subject product. RegWest will act as sole agent in this endeavor.

Attached are the following:

- 1. Application for Pesticide: Other (Notification)
- 2. One **Reference Label** (illustrating the additional product sizes)
- 3. Two **New Labels** (incorporating the additional product sizes)
- 4. Two copies of final printed labeling for each new product size

The additional product sizes are:

1 lb. (8 packs x 2 oz.) and 3.125 lbs. (25 packs x 2 oz.)

Please contact me at (970) 353-0611 or kim@regwest.com if you require additional information or have any questions.

Regards,
RegWest Company, LLC

Kim Davis, CC, APC, Manager
Consultant/Agent

Attachments

cc: SLC – Poché, R.

30856 Rocky Road
Greeley, CO 80631-9375
970-353-0611
970-353-0613 Fax
Kim@regwest.com

Novel Commensal Rodent Pellet #2

Master Label

Master Label Includes:

- Sub-Label A: Box/Pail Label
- Sub-Label B: Place Pack Label (Unit Packaging)
- Sub-Label C: Alternate Place Pack Label (When Sold Individually)

Kills Norway Rats, Roof Rats, House Mice and Voles
 First dead rats, mice and voles will appear 4 to 5 days after feeding begins.
 Rodent Control Pellets
 A Bait Containing a Rodenticide to Kill Norway Rats, Roof Rats, House Mice and Voles and an
 Insecticide to Aid in the Reduction of Flea Populations on Listed Rodent Hosts

Active Ingredients:

Imidacloprid (CAS Number 138261-41-3)	0.020%
Warfarin (CAS Number 81-81-2)	0.025%
Other Ingredients	<u>99.955%</u>
Total	100.000%

Keep Out of Reach of Children

CAUTION

Harmful if swallowed.

See back [side] panel for additional Precautionary Statements.

EPA Reg. No. 72500-13

EPA Est. 72500-CO-1

Net Weight: 2 oz. (56 grams)

{Unit Packaging:}

- 12 oz. (6 packs x 2 oz.)
- [1 lb. (8 packs x 2 oz.)
- [3.125 lb. (25 packs x 2 oz.)
- [9.38 lbs. (75 packs x 2 oz.)
- [12.5 lbs. (100 packs x 2 oz.)
- [22.5 lbs. (180 packs x 2 oz.)
- [25 lbs. (200 packs x 2 oz.)

Manufactured by:

Scimetrics Ltd. Corp

P.O. Box 1045

Wellington, CO 80549

New Solutions to Old Problems

[] Denotes optional/alternate language
 { } Denotes language that does not appear on the market label

NOTIFICATION

JUL 25 2008

Sub-Label A: Box/Pail Label

{Front Panel}

Novel Commensal Rodent Pellet #2
Place Packs

Kills Norway Rats, Roof Rats, House Mice and Voles
First dead rats, mice and voles will appear 4 to 5 days after feeding begins.

Rodent Control Pellets
A Bait Containing a Rodenticide to Kill Norway Rats, Roof Rats, House Mice and Voles and an
Insecticide to Aid in the Reduction of Flea Populations on Listed Rodent Hosts

Active Ingredients:

Imidacloprid (CAS Number 138261-41-3)	0.020%
Warfarin (CAS Number 81-81-2)	0.025%
Other Ingredients	<u>99.955%</u>
Total	100.000%

Keep Out of Reach of Children
CAUTION

Harmful if swallowed.
See back panel for additional Precautionary Statements.

EPA Reg. No. 72500-13

EPA Est. 72500-CO-1

- Net Weight:** _____
 {Unit Packaging:}
 [12 oz. (6 packs x 2 oz.)]
 [1 lb. (8 packs x 2 oz.)]
 [3.125 lb. (25 packs x 2 oz.)]
 [9.38 lbs. (75 packs x 2 oz.)]
 [12.5 lbs. (100 packs x 2 oz.)]
 [22.5 lbs. (180 packs x 2 oz.)]
 [25 lbs. (200 packs x 2 oz.)]

Manufactured by:
Scimetrics Ltd. Corp
P.O. Box 1045
Wellington, CO 80549
New Solutions to Old Problems

{Back Panel}

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not use this product to control any pests, by any application method, or at any use sites, that are not specified on this label.

Use of this product may aid in the reduction of fleas, including by not necessarily limited to *Xenopsylla cheopis*, that commonly infest listed rodent hosts, if ingesting sufficient quantities of this bait.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets or other non-target animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, OR in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooved livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container and unused or unconsumed bait as specified by this label.

RATS AND MICE:

Use Restrictions: This product may be used to control Norway rats, roof rats and house mice at the sites in "Urban Areas" and "Non-urban Areas" that are indicated below. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come into contact with food. Wear gloves when handling bait or animal carcasses. Do not broadcast bait.

Urban Areas: To control Norway rats, roof rats and house mice this product may be used in and around homes, and industrial, commercial and public buildings; in transport vehicles (ship, trains and aircraft) and in and around related port or terminal buildings. Applications "around" buildings must be made in tamper-resistant bait stations placed along walls or other external parts of buildings (e.g. doorways, ramps, loading docks) where rats or mice might seek to gain entrance.

Non-urban Areas: To control Norway rats, roof rats and house mice this product may be used in and around homes and agricultural buildings. All outdoor placements must be in tamper-resistant bait stations.

Selection of Treatment Areas: Determine areas where rodents will most likely find and consume bait. Generally, these areas will be along walls, by gnawed openings, in corners and concealed places, between floors and walls, beside burrow openings and other locations where signs of rodents have been seen. Remove as much alternative food as possible.

Application Directions:

{Only 12-oz size:}

The amount of bait in this 6-pack package may be sufficient to kill 3 to 6 rats.

{All outer packages:}

Norway Rats and Roof Rats: Apply 2 to 8 [2 to 6 for 6-pack] place packs per bait placement. Space bait placements at intervals of 15 to 30 feet in the infested area, at locations consistent with the requirements of this label. Maintain an uninterrupted supply of fresh bait for 10 days or until there no longer are fresh signs of rat activity or signs that rats are feeding on the bait.

House Mice: Open pack and apply 1/4 to 1/2 ounce of bait per typical placement. Larger placement – up to one unopened 2-ounce pack – may be made at points of extremely high mouse activity. Space bait placements at intervals of 8 to 12 feet in the infested area, at locations consistent with the requirements on this label. Maintain an uninterrupted supply of fresh bait for 15 days or until there no longer are fresh signs of house mouse activity or signs that house mice are feeding on the bait.

Follow-up (Rats & Mice): Immediately replace contaminated or spoiled bait. Collect and properly dispose of all visible dead animals and leftover bait. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation occurs, repeat treatment. Where there is a continuous source of reinfestation, establish permanent bait stations and replenish bait as needed.

2/17

VOLES:

Use Restrictions: This product may be used to control meadow voles (*Microtus pennsylvanicus*), pine voles (*M. pinetus*) and montane voles (*M. montanus*) around (within 15 feet of) homes, and industrial, commercial, agricultural and public buildings. Do not use this product within the habitat of threatened or endangered *Microtus* spp. voles, threatened or endangered *Peromyscus* spp. mice, or other threatened or endangered rodents. Do not expose bait on bare ground. Unless children, pets, birds and other wildlife are excluded from the treated areas, all applications for controlling voles must be made in tamper-resistant bait stations. All bait applications must be made by hand. Wear gloves when handling bait or animal carcasses. Do not broadcast bait.

Selection of Treatment Areas: Determine areas near buildings where voles or signs of vole activity (girdling of plants, consumption of bulbs, burrow openings and surface runways) have been seen. These areas generally will be in flower beds or under shrubs, bushes or ground cover.

Application Directions: Open pack and load one ounce of bait (half of a pack) into each bait station. Situate bait stations near runways, burrow openings or plants that voles have been damaging. If, and only if, children, pets, domestic animals and wildlife reliably are excluded from the treated area, bait may be applied in one-ounce amounts directly into vole runways or burrow openings. Cover treated runways or holes with a shingle or board, or ground vegetation, if present. Check placements daily. Remove any fouled or spoiled bait, and replenish if vole activity and interest in bait persist. Maintain an uninterrupted supply of fresh bait for 15 days or until there no longer are fresh signs of vole activity and feeding on bait. Collect and properly dispose of any carcasses and leftover bait.

{The following will be used for the 4 oz., 12 oz. and 1 lb. sizes:}

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container in a cool, dry place inaccessible to children and pets. **Disposal: If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

{The following will be used for the 3.125 through 25 lbs. sizes:}

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container in a cool, dry place inaccessible to children and pets. **Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Disposal:** Non-Refillable Container. Do not reuse or refill this container. **{Plastic:}** Triple rinse (or equivalent) then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or incineration; or if allowed by state and local authorities, by burning; if burned, stay out of smoke. **{Paper:}** Dispose of empty paper box in a sanitary landfill or by incineration; or if allowed by state and local authorities, by burning; if burned stay out of smoke.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Keep away from humans, domestic animals and pets. Harmful if swallowed because this material may reduce the clotting ability of the blood and cause bleeding. Causes moderate eye irritation. Do not get in eyes or on skin or clothing. Wash arms, hands and face with soap and water after applying and before eating, drinking, chewing gum or using tobacco. Applicators must wear long-sleeved shirt, long pants, socks, shoes and gloves.

First Aid

Have the product container or label with you when calling a poison control center or doctor or going for treatment. **If swallowed**, call a poison control center, doctor or 1-800-858-7378 immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. **If in eyes**, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Notice to Physician or Veterinarian

This product reduces the clotting ability of the blood and may cause bleeding. If ingested, administer Vitamin K-1 intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.

Environmental Hazards

This product is highly toxic to aquatic invertebrates and toxic to mammals and birds. Do not apply this product directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwaters.

Warranty: To the extent consistent with applicable law, Manufacturer and Seller make no warranty of merchantability, fitness for any purpose, or otherwise express or implied, concerning this product or its use which extend beyond the statements on this label.

- [] Denotes optional/alternate language
- { } Denotes language that does not appear on the market label

9/17

Sub-Label B: Place Pack Label (Unit Packaging)

{Front Panel}

Novel Commensal Rodent Pellet #2
Place Pack

Kills Norway Rats, Roof Rats, House Mice and Voles
First dead rats, mice and voles will appear 4 to 5 days after feeding begins.

Rodent Control Pellets

A Bait Containing a Rodenticide to Kill Norway Rats, Roof Rats, House Mice and Voles and an Insecticide to Aid in the Reduction of Flea Populations on Listed Rodent Hosts

Active Ingredients:

Imidacloprid (CAS Number 138261-41-3)	0.020%
Warfarin (CAS Number 81-81-2)	0.025%
Other Ingredients	<u>99.955%</u>
Total	100.000%

Keep Out of Reach of Children

CAUTION

Harmful if swallowed.

See back panel for additional precautionary statements.

EPA Reg. No. 72500-13

EPA Est. 72500-CO-1

Net Weight: 2 oz. (56 grams)

Manufactured by:
Scimetrics Ltd. Corp
P.O. Box 1045
Wellington, CO 80549
New Solutions to Old Problems

{Back Panel}

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read the entire label on the outer package before using this product. It is illegal to sell these place packs individually.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container in a cool, dry place inaccessible to children and pets. **Disposal: If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Keep away from humans, domestic animals and pets. Harmful if swallowed because this material may reduce the clotting ability of the blood and cause bleeding. Causes moderate eye irritation. Do not get in eyes or on skin or clothing. Wash arms, hands and face with soap and water after applying and before eating, drinking, chewing gum or using tobacco. Applicators must wear long-sleeved shirt, long pants, socks, shoes and gloves.

First Aid

Have the product container or label with you when calling a poison control center or doctor or going for treatment. **If swallowed**, call a poison control center, doctor or 1-800-858-7378 immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. **If in eyes**, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Notice to Physician or Veterinarian

This product reduces the clotting ability of the blood and may cause bleeding. If ingested, administer Vitamin K-1 intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.

Environmental Hazards

This product is highly toxic to aquatic invertebrates and toxic to mammals and birds. Do not apply this product directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwaters.

Warranty: To the extent consistent with applicable law, Manufacturer and Seller make no warranty of merchantability, fitness for any purpose, or otherwise express or implied, concerning this product or its use which extend beyond the statements on this label.

[] Denotes optional/alternate language

{ } Denotes language that does not appear on the market label

Sub-Label C: Alternate Place Pack Label (When Sold Individually)

{Front Panel}

Novel Commensal Rodent Pellet #2
Place Packs

Kills House Mice and Voles

First dead house mice and voles will appear 4 to 5 days after feeding begins
House Mouse and Vole Control Pellets

A Bait Containing a Rodenticide to Kill House Mice and Voles and an
Insecticide to Aid in the Reduction of Flea Populations on Listed Rodent Hosts

Active Ingredients:

Imidacloprid (CAS Number 138261-41-3)	0.020%
Warfarin (CAS Number 81-81-2)	0.025%
Other Ingredients	<u>99.955%</u>
Total	100.000%

Keep Out of Reach of Children
CAUTION

Harmful if swallowed.

See back panel for additional precautionary statements.

EPA Reg. No. 72500-13

EPA Est. 72500-CO-1

Net Weight: 2 oz. (56 grams)

Manufactured by:
Scimetrix Ltd. Corp
P.O. Box 1045
Wellington, CO 80549
New Solutions to Old Problems

{Back Panel}

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not use this product to control any pests, by any application method, or at any use sites, that are not specified on this label.

Use of this product may aid in the reduction of fleas, including but not necessarily limited to *Xenopsylla cheopsis*, that commonly infest listed rodent hosts, if ingesting sufficient quantities of this bait.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets or other non-target animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.

2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, OR in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooved livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container and unused or unconsumed bait as specified by this label.

HOUSE MICE:

Use Restrictions: This product may be used to control house mice at the sites in "Urban Areas" and "Non-urban Areas" that are indicated below. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come into contact with food. Wear gloves when handling bait or animal carcasses. Do not broadcast bait.

Urban Areas: To control house mice this product may be used in and around homes, and industrial, commercial and public buildings; in transport vehicles (ship, trains and aircraft) and in and around related port or terminal buildings. Applications "around" buildings must be made in tamper-resistant bait stations placed along walls or other external parts of buildings (e.g. doorways, ramps, loading docks) where rats or mice might seek to gain entrance.

Non-urban Areas: To control house mice this product may be used in and around homes and agricultural buildings.

Selection of Treatment Areas: Determine areas where house mice will most likely find and consume bait. Generally, these areas will be along walls, by gnawed openings, in corners and concealed places, between floors and walls, beside burrow openings and other locations where signs of house mice have been seen. Remove as much alternative food as possible.

Application Directions:

Open pack and apply 1/4 to 1/2 ounce of bait per typical placement. Larger placement – up to one unopened 2-ounce pack – may be made at points of extremely high mouse activity. Space bait placements at intervals of 8 to 12 feet in the infested area, at locations consistent with the requirements on this label. Maintain an uninterrupted supply of fresh bait for 15 days or until there no longer are fresh signs of house mouse activity or signs that house mice are feeding on the bait.

Follow-up: Immediately replace contaminated or spoiled bait. Collect and properly dispose of all visible dead animals and leftover bait. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation occurs, repeat treatment. Where there is a continuous source of reinfestation, establish permanent bait stations and replenish bait as needed.

VOLES:

Use Restrictions: This product may be used to control meadow voles (*Microtus pennsylvanicus*), pine voles (*M. pinetus*) and montane voles (*M. montanus*) around (within 15 feet of) homes, and industrial, commercial, agricultural and public buildings. Do not use this product within the habitat of threatened or endangered *Microtus* spp. voles, threatened or endangered *Peromyscus* spp. mice, or other threatened or endangered rodents. Do not expose bait on bare ground. Unless children, pets, birds and other wildlife are excluded from the treated areas, all applications for controlling voles must be made in tamper-resistant bait stations. All bait applications must be made by hand. Wear gloves when handling bait or animal carcasses. Do not broadcast bait.

Selection of Treatment Areas: Determine areas near buildings where voles or signs of vole activity (girdling of plants, consumption of bulbs, burrow openings and surface runways) have been seen. These areas generally will be in flower beds or under shrubs, bushes or ground cover.

Application Directions: Open pack and load one ounce of bait (half of a pack) into each bait station. Situate bait stations near runways, burrow openings or plants that voles have been damaging. If, and only if, children, pets,

domestic animals and wildlife reliably are excluded from the treated area, bait may be applied in one-ounce amounts directly into vole runways or burrow openings. Cover treated runways or holes with a shingle or board, or ground vegetation, if present. Check placements daily. Remove any fouled or spoiled bait, and replenish if vole activity and interest in bait persist. Maintain an uninterrupted supply of fresh bait for 15 days or until there no longer are fresh signs of vole activity and feeding on bait. Collect and properly dispose of any carcasses and leftover bait.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container in a cool, dry place inaccessible to children and pets. **Disposal: If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Keep away from humans, domestic animals and pets. Harmful if swallowed because this material may reduce the clotting ability of the blood and cause bleeding. Causes moderate eye irritation. Do not get in eyes or on skin or clothing. Wash arms, hands and face with soap and water after applying and before eating, drinking, chewing gum or using tobacco. Applicators must wear long-sleeved shirt, long pants, socks, shoes and gloves.

First Aid

Have the product container or label with you when calling a poison control center or doctor or going for treatment. **If swallowed,** call a poison control center, doctor or 1-800-858-7378 immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. **If in eyes,** hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Notice to Physician or Veterinarian

This product reduces the clotting ability of the blood and may cause bleeding. If ingested, administer Vitamin K-1 intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.

Environmental Hazards

This product is highly toxic to aquatic invertebrates and toxic to mammals and birds. Do not apply this product directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwaters.

Warranty: To the extent consistent with applicable law, Manufacturer and Seller make no warranty of merchantability, fitness for any purpose, or otherwise express or implied, concerning this product or its use which extend beyond the statements on this label.

[] Denotes optional/alternate language
{ } Denotes language that does not appear on the market label

Kaput[®]

Kills Norway Rats, Roof Rats, House Mice and Voles.
First dead rats, mice and voles will appear
4 to 5 days after feeding begins.

A Bait Containing a Rodenticide to Kill Norway Rats, Roof
Rats, House Mice and Voles and an Insecticide to Aid in the
Reduction of Flea Populations on Listed Rodent Hosts

COMBO BAIT PELLETS

RODENT CONTROL PELLETS

PLACE PACKS

NET WT. 3.125 LB (1.42 kg) (25 PACKS x 2 oz.)

ACTIVE INGREDIENTS:

Imidacloprid (CAS Number 186261-41-3) 0.020%
Warfarin (CAS Number 81-81-2) 0.025%

OTHER INGREDIENTS

TOTAL 99.955%

100.000%

EPA Est. 72500-CO-1
EPA Reg. No. 72500-13



Keep Out of Reach of Children
CAUTION
Harmful If Swallowed.

See back panel for additional Precautionary Statements.

Manufactured by:
Scimetrics
LTD. CORP.
P.O. Box 1045
Wellington, CO 80549
New Solutions to Old Problems

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

CAUTION

Keep away from humans, domestic animals and pets. Harmful if swallowed because this material may reduce the clotting ability of the blood and cause bleeding. Causes moderate eye irritation. Do not get in eyes or on skin or clothing. Wash arms, hands and face with soap and water after applying and before eating, drinking, chewing gum or using tobacco. Applicators must wear long-sleeved shirt, long pants, socks, shoes and gloves.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor or going for treatment.
If swallowed, call a poison control center, doctor or 1-800-858-7378 immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. If in eyes, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

NOTICE TO PHYSICIAN OR VETERINARIAN

This product reduces the clotting ability of the blood and may cause bleeding. If ingested, administer Vitamin K-1 intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates and toxic to mammals and birds. Do not apply this product directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not use this product to control any pests, by any application method, or at any use sites, that are not specified on this label.

Use of this product may aid in the reduction of fleas, including by not necessarily limited to *Xenopsylla cheopis*, that commonly infest listed rodent hosts, if ingesting sufficient quantities of this bait.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets or other non-target animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, OR in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooved livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container and unused or unconsumed bait as specified by this label.

RATS AND MICE

Use Restrictions: This product may be used to control Norway rats, roof rats and house mice at the sites in "Urban Areas" and "Non-urban Areas" that are indicated below. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come into contact with food. Wear gloves when handling bait or animal carcasses. Do not broadcast bait.

Urban Areas: To control Norway rats, roof rats and house mice this product may be used in and around homes, and industrial, commercial and public buildings; in transport vehicles (ship, trains and aircraft) and in and around related port or terminal buildings. Applications "around" buildings must be made in tamper-resistant bait stations placed along walls or other external parts of buildings (e.g. doorways, ramps, loading docks) where rats or mice might seek to gain entrance.

Non-urban Areas: To control Norway rats, roof rats and house mice this product may be used in and around homes and agricultural buildings. All outdoor placements must be in tamper-resistant bait stations.

Selection of Treatment Areas: Determine areas where rodents will most likely find and consume bait. Generally, these areas will be along walls, by gnawed openings, in corners and concealed places; between floors and walls, beside burrow openings and other locations where signs of rodents have been seen. Remove as much alternative food as possible.

Application Directions:
Norway Rats and Roof Rats: Apply 2 to 8 place

packs per bait placement. Space bait placements at intervals of 15 to 30 feet in the infested area, at locations consistent with the requirements of this label. Maintain an uninterrupted supply of fresh bait for 10 days or until there no longer are fresh signs of rat activity or signs that rats are feeding on the bait.

House Mice: Open pack and apply 1/4 to 1/2 ounce of bait per typical placement. Larger placement - up to one unopened 2-ounce pack - may be made at points of extremely high mouse activity. Space bait placements at intervals of 8 to 12 feet in the infested area, at locations consistent with the requirements on this label. Maintain an uninterrupted supply of fresh bait for 15 days or until there no longer are fresh signs of house mouse activity or signs that house mice are feeding on the bait.

Follow-up (Rats & Mice): Immediately replace contaminated or spoiled bait. Collect and properly dispose of all visible dead animals and leftover bait. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation occurs, repeat treatment. Where there is a continuous source of reinfestation, establish permanent bait stations and replenish bait as needed.

VOLES

Use Restrictions: This product may be used to control meadow voles (*Microtus pennsylvanicus*), pine voles (*M. pinetis*) and montane voles (*M. montanus*) around (within 15 feet of) homes, and industrial, commercial, agricultural and public buildings. Do not use this product within the habitat of threatened or endangered *Microtus* spp. voles, threatened or endangered *Peromyscus* spp. mice, or other threatened or endangered rodents. Do not expose bait on bare ground. Unless children, pets, birds and other wildlife are excluded from the treated areas, all applications for controlling voles must be made in tamper-resistant bait stations. All bait applications must be made by hand. Wear gloves when handling bait or animal carcasses. Do not broadcast bait.

Selection of Treatment Areas: Determine areas near buildings where voles or signs of vole activity (gridding of plants, consumption of bulbs, burrow openings and surface runways) have been seen. These areas generally will be in flower beds or under shrubs, bushes or ground cover.

Application Directions: Open pack and load one ounce of bait (half of a pack) into each bait station. Situate bait stations near runways, burrow openings or plants that voles have been

damaging. If, and only if, children, pets, domestic animals and wildlife reliably are excluded from the treated area, bait may be applied in one-ounce amounts directly into vole runways or burrow openings. Cover treated runways or holes with a shingle or board, or ground vegetation, if present. Check placements daily. Remove any fouled or spoiled bait, and replenish if vole activity and interest in bait persist. Maintain an uninterrupted supply of fresh bait for 15 days or until there no longer are fresh signs of vole activity and feeding on bait. Collect and properly dispose of any carcasses and leftover bait.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container in a cool, dry place inaccessible to children and pets. **Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Disposal:** Non-Refillable Container. Do not reuse or refill this container. Triple rinse (or equivalent) then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill, or incineration; or if allowed by state and local authorities, by burning; if burned, stay out of smoke.

WARRANTY

To the extent consistent with applicable law, Manufacturer and Seller make no warranty of merchantability, fitness for any purpose, or otherwise express or implied, concerning this product or its use which extend beyond the statements on this label.

Scimetrics
LTD. CORP.

New Solutions to Old Problems
P.O. Box 1045 Wellington, CO 80549
(970) 482-1330

Label Version 4/2/08



NOTIFICATION
JUL 25 2008

17

Kaput[®]

Kills Norway Rats, Roof Rats, House Mice and Voles.
First dead rats, mice and voles will appear
4 to 5 days after feeding begins.

A Bait Containing a Rodenticide to Kill Norway Rats, Roof Rats, House Mice and Voles and an Insecticide to Aid in the Reduction of Flea Populations on Listed Rodent Hosts

COMBO BAIT[®] PELLETS

RODENT CONTROL PELLETS

PLACE PACKS

NET WT. 3.125 LB (1.42 kg) (25 PACKS x 2 oz.)

ACTIVE INGREDIENTS:

Imidacloprid (CAS Number 138261-41-3)	0.020%
Warfarin (CAS Number 81-81-2)	0.025%

OTHER INGREDIENTS

99.955%

TOTAL

100.000%

EPA Est. 72500-CO-1
EPA Reg. No. 72500-13



Manufactured by:
Scimetrics
 LTD. CORP.
 P.O. Box 1045
 Wellington, CO 80549
New Solutions to Old Problems

Keep Out of Reach of Children

CAUTION

Harmful If Swallowed.

See back panel for additional Precautionary Statements.

NOTIFICATION

JUL 25 2008

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

CAUTION

Keep away from humans, domestic animals and pets. Harmful if swallowed because this material may reduce the clotting ability of the blood and cause bleeding. Causes moderate eye irritation. Do not get in eyes or on skin or clothing. Wash arms, hands and face with soap and water after applying and before eating, drinking, chewing gum or using tobacco. Applicators must wear long-sleeved shirt, long pants, socks, shoes and gloves.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

If swallowed, call a poison control center, doctor or 1-800-858-7378 immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. **If in eyes**, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

NOTICE TO PHYSICIAN OR VETERINARIAN

This product reduces the clotting ability of the blood and may cause bleeding. If ingested, administer Vitamin K-1 intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates and toxic to mammals and birds. Do not apply this product directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not use this product to control any pests, by any application method, or at any use sites, that are not specified on this label.

Use of this product may aid in the reduction of fleas, including by not necessarily limited to *Xenopsylla cheopis*, that commonly infest listed rodent hosts, if ingesting sufficient quantities of this bait.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets or other non-target animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, OR in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container and unused or unconsumed bait as specified by this label.

RATS AND MICE

Use Restrictions: This product may be used to control Norway rats, roof rats and house mice at the sites in "Urban Areas" and "Non-urban Areas" that are indicated below. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come into contact with food. Wear gloves when handling bait or animal carcasses. Do not broadcast bait.

Urban Areas: To control Norway rats, roof rats and house mice this product may be used in and around homes, and industrial, commercial and public buildings; in transport vehicles (ship, trains and aircraft) and in and around related port or terminal buildings. Applications "around" buildings must be made in tamper-resistant bait stations placed along walls or other external parts of buildings (e.g. doorways, ramps, loading docks) where rats or mice might seek to gain entrance.

Non-urban Areas: To control Norway rats, roof rats and house mice this product may be used in and around homes and agricultural buildings. All outdoor placements must be in tamper-resistant bait stations.

Selection of Treatment Areas: Determine areas where rodents will most likely find and consume bait. Generally, these areas will be along walls, by gnawed openings, in corners and concealed places, between floors and walls, beside burrow openings and other locations where signs of rodents have been seen. Remove as much alternative food as possible.

Application Directions:

Norway Rats and Roof Rats: Apply 2 to 8 place

packs per bait placement. Space bait placements at intervals of 15 to 30 feet in the infested area, at locations consistent with the requirements of this label. Maintain an uninterrupted supply of fresh bait for 10 days or until there no longer are fresh signs of rat activity or signs that rats are feeding on the bait.

House Mice: Open pack and apply 1/4 to 1/2 ounce of bait per typical placement. Larger placement – up to one unopened 2-ounce pack – may be made at points of extremely high mouse activity. Space bait placements at intervals of 8 to 12 feet in the infested area, at locations consistent with the requirements on this label. Maintain an uninterrupted supply of fresh bait for 15 days or until there no longer are fresh signs of house mouse activity or signs that house mice are feeding on the bait.

Follow-up (Rats & Mice): Immediately replace contaminated or spoiled bait. Collect and properly dispose of all visible dead animals and leftover bait. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation occurs, repeat treatment. Where there is a continuous source of reinfestation, establish permanent bait stations and replenish bait as needed.

VOLES

Use Restrictions: This product may be used to control meadow voles (*Microtus pennsylvanicus*), pine voles (*M. pinetus*) and montane voles (*M. montanus*) around (within 15 feet of) homes, and industrial, commercial, agricultural and public buildings. Do not use this product within the habitat of threatened or endangered *Microtus* spp. voles, threatened or endangered *Peromyscus* spp. mice, or other threatened or endangered rodents. Do not expose bait on bare ground. Unless children, pets, birds and other wildlife are excluded from the treated areas, all applications for controlling voles must be made in tamper-resistant bait stations. All bait applications must be made by hand. Wear gloves when handling bait or animal carcasses. Do not broadcast bait.

Selection of Treatment Areas: Determine areas near buildings where voles or signs of vole activity (girdling of plants, consumption of bulbs, burrow openings and surface runways) have been seen. These areas generally will be in flower beds or under shrubs, bushes or ground cover.

Application Directions: Open pack and load one ounce of bait (half of a pack) into each bait station. Situate bait stations near runways, burrow openings or plants that voles have been

damaging. If, and only if, children, pets, domestic animals and wildlife reliably are excluded from the treated area, bait may be applied in one-ounce amounts directly into vole runways or burrow openings. Cover treated runways or holes with a shingle or board, or ground vegetation, if present. Check placements daily. Remove any fouled or spoiled bait, and replenish if vole activity and interest in bait persist. Maintain an uninterrupted supply of fresh bait for 15 days or until there no longer are fresh signs of vole activity and feeding on bait. Collect and properly dispose of any carcasses and leftover bait.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container in a cool, dry place inaccessible to children and pets.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Non-Refillable Container. Do not reuse or refill this container. Triple rinse (or equivalent) then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or incineration; or if allowed by state and local authorities, by burning; if burned, stay out of smoke.

WARRANTY

To the extent consistent with applicable law, Manufacturer and Seller make no warranty of merchantability, fitness for any purpose, or otherwise express or implied, concerning this product or its use which extend beyond the statements on this label.

Scimetrics

LTD. CORP.

New Solutions to Old Problems

P.O. Box 1045 Wellington, CO 80549
(970) 482-1330

Label Version 4/2/08

